



University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): Brita Dahl Jensen, Joana G. Vicente, Hira K. Manandhar and Steven J. Roberts

Article Title: Occurrence and Diversity of *Xanthomonas campestris* pv. *campestris* in Vegetable Brassica Fields in Nepal

Year of publication: 2010

Link to published version:

[http://dx.doi.org/ 10.1094/PDIS-94-3-0298](http://dx.doi.org/10.1094/PDIS-94-3-0298)

Publisher statement: None

1 **Occurrence and Diversity of *Xanthomonas campestris* pv. *campestris* in**
2 **Vegetable Brassica Fields in Nepal**

3

4 **Brita Dahl Jensen**, Department of Plant Biology and Biotechnology & Department of
5 Agriculture and Ecology, The Faculty of Life Sciences, University of Copenhagen,
6 Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark; **Joana G. Vicente**, Warwick HRI,
7 The University of Warwick, Wellesbourne, Warwick CV35 9EF, UK; **Hira K.**
8 **Manandhar**, Plant Pathology Division, Nepal Agricultural Research Council, P. O. Box
9 1126, Khumaltar, Lalitpur, Nepal; and **Steven J. Roberts**, Plant Health Solutions, 20
10 Beauchamp Road, Warwick CV34 5NU, UK

11

12 Corresponding author: B. D. Jensen

13 E-mail: dahl@life.ku.dk

14

15

1 Correspondence to:

2 Brita Dahl Jensen

3 Department of Plant Biology and Biotechnology & Department of Agriculture

4 and Ecology

5 The Faculty of Life Sciences, University of Copenhagen

6 Thorvaldsensvej 40

7 1871 Frederiksberg C

8 Denmark

9

10 E-mail: dahl@life.ku.dk

11 Phone: + 45 35 33 34 71

12 Fax: + 45 35 33 34 60

13

14

1 **ABSTRACT**

2

3 Jensen, B. D., Vicente J. G., Manandhar H. K., and Roberts. S. J. 200X. Occurrence and
4 diversity of *Xanthomonas campestris* pv. *campestris* in vegetable brassica fields in Nepal.
5 Plant Dis. XX: XXX-XXX.

6

7 Black rot caused by *Xanthomonas campestris* pv. *campestris* (*Xcc*) was found in 28
8 sampled cabbage fields in five major cabbage growing districts in Nepal in 2001, and in
9 four cauliflower fields in two districts and a leaf mustard seed bed in 2003. Pathogenic
10 *Xcc* strains were obtained from 39 cabbage plants, four cauliflower plants and one leaf
11 mustard plant with typical lesions. Repetitive DNA polymerase chain reaction-based
12 fingerprinting (rep-PCR) using REP, ERIC and BOX primers, was used to assess the
13 genetic diversity. Strains were also race-typed using a differential series of *Brassica* spp..
14 Cabbage strains belonged to five races (1, 4, 5, 6 and 7) with races 4, 1 and 6 the most
15 common. All cauliflower strains were race 4, and the leaf mustard strain was race 6. A
16 dendrogram derived from the combined rep-PCR profiles showed that the Nepalese *Xcc*
17 strains clustered separately from other *Xanthomonas* species and pathovars. Race 1
18 strains clustered together and strains of races 4, 5 and 6 were each split into at least two
19 clusters. The presence of different races and the genetic variability of the pathogen should
20 be considered when resistant cultivars are bred and introduced into regions in Nepal to
21 control black rot of brassicas.

22 Additional keywords: genetic variation, pathogenicity

23

1 INTRODUCTION

2 Cabbage and cauliflower production increased significantly in Nepal during the 1990's;
3 in 2004/2005, nearly 20,000 ha were cultivated with cabbage, accounting for 11% of the
4 vegetable area in Nepal (3). Cabbage is considered an important cash crop, and in some
5 areas production is almost continuous, with up to three crops per year. The production
6 increase has been accompanied by an increase in the use of commercial hybrid seed (32).

7 Black rot, caused by the vascular bacterium *Xanthomonas campestris* pv. *campestris*,
8 (*Xcc*), is one of the most serious diseases of vegetable Brassicas (4, 43). Typical disease
9 symptoms in the field are V-shaped, chlorotic to necrotic lesions starting from leaf
10 margins and with blackening of the vascular tissue (33, 43). Blight symptoms have also
11 been reported (45). Black rot of crucifers was reported in Nepal in 1977 (30). Adhikari &
12 Basnyat (1) found that 20 strains of *Xcc* from broccoli, cabbage and cauliflower from the
13 Kathmandu and Chitwan valleys, had similar biochemical and physiological
14 characteristics, but strains from Chitwan tended to be more aggressive on cabbage,
15 cauliflower and broccoli, than strains from the Kathmandu Valley.

16 Until 2007, six races of *Xcc* were defined on the basis of their reactions on a differential
17 set that includes genotypes of *Brassica oleracea*, *B. rapa*, *B. juncea*, *B. napus* and *B.*
18 *carinata* (14, 40). A gene-for-gene model has been proposed to explain the interactions
19 between the races and the differential cultivars (40). Races 1, 4 and 6 are the most
20 frequently reported races (11, 39, 40). In 2007, race 7 was reported in Belgium by Fargier
21 & Manceau (8), along with races 8 and 9, which had a narrower host range on differential

1 cultivars. We also found race 7 in this study and a preliminary report of our work
2 appeared in 2007 (13).

3 Conserved, repetitive DNA sequences in bacteria have been targeted for use in
4 distinguishing bacterial species and strains by the polymerase chain reaction (PCR) (7,
5 37, 38), using primers specific to the repetitive extragenic palindromic (REP) sequence
6 (10), the enterobacterial repetitive intergenic consensus (ERIC) sequence (28), and the
7 BOX element (17). Previous studies have shown the potential of repetitive DNA PCR-
8 based fingerprinting (rep-PCR) to differentiate *Xanthomonas* pathovars (20, 21, 26, 41)
9 and strains within pathovars (16, 20), and therefore could be used as a diagnostic tool in
10 plant pathology.

11 Some strains of *Xcc* have been characterised by genetic fingerprinting using rep-PCR.
12 *Xcc* strains from Tanzania tended to cluster within local geographical areas (18) and
13 similarly it was found that *Xcc* strains from different countries to some extent clustered
14 within larger geographical regions (35), whereas this was not the case in a study carried
15 out on Italian strains (46). Results of rep-PCR fingerprinting allowed the separation of
16 *Xcc* from *X. campestris* pv. *raphani* (*Xcr*) strains, and showed that among *Xcc* and *Xcr*
17 strains there was a tendency for strains of the same race to cluster together (41). Rep-
18 PCR, AFLP, PFGE (pulsed-field gel electrophoresis) and integron gene cassette PCR was
19 used to study the diversity of 22 *Xcc* strains, and revealed 11, 12, 13, and 4 differentiated
20 genotypes, respectively, supporting the heterogeneous nature and relatively high level of
21 diversity in this pathovar (36, 46).

22 The aim of this work was: (i) to study the occurrence of black rot in Brassica fields in
23 major cabbage growing regions of Nepal; (ii) to characterise the diversity of the *Xcc*

1 strains by rep-PCR fingerprinting, and race typing on a set of differential *Brassica* spp.
2 genotypes with the aim to suggest efficient control strategies involving breeding,
3 introduction and cultivation of resistant cultivars.

4

5 **MATERIALS AND METHODS**

6 The work was divided into three steps: 1) Collection of black rot infected leaf material in
7 Nepal, 2) isolation of *Xcc* and pathogenicity testing at the Faculty of Life Sciences,
8 Denmark, and 3) further pathogenicity testing, rep-PCR and race typing at Warwick HRI,
9 The University of Warwick, UK.

10 **Collection of black rot infected leaf samples.** Twenty-eight white cabbage (*Brassica*
11 *oleracea* var. *capitata*) fields, each at least 10 x 10 m, were sampled in five major
12 cabbage growing areas (five districts in four zones) in the Eastern, Central and Western
13 parts of Nepal (Fig. 1; Table 1) during the monsoon season in July 2001. According to
14 farmers' information, the cultivar grown was usually Green Coronet F₁. From each field,
15 a leaf with typical black rot symptoms (V-shaped lesions) was sampled from each of ten
16 well-separated plants. In 2003, single plants in four cauliflower (*B. oleracea* var. *botrytis*)
17 fields were sampled, and a single sample was collected from a seedbed of leaf mustard
18 (*B. juncea*). Samples were dried between sheets of paper at room temperature before
19 isolation of the causal agent. The altitude of each field was recorded using a hand held
20 GPS device (eTrex Venture GPS navigator, Garmin Corporation, Taiwan).

21 **Isolations.** One or two leaf samples, representing one or two plants from each field, were
22 selected for isolation of the causal agent from the typical V-shaped lesions. From each
23 leaf sample, 0.5-1 cm² leaf tissue was excised with a sterile scalpel from the margin of a

1 lesion and placed in a drop of sterile 0.85% saline for 5 min. A loopful of saline
2 suspension from a point with bacterial ooze was streaked on nutrient starch
3 cycloheximide agar (NSCA) (23) and incubated at 28°C for 48-72 h and observed for
4 typical *Xcc* colonies (pale yellow, mucoid, starch-hydrolyzing). A typical colony from
5 each isolation plate was sub-cultured to yeast dextrose calcium carbonate agar (YDC)
6 (31), and then transferred to sucrose-peptone agar (SPA) for viscosity testing (19). For
7 most isolations, and additional colony from the same plate was sub-cultured and the
8 strain was kept as a 'duplicate'. All isolates with a flow time of more than 30 seconds
9 (19) were subsequently tested for pathogenicity. Isolates were stored at -80°C in
10 cryovials (Simport cryovial, Canada) containing glass beads and nutrient broth plus 15 %
11 glycerol (9), or in Protect cryovials (Technical Service Consultants Ltd., Lancashire,
12 UK).

13 **Pathogenicity.** Isolates were tested for pathogenicity on six 3 to 4-week-old plants of
14 cabbage cv. Copenhagen Market (L. Daehnfelddt, Denmark), grown in 12 x 13 cm square
15 pots in a peat-based compost (Weibulls Enhetsjord, K-Jord, Svalöf Weibull Trädgård AB,
16 Hammenhög, Sweden) in a glasshouse at 20-25°C. Each isolate was tested twice
17 (different inoculation date). Bacteria were grown on YDC agar at 28°C for 48 h, and
18 inoculum was prepared by suspending a small amount of growth in 0.85% saline. The
19 suspensions were adjusted to a final concentration of approx. 10^8 CFU ml⁻¹ by dilution.
20 Each pot with six plants was sprayed with 10 ml of bacterial suspension using a hand
21 held plastic atomiser (KAB1005, KABI, Denmark) at a distance of approx. 15 cm from
22 the plants. After inoculation, plants were maintained in a transparent polyethylene tent for
23 48 h to increase the humidity. The temperature varied between 25 and 35°C during the

1 experiment with a day length of approx. 16 h. The six plants were assessed for symptoms
2 (V-shaped lesions) 2 and 3 weeks after inoculation; isolates were considered pathogenic
3 when V-shaped lesions occurred on at least three plants. In addition, the pathogenicity of
4 twenty isolates was tested on individual 4-week-old Savoy cabbage plants (*B. oleracea*
5 var. *sabauda*) of cv. Wirosa F₁ (Bejo Zaaden, The Netherlands) using the inoculation
6 method of Vicente et al. (41). Plants were raised from seed sown in 6.5-cm square plastic
7 pots with Levington M2 compost (Scotts, England) in a glasshouse with a minimum
8 temperature of 20/15°C and venting at 22/17°C (day/night), and with supplementary
9 lighting to give a 16 h photoperiod. Isolates were grown on YDC medium for 48 h at
10 30°C. The three youngest leaves of 4-week-old plants were inoculated at six sites near the
11 leaf margin and three sites in the midrib using a sterile pin charged with growth from a
12 YDC plate (41). One plant was inoculated per isolate, but the inoculations were repeated
13 if the first results indicated that the isolate was weak or not pathogenic. The presence of
14 symptoms was recorded two and three weeks after inoculation (40).

15 **rep-PCR fingerprinting.** DNA extraction and rep-PCR amplification was done for all
16 pathogenic strains from Nepal (44 independent strains plus 32 duplicate strains from the
17 same leaf) and two reference strains, representing two commonly found races, HRI
18 1279A (race 4) and HRI 3811 (race 1), under the same laboratory conditions at Warwick
19 HRI as previously reported (41).

20 All samples, 'blank' control lanes using water as template, and the two reference strains
21 were included at the same time in each of the rep-PCR setups. From each strain and blank
22 controls, 6 µl aliquots (REP, and ERIC primer sets, and 2.5 µl aliquots (BOX primer) of
23 amplified PCR products were separated by electrophoresis in 1.2% agarose 22 lane gels

1 (41). A molecular mass marker (1-Kb Plus DNA Ladder, Invitrogen) was loaded on both
2 sides and central lanes, and the two reference strains were loaded on all gels. Gels were
3 visualized on a UV transilluminator and photographed on an imaging system (ImaGo, B
4 & L Systems, The Netherlands). A digital image of each gel was analysed using the
5 GelCompar II Software (Applied Maths, Kortrijk, Belgium). Gels were normalised using
6 the molecular mass markers; optimisation settings were chosen which maximised the
7 similarity of the strains, which were repeated on each gel. Bands less than 500 bp (REP),
8 400 bp (ERIC), 300 bp (BOX) and larger than 4000 bp were excluded from the analysis.
9 The normalized data generated from BOX, ERIC and REP fingerprinting profiles were
10 combined. The similarity between profiles was calculated using the Pearson product-
11 moment correlation coefficient. Cluster analysis of the pairwise similarity values was
12 performed using the UPGMA algorithm. For comparative purposes, profiles of additional
13 strains, representing all six races of *Xcc*, three races of *Xcr*, three strains of other
14 *Xanthomonas campestris* pathovars and two strains of other *Xanthomonas* species
15 generated previously under the same laboratory conditions as used here (41) were
16 included in the analysis.

17 **Race typing.** Forty-four independent strains from Nepal and seven duplicate strains
18 (obtained from the same plants), were race typed. Race typing was carried out using a set
19 of differential *Brassica* spp. using the method and experimental conditions at Warwick
20 HRI as previously described (40) with the following modifications: the doubled haploid
21 lines Cob60 and FBLM2 and the inbred line Bo99075 derived respectively from *B. napus*
22 cv. Cobra line 14R, *B. juncea* cv. Florida Broad Leaf Mustard and *B. carinata* accession
23 PI 199947 replaced the lines from which they were originally derived. In addition, the

1 doubled-haploid line SxD1 derived from a cross between *B. oleracea* cv.
2 Böhmerwaldkohl and a rapid cycling line was also tested to confirm the results obtained
3 with Miracle F₁. The differentials used are included in Table 2. Three plants of ‘Seven
4 Top Turnip’ and one plant of each of the other genotypes were inoculated with each
5 strain. One control strain of race 5 (HRI 3880) was also race typed for comparison.
6 Plants were raised in the glasshouse as described above for the pathogenicity test on
7 Wirosa F₁. Strains were grown on YDC medium for 48 h at 30°C, and suspended in
8 0.85% saline to produce a turbid suspension containing approx. 10⁸-10⁹ CFU ml⁻¹. The
9 three youngest leaves of 4-week-old plants were inoculated by clipping the veins near the
10 leaf margins with mouse-tooth forceps with teeth wrapped in absorbent cotton wool and
11 dipped in the bacterial suspension (40). Disease assessments were carried out two and
12 three weeks after inoculation (40).

13

14 **RESULTS**

15 **Occurrence of black rot and symptoms in the field.** Black rot symptoms were
16 commonly observed in all 28 cabbage fields visited and sampled in the five major
17 cabbage growing districts in Nepal during the monsoon season in July 2001 (Fig. 1, Table
18 1). The altitude of the fields ranged from 1296 to 2139 m. Most samples were collected
19 from fields in Lalitpur, Dhankuta, and Makawanpur districts, as cabbage production was
20 at its peak at the time of sampling, whereas most fields had already been harvested in
21 Bhaktapur and Baglung districts. Typical V-shaped, chlorotic lesions with black veins
22 were found on leaves in all fields, irrespective of plant maturity. In some fields, mature
23 heads exhibited blackening of the vascular stem tissue when cut in half. In 2003, the four

1 cauliflower fields and the seedbed of leaf mustard visited, also exhibited typical
2 symptoms: V-shaped, chlorotic lesions with dark veins. Blight symptoms were not
3 observed.

4 **Isolations and pathogenicity.** Isolates from 39 cabbage plants, four cauliflower plants
5 and one leaf mustard plant produced yellow mucoid growth on YDC and a positive
6 viscosity test (Table 1, Table 3). Thirty duplicate isolates from the same cabbage plants
7 and two duplicate isolates from the cauliflower plants also had the above typical
8 characteristics. The isolates all produced clear and distinct V-shaped lesions on leaves of
9 *B. oleracea* cv. Copenhagen Market and/or Wirosa F₁ and were considered to be *Xcc*.

10 **rep-PCR fingerprinting.** All 39 cabbage strains, the four cauliflower strains and the leaf
11 mustard strain together with the 32 duplicate strains from Nepal and the two reference
12 strains, HRI 1279A (race 4) and HRI 3811 (race 1), were fingerprinted using rep-PCR
13 (Table 1). The rep-PCR fingerprint profiles, obtained with the BOX, ERIC and REP
14 primers, confirmed that all isolates were strains of *Xcc*. These profiles were complex and
15 revealed polymorphic bands among *Xcc* strains. The combined dendrogram had a
16 cophenetic correlation coefficient of 0.95. The separate dendrograms obtained with REP,
17 ERIC and BOX had cophenetic correlations of 0.93, 0.94 and 0.88 respectively (data not
18 shown).

19 The profiles of the two reference strains (HRI 1279A and 3811) that were repeated in five
20 different gels had similarities ranging from 88.6 to 97.7% and 81.1 (in one gel) to 95.3%
21 respectively, with most profiles of the same strain having similarities greater than 93%.
22 Thus, the separation of strains with similarities greater than 80% might not be reliable,
23 and strains with a similarity greater than approximately 90% might be identical.

1 All, except one, of the pairs of strains originating from a single lesion (30 pairs from
2 cabbage, two from cauliflower plants), produced fingerprint profiles that were identical
3 (i.e. with a similarity $\geq 90\%$). The pair of strains (N47 and N48) obtained from the same
4 lesion had distinct profiles (similarity of 81.9%, with profiles obtained on the same gel
5 runs).

6 Figure 2 presents a dendrogram constructed with the results from Nepalese strains from
7 39 individual cabbage plants, of which one plant is represented by two strains (N47 and
8 N48), four strains from cauliflower, one strain from leaf mustard, fifteen strains
9 representative of the six races of *Xcc*, and eight strains of other *Xanthomonas* species and
10 pathovars. All *Xcc* strains from Nepal and all reference *Xcc* strains, except race 2, formed
11 a single cluster at 44.8% similarity. This group was differentiated from the other *X.*
12 *campestris* pathovars and other *Xanthomonas* species. *Xcc* race 2, of which only a single
13 strain has been obtained (39), is clearly distinct from other *Xcc* strains. At 70% similarity,
14 the *Xcc* strains separated into a large group of 29 strains (containing 21 strains from four
15 districts of Nepal), a group of 12 Nepalese strains (from four districts of Nepal), a group
16 of 11 strains (containing 9 Nepalese strains from three districts of Nepal), two strains
17 from Nepal (N68 and N70), two strains from Portugal and Nepal (HRI 6185 and N8),
18 respectively, and three individual strains (HRI 6181, HRI 3880, HRI 6312A). At 80%
19 similarity, the group of 29 strains split into two major groups (containing 9 and 13
20 strains), three minor groups containing two strains and an individual strain (HRI 6412);
21 the group of 12 strains split into a group of 8 strains, a group of two strains (N35 and
22 N47) and two single strains (N55 and N34); the group of 11 strains remained clustered.

1 **Race-typing.** Thirty-nine cabbage *Xcc* strains, seven duplicate strains, four cauliflower
2 strains, and one leaf mustard strain from Nepal plus the race 5 type-strain (HRI 3880)
3 were inoculated into the differential set (Table 2). Four of the six previously described
4 races, namely 1, 4, 5, and 6 were identified (Table 3; Table 4). In cabbage, on an
5 individual plant basis, 14 strains were identified as race 4, 12 strains as race 1, nine
6 strains as race 6, and three as race 5. Races 2 and 3 were not found. Race 6 strains were
7 generally pathogenic on all differentials, but weakly pathogenic on *B. oleracea* cv.
8 Miracle F₁ and line SxD1. Two strains, N35 and N47, were pathogenic and very
9 aggressive on all genotypes except on *B. juncea* line FBLM2. This was an unknown
10 differential reaction at the time of the experiments, and these strains were assigned to race
11 7 (13). Seven plants were represented by seven pairs (duplicates) of strains from the same
12 lesion (Table 3). These pairs of strains were the same race-type, except for one pair (N47
13 and N48), which were identified as races 6 and 7, respectively (Table 3). Races 4, 1 and
14 6 were the most common in cabbage and were all found in four of the five districts (Table
15 4). Races 5 and 7 were represented by three and two of the strains, respectively (Table 4).
16 Race 5 was identified in two of the districts, and race 7 in only one district. The four
17 strains from cauliflower were all race 4, and the leaf mustard strain was race 6 (Table 3,
18 Table 4). Cabbage production in the Dhankuta district was intensive at the time of
19 sampling, and all five races identified in this study were present. Three races were
20 identified amongst the 13 strains from Lalitpur (Table 4) whereas 2 or 3 races were
21 identified on the basis of a more limited number of strains in the other districts. More
22 than one race was present in 11 out of 17 tested cabbage fields, on the basis of tests of
23 strains from two to three plants per field (Table 3).

1 Typical black rot symptoms were frequent on inoculated plants, and dark necrotic, blight
2 symptoms were observed in some cases. Blight-like symptoms were frequently produced
3 by race 7 strains and by some strains of races 5 and 6, and a few strains of race 1, but not
4 race 4 strains.

5 **Combining rep-PCR fingerprinting, race affiliation and geographical origin.** The
6 geographical origin of the Nepalese strains does not seem to be correlated with the rep-
7 PCR clusters shown in the combined dendrogram (Fig. 2). The clusters seem to be more
8 related to the race of the strains. All 12 Nepalese *Xcc* race 1 strains formed a single sub-
9 cluster at 85% similarity jointly with the race 1 reference strain (HRI 3811) from USA.
10 The *Xcc* race 4 strains were divided amongst two distinct clusters, containing nine and 11
11 strains, at 81% and 85% similarity, respectively. The latter group included the race 4
12 reference strain (HRI 1279A) with rep-PCR profiles from the present study and a race 4
13 strain (HRI 6189) from a previous study (40, 41). Strains from the Lalitpur district were
14 included in both race 4 clusters, even at the field level (strains N20/21 and N27). Most of
15 the race 1 strains and strains in each of the race 4 clusters might be identical. Two
16 Nepalese race 5 strains (N68 and N70) were identical and distinct from another race 5
17 Nepalese strain (N34). Nine race 6 and the two race 7 Nepalese *Xcc* cabbage strains were
18 included in the same cluster at 77% similarity. One Nepalese race 6 strain (N8) clustered
19 with the race 6 reference strain from Portugal.

20

1 **DISCUSSION**

2 The present study supports previous findings related to the occurrence of black rot in
3 Nepal and the diversity of the causal bacterium, *Xcc* (1, 27), and it confirms that the
4 disease is widespread in the vegetable Brassica growing areas in the country. Symptoms
5 of black rot were observed in all visited fields, in seedbeds, in newly transplanted crops
6 and in mature crops. The disease is frequently overlooked by farmers and extension
7 workers, who observe symptoms, but mistakenly think they are related to natural plant
8 senescence. The increased and almost continuous production of Brassica crops in some
9 areas, the ability of *Xcc* to spread by seed movement (42) and rain splash (43) and to
10 survive in plant debris left in the field (24), may cause build up and maintenance of
11 inoculum, thereby increasing the impact of the disease in the future, unless appropriate
12 control measures are adopted.

13 A significant proportion of cabbage seed is imported to Nepal (32), and it is important to
14 ensure that such seed is tested and free of the pathogen to minimise the risk of
15 introduction of additional new races or pathotypes. Selection of cultivars with disease
16 resistance is another means of control (12, 34, 43), which should be considered in
17 combination with crop rotation and sanitation schemes.

18 V-shaped leaf lesions with black veins, typical of black rot (6, 44), were frequently
19 observed in the fields, and *Xcc* was isolated from these lesions regardless of the crop
20 growth stage. Blight symptoms, characterised by a sudden collapse of interveinal tissues
21 and lack of veinal necrosis at earlier stages of infection, similar to those observed in
22 Hawaii (29, 45) and in the UK (S.J. Roberts, unpublished), were not observed in the
23 Nepalese fields. However, during race-typing under controlled conditions, we observed

1 both typical black rot symptoms and blight-like symptoms especially in plants inoculated
2 with strains of races 5, 6 and 7. A previous study has also shown that inoculation of *B.*
3 *oleracea* accessions with race 5 and 6 strains more commonly produced blight symptoms
4 than when inoculated with other races (40). The blight symptoms appeared similar to
5 those previously described (2) after artificial inoculation under greenhouse conditions.

6 In this study, duplicate isolates obtained from the same lesion produced similar rep-PCR
7 profiles, except in one case where two isolates belonged to different races. This suggests
8 that rep-PCR can be used for rapid initial screening of isolates to select non-identical
9 ones for further analysis, including race-typing. The race affiliation of the two isolates
10 from the same or adjacent lesions was tested twice to ensure that the result was
11 reproducible. The results in general indicate that most leaf lesions, from which isolations
12 were made, were a result of infection by a single race of the pathogen. However, the two
13 races obtained from the same leaf indicate that mixed infections, or leaf lesions with a
14 race mixture produced as the result of merged lesions from several infection points, may
15 develop if more than one race is present in the field. This is not unlikely as different races
16 were identified at the field level in more than half of the cabbage fields sampled in this
17 study.

18 Rep-PCR of strains of *Xcc* from different geographic regions of Nepal produced
19 fingerprinting profiles for races 1, 4 and 6 that clustered according to races and different
20 lineages therein. Nepalese race 1 strains formed a single cluster, and strains of races 4, 5
21 and 6 were split into at least two clusters. All strains of race 4 and 6, except one strain of
22 each race, clustered in two separate clusters, respectively. Previously, it has also been
23 found that *X. campestris* pv. *raphani* and a selection of *Xcc* strains clustered in

1 accordance with races (40). In contrast, it was concluded on the basis of rep-PCR (with
2 REP and BOX primers only) that fingerprints of *Xcc* strains from Tanzania (18) and
3 Nepalese strains (27) were related to geographical origin, but the races of these strains
4 were not determined and it is possible that different races were present in different areas.
5 We found (data not shown) that the ERIC primers, which were not included in that study
6 (18), are the most important for discriminating between *Xcc* races. ERIC-PCR also
7 revealed a higher level of genetic diversity than REP and BOX primers in a previous
8 study of *X. axonopodis* pv. *phaseoli* and pv. *phaseoli* var. *fuscans* (15). Another study
9 showed that two *Xcc* race 3 strains had different rep-PCR fingerprinting profiles (or were
10 grouped in different clusters) and suggested that race affiliation cannot be inferred by
11 DNA fingerprinting (36). However, the 14 Israeli strains were not race-typed. More
12 recently, results of fingerprints using BOX-PCR and M13-PCR of *Xcc* strains from Italy
13 were not correlated with geographical and host origin (45). Other rep-PCR studies of
14 other *Xanthomonas* species and pathovars have shown grouping of strains affiliated with
15 distinct symptom groups (22), geographical origin (25) and year of isolations (26). This
16 indicates that the potential to use rep-PCR to differentiate pathogenic variants within a
17 pathovar may depend on several parameters.

18 Five races of *Xcc* were found in Nepal with races 1, 4 and 6 dominating. The dominance
19 of races 1 and 4 has also been reported from other countries and the other races were
20 considered rare (40). A previous study (39), also showed the presence and dominance of
21 race 4, followed by race 1 and 6 amongst 51 strains from Portugal, mainly from *B.*
22 *oleracea*; in that study race 4 accounted for 53% of the strains, race 6 for 29% and race 1
23 for 18%. In contrast, race-typing of 102 strains from the UK mainly from *B. oleracea*

1 (including 68 strains from cauliflower) showed that race 1 dominated (69%) followed by
2 race 4 (30%); a single strain of race 3 was also identified in the UK collection (40). The
3 presence of race 5 has previously been reported for only three strains of older origin
4 (1953-1975) (40), which suggested that these races might not exist anymore. However,
5 our study revealed that race 5 still exists, although at low frequency (7%). The two
6 lineages of races 4 and 6 may suggest two different origins for these races, or that races 4
7 and 6 have evolved into two distinct lineages, which might even be different races, not
8 yet identified by lack of discerning differentials.

9 Two strains, N35 and N47, were pathogenic and very aggressive on all genotypes except
10 on *B. juncea* line FBLM2, and were assigned to race 7. Differences in the ability of the
11 strains to cause disease in artificial inoculation tests were also revealed previously with
12 Nepalese strains from the Katmandu and Chitwan valleys (1). Florida Broad Leaved
13 Mustard (FBLM2) is compatible with (i.e. susceptible to) *Xcc* race 6, but not with the
14 new race 7 (8, 13). It has been shown that mutants of *Xcc* race 1 with insertions or
15 deletions in *avrXccFM* become virulent on Florida Broad Leaf Mustard (5). Similarly, a
16 single gene difference may in this way determine the difference in compatibility between
17 races 6 and 7 on Florida Broad Leaf Mustard. The rep-PCR clustering of the two races
18 indicate a close relatedness between these two races, so it is possible that they may differ
19 only in the presence of an *avr* gene.

20 More than one race was identified in all districts, and there were no obvious differences
21 in the race composition within the geographically isolated regions sampled. Most
22 cabbage seed is imported, and during sampling it became evident that one cultivar
23 dominated in all growing areas. This may imply that the pathogen was introduced into the

1 areas with the seed. However, many factors play a role in the occurrence and diversity of
2 the pathogen, for instance tight crop rotation schemes with cabbage and other Brassica
3 crop species, and may together contribute to the occurrence and diversity of the pathogen.
4 Race-typing of strains is time, space and labour intensive. This study has shown that
5 fingerprinting of strains using the combined rep-PCR profiles (REP, ERIC, and BOX
6 primer sets) has potential to differentiate species, pathovars, pathotypes and lineages
7 within *Xcc*. The presence of several races of the bacterium within Nepal, including the
8 new race 7, emphasises the need for local surveys of *Xcc* to gain insight in the population
9 composition and change. The results may be of value in relation to regulatory issues, such
10 as seed import, but also highlight the need to account for pathogenic diversity when
11 breeding, introducing and cultivating resistant cultivars to control the disease.

12

13 **ACKNOWLEDGEMENTS**

14 This study was supported by Danida, Denmark (RUF, grant no. 91013, and 91116) and
15 by the Department for Environment, Food and Rural Affairs (Defra), UK. Some of the
16 work in the UK was done under a Defra Plant Health Licence (PHL 166C/4499). We
17 thank Liz Bailey and Sandy McClement from WHRI, University of Warwick for
18 providing doubled haploid accessions. In Nepal, we are greatly indebted to Mr.
19 Purushottam Prasad Kathiwada, Agriculture Research Station, Pakhribas, Mr. Birendra
20 A. C., Dhaulagiri Community Resource and Development Centre, Baglung, and Dr.
21 Kedar Budhatoki, Horticulture Research Division, Nepal Agricultural Research Council,
22 Khumaltar, for locating the crops, and to Gopal P. Parajuli, Plant Pathology Division,
23 Nepal Agricultural Research Council, Khumaltar for assistance with the collection of

- 1 infected leaf samples. We thank all farmers for welcoming us onto their land and
- 2 allowing collection of diseased material.

Table 1. Cabbage growing areas in the lower hills of the Himalayas and the Kathmandu valley visited July 2001, the altitude, number of fields sampled with black rot symptoms, number of fields from which *Xanthomonas campestris* pv. *campestris* (*Xcc*) isolation was attempted, and the number of strains included in rep-PCR and race typing.

Region	Zone	District	Altitude (m)	Sampled fields with black rot symptoms	Fields from which isolation was carried out	Number of strains ^a	
						Pathogenicity test positive and rep-PCR	race-typed ^b
Western	Dhawalagiri	Baglung	1521-69	3	2	4 (+3)	4 (+1)
Central	Bagmati	Lalitpur	1430-84	7	5	13 (+7)	13 (+2)
Central	Bagmati	Bhaktapur	1296-1304	2	2	5 (+4)	5 (+1)
Central	Narayani	Makawanpur	1710-1950	7	3	6 (+6)	6 (+1)
Eastern	Koshi	Dhankuta	1734-2139	9	6	11 (+10)	11 (+2 ^c)
Total				28	18	39 (+30)	39 (+7 ^c)

^aNumber of strains obtained from individual plants; parentheses indicate the number of duplicate strains that were also tested (+x).

^bRace typing included one pair of strains from the same plant that had different rep-PCR profiles^c.

1

2

Table 2. Brassica accessions used in pathogenicity tests to identify *Xanthomonas campestris* pv. *campestris* races

Genotype	Species	Seed source ^a	Type	Interaction with race ^b						
				1	2	3	4	5	6	7
Wirosa F ₁	<i>B. oleracea</i>	Bejo	Hybrid	+	+	+	+	+	+	+
Just Right Hybrid Turnip	<i>B. rapa</i>	Otis	Hybrid	+	+	+	-	+	+	+
Cob60	<i>B. napus</i>	WHRI	Doubled haploid	+	+	+	-	+	+	+
Seven Top Turnip	<i>B. rapa</i>	Otis	Open pollinated	+	-	+	-/+	-/+	+	+
Bo99075 (PI 199947)	<i>B. carinata</i>	WHRI	Inbred line	-	+	-	-	+	+	+
FBLM 2	<i>B. juncea</i>	Otis	Doubled haploid	-	+	-	-	-/(+)	+	-
Miracle F ₁	<i>B. oleracea</i>	Bejo	Hybrid	+	-	-	+	-	(+)/+	+
SxD1	<i>B. oleracea</i>	WHRI	Doubled haploid	+	-	-	+	-	(+)/+	+

^a Bejo= Bejo Zaden, Warmenhuizen, Holland; WHRI= Warwick HRI, The University of Warwick, UK; Otis=Otis S. Twilley Seed Co., US.

^b +, compatible interaction (susceptible host); -, incompatible interaction (resistant host); (+), weakly pathogenic; -/+, variable result within accession. Adapted from Vicente *et al.* (2001) based on unpublished results and results of the present study.

1

2

Table 3. The origin of *Xanthomonas campestris* pv. *campestris* strains and the results of pathogenicity testing and race-typing.

HRI number	Original number ^a	Crop	District/ Field number	Pathogenicity Cph. Market ^b	Pathogenicity Wirosa	Race
<i>Collected in 2001</i>						
8429 A	N1	Cabbage	Lalitpur/2	+	nt	6 ^c
8430	N5	Cabbage	Lalitpur/2	+	nt	4
8431 A	N6	Cabbage	Lalitpur/3	+	nt	4 ^c
8432	N8	Cabbage	Lalitpur/3	+	nt	6
8434	N11	Cabbage	Lalitpur/4	+	nt	1
8435 A	N12	Cabbage	Lalitpur/4	+	nt	4
8388 A/B	N14/N15	Cabbage	Lalitpur/4	+	nt	4
8389	N16	Cabbage	Lalitpur/4	+	nt	4
8436 B	N18	Cabbage	Lalitpur/5	+	nt	4
8437 A	N25	Cabbage	Lalitpur/5	+	+	1
8390 A/B	N20/N21	Cabbage	Lalitpur/6	+	+	4 ^c
8439 A	N22	Cabbage	Lalitpur/6	+	nt	1
8440	N27	Cabbage	Lalitpur/6	+	nt	4
8441 A	N29	Cabbage	Dhankuta/3	+	nt	1
8442 A	N31	Cabbage	Dhankuta/3	+	nt	1
8443 B	N34	Cabbage	Dhankuta/5	+	nt	5 ^c
8444A	N35	Cabbage	Dhankuta/5	+	nt	7 ^c
8445 A/B	N37/N38	Cabbage	Dhankuta/6	+	nt	1

8446	N40	Cabbage	Dhankuta/7	+	nt	6 ^c
8447 A	N41	Cabbage	Dhankuta/7	+	nt	1
8448 A	N43	Cabbage	Dhankuta/8	+	nt	6 ^c
8449 A	N45	Cabbage	Dhankuta/8	+	nt	4
8450 A	N47	Cabbage	Dhankuta/9	+	nt	7 ^c
8450 B	N48	Cabbage	Dhankuta/9	+	nt	6 ^c
8451 A	N49	Cabbage	Dhankuta/9	+	nt	6
8452 A/B	N51/N52	Cabbage	Makawanpur/1	+	nt	1
8453 B	N54	Cabbage	Makawanpur/1	+	nt	4
8454 A	N55	Cabbage	Makawanpur/4	+	nt	6 ^c
8455 A	N57	Cabbage	Makawanpur/4	+	nt	1
8456 A	N59	Cabbage	Makawanpur/7	+	nt	1
8457 A	N61	Cabbage	Makawanpur/7	+	nt	1
8458 A/B	N63/N64	Cabbage	Baglung/1	+	nt	6 ^c
8459 B	N66	Cabbage	Baglung/1	+	+	6
8460 B	N68	Cabbage	Baglung/2	+	nt	5 ^c
8461 B	N70	Cabbage	Baglung/2	+	+	5
8462 A/B	N71/N72	Cabbage	Bhaktapur/1	+	+	1
8463 A	N77	Cabbage	Bhaktapur/1	nt	+	4
8464 B	N74	Cabbage	Bhaktapur/3	+	nt	4
8465 A	N75	Cabbage	Bhaktapur/3	+	nt	4
8466	N79	Cabbage	Bhaktapur/3	nt	+	4

Collected in 2003

8467 A	N81	Cauliflower	Dhankuta/10	nt	+	4
8468 A	N83	Cauliflower	Lalitpur/7	nt	+	4
8469	N85	Cauliflower	Lalitpur/8	+	+	4

8471	N87	Leaf mustard	Lalitpur/10	+	+	6 ^c
8472	N88	Cauliflower	Lalitpur/11	nt	+	4

^a Two numbers indicate that duplicate strains from the same plant were tested with the same results.

^b +, compatible interaction; nt, not tested.

^c The duplicate strains or one strain were tested twice in two separate experiments.

Table 4. Frequency of occurrence of *Xanthomonas campestris* pv. *campestris* races among 40 strains obtained from cabbage plants in five districts in Nepal.

District	Number of strains of race					Total
	1	4	5	6	7	
Baglung			2	2		4
Lalitpur	3	8		2		13
Bhaktapur	1	4				5
Makawanpur	4	1		1		6
Dhankuta	4	1	1	4 ^a	2 ^a	12
Total	12	14	3	9	2	40
	(30.0%)	(35.0%)	(7.5%)	(22.5%)	(5%)	

^a In one case, two races were obtained from the same lesion (plant).

Fig. 1. Cabbage growing areas (districts) from which leaves with typical black rot symptoms were collected in 2001 and confirmed to be caused by *Xanthomonas campestris* pv. *campestris* in the present study. (Map courtesy of Matt Rosenberg, Geography at About.com: <http://geography.about.com>).

Fig. 2. Dendrogram of the genetic similarity of 40 race-typed Nepalese *Xcc* strains from cabbage, four from cauliflower and one from leaf mustard, analysed together with two control strains of *Xcc* (HRI 1279A and 3811), and combined with previous fingerprinting data from 13 *Xcc* and eight other *Xanthomonas* sp. strains (including representative strains of *X. campestris* pvs. *raphani* and *incanae*, *X. campestris* strains from wallflower and candytuft, *X. vesicatoria* and *X. axonopodis* pv. *vesicatoria*) from Vicente *et al.* (2006). Race type strains of *Xcc* and *X. campestris* pv. *raphani* and type strain of *X. vesicatoria* are indicated with a T. Similarities were calculated from the combined data of PCR fingerprinting using repetitive extragenic palindromic (REP), enterobacterial repetitive intergenic consensus (REP) and BOX primer sets. Strain numbers without a prefix are HRI numbers. Solid and dashed lines delimit the most important groups of strains at 70 and 80% similarity, respectively.

Literature Cited

1. Adhikari, T. B., and Basnyat, R. 1999. Phenotypic characteristics of *Xanthomonas campestris* pv. *campestris* from Nepal. Eur. J. Plant Pathol. 105:303-305.

2. Alvarez, A. M., Benedict, A. A., Mizumoto, C. Y., Hunter, J. E., and Gabriel, D. W. 1994. Serological, pathological, and genetic diversity among strains of *Xanthomonas campestris* infecting crucifers. *Phytopathology*. 84:1449-1457.
3. Anonymous. 2006. Annual Report. Vegetable Development Directorate, Department of Agriculture, Khumaltar, Lalitpur, Nepal.
4. Cab International. 2006. Crop Protection Compendium, 2006 Edition. CAB International, Wallingford, UK.
5. Castañeda, A., Reddy, J. D., El-Yacoubi, B., and Gabriel, D. W. 2005. Mutagenesis of all eight *avr* genes in *Xanthomonas campestris* pv. *campestris* had no detected effect on pathogenicity, but one *avr* gene affected race specificity. *Mol. Plant Microbe Interact.* 18:1306-1317.
6. Cook, A. A., Walker, J. C., and Larson, R. H. 1952. Studies on the disease cycle of black rot of crucifers. *Phytopathology* 42:162-167.
7. De Bruijn, F. J. 1992. Use of repetitive (repetitive extragenic palindromic and enterobacterial repetitive intergeneric consensus) sequences and the polymerase chain reaction to fingerprint the genomes of *Rhizobium meliloti* isolates and other soil bacteria. *Appl. Environ. Microbiol.* 58:2180-2187.
8. Fargier, E., and Manceau, C. 2007. Pathogenicity assays restrict the species *Xanthomonas campestris* into three pathovars and reveal nine races within *X. campestris* pv. *campestris*. *Plant Pathol.* 56:805-818.
9. Feltham, R. K. A., Power, A. K., Pell, P. A., and Sneath, P. H. A. 1978. A simple method for storage of bacteria at -76°C. *J. Appl. Bacteriol.* 44:313-316.

10. Higgins, C. F., Ames, G. F.-L., Barnes, W. M., Clement, J. M., and Hofnung, M. 1982. A novel intercistronic regulatory element of prokaryotic operons. *Nature* 298:760-762.
11. Ignatov, A., Hida, K., and Kuginuki, Y. 1998. Black rot of crucifers and sources of resistance in brassica crops. *Jpn. Agric. Res. Q.* 32:167-172.
12. Jensen, B. D., Massomo, S. M. S., Swai, I. S., Hockenhull, J., and Andersen, S. B. 2005. Field evaluation for resistance to the black rot pathogen *Xanthomonas campestris* pv. *campestris* in cabbage (*Brassica oleracea*). *Eur. J. Plant Pathol.* 113:297-308.
13. Jensen, B. D., Vicente, J. G., Manandhar, H. K., and Roberts, S. J. 2007. An increasing risk for bacterial plant diseases in Northern Europe? Occurrence and diversity of *Xanthomonas campestris* pv. *campestris* in vegetable Brassica fields in Nepal and identification of a new race. Abstract Book. Plant Biotech Denmark, Annual meeting, Faculty of Life Sciences, University of Copenhagen, Denmark, 23-24th January.
14. Kamoun, S., Kamdar, H. V., Tola, E., and Kado, C. I. 1992. Incompatible interactions between crucifers and *Xanthomonas campestris* involve a vascular hypersensitive response: role of the hrpx locus. *Mol. Plant Microbe Interact.* 5:22-33.
15. López, R., Asensio, C., and Gilbertson, R. L. 2006. Phenotypic and genetic diversity in strains of common blight bacteria (*Xanthomonas campestris* pv. *phaseoli* and *x. campestris* pv. *phaseoli* var. *fuscans*) in a secondary center of diversity of the common bean host suggests multiple introduction events. *Phytopathology* 96:1204-1213.
16. Louws, F. J., Fulbright, D. W., Stephens, C. T., and De Bruijn, F. J. 1994. Specific genomic fingerprints of phytopathogenic *Xanthomonas* and *Pseudomonas* pathovars and strains generated with repetitive sequences and PCR. *Appl. Environ. Microbiol.* 60:2286-2295.
17. Martin, C., Briese, T., and Hakenbeck, R. 1992. Nucleotide-sequences of genes encoding penicillin-binding proteins from *Streptococcus pneumoniae* and *Streptococcus oralis* with

- high homology to *Escherichia coli* penicillin-binding proteins 1A and 1B. *J. Bacteriol.* 174:4517-4523.
18. Massomo, S. M. S., Nielsen, H., Mabagala, R. B., Mansfeld-Giese, K., Hockenhull, J., and Mortensen, C. N. 2003. Identification and characterisation of *Xanthomonas campestris* pv. *campestris* strains from Tanzania by pathogenicity tests, Biolog, rep-PCR and fatty acid methyl ester analysis. *Eur. J. Plant Pathol.* 109:775-789.
 19. Pierce, L., Schroth, M. N., and McCain, A. H. 1990. Viscosity test for preliminary identification of strains of *Xanthomonas campestris*. *Plant Dis.* 74:646-647.
 20. Rademaker, J. L. W., Louws, F. J., Schultz, M. H., Rossbach, U., Vauterin, L., Swings, J., and De Bruijn, F. J. 2005. A comprehensive species to strain taxonomic framework for *Xanthomonas*. *Phytopathology* 95:1098-1111.
 21. Rademaker, J. L. W., Norman, D. J., Forster, R. L., Louws, F. J., Schultz, M. H., and De Bruijn, F. J. 2006. Classification and identification of *Xanthomonas translucens* isolates, including those pathogenic to ornamental asparagus. *Phytopathology* 96:876-884.
 22. Sahin, F., Abbasi, P. A., Ivey, M. L. L., Zhang, J., and Miller, S. A. 2003. Diversity among strains of *Xanthomonas campestris* pv. *vitians* from lettuce. *Phytopathology* 93:64-70.
 23. Schaad, N. W., and Kendrick, R. 1975. A qualitative method for detecting *Xanthomonas campestris* in crucifer seed. *Phytopathology* 65:1034-1036.
 24. Schaad, N. W., and White, W. C. 1974. Survival of *Xanthomonas campestris* in soil. *Phytopathology* 64:1518-1520.
 25. Scortichini, M., Marchesi, U., and Di Prospero, P. 2001. Genetic diversity of *Xanthomonas arboricola* pv. *juglandis* (synonyms: *X. campestris* pv. *juglandis*; *X. juglandis* pv. *juglandis*)

- strains from different geographical areas shown by repetitive polymerase chain reaction genomic fingerprinting. *J. Phytopathol.* 149:325-332.
26. Scortichini, M., and Rossi, M. P. 2003. Genetic diversity of *Xanthomonas arboricola* pv. *fragariae* strains and comparison with some other *X. arboricola* pathovars using repetitive PCR genomic fingerprinting. *J. Phytopathol.* 151:113-119.
 27. Shakya, D. D., Louws, F. J., and Alvarez, A. M. 2000. Diversity of *Xanthomonas campestris* pv. *campestris* (*Xcc*) populations in Nepal. *Phytopathology* 90:S71.
 28. Sharples, G. J., and Lloyd, R. G. 1990. A novel repeated DNA sequence located in the intergenic regions of bacterial chromosomes. *Nucleic Acids Res.* 18:6503-6508.
 29. Shigaki, T., Nelson, S. C., and Alvarez, A. M. 2000. Symptomless spread of blight-inducing strains of *Xanthomonas campestris* pv. *campestris* on cabbage seedlings in misted seedbeds. *Eur. J. Plant Pathol.* 106:339-346.
 30. Shrestha, K., Mathur, S. B., and Neergaard, P. 1977. Seed-borne organisms in some crops of Nepal. *Seed Sci. & Technol.* 5:111-121.
 31. Stolp, H., and Starr, M. P. 1964. Bacteriophage reactions and speciation of phytopathogenic xanthomonads. *J. Phytopathol.* 51:442-478.
 32. Subedi, G. D., and K.C, R. B. 2004. Scenario of hybrid vegetable seed use in Nepal. Pages 275-281. B. B. Khatri, B. P. Sharma, P. P. Khatiwada, K. P. Paudyal, B. R. Khadge, and H. N. Regmi, eds. in: *Advances of Horticultural Research in Nepal. Proceedings of the Fourth National Workshop on Horticulture, March 2-4.* Nepal Agricultural Research Council, National Agricultural Research Institute and Horticulture Research Division, Khumaltar, Lalitpur, Nepal.

33. Sutton, J. C., and Williams, P. H. 1970. Relation of xylem plugging to black rot lesion development in cabbage. *Can. J. Bot.* 48:391-401.
34. Taylor, J. D., Conway, J., Roberts, S. J., Astley, D., and Vicente, J. G. 2002. Sources and origin of resistance to *Xanthomonas campestris* pv. *campestris* in Brassica genomes. *Phytopathology* 92:105-111.
35. Tsygankova, S. V., Ignatov, A. N., Boulygina, E. S., Kuznetsov, B. B., and Korotkov, E. V. 2004. Genetic relationships among strains of *Xanthomonas campestris* pv. *campestris* revealed by novel rep-PCR primers. *Eur. J. Plant Pathol.* 110:845-853.
36. Valverde, A., Hubert, T., Stolov, A., Dagar, A., Kopelowitz, J., and Burdman, S. 2007. Assessment of genetic diversity of *Xanthomonas campestris* pv. *campestris* isolates from Israel by various DNA fingerprinting techniques. *Plant Pathol.* 56:17-25.
37. Versalovic, J., Koeuth, T., and Lupski, J. R. 1991. Distribution of repetitive DNA-sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Res.* 19:6823-6831.
38. Versalovic, J., Schneider, M., De Bruijn, F. J., and Lupski, J. R. 1994. Genomic fingerprinting of bacteria using repetitive sequence-based polymerase chain reaction. *Methods Mol. Cell Biol.* 5:25-40.
39. Vicente, J. G. 2004. A Podridão Negra das Crucíferas. Centro Operativo e Tecnológico Hortofrutícola Nacional, Alcobaça, Portugal. 102 pp.
40. Vicente, J. G., Conway, J., Roberts, S. J., and Taylor, J. D. 2001. Identification and origin of *Xanthomonas campestris* pv. *campestris* races and related pathovars. *Phytopathology* 91:492-499.

41. Vicente, J. G., Everett, B., and Roberts, S. J. 2006. Identification of isolates that cause a leaf spot disease of brassicas as *Xanthomonas campestris* pv. *raphani* and pathogenic and genetic comparison with related pathovars. *Phytopathology* 96:735-745.
42. Walker, J. C., and Tisdale, W. B. 1920. Observations on seed transmission of the cabbage black rot organism. *Phytopathology* 10:175-177.
43. Williams, P. H. 1980. Black rot: a continuing threat to world crucifers. *Plant Dis.* 64:736-742.
44. Williams, P. H., Staub, T., and Sutton, J. C. 1972. Inheritance of resistance in cabbage to black rot. *Phytopathology* 62:247-252.
45. Yuen, G. Y. K., and Alvarez, A. M. 1985. Aberrant symptoms on cabbage caused by strains of *Xanthomonas campestris*. *Phytopathology* 75:1382.
46. Zaccardelli, M., Campanile, F., Moretti, C., and Buonauro, R. 2008. Characterization of Italian populations of *Xanthomonas campestris* pv. *campestris* using primers based on DNA repetitive sequences. *J. Plant Pathol.* 90:375-381.