



The OPERA experimental program with CNGS

D. Duchesneau

► **To cite this version:**

D. Duchesneau. The OPERA experimental program with CNGS. International Workshop on Topics in Astroparticle and Underground Physics 8 TAUP2003, Sep 2003, Seattle, United States. 138, pp.411-414, 2005, <10.1016/j.nuclphysbps.2004.11.094>. <in2p3-00014170>

HAL Id: in2p3-00014170

<http://hal.in2p3.fr/in2p3-00014170>

Submitted on 25 Nov 2003

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LAPP-EXP 2003-25
November 2003

The OPERA experimental program with CNGS

D. Duchesneau

LAPP-IN2P3-CNRS
9 chemin de Bellevue - BP. 110
F-74941 Annecy-le-Vieux Cedex

**Invited talk at the 8th International Workshop on Topics in Astroparticle
and Underground Physics (TAUP2003), Seattle, Etats-Unis, September 5-9, 2003**

The OPERA Experimental Program with CNGS

D. Duchesneau^a *

^aLAPP, IN2P3-CNRS,
9 Chemin de Bellevue, BP110, F-74941, Annecy-le-Vieux, France
(email: duchesneau@lapp.in2p3.fr)

This paper reviews the current status of the OPERA detector preparation which is designed to investigate the neutrino oscillation properties using the future CERN neutrino beam called CNGS. The physics potential and performances for neutrino oscillation studies including $\nu_\mu \rightarrow \nu_e$ search will also be presented.

1. Introduction

OPERA [1] is a first generation long baseline neutrino experiment to be located in the Gran Sasso underground laboratory. The detector is a massive hybrid detector with nuclear emulsions used as very precise tracking devices and electronic detectors to locate the neutrino interaction events in the emulsions.

It is designed to primarily search for ν_τ appearance in the CERN high energy ν_μ beam CNGS [2] at 730 km from the neutrino source, in order to establish unambiguously the origin of the neutrino oscillations observed at the atmospheric Δm^2 scale. The best fit of the $\nu_\mu \rightarrow \nu_\tau$ oscillation hypothesis to the Super-Kamiokande atmospheric data gives the oscillation parameters $\Delta m^2 = 2.0 \times 10^{-3} \text{eV}^2$ with $\sin^2 2\theta = 1.0$ [3]. The range of allowed values at 90% CL correspond to $1.3 \times 10^{-3} < \Delta m^2 < 3.0 \times 10^{-3} \text{eV}^2$ and $\sin^2 2\theta > 0.90$.

The CNGS neutrino beam is a high energy beam optimised for ν_τ appearance with a mean neutrino energy of about 17 GeV. The beamline characteristics are summarised in Ref. [4] and the construction status of the CNGS project can be found in Ref. [5]. The possibility of an increase of the neutrino beam intensity by a factor 1.5 compared to the original design is under study [6]. However the feasibility for this upgrade should await results of tests performed in 2003.

Using the CERN SPS accelerator in a shared

*Representing the OPERA Collaboration.

mode, 6.76×10^{19} protons on target (pot) can be delivered during one year, assuming 200 days of operation. The number of charged current and neutral current interactions expected in the Gran Sasso laboratory from ν_μ are about 4000 /kton/year and 1240 /kton/year respectively. If the $\nu_\mu \rightarrow \nu_\tau$ oscillation hypothesis is confirmed, the number of τ 's produced via charged current interaction at the Gran Sasso should be of the order of 16 /kton/year for $\Delta m^2 = 2.0 \times 10^{-3} \text{eV}^2$ at full mixing.

2. Status of the detector construction

The ν_τ appearance search is based on the observation of events produced by charged current interaction (CC) with the τ decaying in all possible decay modes. The principle of the OPERA experiment is to observe the τ trajectories and the decay products in emulsion films composed of two thin emulsion layers (50 μm thick) put on either side of a plastic base (200 μm thick). To achieve a large target mass, the emulsion films are interleaved with 1 mm thick lead plates. The basic detector unit, called ECC brick, is obtained by stacking 56 lead plates with 57 emulsion films. In order to reach 1.8 kton target mass, 206336 bricks will be installed into walls separated from each other by vertical planes of electronic target trackers. A more complete description of the detector setup can be found in Ref [4].

A brief summary of the construction status of the different detector parts follows.

2.1. The Emulsion Films and Lead

The emulsion films are produced in Japan by FUJI company and the mass production started in April 2003 at a rate of about 8000 m²/month. The entire production will correspond to a total of 150 000 m². To erase the cosmic ray tracks recorded during film production and subsequent manipulation, a refreshing process, consisting of leaving the emulsion in a 95% relative humidity during 3 days at a temperature of 30°C, is performed in the Tono mine in Japan. This procedure allows to erase more than 98% of the tracks before transportation to Gran Sasso.

The lead used for the bricks consists of low radioactivity lead (Boliden) with 0.7% Ca added for rigidity purpose. The lead thickness is controlled with a 10 μm accuracy. A prototype production is scheduled with the Goslar firm in Germany.

2.2. The Target Tracker

The electronic target tracker is composed of X and Y planes of 256 AMCRYS-H plastic scintillator strips (6.7 m x 2.5 cm x 1 cm). Each plane is divided in 4 modules. Each strip is read out on both ends by a Kuraray wave length shifting optical fiber connected to 64 channel Hamamatsu photomultiplier tubes. The main goal of the electronic detector is to provide a trigger with an efficiency greater than 99% for neutrino interactions and an efficient localisation of the bricks where the events occur.

The construction and assembly of the target tracker modules take place in Strasbourg. The production rate should be of 8 modules per week. Delivery and installation at the Gran Sasso laboratory is planned to start beginning of 2004.

2.3. The Muon spectrometer

The muon spectrometer allows a determination of the charge and momentum of muons going through by measuring their curvature in a dipolar magnet which provides 1.6 Tesla transverse to the neutrino beam axis. Each spectrometer is equipped with six vertical planes of drift tubes as precision tracker together with 22 planes (8x8 m²) of RPC bakelite chambers. The RPC's are located in the magnets between the 5 cm thick vertical iron slabs. The installation of the final

spectrometers in the Hall C of the Gran Sasso underground laboratory has already started since May 2003. The iron slabs, the yokes, the coils and the power supplies have been ordered in the mean time. The first batch of RPC chambers are expected to arrive in November 2003 at Gran Sasso to be installed in the first spectrometer.

The precision tracker planes are composed of 4 staggered layers of 168 aluminium tubes, 8 m long with 38 mm outer diameter. A full scale prototype module has been built and tested in Hamburg. The spatial resolution of this detector is better than 500 μm.

The physics performance of the complete spectrometer reduces the charge confusion to less than 0.3% and gives a momentum resolution better than 20% for momentum less than 50 GeV. The muon identification efficiency reaches 95% using the target tracker information for the cases where the muons stop inside the target.

3. Physics performance: $\nu_\mu \rightarrow \nu_\tau$ search

The τ decay channels investigated by OPERA are the e, μ and hadron. Table 1 summarises the OPERA performance after 5 years of running with the 50% increased CNGS intensity (3.35x10²⁰ pot). The numbers in parenthesis are obtained from the nominal CNGS intensity delivering 2.25x10²⁰ pot in 5 years. The number of expected signal events from $\nu_\mu \rightarrow \nu_\tau$ oscillation is given as a function of the studied channel for three different values of Δm^2 at full mixing. The total efficiency including the branching ratios amounts to 9.1% and the total background is estimated to be less than 1.06 event. The main background sources are charm decays (54%), large angle muon scattering (16%) and hadron reinteractions (29%). A probability larger than 95% to have a 4 σ significant effect after 5 years is reached for $\Delta m^2 > 2.0 \times 10^{-3} \text{eV}^2$. This probability drops to 45% for $\Delta m^2 = 1.3 \times 10^{-3} \text{eV}^2$.

4. Search for $\nu_\mu \rightarrow \nu_e$ appearance

Having excellent electron identification capabilities, OPERA has estimated its sensitivity in searching for $\nu_\mu \rightarrow \nu_e$ appearance [7] with the

Table 1

Summary of the expected numbers of τ events in 5 years for different Δm^2 with the expected background and detection efficiencies per decay channel for OPERA. The numbers in parenthesis are obtained with the initial CNGS intensity design

channel	signal for Δm^2 (eV^2)			$\epsilon \times \text{Br}$	Background
	1.3×10^{-3}	2.0×10^{-3}	3.0×10^{-3}		
$\tau \rightarrow e$	1.8 (1.2)	4.1 (2.7)	9.2 (6.1)	3.4%	0.31 (0.21)
$\tau \rightarrow \mu$	1.4 (0.9)	3.4 (2.3)	7.6 (5.1)	2.8%	0.33 (0.22)
$\tau \rightarrow h$	1.5 (1.0)	3.5 (2.3)	7.8 (5.2)	2.9%	0.42 (0.28)
Total	4.7 (3.1)	11.0 (7.3)	24.6 (16.4)	9.1%	1.06 (0.71)

CNGS beam. The analysis principle is based on a search for an excess of ν_e CC events at low neutrino energies. The main background comes from the electron neutrino contamination (0.8%) present in the beam. The analysis takes into account the electron events coming from $\nu_\mu \rightarrow \nu_\tau$ events where $\tau \rightarrow e \nu_\tau \nu_e$ since both oscillations would occur at the atmospheric Δm^2 scale. These events distort the kinematical distributions where the low energy events contribute. The sensitivity to θ_{13} is obtained by doing a χ^2 minimisation using the visible energy, the missing transverse energy and the electron energy distributions in which the oscillation parameter varies. Table 2 summarises the expected number of selected events, assuming 5 years running and still the original CNGS design intensity (2.25×10^{20} pot), from $\nu_\mu \rightarrow \nu_e$ and $\nu_\mu \rightarrow \nu_\tau$ oscillations and from background ν_e at 3 values of θ_{13} . Background events coming from ν_μ CC and ν_μ NC interactions amount to 1.0 and 5.2 respectively.

Table 2

Expected number of signal and background events in 5 years obtained in the search of $\nu_\mu \rightarrow \nu_e$ oscillation at $\Delta m^2 = 2.5 \times 10^{-3} \text{eV}^2$.

θ_{13} (deg)	$\sin^2 2\theta_{13}$	ν_e CC	$\nu_\mu \rightarrow \nu_\tau$ $\tau \rightarrow e$	signal $\nu_\mu \rightarrow \nu_e$
9	0.095	18	4.5	9.3
7	0.058	18	4.6	5.8
5	0.030	18	4.6	3.0

The limit obtained by OPERA at 90% CL on θ_{13} is 7.1° after 5 years, which leads to significant improvement over the actual CHOOZ limit [8] and opens an important window on the third mixing angle.

5. Conclusion

The CNGS construction is progressing well. The project is on schedule and a startup is expected for June 2006. At the same time OPERA enters the construction phase and should be ready to take data by 2006.

The detector performances are such that the first evidence for $\nu_\mu \rightarrow \nu_\tau$ appearance signal could be seen after only a few years of data taking. The very good electron identification and measurement of the detector give the possibility to explore the $\nu_\mu \rightarrow \nu_e$ appearance channel pushing down the θ_{13} limit.

REFERENCES

1. OPERA Collaboration, M. Guler *et al.*, Experiment proposal, CERN-SPSC-2000-028 and LNGS P25/2000 (2000); Status Report on the OPERA experiment, CERN-SPSC-2001-025 and LNGS-EXP 30/2001 (2001).
2. G. Acquistapace *et al.*, CERN 98-02 and INFN/AE-98/105 (1998); R. Bailey *et al.*, "Addendum to Report CERN 98-02", CERN-SL 99-034 and INFN/AE-99/05 (1999); A.E. Ball *et al.*, SL-Note 2000-063 (2000).
3. Y. Hayato, contribution to the EPS conference, July 2003, Aachen, Germany.
4. D. Duchesneau, Nucl. Phys. B (Proc. Suppl.) 123 (2003) 279 [hep-ex/0209082].
5. F. Arneodo, these proceedings.
6. R. Cappi *et al.*, CERN-SL-2001-032 (2001).
7. M. Komatsu, P. Migliozzi and F. Terranova, J. Phys. G29 (2003) 443 [hep-ph/0210043].
8. CHOOZ Collaboration, M. Apollonio *et al.*, Phys. Lett. B 466 (1999) 415.