



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Acquisition of extended-spectrum beta-lactamase-producing Enterobacteriaceae (ESBL-PE) carriage after exposure to systemic antimicrobials during travel: systematic review and meta-analysis

Citation for published version:

Wuerz , TC, Kassim, SS & Atkins, KE 2020, 'Acquisition of extended-spectrum beta-lactamase-producing Enterobacteriaceae (ESBL-PE) carriage after exposure to systemic antimicrobials during travel: systematic review and meta-analysis', *Travel medicine and infectious disease*.
<https://doi.org/10.1016/j.tmaid.2020.101823>

Digital Object Identifier (DOI):

[10.1016/j.tmaid.2020.101823](https://doi.org/10.1016/j.tmaid.2020.101823)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Travel medicine and infectious disease

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Acquisition of extended-spectrum beta-lactamase-producing Enterobacteriaceae (ESBL-PE) carriage
after exposure to systemic antimicrobials during travel: systematic review and meta-analysis.

Terence C. Wuerz MD, MSc^{1,2,*}, Sameer S. Kassim MBBChBAO, MSc³, Katherine E. Atkins, Ph.D^{4,5}

¹St. Boniface General Hospital, Winnipeg, Manitoba, Canada

²Departments of Internal Medicine and Community Health Sciences, University of Manitoba, Winnipeg, Manitoba, Canada

³Department of Family Medicine, University of Manitoba, Winnipeg, Manitoba, Canada

⁴Centre for Mathematical Modelling of Infectious Diseases, London School of Hygiene and Tropical Medicine, London, United Kingdom

⁵Centre for Global Health Research, Usher Institute for Population Health Sciences and Informatics, Edinburgh Medical School, The University of Edinburgh, UK

*Corresponding Author: Dr Terence C. Wuerz, Department of Internal Medicine, University of Manitoba, C5002-1
– 409 Tache Ave, Winnipeg, MB, R2H 2A6, Canada. Telephone 204-237-2948, fax 204-233-7125

ABSTRACT

Background: International travel is an important risk factor for colonization with extended-spectrum beta-lactamase-producing Enterobacteriaceae (ESBL-PE). Antimicrobial use during travel likely amplifies this risk, yet to what extent, and whether it varies by antimicrobial class, has not been established.

Methods: We conducted a systematic review that included prospective cohorts reporting both receipt of systemic antimicrobials and acquired ESBL-PE isolated from stool or rectum during international travel. We performed a random effects meta-analysis to estimate odds of acquiring ESBL-PE due to antimicrobials during travel, overall and by antimicrobial class. **Results:** Fifteen studies were included. The study population was mainly female travellers from high income countries recruited primarily from travel clinics. Participants travelled most frequently to Asia and Africa with 10% reporting antimicrobial use during travel. The combined odds ratio (OR) for ESBL-PE acquisition during travel was 2.37 for antimicrobial use overall (95% confidence interval [CI], 1.69 to 3.33), but there was substantial heterogeneity between studies. Fluoroquinolones were the antibiotic class associated with the highest combined OR of ESBL-PE acquisition, compared to no antimicrobial use (OR 4.68, 95% CI, 2.34 to 9.37). **Conclusions:** The risk of ESBL-PE colonization during travel is increased substantially with exposure to antimicrobials, especially fluoroquinolones. While a small proportion of colonized individuals will develop a resistant infection, there remains the potential for onward spread among returning travellers. Public health efforts to decrease inappropriate antimicrobial usage during travel are warranted.

Research in context

Evidence before this study

Antimicrobial resistance (AMR) among bacteria that commonly cause human infection is of increasing public health concern. International travel has recently been associated with colonization with Extended-Spectrum Beta-Lactamase Producing-Enterobacteriaceae (ESBL-PE), increasing the spread of drug resistance among these important pathogens. We searched Pubmed, Embase, MEDLINE, Web of Science, SCOPUS, and the Cochrane Library for prospective cohort studies published between January 2000 and June 2018, reporting on acquisition of ESBL-PE among travellers, which reported on antimicrobial use during travel. 15 studies were included, which were at moderate risk of bias. The pooled odds ratio for acquisition of ESBL-PE during travel was 2.37 among antimicrobial users, compared to non-users (95% CI, 1.69 to 3.33). The magnitude of this association was stronger among travellers reporting fluoroquinolone use (OR 4.68, 95% CI 2.34 to 9.37).

Added value of this study

This is the first study to quantify the association between antimicrobial use during travel, overall and by specific antimicrobial class, with ESBL-PE acquisition across broad populations of travellers and destination countries.

Implications of all the available evidence

Further study into the mechanisms by which antimicrobials, such as fluoroquinolones, contribute to AMR may identify protective measures. Meanwhile, antimicrobial use during travel for prevention or treatment of mild-to-moderate traveller's diarrhea should not be recommended routinely. Where indicated, alternatives to fluoroquinolone antimicrobials should be considered.

INTRODUCTION

In 2014 the World Health Organization declared that a “post-antibiotic era” is within sight if urgent action was not taken¹. At the forefront of this growing public health threat is drug resistance in Enterobacteriaceae, a family of gram-negative bacteria which make up a large part of normal human gut flora. Extended-spectrum beta-lactamase (ESBL) enzymes are an important cause of increasing bacterial resistance globally^{2,3} with ESBL-producing Enterobacteriaceae (ESBL-PE) colonizing humans either de novo or through fecal-oral transmission⁴. ESBL enzymes render all penicillin, monobactam and expanded-spectrum cephalosporin antimicrobials^{2,5} ineffective. Carriers of ESBL-PE are usually asymptomatic⁴, but are at risk to develop clinical infection due to these organisms, resulting in increased cost, morbidity, and mortality compared to infections due to drug-susceptible Enterobacteriaceae^{4,6,7}.

The prevalence of ESBL-PE varies worldwide and by epidemiologic setting^{1,3,4}. For example, the frequency of ESBL-PE is relatively low in North America and Europe¹, and comparatively much higher in many lower income countries, especially in South Asia⁴. Accordingly, observational studies have established a strong association between asymptomatic ESBL-PE acquisition and international travel⁸⁻²³. The risk of acquiring carriage with ESBL-PE is over 20% with any international travel, and higher with travel to areas of particularly high prevalence such as South Asia^{15,24}. Few healthy travellers colonized with ESBL-PE will develop infection,^{25,26} however, an estimated 12% will transmit these bacteria to other household members⁹.

Antimicrobial use has been inconsistently reported as an independent risk factor for ESBL-PE acquisition among international travellers^{15,24}. Estimates around the degree of risk antimicrobial use poses, as well as the relative role that different antimicrobial classes play, have not been well established.

We conducted a systematic review and meta-analysis to determine the extent to which receipt of antimicrobials during travel increase the odds of faecal carriage with ESBL-PE compared to those that do not receive antimicrobials, as well as how much these odds vary between antimicrobial classes.

METHODS

We included only prospective cohort studies, related to travel across an international boundary, reporting screening for faecal Enterobacteriaceae carriage (in asymptomatic participants) both prior to and after travel, in our review. Studies must have presented phenotypic antimicrobial susceptibility data (or molecular equivalent) which is adequate to ascertain ESBL-PE presence or absence, and were excluded if ESBL-PE acquisition status by presence or absence of receipt of systemic antimicrobials taken during travel was not reported.

We searched Embase, MEDLINE, Web of Science, SCOPUS, Cochrane Library, and PubMed for studies published in peer-reviewed English journals from January 2000 to June 2018 (Box 1). The reference lists, and citations of included studies, were inspected to identify further potentially eligible studies. Citations were entered and housed in Covidence™, and abstracts were screened for eligibility by 2 independent reviewers (TW, SK). Studies selected for full text review were read in full to ascertain whether inclusion criteria were met and selected for data extraction as appropriate. Conflicts were resolved by consensus by 2 authors (TW, SK). Each included study was assessed using tools for assessing bias in observational studies as recommended in the STROBE statement²⁷ and modified to suit the study design.

Data were extracted by a single author (TW) from each included article and exported to Stata 13 for analysis. Data gathered included details on study population and travel characteristics, potential confounders, antimicrobials taken during travel, method of determining ESBL-PE pre- and post-travel, the prevalence of ESBL-PE pre- and post-travel, and study quality metrics. Odds ratios for acquisition of ESBL-PE, by presence or absence of antimicrobial exposure during travel, with 95% confidence intervals

of all included studies were displayed as a Forrest plot. Data were visually inspected for heterogeneity of results; Cochrane's Q and I^2 were calculated to quantify this heterogeneity. As there was substantial evidence of heterogeneity, a random effects model of meta-analysis, representing the average odds ratio of ESBL-PE due to antimicrobial exposure during travel, was built. For the secondary analysis, a random effects meta-analysis was undertaken of rates of ESBL-PE acquisition by antimicrobial class (beta-lactam, fluoroquinolone, macrolide or tetracycline) received during travel, compared to no antimicrobial receipt (the baseline group).

All antimicrobials including doxycycline were included in the primary analysis, however, doxycycline was excluded from this analysis in 2 studies, as it was counted as an antimalarial drug and the data grouping did not permit analysis compared no antibiotic receipt for these studies. We therefore also performed a sensitivity analysis by comparing the random effects meta-analysis which excluded doxycycline as an antimicrobial to the primary analysis which included. For all studies, we plotted effect size (log OR for ESBL-PE acquisition during travel related antimicrobial exposure) against log standard error of each study to create a funnel plot. The quality of the studies was assessed using the modified STROBE²⁷ tool.

Studies were excluded from the secondary analysis if no information was available on proportion of ESBL-PE acquisition stratified by antimicrobial class. Authors of the 10 studies missing crucial data for our secondary analysis were contacted to obtain it; 6 replies were received.

RESULTS

The search strategy identified 5323 journal articles in published peer-reviewed journals (Figure 1). 1893 duplicates were identified and removed, leaving 3430 study titles and abstracts, of which 3372 were excluded on screening. 58 published manuscripts were reviewed in full and 15 studies were included for the systematic review.

The 15 included prospective cohort studies were published from 2010 to 2017, and enrolled travellers from 2007 to 2015^{8-14,16-23}. The studies included 5283 participants (median, 205; range, 58 to 1965) and in all but 2 instances were conducted in Northern or Western Europe (Supplementary Table 1). The median age of participants varied substantially, but in all studies, fewer than 50% were male. 13 studies used a phenotypic approach for determination of ESBL-PE status (with polymerase chain reaction [PCR] for confirmation) whereas 2 studies adopted a molecular approach, using PCR to identify the gene encoding ESBL production within Enterobacteriaceae, bla_{CTX-M}, without culture^{14,20}.

Asia, including South Asia region to which travel carries the highest risk of ESBL-PE acquisition, was the most common region travelled to, followed by Africa (Table 1a). Median travel time ranged from 14-21 days in most studies, and the median percentage of travellers reporting diarrhea or gastroenteritis was 38% (range: 12 to 69%). In total, 550 participants (10% of the total) reported systemic antimicrobial usage (range by study: 4-49%). The most common indications were lower respiratory tract infection and traveller's diarrhea. Of participants reporting a specific class of antimicrobial use, beta-lactams (30%), fluoroquinolones (25%), and doxycycline (20%) were more common than macrolides (8%). A total of 1748 participants acquired ESBL-PE carriage during travel; this also varied significantly by study, from 9% of to 69% (median 31%).

All studies used reliable laboratory phenotypic or genotypic methodology; however, all enrolment was from non-random samples of travellers. All but 2 studies^{10,23} did not exclude co-travellers (groups of travellers) from enrolment. The exposure of interest, antimicrobial use during travel, was ascertained via post-travel questionnaire in all studies. While important potential confounding was reported in all studies, duration of antimicrobial usage was not reported by any. 4 studies^{14,16,23,28} provided no information about losses to follow-up; for the remainder, the median number lost to follow-up was low (7.5%).

The point estimates of the odds ratio (OR) for ESBL-PE acquisition by any or no systemic antimicrobial exposure varied between 0.53 to 8.05 for individual studies (Table 1b).

We included all studies in a random effects meta-analysis (Figure 2). Our primary analysis found that the combined OR for effect of all antimicrobials on ESBL-PE acquisition was 2.37 (95% CI 1.69 to 3.33, $p < 0.01$, $I^2=57\%$, which indicates strong evidence for heterogeneity). A sensitivity analysis excluding doxycycline (all but 2 studies^{21,22}), yielded very similar results (OR 2.48, 95% CI 1.76 to 3.50, $p < 0.01$, $I^2 54\%$). Visual inspection of the funnel plot (Supplementary Figure 1) found no strong evidence of bias across the studies.

Excluding 4 studies with no information on ESBL-PE acquisition by class of antimicrobial received^{13,14,17,22}, we found fluoroquinolone exposure was associated with a combined OR of 4.68 compared to no antibiotics (95% CI, 2.34 to 9.37), tetracyclines were associated a combined OR of 1.68 (95% CI, 1.03 to 2.72) (Table 2). On average, there was no evidence of a combined increased odds of ESBL-PE acquisition by exposure to beta-lactams or macrolides.

DISCUSSION

International travel has been previously identified as an important risk factor for both colonization and infection with ESBL-PE^{15,25,26}, and likely plays an important role in the spread of ESBL-PE within high income countries. However, the risk has not been systematically quantified. Our systematic review and meta-analysis found that systemic antimicrobial use during travel was associated with an odds ratio of 2.37 for acquiring ESBL-PE. Given a baseline risk of ESBL-PE acquisition of 20% or higher among international travellers^{9,15}, and antimicrobial use in an average of 10% of such travellers, the burden of this additional risk is substantial. Our analysis found that fluoroquinolones and tetracyclines (doxycycline) were associated with the increased risk of ESBL-PE acquisition, with a combined OR of 4.68 and 1.68 respectively.

While antimicrobial use is widely recognized to be a major driver of antimicrobial resistance (AMR), quantifying its impact can be challenging. Ecologic data comparing antimicrobial consumption in the community and hospital has been positively correlated to AMR (from hospital bacterial isolates) over time and across countries^{29,30}. Previous systematic reviews^{15,24} have examined the risk of ESBL-PE acquisition in travellers. Unlike the previous studies, we used a meta-analysis to independently estimate the degree to which antimicrobial usage during travel increases the odds of ESBL-PE acquisition across populations of international travellers from high income countries, and to compare specific classes of antimicrobials. Doxycycline has not previously been associated with an increased risk of MRE acquisition²⁴, possibly due to lower numbers exposed and a less strong association compared to fluoroquinolones.

There exist several pathways through which antimicrobial use might cause an increase in acquisition in ESBL-PE during travel. First, selection pressure due to antimicrobial effect on more drug-sensitive strains of Enterobacteriaceae can select for increased resistance within an individual. This selection could be due to antimicrobials taken in response to disease causing Enterobacteriaceae or, substantially more likely, due to the by-stander effect whereby non-invasive bacteria are exposed to antimicrobials³¹. This pathway would predict beta-lactams might be causatively associated with increased risk of ESBL-PE acquisition, which we did not observe; additionally, ESBL-PE are commonly co-resistant to other antibiotic classes such as fluoroquinolones³. An alternate pathway involves the ability of the healthy gut microbiota to prevent expansion of potential pathogens including resistant bacteria such as ESBL-PE, a phenomenon termed 'colonization resistance'³². Disruption of microbial composition due to exposure to antimicrobials decreases colonization resistance³³, reducing the normally protective capabilities of an intestine with fully diverse and intact microbial composition. Proposed mechanisms for the normal flora-mediated resistance to colonization include direct microbial competition for nutrients, production of bacteriocin peptides which inhibit the growth of specific types of bacteria, and more

complex indirect mechanisms involving interaction between bacterial communities to help maintain host immune responses^{33,34}. Fluoroquinolones, in contrast with narrower spectrum antimicrobials, have a relatively large impact on the composition of normal gut microbiota composition due to their broad spectrum of activity and high local concentrations achieved³⁵. In addition to high rates of co-resistance to fluoroquinolones among ESBL-PE^{2,3}, this may in part explain their large effect on ESBL-PE acquisition seen in this study.

An estimated 30% of outpatient antimicrobial prescriptions in the US are considered inappropriate³⁶. It is likely that inappropriate antimicrobial use is even higher in lower income countries, where most consumption is attributable to over-the-counter use. Moreover, travellers from high income countries to tropical settings have historically been counselled to bring antimicrobials, including commonly fluoroquinolones, as preventative or abortive treatment of diarrhea^{37,38}. While more recent guidelines have scaled back the uniform recommendation of traveller antimicrobial use, partly in response to increased recognition of the risk of faecal drug resistant carriage, some still suggest antimicrobials may be used as therapy for cases of moderate or severe traveller's diarrhea (TD)³⁷. Moreover, antibiotics are not commonly prescribed pre-travel as 'stand-by' therapy, a practice which increases their inappropriate use in milder TD cases³⁸. This study supports calls³⁹ to avoid unnecessary antimicrobial consumption during travel, including those taken for prevention or treatment of mild-to-moderate severity TD in healthy travellers. Stand-by antibiotic prescriptions before travel should be limited to those who are at increased risk (such as immunocompromised travellers). When they cannot be avoided altogether, our results suggest using alternative antimicrobials to fluoroquinolones, in view of the higher odds of ESBL-PE acquisition associated with this class.

This meta-analysis included only prospective cohort studies with robust microbiologic assessments of ESBL-PE carriage. The strength of excluding ESBL-PE carriers pre-travel is that any ESBL-PE carrier can be shown to have acquired the resistant bacteria during travel, which can in turn may

generally be attributed to exposures and behaviours taken during the trip. Additionally, while each study recruited travellers from a single country in North America, Europe or Australia, the aggregate analysis supports a broader generalization of healthy tourists from high income regions of the world travelling internationally to tropical, lower income countries.

However, there are limitations to the analysis. First, antimicrobial effect on ESBL-PE acquisition may be mediated partially through an indication of gastroenteritis, which is both an independent risk factor for ESBL-PE acquisition^{9,17} and also a condition for which fluoroquinolones are commonly recommended for international travellers⁴⁰. That is, there is a question about the direction of association between fluoroquinolone use and carriage of ESBL-PE. While an analysis accounting for antimicrobial indication was not possible in this review, several prospective studies have adjusted for diarrhea or gastroenteritis as a confounder^{9-11,13,17}. In these studies, a strong and significant effect of antimicrobial on increased rates of ESBL-PE acquisition during travel persisted. Similarly, many studies documented the presence or absence of proven confounders in individual participants, including age, sex, length of travel, and travel destination which are potentially related to both rates of antimicrobial use, and risk of ESBL-PE acquisition. However, individual studies including the single largest study included in this review⁹ were able to account for these variables in the analysis, and found a persistent effect on ESBL-PE acquisition related to antimicrobial exposure during travel.

Additionally, the designs of the included studies present potential limitations. For instance, losses to follow-up after travel, while not large where provided, were not reported in 4 studies. Selection bias may occur if those lost had different rates of antimicrobial use and rates of ESBL-PE acquisition; however, it seems unlikely the magnitude of this effect would be large enough to substantially change the observed odds ratio. Participants in all studies were non-random volunteers from higher income countries which may limit generalizability of the results to travellers not attending such clinics or originating from lower income countries. Meanwhile, antimicrobial exposure was self-

reported after travel, and data on antimicrobial class was limited to 11 studies, which might underestimate the effect of specific classes during travel on ESBL-PE acquisition.

CONCLUSION

The odds of acquiring ESBL-PE are substantially increased when antimicrobials are consumed during travel. This risk is shared unevenly among antimicrobial classes, with fluoroquinolones posing a substantial risk compared to others. Incorporation of this analysis in decisions and guidelines addressing whether to use antimicrobials during travel will allow for a better realization of the true risks versus benefits. We call for further study into the mechanisms by which antimicrobials, and in particular fluoroquinolones, contribute to AMR, including indirect mechanisms mediated by disruption of the gut microbiome. This may identify fruitful protective factors which may ameliorate the effect of antimicrobials on AMR, which are needed. In the meantime, it seems prudent for travel health practitioners to avoid prescribing stand-by antibiotics for travellers who are not at increased risk, and to emphasize limiting the consumption of antimicrobials for these individuals to the treatment of severe cases of TD only.

Funding

No outside funding supported this work.

Acknowledgements

We would like to offer special thanks to corresponding authors of studies who responded to our requests for further data, which made an invaluable contribution to the precision of our analysis: Drs. Christoph Lübbert, Sunita Paltansing, Thongpan Leangapichart, John Penders, Martin Angelin, Esther Künzli, Karin Ellen Veldkamp, and Marguerite Bruijning.

BOX 1. DETAILED SEARCH STRATEGY USED IN PUBMED.

PubMed

(Enterobacteriaceae[MeSH Major Topic] OR (Enterobacteriaceae*[tiab] OR Gram negative bacteri*[tiab] OR E. coli [tiab] OR Escherischia*[tiab] OR Klebsiella*[tiab] OR Salmonella[tiab] OR Proteus[tiab] OR Enterobacter[tiab] OR Shigella[tiab] OR Yersinia[tiab] OR gut flora[tiab])) AND (Drug Resistance, Bacterial[MeSH Major Topic] OR (Drug resistan*[tiab] OR Extended-spectrum beta-lactamase[tiab] OR ESBL[tiab] OR Amp C[tiab] OR Carbapenemase*[tiab] OR CPE[tiab] OR CRE[tiab] OR Cephalosporinase[tiab] OR Penicillinase[tiab] OR beta-lactamase[tiab] OR CTX-M[tiab] OR ((Cephalosporin[tiab] OR Cefepime[tiab] OR ceftriaxone[tiab] OR Cefotaxime[tiab] OR Ceftazidime[tiab] OR antibiotic*[tiab] OR antimicrob*[tiab]) AND (sensitivit*[tiab] OR susceptibil*[tiab] OR resistan*[tiab]))) AND (Travel[MeSH Major Topic] OR (Travel*[tiab] OR International[tiab] OR Trip[tiab] OR Voyage[tiab] OR Air Transport[tiab] OR Post-Travel[tiab] OR Foreign[tiab] OR Touris*[tiab] OR Aviation[tiab] OR Airport[tiab]))

Figure 1: Flowchart depicting studies screened, reviewed and finally included for systematic review.




Table 1a: Characteristics of travel for studies included for systematic review of the risk of acquiring Extended-Spectrum Producing-Enterobacteriaceae after exposure to antimicrobials during travel.

First Study Author	Top 3 Regions Travelled to ² (Percentage of Participants ¹)	Travel Time Median in Days (range)	Percentage of all travellers who:			Top 2 Classes of Antimicrobials Used (% Total)
			Had Diarrhea	Used Any Abtx	Acquired ESBL-PE	
Angelin	South Asia(40), SSA (39), Asia(8)	45 (13-365)	69	20	36	Beta-lactams (30), FQs (20)
Arcilla	Asia (41), SSA (21), South/Central America (18)	20 (15-25‡)	39	7	34	FQs (31), Beta-lactams (20)
Blyth	South/Central America (33), Asia & South Asia (27), Africa (27)	12* (6-105)	12	10	9	Tetracyclines (50), FQs (33%)
Kantele	SSA (45), Asia (25), South Asia (14)	19 (4-133)	67	15	21	FQs (70), Macrolides (14)
Kennedy	Asia (49), Europe (16), South Asia (11)	21 (9-135)	37	27	49	Tetracyclines (54), FQ (7)
Kuenzli	South Asia (100)	18 (5-35)	37	4	69	FQs (43), Beta-lactams (29)
Leanga-pichart	Middle East (100)	28 (24-32)	19	49	37	Beta-lactams (72) Macrolides (29)
Lübbert	SSA (31), Asia (30), South/Central America (27)	21 (3-218)	38	13	30	Tetracyclines (38), FQs (17)
Osthalm-Balkhed	Africa (34), Asia (26) South & Central America (13)	16 (4-119)	42	9	30	NR
Paltansing	Asia (43), South Asia (25), SSA (25)	21 (6-90)	38	6	33	Beta-lactams (32), FQs (21)
Reuland	Asia & South Asia (56), Africa (22), South/Central America (18)	NR	46	5	23	FQs (27), Beta-lactams (18)
Ruppé	Asia & South Asia (34), SSA (34), South/Central America (31)	20 (15-30)	40	10	51	Beta-lactams (42), FQs (22)
Tängdén	Asia (30), SSA (24), Europe (15)	14 (1-26)	30	10	24	Beta-lactams (50), FQs (30)
Vading	Asia (39), South Asia (35), Northern Africa (14)	14* (8-20‡)	29	9	24	FQs (33), Beta-lactams (20)
von Wintersdorff	Africa (27), South Asia (25), Asia (23)	21 (5-240)	37	14	31	NR

¹Who provided both pre-travel and post-travel samples for analysis. ²Unless otherwise indicated, South Asia (including India, Pakistan, Nepal, Sri Lanka, and Bangladesh) included separately from the rest of Asia, due to higher rates of ESBL-PE. *Median is presented. †Interquartile range is presented. Definitions: NR not reported, ESBL-PE Extended-spectrum beta-lactamase-producing Enterobacteriaceae, SSA Sub-Saharan Africa, FQ fluoroquinolone


Table 1b: Rates extended-spectrum beta-lactamase acquisition by antimicrobial exposure during travel, with crude odds ratio and 95% confidence interval, for included studies.

First Study Author	Participants ¹ who acquired ESBL-PE (% all participants)	Number exposed to antibiotics who acquired ESBL-PE ² (% of those exposed)	Number not exposed to antibiotics who acquired ESBL-PE ² (% of those not exposed)	Odds ratio for acquisition of ESBL-PE by antibiotic exposure (95% CI)
Angelin	35 (36)	20 (25)	30 (38)	0.53 (0.18, 1.62)
Arcilla	633 (34)	132 (55)	553 (33)	2.56 (1.79, 3.66)
Blyth	5 (9)	6 (33)	3 (6)	8.50 (1.09, 66.6)
Kantele	90 (21)	66 (42)	62 (17)	3.53 (2.02, 6.18)
Kennedy	50 (49)	28 (68)	31 (42)	2.93 (1.17, 7.33)
Kuenzli	118 (69)	7 (71)	101 (69)	1.14 (0.21, 6.09)
Leangapichart	73 (37)	107 (36)	35 (40)	0.83 (0.47, 1.49)
Lübbert	58 (30)	24 (38)	15 (9)	6.08 (2.28, 16.23)
Ostholm-Balkhed	68 (30)	20 (25)	15 (7)	4.24 (1.36, 13.28)
Paltansing	113 (33)	19 (47)	104 (33)	1.86 (0.73, 4.72)
Reuland	95 (23)	22 (50)	87 (22)	3.46 (1.45, 8.25)
Ruppé	292 (51)	59 (73)	249 (48)	2.87 (1.58, 5.23)
Tängdén	24 (24)	10 (30)	21 (23)	1.41 (0.33, 5.93)
Vading	56 (24)	15 (67)	46 (29)	4.91 (1.59, 15.16)
Von Wintersdorff	38 (31)	15 (40)	32 (30)	1.56 (0.51, 4.75)

¹Who provided both pre-travel and post-travel samples for analysis.

²Excluding participants with missing antibiotic exposure information.

Figure 2: Forest plot of random effects model, for odds ratio (OR) of extended-spectrum beta-lactamase-producing Enterobacteriaceae (ESBL-PE) among participants exposed to, compared to those not exposed to, antimicrobials during travel



Definitions: OR odds ratio, CI confidence intervals.

Table 2: Summary effect measures, with 95% confidence intervals and measures of heterogeneity, for odds of acquiring extended-spectrum beta-lactamase-producing Enterobacteriaceae (ESBL-PE) during travel, with exposure to specific classes of antimicrobials during travel across all included studies¹

Antimicrobial class	Odds ratio, ESBL-PE acquisition (95% Confidence Interval) ²	P value for chi-squared test of heterogeneity from RE model (Cochran's Q)	I ² statistic (variation in the OR attributable to heterogeneity)
Beta lactam ³	1.57 (0.86, 2.87)	0.097	37.8%
Fluoroquinolone ³	4.68 (2.34, 9.37)	0.149	31.3%
Macrolide ⁴	0.64 (0.17, 2.43)	0.030	57.0%
Tetracycline (doxycycline) ³	1.68 (1.03, 2.72)	0.890	0.0%
Any antimicrobials ⁵	2.37 (1.69, 3.33)	0.004	56.8%

¹From random effects model

²Baseline category is to participants with no antimicrobial exposures during travel

³In all studies, doxycycline was the only tetracycline reported, most frequently for malaria prophylaxis. 4 studies were excluded from this analysis.

⁴8 studies were excluded from this analysis: 4 studies provided no data on rates of ESBL-PE acquisition by macrolide exposure, 4 studies reported no macrolide exposure among participants.

⁵Tetracyclines included here as antimicrobial exposure, in all but 2 studies for which doxycycline was excluded from overall antimicrobial exposure category (considered an antimalarial).

References

1. WHO. *ANTIMICROBIAL RESISTANCE Global Report on Surveillance 2014*. Geneva; 2014. <http://www.who.int/drugresistance/documents/surveillancereport/en/>.
2. Pitout JDD, Laupland KB. Extended-spectrum β -lactamase-producing Enterobacteriaceae: an emerging public-health concern. *Lancet*. 2008;8(3):159-166. doi:10.1016/S1473-3099(08)70041-0
3. Falagas ME, Karageorgopoulos DE. Extended-spectrum β -lactamase-producing organisms. *J Hosp Infect*. 2009;73(4):345-354. doi:10.1016/j.jhin.2009.02.021
4. Woerther PL, Burdet C, Chachaty E, Andremont A. Trends in human fecal carriage of extended-spectrum beta-lactamases in the community: Toward the globalization of CTX-M. *Clin Microbiol Rev*. 2013;26(4):744-758. doi:10.1128/CMR.00023-13
5. Jacoby GA, Silvia Munoz-Price L. The New β -Lactamases. *N Engl J Med*. 2005;352(4):380-391. doi:10.1056/NEJMr041359
6. Denis B, Lafaurie M, Donay J-L, et al. Prevalence, risk factors and impact on clinical outcome of ESBL producing *Escherichia coli* bacteraemia: a 5 year study. *Int J Infect Dis*. 2015; epub(2015):1-6. doi:10.1016/j.ijid.2015.07.010
7. Cornejo-Juárez P, Suárez-Cuenca JA, Volkow-Fernández P, et al. Fecal ESBL *Escherichia coli* carriage as a risk factor for bacteremia in patients with hematological malignancies. *Support Care Cancer*. 2015; Epub ahead of print. doi:10.1007/s00520-015-2772-z
8. Angelin M, Forsell J, Granlund M, Evengard B, Palmgren H, Johansson A. Risk factors for colonization with extended-spectrum beta-lactamase producing Enterobacteriaceae in healthcare students on clinical assignment abroad: A prospective study. *Travel Med Infect Dis*. 2015;13(3):223-229. doi:10.1016/j.tmaid.2015.04.007
9. Arcilla MS, van Hattem JM, Haverkate MR, et al. Import and spread of extended-spectrum β -lactamase-producing Enterobacteriaceae by international travellers (COMBAT study): a prospective, multicentre cohort study. *Lancet Infect Dis*. 2017;17(1):78-85. doi:10.1016/S1473-3099(16)30319-X
10. Reuland EA, Sonder GJB, Stolte I, et al. Travel to Asia and traveller's diarrhoea with antibiotic treatment are independent risk factors for acquiring ciprofloxacin-resistant and extended spectrum β -lactamase-producing Enterobacteriaceae—a prospective cohort study. *Clin Microbiol Infect*. 2016;22:731.e1-731.e7. doi:10.1016/j.cmi.2016.05.003
11. Ruppé E, Armand-Lefèvre L, Estellat C, et al. High rate of acquisition but short duration of carriage of Multidrug-resistant Enterobacteriaceae after travel to the tropics. *Clin Infect Dis*. 2015;61(4):593-600. doi:10.1093/cid/civ333
12. Tängdén T, Cars O, Melhus Å, Löwdin E. Foreign travel is a major risk factor for colonization with *Escherichia coli* producing CTX-M-type extended-spectrum β -lactamases: A prospective study with Swedish volunteers. *Antimicrob Agents Chemother*. 2010;54(9):3564-3568. doi:10.1128/AAC.00220-10
13. Vading M, Kabir MH, Kalin M, et al. Frequent acquisition of low-virulence strains of ESBL-producing *Escherichia coli* in travellers. *J Antimicrob Chemother*. 2016;71:3548-3555. doi:10.1093/jac/dkw335
14. von Wintersdorff CJH, Penders J, Stobberingh EE, et al. High rates of antimicrobial drug resistance gene acquisition after international travel, The Netherlands. *Emerg Infect Dis*. 2014;20(4):649-657. doi:10.3201/eid2004.131718
15. Hassing RJ, Alsmas J, Arcilla MS, Genderen PJ Van, Stricker BH, Verbon A. International travel and acquisition of multidrug-resistant Enterobacteriaceae: a systematic review. *Euro Surveill*. 2015;20(47):1-14. doi:10.2807/1560-7917.ES2015.20.47.30074
16. Blyth DM, Mende K, Maranich AM, et al. Antimicrobial resistance acquisition after international travel in U.S.

- travelers. *Trop Dis Travel Med Vaccines*. 2016;2(4):1-8. doi:10.1186/s40794-016-0020-2
17. Kantele A, Lääveri T, Mero S, et al. Antimicrobials Increase Travelers' Risk of Colonization by Extended-Spectrum Betalactamase-Producing Enterobacteriaceae. *Clin Infect Dis*. 2015;60(6):837-846. doi:10.1093/cid/ciu957
 18. Kennedy K, Collignon P. Colonisation with Escherichia coli resistant to "critically important" antibiotics: A high risk for international travellers. *Eur J Clin Microbiol Infect Dis*. 2010;29(12):1501-1506. doi:10.1007/s10096-010-1031-y
 19. Künzli E, Jaeger VK, Frei R, et al. High colonization rates of extended-spectrum β -lactamase (ESBL)-producing Escherichia coli in Swiss Travellers to South Asia—a prospective observational multicentre cohort study looking at epidemiology, microbiology and risk factors. *BMC Infect Dis*. 2014;14:528-537. doi:doi: 10.1186/1471-2334-14-528.
 20. Leangapichart T, Tissot-Dupont H, Raoult D, Memish ZA, Rolain JM, Gautret P. Risk factors for acquisition of CTX-M genes in pilgrims during Hajj 2013 and 2014. *J Antimicrob Chemother*. 2017;72:2627-2635. doi:10.1093/jac/dkx155
 21. Lübbert C, Straube L, Stein C, et al. Colonization with extended-spectrum β -lactamase-producing and carbapenemase-producing Enterobacteriaceae in international travelers returning to Germany. *Int J Med Microbiol*. 2015;305(1):148-156. doi:10.1016/j.ijmm.2014.12.001
 22. Ostholm-Balkhed A, Tärnberg M, Nilsson M, Nilsson L, Hanberger H, Hällgren A. Travel-associated faecal colonization with ESBL-producing Enterobacteriaceae: incidence and risk factors. *J Antimicrob Chemother*. 2013;68(9):2144-2153. doi:doi: 10.1093/jac/dkt167
 23. Paltansing S, Vlot JA, Kraakman MEM, et al. Extended-spectrum β -lactamase-producing enterobacteriaceae among travelers from the Netherlands. *Emerg Infect Dis*. 2013;19(8):1206-1213. doi:10.3201/eid1908.130257
 24. Ruppé E, Andremont A, Armand-Lefèvre L. Digestive tract colonization by multidrug-resistant Enterobacteriaceae in travellers: An update. *Travel Med Infect Dis*. 2018;21:28-35. doi:10.1016/j.tmaid.2017.11.007
 25. Laupland KB, Church DL, Vidakovich J, Mucenski M, Pitout JDD. Community-onset extended-spectrum β -lactamase (ESBL) producing Escherichia coli: importance of international travel. *J Infect*. 2008;57:441-448. doi:10.1016/j.jinf.2008.09.034
 26. Epelboin L, Robert J, Tsyryna-Kouyoumdjian E, Laouira S, Meyssonier V, Caumes E. High rate of multidrug-resistant gram-negative bacilli carriage and infection in hospitalized returning travelers: a cross-sectional cohort study. *J Travel Med*. 2015;22(5):292-299. doi:10.1111/jtm.12211
 27. Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *PLoS Med*. 2007;4(10):296-1623. doi:10.1371/journal.pmed
 28. Vading M, Kabir MH, Kalin M, et al. Frequent acquisition of low-virulence strains of ESBL-producing Escherichia coli in travellers. *J Antimicrob Chemother*. 2016;71:3548-3555. doi:10.1093/jac/dkw335
 29. Shrestha P, Cooper BS, Coast J, et al. Enumerating the economic cost of antimicrobial resistance per antibiotic consumed to inform the evaluation of interventions affecting their use. *Antimicrob Resist Infect Control*. 2018;7(98):1-9. doi:10.1186/s13756-018-0384-3
 30. Goossens H, Ferech M, Stichele R Vander, Elseviers M. Outpatient antibiotic use in Europe and association with resistance: a cross-national database study. *Lancet*. 2005;365(9459):579-587. doi:10.1016/S0140-6736(05)17907-0
 31. Tedijanto C, Olesen SW, Grad YH, Lipsitch M. Estimating the proportion of bystander selection for antibiotic resistance among potentially pathogenic bacterial flora. *Proc Natl Acad Sci U S A*. 2018;115(51):E11988-E11995.

doi:10.1073/pnas.1810840115

32. Sorbara MT, Pamer EG. Interbacterial mechanisms of colonization resistance and the strategies pathogens use to overcome them. *Mucosal Immunol.* 2018;(March):1-9. doi:10.1038/s41385-018-0053-0
33. Kim S, Covington A, Pamer EG. The intestinal microbiota: Antibiotics, colonization resistance, and enteric pathogens. *Immunol Rev.* 2017;279(1):90-105. doi:10.1111/imr.12563
34. Ubeda C, Djukovic A, Isaac S. Roles of the intestinal microbiota in pathogen protection. *Clin Transl Immunol.* 2017;6(2):e128-10. doi:10.1038/cti.2017.2
35. Stewardson AJ, Gaïa N, François P, et al. Collateral damage from oral ciprofloxacin versus nitrofurantoin in outpatients with urinary tract infections: A culture-free analysis of gut microbiota. *Clin Microbiol Infect.* 2015;21(4):344.e1-344.e11. doi:10.1016/j.cmi.2014.11.016
36. Fleming-Dutra KE, Hersh AL, Shapiro DJ, et al. Prevalence of inappropriate antibiotic prescriptions among us ambulatory care visits, 2010-2011. *JAMA.* 2016;315(17):1864-1873. doi:10.1001/jama.2016.4151
37. Riddle MS, Connor BA, Beeching NJ, et al. Guidelines for the prevention and treatment of travelers' diarrhea: a graded expert panel report. *J Travel Med.* 2017;24(Suppl 1):S63-S80. doi:10.1093/jtm/tax026
38. Vilkin K, Lääveri T, Pakkanen SH, Kantele A. Stand-by antibiotics encourage unwarranted use of antibiotics for travelers' diarrhea: A prospective study. *Travel Med Infect Dis.* 2019 Jan - Feb;27:64-71. doi:10.1016/j.tmaid.2018.06.007
39. Hill DR, Ericsson CD, Pearson RD, et al. The practice of travel medicine: guidelines by the Infectious Diseases Society of America. *Clin Infect Dis.* 2006;43:1499-1539. doi:10.1086/508782
40. McDonald LC. Effects of short- and long-course antibiotics on the lower intestinal microbiome as they relate to traveller's diarrhea. *J Travel Med.* 2017;24(1):S35-S38. doi:10.1093/jtm/taw084