

U. Albert¹, L. Pellegrini¹, G. Maina²,
A.-R. Atti¹, D. De Ronchi¹, Z. Rhimer³

¹ Department of Medicine, Surgery, and Health Sciences, University of Trieste, Italy; ² Rita Levi Montalcini Department of Neuroscience, University of Torino, Italy; ³ Department of Clinical and Theoretical Mental Health, Kútvölgyi Clinical Center, Semmelweis University, Budapest, Hungary

Suicide in obsessive-compulsive related disorders: prevalence rates and psychopathological risk factors

Summary

Objectives

To estimate prevalence rates of suicide attempts and suicidal ideation in individuals with a principal diagnosis of obsessive-compulsive related disorders (OCRDs); 2. to identify predictors of suicide risk among subjects with OCRDs (where available).

Methods

The systematic review was conducted by searching PubMed from the date of the first available article to December 31, 2018. The search terms [suicide] OR [suicidality] OR [suicide attempts] OR [suicidal ideation] OR [suicidal thoughts] were combined with the following: [BDD] OR [body dysmorphic disorder]; [HD] OR [hoarding disorder]; [trichotillomania] OR [hair pulling disorder]; [excoriation disorder] OR [skin picking disorder].

Results

In BDD, data concerning lifetime suicide attempts are consistent across studies: mean rate is 21.5% (range 9-30.3%). Mean rate of current suicidal ideation is 37.4% (range 26.5-49.7%) and mean rate of lifetime suicidal ideation is 74.5% (range 53.5-85%). BDD-specific factors such as early onset, severity, poor insight and muscle dysmorphia and comorbid disorders increase the risk of suicide attempts or suicidal ideation. Only 2 studies recruited individuals with DSM-5 HD: suicidality appears to be low, with rates of current suicidal ideation comprised between 5% and 10%, although 19% of individuals attempted suicide during their lifetime. Concerning the grooming disorders, lifetime rates of suicide attempts are low as compared to rates in other OCRDs; approximately 40% of individuals, however, reported lifetime suicidal ideation.

Conclusions

OCRDs taken together may be at risk for suicide attempts and suicidal ideation independently from comorbid disorders (and specifically independently from comorbid OCD); BDD remains the disorder more strongly associated with an increased risk for suicide, followed by HD and then the grooming disorders.

Key words

Suicide attempts • Suicidal ideation • BDD • HD • Trichotillomania • Skin Picking Disorder

© Copyright by Pacini Editore Srl



OPEN ACCESS

Received: March 13, 2019

Accepted: June 20, 2019

Correspondence

Umberto Albert
Department of Medicine, Surgery,
and Health Sciences, University of Trieste,
via G. de Pastovich 3, 34128 Trieste, Italy
• Tel. +39 040 3997327 • E-mail: ualbert@units.it

Introduction

Recent systematic reviews and meta-analyses confirmed that Obsessive-Compulsive Disorder (OCD), historically considered to be associated with a relatively low risk of suicide, is actually in itself associated with considerable risk for lifetime suicide attempts and suicidal ideation^{1,2}. Data from recent large epidemiological studies performed on National Registers, providing data on the longitudinal association between OCD and death by suicide and lifetime suicide attempts over a follow-up of several years^{3,4}, confirmed that individuals with OCD are at greater risk for committing suicide as compared to the general population.

Less is known about suicidality and other DSM-5 Obsessive-Compulsive Related Disorders (OCDs); in the *new* chapter, *new* disorders such as Hoarding Disorder (HD) and Skin Picking Disorder (SPD) and disorders once classified elsewhere (Body Dysmorphic Disorder – BDD – previously in the chapter of Somatoform Disorder, and Trichotillomania – TTM – previously classed among the Impulse Control Disorders) have been grouped together with the *nosological organizer* OCD⁵. All disorders included in this chapter share similarities with OCD, although some appear to have a stronger cognitive component – and thus are closer to OCD – while others mainly consist of body-focused repetitive behaviors.

While several issues concerning phenomenological characteristics of these disorders have been studied, less attention has been devoted to suicidality. A recent systematic review and meta-analysis examined the strength and patterns of the association between suicidality and BDD, concluding that BDD is actually associated with increased odds for both suicide attempts and suicidal ideation⁶. No similar studies are available for the other disorders of the OCDs chapter.

Given the prevalence of these disorders in the general population and the impact in terms of psychosocial impairment associated with these disorders, the investigation of suicidality and the identification of potential socio-demographic and clinical factors that could increase the risk for suicide is, to our opinion, of particular clinical relevance.

The aims of the present systematic review were: 1) to estimate prevalence rates of suicide attempts and suicidal ideation in individuals with a principal diagnosis of obsessive-compulsive related disorders; 2) to identify predictors of suicide risk among subjects with OCDs (where available).

Methods

Search strategy

The systematic review was conducted using the PRISMA guidelines by searching PubMed from the date of the first available article to December 31, 2018. The search terms [suicide] OR [suicidality] OR [suicide attempts] OR [suicidal ideation] OR [suicidal thoughts] were combined with the following: [BDD] OR [body dysmorphic disorder]; [HD] OR [hoarding disorder]; [trichotillomania] OR [hair pulling disorder]; [excoriation disorder] OR [skin picking disorder].

Article selection and review strategy

Articles were identified and assessed for eligibility by two independent reviewers (UA and LP), who independently decided which identified articles to include according to clinical importance and eligibility criteria. In case of disagreement, a third author (GM) was consulted to mediate consensual decisions. Duplicate studies were excluded. Cross-references from the articles identified were also

examined. Unpublished studies, conference abstracts or poster presentations were not included. The database search was restricted to English language papers.

Eligibility criteria

The inclusion criteria for the studies were the following: 1) studies with appropriate definition of the obsessive-compulsive related disorder (diagnosis made through specific structured interviews and/or established international criteria); 2) adolescents and/or adults; 3) cross-sectional or prospective designs; 4) performed in clinical samples or in the general population (epidemiological studies); 5) employed a quantitative measure of suicidality in order to derive prevalence rates of current/lifetime suicide attempts, suicidal ideation and/or family history of suicide attempts/completed suicide; and/or 6) reported an outcome measure of the association between suicidality and OCD (e.g. odds ratios) or examined factors associated with suicidality.

Results

Search results

The flowchart of studies selected and included in the systematic review for BDD is provided in Figure 1. In total, 24 studies were included in the qualitative synthesis (providing data on prevalence rates of suicide attempts, suicidal ideation). Additional 17 studies were retrieved from PubMed search and manual search providing data on suicidality and HD (N = 8), TTM (N = 4) and SPD (N = 5).

Body dysmorphic disorder

Table I reports prevalence rates of suicide attempts and suicidal ideation in individuals with BDD as from clinical studies; 17 studies provided information, although some of the studies included partially overlapping samples. Additional studies included exactly the same sample of Phillips et al. 2005⁷ and thus were excluded from the table (see Appendix 1).

Concerning lifetime suicide attempts, data are consistent across studies: mean rate is 21.5% (range 9-30.3%; median value: 22.4%). Higher rates are reported in individuals with comorbid OCD (OCD+BDD: 40%)⁸, in Veterans with comorbid MDD (92% of the sample) (58.3%)⁹ and among inpatients (75% had comorbid Substance Use Disorder) (93.8%)¹⁰. Mean rate of current suicidal ideation is 37.4% (range 26.5-49.7%; median value: 36.9%) and mean rate of lifetime suicidal ideation is 74.5% (range 53.5-85%; median value: 77.9%).

Studies performed in the general population confirmed that BDD is associated with a significantly higher risk of suicide attempts and suicidal ideation as compared to individuals without that diagnosis (Tab. II), although reported prevalence rates somewhat lower than those in clinical settings. When suicide risk was estimated in the general population using specific instruments, such as the




FIGURE 1. Flow chart showing the selection of BDD studies.

SBQ-R (Suicide Behaviors Questionnaire Revised) or the ACSS (Acquired Capability Suicide Scale), it was found elevated¹¹⁻¹³.

Twenty-six studies provided information on predictors of suicidality. Table III presents results of the analysis of socio-demographic and clinical factors potentially associated with increased risk for suicide.

Hoarding disorder

We could find only two studies that recruited individuals with DSM-5 criteria HD^{14 15}. All the other studies shown in Table IV investigated suicidality in individuals with a diagnosis of OCD and hoarding symptoms (in one study only subjects with hoarding as the primary problem were included, thus suggesting that these individuals could have been diagnosed with HD provided that DSM-5 criteria were available at that moment). Interpretation of data concerning suicidality in HD is then compromised by these limits.

Suicidality in individuals with HD appears to be low, with rates of current suicidal ideation comprised between 5% and 10%, although 19% of individuals attempted suicide during their lifetime^{14 15}. Suicidality appears to be higher in samples composed of individuals with OCD and hoarding symptoms (Table IV).

Only one recent study¹⁴ specifically examined factors associated with increased suicide risk in individuals diagnosed with HD according to DSM-5 criteria: severity of HD (as measured by the SI-R total score), hoarding-related impairment (as measured by the ADL-H total score), higher number of psychiatric comorbidities, and specifically MDD and BD, all predicted suicidality¹⁴.

Trichotillomania (hair pulling disorder) and skin picking disorder

Only four studies provided information concerning suicidality in individuals with TTM. These studies, moreover, suffered from the inclusion of few subjects whose diagnosis was made without structured interviews. In one study¹⁶, data concerning suicidality in individuals with OCD and grooming disorders were provided without specifying whether it was TTM or SPD. Table V presents results of our review: lifetime rates of suicide attempts are low (3.7-12%) as compared to rates in other OCRDs; approximately 40% of individuals, however, reported lifetime suicidal ideation. Concerning SPD, five studies provided prevalence rates of suicide attempts and suicidal ideation; however, only two studies included samples made of non-comorbid SPD (Tab. VI). The only study that investigated lifetime suicide attempts rate in individuals with SPD without comorbid disorders found a low prevalence (5.7%)¹⁷. Approximately 40% of individuals reported lifetime suicidal ideation.

Discussion

Obsessive-compulsive disorder (OCD) has long been considered a disorder which did not carry a notable risk for suicide. Recent meta-analyses and systematized reviews, however, challenged this opinion and found that OCD may actually be considered at risk for suicidal ideation, suicide attempts and committed suicide. A recent systematic review from our research group² found a mean prevalence of current suicidal ideation in OCD of 25.9% (median: 15.6%); lifetime suicidal ideation of 44.1% (median: 36.4%) and lifetime suicide attempts of 14.2% (median

TABLE I. *Suicidality in BDD: studies in clinical samples.*

Author	Country	Design	BDD diagnosis	Screening for suicidality	Mode of suicidality	Sample N		Suicidality (%)	
						Mean age % males	Suicide attempts	Suicidal ideation	
Veale et al., 1996 ²⁰	UK	Cross-sectional	BDDE BDD-YBOCS	n/r	Lifetime suicide attempts	50 32.6 (range 19-58) 24	24 (1 completed suicide)	-	
Perugi et al., 1997 ²¹	Italy	Cross-sectional	DID DSS HSCL-90	BDSS for suicidal ideation	Current suicidal ideation	58 25 (SD 5.9) 59	-	45	
Zimmerman & Mattia, 1998 ²²	USA	Cross-sectional	SCID-I	n/r	Lifetime suicide attempts	16 31.6 (SD 10.8) 25	18.8	-	
Albertini & Phillips, 1999 ²³	USA	Cross-sectional	SCID-I BDD Form BDD-YBOCS	BDD data form	Lifetime suicidal ideation & lifetime suicide attempts	33 (adolescents) 14.9 (SD 2.2) 9	21	67	
Altamura et al., 2001 ²⁴	Italy	Cross-sectional	SCID-I BDD-YBOCS	SCID-I	Current suicidal ideation	30 28.5 (SD 2.3) 13	-	49.7	
Grant et al., 2002 ²⁵	USA	Cross-sectional	BDD-Q Self-report screening measure for BDD SCID-I	SCID-I	Lifetime suicide attempts	AN+BDD: 16 27.4 (SD 9.7) 0	62.5	-	
Frare et al., 2004 ²⁶	Italy	Cross-sectional	SCID-I	ADPI	Current suicidal ideation	BDD: 34 24.7 (SD 5.6) 55.9 BDD + OCD: 24 16.4 (SD 2.3) 62.5	- -	26.5 29.1	
Phillips et al., 2005 ⁷	USA	Cross-sectional	SCID-I BDD-YBOCS	BDD Form	Lifetime suicidal ideation & lifetime suicide attempts	200 32.6 (SD 12.1) 31.5	27.5 (2 completed suicide)	78	
Fontanelle et al., 2006 ²⁷	Brazil	Cross-sectional	SCID-I	Specific questionnaire	Current suicidal ideation & lifetime suicide attempts	20 29.2 (SD 8.6) 45	15	35	
Phillips et al., 2007 ^{* 8}	USA	Cross-sectional	SCID-I BDD-YBOCS	SCID-I HAM-D	Lifetime suicidal ideation & lifetime suicide attempts	BDD: 45 36.5 (SD 12.7) 62.3 BDD + OCD: 40 36.5 (SD 11.7) 55	13.3 40	77.8 85	
Conroy et al., 2008 ¹⁰	USA	Cross-sectional	BDD-Q SCID-I/P	A brief version of the BDD data form	Lifetime suicidal ideation & lifetime suicide attempts	16 31.9 (SD 11) 31.2	93.8	100	
Phillips & Kelly, 2009 ²⁸	USA	Prospective	BDD-YBOCS	HAM-D	Current suicidal ideation	67 32.1 (SD: 10.5) 32.3%	-	38.8	
Costa et al., 2012 ²⁹	Brazil	Cross-sectional	SCID-I	Questionnaire assessing suicidality	Lifetime suicidal ideation & lifetime suicide attempts	OCD + BDD: 109 31.3 (SD 10.2) 44	23.8	53.5 (suicidal plans 36.6)	
Bjornsson et al., 2013 ^{* 30}	Iceland	Cross-sectional	BDD-YBOCS BDD-PSR BDDE	Suicidality items from the HAM-D	Lifetime suicidal ideation & lifetime suicide attempts	Sample 1: 184 16.7 (SD 7.3) 33.15 Sample 2: 244 16.7 (7.2) 46.7	30.3-14.5 [*] 29.9-16.7 [*]	81-69.4 [#] 83.9-79.7 [*]	

(continues)

TABLE I (follows). *Suicidality in BDD: studies in clinical samples.*

Author	Country	Design	BDD diagnosis	Screening for suicidality	Mode of suicidality	Sample N		Suicidality (%)	
						Mean age % males	Suicide attempts	Suicidal ideation	
Hart et al., 2013* ³¹	USA	Cross-sectional	SCID-I BDD-YBOCS BDD form	SCID-I	Lifetime suicidal ideation & lifetime suicide attempts	Sample 1: 160 28.80 (SD 11.04) 41	29.4	78.0	
						Sample 2: 115 32.93 (SD 11.83) 30	28.2	74.1	
De Brito et al., 2015 ³²	Brazil	Cross-sectional	BDDE Clinical assessment	n/r	Lifetime suicidal ideation & lifetime suicide attempts	300 n/r 14.6	9	18.4	
Kelly et al., 2015 ⁹	USA	Cross-sectional	BDD-Q SCID-P	n/r	Lifetime suicidal ideation & lifetime suicide attempts	12 49.6 (SD 13.7) 83.3	58.3	66.7	

*n/r: not reported; *: partially overlap with Phillips 2005. The additional studies which used exactly the same cohort of participants are included in Appendix (not shown in the table); #: age of onset before 18 – age of onset after 18; BDDE: Body Dysmorphic Disorder Examination; BDD-YBOCS: Yale-Brown Obsessive-Compulsive Scale adapted for BDD; DID: Diagnostic Interview for Body Dysmorphobia; DSS: Body Dysmorphic Symptom Scale; HSCL 90: Hopkins Symptom Checklist 90; SCID-I: Structured Clinical Interview for DSM-IV Axis-I Disorders; BDD-Q: The Body Dysmorphic Disorder Questionnaire; BDD-PSR: Psychiatric Status Rating Scale for Body Dysmorphic Disorder; SCID/P: Structured Clinical Interview for DSM-IV Patient Edition; ADPI: Adult Demographic and Personal Inventory; HAM-D: Hamilton Rating Scale for Depression.*

TABLE II. *Suicidality in BDD: epidemiological studies.*

Authors	Country	Design	BDD diagnosis	Screening for suicidality	Mode of suicidality	N BDD-sample§		Suicidality	
						Mean age % males	Suicide attempts	Suicidal ideation	
Rief et al., 2006 ³³	Germany	Cross-sectional	DSM criteria Clinical assessment SOMS-7	n/r	Current suicidal ideation & lifetime suicide attempts	42 44.3 (SD 17.2) 40	7.2 (1.0 in no-BDD, p < .001)	19.1 (3.4 in no-BDD, p < .001)	
Buhlmann et al., 2010 ³⁴	Germany	Cross-sectional	DSM criteria	Specific questionnaire	Lifetime suicidal ideation & lifetime suicide attempts	45 48.9 (SD: 17.1) 37.7	22.2 (3.5 in no-BDD, p = 0.02)	31.0 (2.1 in no-BDD, p = 0.02)	
Schieber et al., 2015 ³⁵	Germany	Cross-sectional	DSM-IV and DSM-5 criteria	PHQ-9	Current suicidal ideation	62 42.1 (SD 13.6) 21	-	31.1 (7.3 in no-BDD)	
Moolman et al., 2017 ³⁶	Germany	Cross-sectional	DSM-5 criteria BDSI	Items 10, 16, 17, and 18 from the FKS	Current suicidal ideation	11* n/r 18.2	-	36.4 (8.8 in no-BDD, p < .001)	
Shaw et al., 2016 ¹²	USA	Cross-sectional	BDD-SS	BDD-SS ACSS	Current suicide risk	235 32.1 (SD 9.9) 43	INQ-burd: 2.71 (SD 1.50) INQ-belong: 3.37 (SD 1.62) ACSS: 8.01 (SD 4.67)		
Weingarden et al., 2016 ¹³	USA	Cross-sectional	BDD-Q BDD-YBOCS	SBQ-R	Current suicide risk	114 30.2 (SD 10.9) 8%	SBQ-R: 8.18 (SD 3.70)		
Weingarden et al., 2017 ¹¹	USA	Cross-sectional	BDD-Q BDD-YBOCS	SBQR BDD-SS	Current suicide risk	184 29.7 (SD 10.1) 7.6	SBQ-R: 9.69 (SD 4.25)		

*§: N of BDD patients from the general population; *: adolescent population; SOMS-7: Somatoform Disorders Screening Symptoms-7; BDSI: Body Dysmorphic Symptoms Inventory; BDD-SS: Body Dysmorphic Disorder Symptoms Scale; BDD-Q: The Body Dysmorphic Disorder Questionnaire; BDD-YBOCS: Yale-Brown Obsessive-Compulsive Scale adapted for BDD; PHQ-9: Patient Health Questionnaire-9; FKL: Fragebogen körperdysmorpher Symptome; ACSS: Acquired Capability Suicide Scale; SBQ-R: Suicide Behaviors Questionnaire Revised; INQ: Interpersonal Needs Questionnaire - INQ-burd: INQ perceived burdensomeness subscale; INQ-belong: INQ thwarted belongingness subscale.*

TABLE III. Studies with data on associated factors or predictors of suicidality in body dysmorphic disorder.

Predictors		Current/lifetime suicidal ideation	Lifetime suicide attempts	Deaths by suicide
Socio-demographic or personal factors	Lifetime academic/occupational/role impairment	Didie et al., 2008 ⁴⁹ Witte et al., 2012 ⁵⁰	Didie et al., 2008 ⁴⁹ Witte et al., 2012 ⁵⁰	Philipps et al. 2005 ⁷
	Lifetime/current social and functional impairment	Witte et al., 2012 (lifetime) ⁵⁰ Philipps et al., 2005 (lifetime and current) ⁷	Philipps et al., 2005 (lifetime and current) ⁷	-
	Age 20 or younger	Philipps et al., 2006 ⁵¹	-	-
	Being single or divorced	Philipps et al., 2005 ⁷	-	-
Disorder-specific (BDD-related) variables	Early onset (< 18 yrs)	Bjornsson et al., 2013 ³⁰	Bjornsson et al., 2013 ³⁰	-
	Appearance related symmetry concerns	Hart & Philipps, 2013 ³¹	-	-
	BDD-SS Severity	Philipps et al., 2005 ⁷ Philipps & Menard, 2006 ⁵² Shaw et al., 2016 ¹²	-	Philipps et al. 2005 ⁷
	Delusional form of BDD	-	Philipps et al., 2005 ⁷ Philipps & Menard, 2006 ⁵²	-
	Muscle dysmorphia	-	Pope et al., 2005 ⁵³	-
	BDD-related restrictive food intake	-	Witte et al., 2012 ⁵⁰	-
	BDD-related excessive exercise (protective)	-	Witte et al., 2012 ⁵⁰	-
	Type of plastic surgery performed for correction of apparent defects: rhytidectomy ¹	De Brito et al., 2016 ³²	De Brito et al., 2016 ³²	-
Comorbidities	Comorbid OCD	Frare et al., 2004 ²⁶ Philipps et al., 2007 ⁸	Philipps et al., 2007 ⁸	-
	Comorbid social phobia	Coles et al., 2006 ⁵⁴	-	Philipps et al. 2005 ⁷
	Comorbid panic attacks	Philipps et al., 2013 ⁵⁵	-	-
	Lifetime history of PTSD	-	Philipps et al., 2005 ⁷ Witte et al., 2012 ⁵⁰	-
	History of psychiatric hospitalization	-	Philipps et al., 2005 ⁷	-
	Substance/alcohol use disorders	Grant et al., 2005 ⁵⁶ Witte et al., 2012 ⁵⁰	Philipps et al., 2005 ⁷	Philipps et al. 2005 ⁷
	MDD (current and lifetime)	Philipps et al., 2005 ⁷ Philipps et al., 2007 ⁸ Witte et al., 2012 ⁵⁰ Shaw et al., 2016 ¹²	-	Philipps et al. 2005 ⁷
	Lifetime bipolar disorder	Philipps et al., 2005 ⁷	Philipps et al., 2005 ⁷	-
	Comorbid eating disorder	-	Philipps et al., 2005 ⁷	-
	Any personality disorder	Philipps et al., 2005 ⁷	Philipps et al., 2005 ⁷	-
Borderline personality disorder	Philipps et al., 2005 ⁷	Philipps et al., 2005 ⁷	-	
Emotion-cognitive factors	Childhood trauma (emotional, physical and sexual abuse) ²	-	Didie et al., 2006 ⁵⁷	-
	Shame/defectiveness beliefs	Weingarten et al., 2016 ¹³	-	-
	Anxiety (considered as emotion)	Weingarten et al., 2016 ¹³	-	-
	Weight concerns	-	Kittler, 2007 ⁵⁸	-
	High levels of impulsivity	Phillips & Menard, 2006 ⁵⁹	Phillips & Menard, 2006 ⁵⁹	-

1) Compared to other types of surgery: abdominoplasty and rhinoplasty; 2) Measured by Childhood Trauma Questionnaire (CTQ).

10.8%). Specific factors are more strongly associated with suicide in OCD patients; the severity of OCD, the unacceptable thoughts symptom dimension (aggressive, sexual, religious obsessions), comorbid Axis I disorder (bipolar disorder or major depressive disorder but also substance use disorder), the severity of comorbid depressive and anxiety symptoms, a previous history of suicide attempts, and some emotion-cognitive factors, such as alexithymia

and hopelessness, all increase the risk of having suicidal ideation or attempting suicide ¹⁸. Our systematic review clearly showed that OCD is at a greater suicide risk, compared to the general population. Hence, clinicians should actively inquire about suicidal thoughts and attempts when interviewing a patient with OCD, keeping in mind that risk identification remains a crucial factor for establishing preventive strategies. The recognition that specific risk factors

TABLE IV. *Suicidality in HD (or in OCD subjects with hoarding symptoms).*

Author	Country	Design	Hoarding disorder symptoms diagnosis	Screening for suicidality	Mode of suicidality	Sample N	Suicidality (%)	
							Mean age	% males
Balci & Sevincok, 2010 ³⁷	Turkey	Cross-sectional	YBOCS	SSI	Current suicidal ideation	11* n/r n/r	-	36.4
Matsunaga et al., 2010 ³⁸	Japan	Cross-sectional	YBOCS	Self-report questionnaire	Lifetime impulsive behaviors (including suicide attempts)	54* 30.8 (SD 8.9) 44.4	41 (vs 18 non-hoarders, p < 0.01)	-
Alonso et al., 2010 ³⁹	Spain	Prospective	YBOCS	Beck suicide intent scale	Lifetime suicide attempts	62* n/r n/r	6.45 (vs 5.91 in total OCD population)	-
Torres et al., 2011 ⁴⁰	Brazil	Cross-sectional	DYBOCS	Specifically created questionnaire	Lifetime and current suicidal ideation, lifetime suicidal plans & lifetime suicide attempts	297* n/r n/r	13.1	11.5 (current) 39.4 (lifetime) 24.2 lifetime suicidal plans
Chakraborty et al., 2012 ⁴¹	UK	Cross-sectional	SI-R	Clinical interview	Current suicidal ideation & lifetime suicide attempts	20# 31.5 (SD 9.98) 50	40	20
Torres et al., 2012 ⁴²	Brazil	Cross-sectional	DYBOCS	Specific questions	Current and lifetime suicidal ideation, lifetime suicidal plans & lifetime suicide attempts	528 (4 of them only HD)* 35.9 (SD 13.2) 40.2	12.7	12.3 (current) 38.9 (lifetime) 23.7 lifetime suicidal plans
Ayers et al., 2015 ¹⁵	USA	Cross-sectional	DSM-5 criteria UHSS SI-R CIR	n/r	Current suicidal ideation	71 67 (SD 5.8) 31	-	4.69
Archer et al., 2018 ¹⁴	USA	Cross-sectional	SIHD	MINI	Current suicidal ideation & lifetime suicide attempts	313 59 (SD 11.8) 25.9	19	10

*: patients with OCD and hoarding symptoms as measured by the Dimensional YBOCS; #: patients with OCD and hoarding symptoms. In all cases, hoarding was a primary problem, that is, not secondary to other OCD symptoms. YBOCS: Yale Brown Obsessive Compulsive Scale; DYBOCS: Dimensional Yale Brown Obsessive Compulsive Scale; SI-R: Saving Inventory Revised; UHSS: UCLA Hoarding Severity Scale; CIR: Clutter Image Rating Scale; SIHD: The Structured Interview for Hoarding Disorder; SSI: Scale for Suicidal Ideation; MINI: Mini International Neuropsychiatric Interview.

are associated with suicidal ideation and attempts among individuals with OCD could potentially lead to saving lives in the future.

Less research has been devoted to understanding suicidality among individuals with obsessive-compulsive related disorders; this group of disorders may share with OCD the high risk for suicide attempts and suicidal ideation. However, only for BDD systematic reviews and a meta-analysis are available on the topic^{6,19}. This lead us to perform the present systematic review including all papers on suicidality among individuals with OCRDs.

Concerning suicidality among individuals with BDD, our results are consistent with those of a previous systematic review and meta-analysis which, however, included only seventeen studies⁶: a positive and statistically significant association was found between BDD and suicidality (attempts and ideation together, without differentiating between current and lifetime rates): OR = 3.63 (CI 2.62-4.63).

Our systematic review found that approximately 20% of individuals with a primary diagnosis of BDD attempted suicide during their lifetime and 75% had suicidal ideation. Studies performed in the general population confirmed that BDD is in itself at greater risk for suicide as compared to the general population. Suicidality in BDD appears, then, even higher than among patients with OCD. Clinicians, then, should not overlook BDD as being not at risk for suicide and should actively inquire about past suicide attempts and current suicidal ideation in each patient with a diagnosis of BDD, independently from other comorbid disorders eventually present.

However, we found that BDD-specific factors such as early onset, severity, poor insight and muscle dysmorphia and comorbid disorders (mainly MDD, anxiety disorders or OCD) increase the risk of suicide attempts or suicidal ideation, and thus may constitute specific predictors of suicidality to be actively inquired and that clinicians could

TABLE V. *Suicidality in trichotillomania.*

Author	Country	Design	TTM diagnosis	Screening for suicidality	Mode of suicidality	Sample N	Suicidality (%)	
						Mean age % males	Suicide attempts	Suicidal ideation
Streichenwein et al., 1995 ⁴³	USA	Clinical trial	DSM-IV criteria	n/r	Lifetime suicide attempts	16 39 (SD 12) 12.5	6.3	-
Seedat & Stein, 1998 ⁴⁴	South Africa	Cross-sectional	Specific questionnaire DSM-IV criteria YBOCS	Specific questionnaire	Current suicidal ideation & lifetime suicide attempts	27 29.8 (SD 10.7) 37	3.7	44.4
Lejoyeux et al., 2002 ⁴⁵	France	Cross-sectional	MIDI	Specific questionnaire	Lifetime suicide attempts	3* 30 (SD 5) 0	66.6	-
Lovato et al., 2012 ¹⁶	Brazil	Cross-sectional	SCID-I DYBOCS	n/r	Current suicidal ideation & lifetime suicide attempts	121** 31.0 (SD 11.8) 27.3	12.0	39.3

YBOCS: Yale Brown Obsessive Compulsive Scale; MIDI: Minnesota Impulsive Disorders Interview; SCID-I=Structured Clinical Interview for DSM-IV Axis-I Disorders; DYBOCS: Dimensional Yale Brown Obsessive Compulsive Scale; *: patients diagnosed with TTM taken from a sample of depressed patients; **: population with OCD and GD (grooming disorders, either skin-picking disorder or trichotillomania).

consider when planning treatment and visit schedules. Results of our systematic review are in accordance to those of previous reviews on the same topic^{6,19}. It has to be stated, however, that few studies specifically examined potential predictors of suicidality in BDD, and many of those studies are flawed by methodological biases, suggesting caution in the interpretation of these findings. It has to be noted, moreover, that some of the factors found to increase suicidality among subjects with BDD do also increase suicide risk in individuals with OCD². Risk identification and stratification of risk remain essential components of suicide prevention and should guide the clinical approach to subjects with OCD. Whether and how these risk factors for suicide work together, and whether the spe-

cific factors act as moderators or mediators, remains to be fully elucidated.

The evaluation of suicide risk in individuals with HD, Trichotillomania and SPD is hampered by the very low number of studies investigating suicidality specifically in samples of individuals with these DSM-5 disorders. The revision of the classification made by DSM-5 (with the creation of the OCD and related disorders new category and with the new disorders – HD and SPD) surely represented an advance for clinicians and researchers, but unfortunately very few studies at yet investigated whether individuals with these new disorders are at risk for committing suicide. When examining suicide risk and hoarding, moreover, we have still to rely on data gathered from samples of subjects

TABLE VI. *Suicidality in skin picking disorder.*

Author	Country	Design	SPD diagnosis	Screening for suicidality	Mode of suicidality	Sample N	Suicidality (%)	
						Mean age % males	Suicide attempts	Suicidal ideation
Philipps et al., 1995 ⁴⁶	USA	Cross-sectional	n/r	n/r	Lifetime suicide attempts	33* - 42.4	33	-
Grant et al., 2006 ⁴⁷	USA	Cross-sectional	SCID-I YBOCS	n/r	Lifetime suicide attempts	79* 30.5 (SD 11.3) 17.7	25.3	-
Grant et al., 2010 ¹⁷	USA	Cross-sectional	proposed DSM criteria	n/r	Lifetime suicide attempts	53 34.2 (SD 13.11) 13.2	5.7 (vs 13.7 in OCD)	-
Lovato et al., 2012 ¹⁶	Brazil	Cross-sectional	SCID-I DYBOCS	n/r	Current suicidal ideation & lifetime attempts	121** 31.0 (SD 11.8) 27.3	12.0	39.3
Machado et al., 2018 ⁴⁸	Brazil	Cross-sectional Epidemiological	SPSQ	PHQ-9	Current suicidal ideation	259 27.8 (8.4) 17.8	-	41.3

*: individuals with BDD and SPD; **: population with OCD and GD (grooming disorders, either skin-picking disorder or trichotillomania); SCID-I: Structured Clinical Interview for DSM-IV Axis-I Disorders; YBOCS: Yale Brown Obsessive Compulsive Scale; DYBOCS: Dimensional Yale Brown Obsessive Compulsive Scale; OCD: Obsessive Compulsive Disorder; SPSQ: Skin Picking Stanford Questionnaire; PHQ-9: Patient Health Questionnaire-9.

with OCD as the primary diagnosis and prominent hoarding symptoms; it is not clear, then, whether suicidality is associated with HD in itself or whether it is associated with the specific subtype of OCD with hoarding symptoms. It is possible that HD, being usually associated with poor insight and then with a long delay from onset to help-seeking, is not at risk for suicide at the beginning of its history, and subsequently becomes associated with a higher risk when comorbid with OCD or MDD, or when the impairment associated with HD is huge. However, this is only a hypothesis that needs to be confirmed.

Concerning the grooming disorders (TTM and SPD), too few studies are available to draw some conclusions; very preliminary data seem to suggest that suicide attempts are low in TTM and SPD as compared to other OCRDs. The inclusion of these disorders among the chapter of OCD and related disorders will for sure draw attention on

such neglected disorders, and we will expect that more reliable data on suicidality will appear in the next future. In conclusion, our present systematic review showed that like in the case of pure OCD², 3 OCRDs taken together may be at risk for suicide attempts and suicidal ideation independently from comorbid disorders (and specifically independently from comorbid OCD); BDD remains the disorder more strongly associated with an increased risk for suicide, followed by HD and then the grooming disorders. A greater awareness of such suicide risk should prompt clinicians to actively inquire about past suicide attempts and current suicidal ideation whenever a patient with one of the obsessive-compulsive related disorders presents for a visit.

Conflict of interest

The Authors have no conflict of interest to declare.

References

- Angelakis I, Gooding P, Tarrier N, et al. *Suicidality in obsessive compulsive disorder (OCD): a systematic review and meta-analysis*. Clin Psychol Rev 2015;39:1-15.
- Albert U, De Ronchi D, Maina G, et al. *Suicide risk in obsessive-compulsive disorder and exploration of risk factors: a systematic review*. Curr Neuropharmacol 2018 Jun 20.
- Fernández de la Cruz L, Rydell M, Runeson B, et al. *Suicide in obsessive-compulsive disorder: a population-based study of 36 788 Swedish patients*. Mol Psychiatry 2017;22:1626-32.
- Meier SM, Mattheisen M, Mors O, et al. *Mortality among persons with obsessive-compulsive disorder in Denmark*. JAMA Psychiatry 2016;73:268-74.
- Albert U, De Cori D, Barbaro F, et al. *Hoarding disorder: a new obsessive-compulsive related disorder in DSM-5*. Off J Ital Soc Psychopathol 2015;21:354-64.
- Angelakis I, Gooding PA, Panagioti M. *Suicidality in body dysmorphic disorder (BDD): a systematic review with meta-analysis*. Clin Psychol Rev 2016;49:55-66.
- Phillips KA, Coles ME, Menard W, et al. *Suicidal ideation and suicide attempts in body dysmorphic disorder*. J Clin Psychiatry 2005;66:717-25.
- Phillips KA, Pinto A, Menard W, et al. *Obsessive-compulsive disorder versus body dysmorphic disorder: a comparison study of two possibly related disorders*. Depress Anxiety 2007;24:399-409.
- Kelly MM, Zhang J, Phillips KA. *The prevalence of body dysmorphic disorder and its clinical correlates in a VA primary care behavioral health clinic*. Psychiatry Res 2015;228:162-5.
- Conroy M, Menard W, Fleming-Ives K, et al. *Prevalence and clinical characteristics of body dysmorphic disorder in an adult inpatient setting*. Gen Hosp Psychiatry 2008;30:67-72.
- Weingarden H, Renshaw KD, Davidson E, et al. *Relative relationships of general shame and body shame with body dysmorphic phenomenology and psychosocial outcomes*. J Obsessive Compuls Relat Disord 2017;14:1-6.
- Shaw AM, Arditte Hall KA, Rosenfield E, et al. *Body dysmorphic disorder symptoms and risk for suicide: the role of depression*. Body Image 2016;19:169-74.
- Weingarden H, Renshaw KD, Wilhelm S, et al. *Anxiety and shame as risk factors for depression, suicidality, and functional impairment in body dysmorphic disorder and obsessive compulsive disorder*. J Nerv Ment Dis 2016;204:832-9.
- Archer CA, Moran K, Garza K, et al. *Relationship between symptom severity, psychiatric comorbidity, social/occupational impairment, and suicidality in hoarding disorder*. J Obsessive Compuls Relat Disord. 2018; Nov. [Epub ahead of print]
- Ayers CR, Dozier ME. *Predictors of hoarding severity in older adults with hoarding disorder*. Int Psychogeriatr 2015;27:1147-56.
- Lovato L, Ferrão YA, Stein DJ, et al. *Skin picking and trichotillomania in adults with obsessive-compulsive disorder*. Compr Psychiatry 2012;53:562-8.
- Grant JE, Odlaug BL, Kim SW. *A clinical comparison of pathologic skin picking and obsessive-compulsive disorder*. Compr Psychiatry 2010;51:347-52.
- Albert U, Speciani M, De Ronchi D, et al. *Factors associated with increased suicide risk in obsessive-compulsive disorder*. Off J Ital Soc Psychopathol 2018;24:154-63.
- De La Vega D, Giner L, Courtet P. *Suicidality in subjects with anxiety or obsessive-compulsive and related disorders: recent advances*. Curr Psychiatry Rep 2018;20:26.
- Veale D, Boocock A, Gournay K, et al. *Body dysmorphic disorder. A survey of fifty cases*. Br J Psychiatry J Ment Sci 1996;169:196-201.
- Perugi G, Giannotti D, Frare F, et al. *Prevalence, phenomenology and comorbidity of body dysmorphic disorder (dysmorphophobia) in a clinical population*. Int J Psychiatry Clin Pract 1997;1:77-82.
- Zimmerman M, Mattia JI. *Body dysmorphic disorder in psychiatric outpatients: recognition, prevalence, comorbidity, demographic, and clinical correlates*. Compr Psychiatry 1998;39:265-70.
- Albertini RS, Phillips KA. *Thirty-three cases of body dysmorphic disorder in children and adolescents*. J Am Acad Child Adolesc Psychiatry 1999;38:453-9.
- Altamura C, Paluello MM, Mundo E, et al. *Clinical and subclinical body dysmorphic disorder*. Eur Arch Psychiatry Clin Neurosci 2001;251:105-8.
- Grant JE, Kim SW, Eckert ED. *Body dysmorphic disorder in patients with anorexia nervosa: prevalence, clinical features, and delusional features of body image*. Int J Eat Disord 2002;32:291-300.
- Frare F, Perugi G, Ruffolo G, et al. *Obsessive-compulsive disorder and body dysmorphic disorder: a comparison of clinical features*. Eur Psychiatry 2004;19:292-8.
- Fontenelle LF, Telles LL, Nazar BP, et al. A

- sociodemographic, phenomenological, and long-term follow-up study of patients with body dysmorphic disorder in Brazil. *Int J Psychiatry Med* 2006;36:243-59.
- 28 Phillips KA, Kelly MM. *Suicidality in a placebo-controlled fluoxetine study of body dysmorphic disorder*. *Int Clin Psychopharmacol* 2009;24:26-8.
- 29 Conceição Costa DL, Chagas Assunção M, Arzeno Ferrão Y, et al. *Body dysmorphic disorder in patients with obsessive-compulsive disorder: prevalence and clinical correlates*. *Depress Anxiety* 2012;29:966-75.
- 30 Bjornsson AS, Didie ER, Grant JE, et al. *Age at onset and clinical correlates in body dysmorphic disorder*. *Compr Psychiatry* 2013;54:893-903.
- 31 Hart AS, Phillips KA. *Symmetry concerns as a symptom of body dysmorphic disorder*. *J Obsessive Compuls Relat Disord* 2013;2:292-8.
- 32 De Brito MJA, Nahas FX, Cordás TA, et al. *Body dysmorphic disorder in patients seeking abdominoplasty, rhinoplasty, and rhytidectomy*. *Plast Reconstr Surg* 2016;137:462-71.
- 33 Rief W, Buhlmann U, Wilhelm S, et al. *The prevalence of body dysmorphic disorder: a population-based survey*. *Psychol Med* 2006;36:877-85.
- 34 Buhlmann U, Glaesmer H, Mewes R, et al. *Updates on the prevalence of body dysmorphic disorder: a population-based survey*. *Psychiatry Res* 2010;178:171-5.
- 35 Schieber K, Kollai I, de Zwaan M, et al. *Classification of body dysmorphic disorder – what is the advantage of the new DSM-5 criteria?* *J Psychosom Res* 2015;78:223-7.
- 36 Möllmann A, Dietel FA, Hunger A, et al. *Prevalence of body dysmorphic disorder and associated features in German adolescents: a self-report survey*. *Psychiatry Res* 2017;254:263-7.
- 37 Balci V, Sevincok L. *Suicidal ideation in patients with obsessive-compulsive disorder*. *Psychiatry Res* 2010;175:104-8.
- 38 Matsunaga H, Hayashida K, Kiriike N, et al. *Clinical features and treatment characteristics of compulsive hoarding in Japanese patients with obsessive-compulsive disorder*. *CNS Spectr* 2010;15:258-66.
- 39 Alonso P, Segalàs C, Real E, et al. *Suicide in patients treated for obsessive-compulsive disorder: a prospective follow-up study*. *J Affect Disord* 2010;124:300-8.
- 40 Torres AR, Ramos-Cerqueira ATA, Ferrão YA, et al. *Suicidality in obsessive-compulsive disorder: prevalence and relation to symptom dimensions and comorbid conditions*. *J Clin Psychiatry* 2011;72:17-26.
- 41 Chakraborty V, Cherian AV, Math SB, et al. *Clinically significant hoarding in obsessive-compulsive disorder: results from an Indian study*. *Compr Psychiatry* 2012;53:1153-60.
- 42 Torres AR, Fontenelle LF, Ferrão YA, et al. *Clinical features of obsessive-compulsive disorder with hoarding symptoms: a multi-center study*. *J Psychiatr Res* 2012;46:724-32.
- 43 Streichenwein SM, Thornby JI. *A long-term, double-blind, placebo-controlled crossover trial of the efficacy of fluoxetine for trichotillomania*. *Am J Psychiatry* 1995;152:1192-6.
- 44 Seedat S, Stein DJ. *Psychosocial and economic implications of Trichotillomania: a pilot study in a South African sample*. *CNS Spectr* 1998;3:40-3.
- 45 Lejoyeux M, Arbaretaz M, McLoughlin M, et al. *Impulse control disorders and depression*. *J Nerv Ment Dis* 2002;190:310-4.
- 46 Phillips KA, Taub SL. *Skin picking as a symptom of body dysmorphic disorder*. *Psychopharmacol Bull* 1995;31:279-88.
- 47 Grant JE, Menard W, Phillips KA. *Pathological skin picking in individuals with body dysmorphic disorder*. *Gen Hosp Psychiatry* 2006;28:487-93.
- 48 Machado MO, Köhler CA, Stubbs B, et al. *Skin picking disorder: prevalence, correlates, and associations with quality of life in a large sample*. *CNS Spectr* 2018;23:311-20.
- 49 Didie ER, Menard W, Stern AP, et al. *Occupational functioning and impairment in adults with body dysmorphic disorder*. *Compr Psychiatry* 2008;49:561-9.
- 50 Witte TK, Didie ER, Menard W, et al. *The relationship between body dysmorphic disorder behaviors and the acquired capability for suicide: BDD and the acquired capability for suicide*. *Suicide Life Threat Behav* 2012;42:318-31.
- 51 Phillips KA, Didie ER, Menard W, et al. *Clinical features of body dysmorphic disorder in adolescents and adults*. *Psychiatry Res* 2006;141:305-14.
- 52 Phillips K, Menard W, Pagano M, et al. *Delusional versus nondelusional body dysmorphic disorder: clinical features and course of illness*. *J Psychiatr Res* 2006;40:95-104.
- 53 Pope CG, Pope HG, Menard W, et al. *Clinical features of muscle dysmorphia among males with body dysmorphic disorder*. *Body Image* 2005;2:395-400.
- 54 Coles ME, Phillips KA, Menard W, et al. *Body dysmorphic disorder and social phobia: cross-sectional and prospective data*. *Depress Anxiety* 2006;23:26-33.
- 55 Phillips KA, Menard W, Bjornsson AS. *Cued panic attacks in body dysmorphic disorder*. *J Psychiatr Pract* 2013;19:194-203.
- 56 Grant JE, Menard W, Pagano ME, et al. *Substance use disorders in individuals with body dysmorphic disorder*. *J Clin Psychiatry* 2005;66:309-16.
- 57 Didie ER, Tortolani CC, Pope CG, et al. *Childhood abuse and neglect in body dysmorphic disorder*. *Child Abuse Negl* 2006;30:1105-15.
- 58 Kittler JE, Menard W, Phillips KA. *Weight concerns in individuals with body dysmorphic disorder*. *Eat Behav* 2007;8:115-20.
- 59 Phillips KA, Menard W. *Suicidality in body dysmorphic disorder: a prospective study*. *Am J Psychiatry* 2006;163:1280-2.
- 60 Phillips KA, Didie ER, Menard W. *Clinical features and correlates of major depressive disorder in individuals with body dysmorphic disorder*. *J Affect Disord* 2007;97:129-35.
- 61 S. Ruffolo J, Phillips KA, Menard W, et al. *Comorbidity of body dysmorphic disorder and eating disorders: severity of psychopathology and body image disturbance*. *Int J Eat Disord* 2006;39:11-9.
- 62 Phillips KA, Menard W, Fay C, et al. *Demographic characteristics, phenomenology, comorbidity, and family history in 200 individuals with body dysmorphic disorder*. *Psychosomatics* 2005;46:317-25.
- 63 Phillips KA, Menard W, Fay C. *Gender similarities and differences in 200 individuals with body dysmorphic disorder*. *Compr Psychiatry* 2006;47:77-87.
- 64 Phillips KA. *Suicidality in body dysmorphic disorder*. *Prim Psychiatry* 2007;14: 58-66.

Appendix 1

16 studies referring to the same sample of patients of Phillips 2005.

How to cite this article: Albert U, Pellegrini L, Maina G, et al. *Suicide in obsessive-compulsive related disorders: prevalence rates and psychopathological risk factors*. *Journal of Psychopathology* 2019;25: 139-48.

This is an open access Journal distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.