

ISSN 1848-2554

10<sup>th</sup> INTERNATIONAL CONGRESS

### FLOUR-BREAD '19

12<sup>th</sup> CROATIAN CONGRESS OF CEREAL TECHNOLOGISTS

## BRAŠNO–KRUH '19.

OSIJEK, CROATIA

June 11-14, 2019

# **BOOK OF ABSTRACTS**

ORGANISED BY

JOSIP JURAJ STROSSMAYER UNIVERSITY OF OSIJEK FACULTY OF FOOD TECHNOLOGY OSIJEK - PTFOS

CROATIAN AGENCY FOR AGRICULTURE AND FOOD - HAPIH

INTERNATIONAL ASSOCIATION FOR CEREAL SCIENCE AND TECHNOLOGY - ICC

ISEKI-FOOD ASSOCIATION - IFA

GLOBAL HARMONIZATION INITIATIVE - GHI

#### Published by JOSIP JURAJ STROSSMAYER UNIVERSITY OF OSIJEK FACULTY OF FOOD TECHNOLOGY OSIJEK

*Editors* Antun Jozinović Sandra Budžaki Ivica Strelec Ante Lončarić

*Proofreading* Antonija Šarić Lahorka Budić

*Cover page* Grafika d.o.o., Osijek

Printed by Grafika d.o.o., Osijek

*Edition* 200 copies

#### **ORGANIZING COMMITTEE**

President: Ivica Strelec, Croatia Vice-President: Michaela Pichler, Austria Vice-President: Huub Lelieveld, The Netherlands Vice-President: Sandra Budžaki, Croatia Secretary: Antun Jozinović, Croatia

#### MEMBERS

Jurislav Babić, Croatia Ante Lončarić, Croatia Krunoslav Dugalić, Croatia Marija Stjepanović, Croatia Andrea Gross-Bošković, Croatia Antonija Šarić, Croatia Lidija Jakobek, Croatia Zdenko Lončarić, Croatia Nada Knežević, Croatia

#### SCIENTIFIC COMMITTEE

President: Hamit Köksel, Turkey Vice-President: Antun Jozinović, Croatia Vice-President: Gerhard Schleining, Austria Vice-President: Darja Sokolić, Croatia

#### MEMBERS

Artur Gryszkin, Poland Jasna Mastilović, Serbia
Vlado Guberac, Croatia
Daliborka Koceva Komlenić, Croatia
Mirela Planinić, Croatia
Ana Bucić-Kojić, Croatia
Biljana Pajin, Serbia
Marina Tišma, Croatia
Drago Šubarić, Croatia
Daniela Horvat, Croatia
Borislav Miličević, Croatia
Jelka Pleadin, Croatia

#### INDUSTRIAL ADVISORY BOARD

Dragan Kovačević (HGK), Croatia Darinka Andrašek (Kraš d.d.), Croatia Ivan Sarić (HUP), Croatia Zvonimir Zdunić (POLJINOS), Croatia Katarina Jukić (BC Institut), Croatia Dominka Maričić (Inspecto d.o.o.), Croatia

#### HONOURABLE COMMITTEE

Žaneta Ugarčić, Croatia Vlasta Piližota, Croatia Vladimir Andročec, Croatia

**Reviewers**: Đurđica Ačkar (Croatia), Jurislav Babić (Croatia), Elene Bartkiene (Lithuania), Paola Battilani (Italy), Ana Bucić-Kojić (Croatia), Sandra Budžaki (Croatia), Daniela Čačić Kenjerić (Croatia), Duška Ćurić (Croatia), Bojana Filipčev (Serbia), Ivana Flanjak (Croatia), Daniela Horvat (Croatia), Elizabet Janić Hajnal (Serbia), Antun Jozinović (Croatia), Katarina Jukić (Croatia), Nada Knežević (Croatia), Hamit Köksel (Turkey), Tihomir Kovač (Croatia), Chathudina Liyanage (Sri Lanka), Ante Lončarić (Croatia), Jasna Mastilović (Serbia), Mladenka Pestorić (Serbia), Mirela Planinić (Croatia), Jelka Pleadin (Croatia), Lea Pollak (Croatia), Cristina Molina Rosell (Spain), Alain Sommier (France), Ivica Strelec (Croatia), Bojan Šarkanj (Croatia), Gordana Šimić (Croatia), Ivan Švec (Czech Republic), Marina Tišma (Croatia), Aleksandra Torbica (Serbia), Natalija Velić (Croatia), Dragan Živančev (Serbia).

#### COMPARISON OF THE $\beta$ -GLUCAN CONTENT IN DIFFERENT BARLEY GENOTYPES

Nevena Djukić<sup>1\*</sup>, Stefan Marković<sup>1</sup>, Dragan Živančev<sup>2</sup>, Aleksandra Torbica<sup>3</sup>, Simin Hagh Nazari<sup>4</sup>

<sup>1</sup>University of Kragujevac, Faculty of Science, Radoja Domanovića 12, 34000 Kragujevac, Serbia <sup>2</sup>Institute of Field and Vegetable Crops, Maksima Gorkog 30, 21000 Novi Sad, Serbia <sup>3</sup>University of Novi Sad, Institute for food technology, Bulevar cara Lazara 1, 21000 Novi Sad, Serbia <sup>4</sup>Department Food Science and Technology, Faculty of Agriculture, University of Zanjan, Zanjan, Iran <sup>\*</sup>corresponding author: nevena@kg.ac.rs

poster presentation

Among cereals, the largest amount of  $\beta$ -glucan is contained in oat and barley. The content of  $\beta$ -glucan varies from 2.5% to 11.3% in the whole barley seed.  $\beta$ -glucans have a number of positive effects on human health and a wide range of potential applications in the food and pharmaceutical industry.

The aim of our research was to analyze the variability of  $\beta$ -glucan content in ten varieties of barley. For determining the content of  $\beta$ -glucan ICC Standard method No. 168 was used.

Based on the identified variability of  $\beta$ -glucan content, a similarity between barley varieties was established and was shown by dendrogram. Three clusters of similar varieties were noticed. Within the Cluster 1 with the highest content of  $\beta$ -glucan, there are two varieties - Oplenac (7.08%) and Pek (6.93%) with a degree of similarity 91. This pair shows the distance from the other cluster of 35, and is in relation to the third cluster 51. The other cluster consists of four varieties (the distance between them is 12 to 18), with a slightly lower concentration of  $\beta$ -glucan: Orion (6.35%), Jastrebac (6.06%), Dinarac (5.80%) and Kraguj (5.65%). In the third cluster, varieties with the largest distance from the previous two clusters are grouped: Novosadski 488, Profit, Dunavac and Midžor, with a concentration of  $\beta$ -glucan less than 5%.

The results indicate a genetic divergence in  $\beta$ -glucan content between the investigated genotypes and the ability to select the appropriate barley genotypes for breeding programmes.

**Keywords**:  $\beta$ -glucan, barley, genotypes