



Article

Periodontitis and dementia: A bidirectional relationship?

Harding, Alice and Singhrao, Simarjit Kaur

Available at <http://clock.uclan.ac.uk/39019/>

Harding, Alice and Singhrao, Simarjit Kaur ORCID: 0000-0001-9573-5963 (2022) Periodontitis and dementia: A bidirectional relationship? Journal of Dental Research, 101 (3). pp. 245-246. ISSN 0022-0345

It is advisable to refer to the publisher's version if you intend to cite from the work.
<http://dx.doi.org/10.1177/00220345211043461>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

Periodontitis and Dementia: A Bidirectional Relationship?

A. Harding¹ and S.K. Singhraj¹

Keywords: Alzheimer's disease, risk factor, causal relationship, periodontal disease, oral health, apolipoprotein E allele 4 susceptibility gene

A body of scientific evidence supports the view that periodontal disease and Alzheimer's disease (AD) are comorbid. While periodontal disease affects tooth-supporting tissues and the host's immune responses, leading to eventual tooth loss, AD is characterized by 2 histologic diagnostic markers at autopsy: the extraneuronal amyloid plaques and the intraneuronal neurofibrillary tangles (Hyman et al. 2012). Other lesions without a role in the neuropathologic diagnosis of AD include neuronal and synaptic loss, neuroinflammation, and cerebral amyloid angiopathy (Dugger and Dickson 2017), which are of importance in understanding the disease process. Understanding the relationship between AD and periodontitis is hindered by the long-standing dogma that those with dementia or AD-associated dementia are at greater incidence of longitudinally manifesting periodontal disease than those without it. This view clearly assumes that AD-associated dementia is a risk factor for periodontal disease. The assumption would be that the behavioral changes associated with the onset of dementia, namely poorer levels of oral hygiene, are the predominant cause rendering an individual with AD more susceptible to periodontal disease. This currently held dogmatic view is behind the proposal of a bidirectional relationship between AD and periodontitis! Given that a growing body of literature (Stein et al. 2007; Sparks Stein et al. 2012; Farhad et al. 2014; Demmer et al. 2020; Nadim et al. 2020) suggests that periodontal disease can precede AD-associated dementia manifestation, the suggestion is that a converse relationship also exists. This was something that Ma et al. (2021) also identified in their cases over the course of their study, but they correctly rejected these cases due to the possibility of confounding factor interferences. To this end, Ma et al. assessed the effect of AD without diverse pathologies that may act as confounding factors on conditions that develop because of the human aging process.

Ma et al. (2021) were able to clarify the plausibility of a bidirectional relationship in their population-based cohort study. It is interesting that the incidence of the bidirectional relationship that Ma et al. identified appears to be associated with the younger patients with AD (<60 y) and in the 61- to 70-y cohort with AD, who are likely to have a longer duration of the mental illness. The bidirectional relationship also exists in the higher age group (>80 y) with AD-associated dementia, but this group shows less severe periodontal disease. A reason behind this could have been the result of lifestyle factors—for example, insomnia, stress, or other confounding factors that

were not identified or considered in this study. Although Ma et al. did not report whether their younger cohorts had known genetic susceptibilities or whether there were mutated genes in the selected AD cases, it is plausible to suggest that genetics played a role in the early-onset AD or inherited form of AD and not in AD cases with as-yet-unidentified genetic vulnerabilities that are categorized into the sporadic or late-onset AD form.

This brings us to the apolipoprotein E allele 4 susceptibility gene inheritance (Corder et al. 1993), as it 1) represents the second-highest risk factor for AD-associated dementia after advancing age, 2) has the implication for an early clinical manifestation of the disease, and 3) has potential relevance for the interaction between this susceptibility gene and infection. In addition, there are known genetic vulnerabilities associated with the inherited or early-onset form of AD, and these include the amyloid precursor protein and presenilin 2. These act to enhance amyloid precursor protein gene processing to promote enhanced amyloid deposition (Selkoe 2001), which again may have played a role for the severity of periodontal disease in the younger patients, as often seen in the dental clinic. It is well documented that, although less common, there are groups of younger patients who have far more aggressive periodontal disease, and this has genetic, racial, and immunologic associations (Merchant et al. 2014).

An alternative view would be that a subset of periodontal patients with subsequent AD development denotes a subset of AD cases where the relationship is likely to be the reverse. In a letter to the *British Dental Journal*, Harding and Singhraj (2019) argued that the evidence is lacking to support a generalization in terms of a bidirectional relationship in all AD cases. It is encouraging that Ma et al. (2021) finally made significant progress in their study to find the actual subset of AD-associated dementia cases with longitudinal periodontal disease, in which they identified a plausible bidirectional relationship. A similar

Journal of Dental Research
1–2

© International Association for Dental Research and American Association for Dental, Oral, and Craniofacial Research 2021



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/00220345211043461

journals.sagepub.com/home/jdr

¹Dementia and Neurodegenerative Diseases Research Group, Faculty of Clinical and Biomedical Sciences, School of Dentistry, University of Central Lancashire, Preston, UK

Corresponding Author:

S.K. Singhraj, Dementia and Neurodegenerative Diseases Research Group, Faculty of Clinical and Biomedical Sciences, School of Dentistry, University of Central Lancashire, Harrington Building, Room 132, Preston PR1 2HE, UK.

Email: sksinghraj@uclan.ac.uk

study will benefit from revealing whether periodontal disease leads to AD-associated dementia development and results in an equivalent subset of AD cases, as postulated by Harding and Singhrao.

Long-standing poor oral hygiene/pathogen load (as reflected by increased pocket depth) is a likely risk for exacerbating early-onset AD and rapid deterioration in cognition due to peripheral and intracerebral inflammation following periodontal disease. Ma et al. (2021) concluded that a convincing causal relationship of AD-associated dementia with periodontal disease exists. However, determining causality in studies on oral-systemic (Raittio and Farmer 2021) and oral-chronic cognitive diseases such as AD-associated dementia is complex due to the nature of human pathophysiology and an individual's lifestyle, genes, and environmental influences—all of which make reaching this simplistic conclusion a challenge. Although current research does indicate a potential dependency/relationship between periodontal disease and AD-associated dementia, further research is required to clarify the directionality of this association.

Author Contributions

A. Harding, S.K. Singhrao, contributed to conception, design, and data acquisition, drafted and critically revised the manuscript. Both authors gave final approval and agree to be accountable for all aspects of the work.

Acknowledgments

We thank the University of Central Lancashire, Preston, UK, from which all funding and support were provided.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

References

- Corder EH, Saunders AM, Strittmatter WJ, Schmechel DE, Gaskell PC, Small GW, Roses AD, Haines JL, Pericak-Vance MA. 1993. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science*. 261(5123):921–923.
- Demmer RT, Norby FL, Lakshminarayan K, Walker KA, Pankow JS, Folsom AR, Mosley T, Beck J, Lutsey PL. 2020. Periodontal disease and incident dementia: the Atherosclerosis Risk in Communities Study (ARIC). *Neurology*. 95(12):e1660–e1671.
- Dugger BN, Dickson DW. 2017. Pathology of neurodegenerative diseases. *Cold Spring Harb Perspect Biol*. 9(7):a028035.
- Farhad SZ, Amini S, Khalilian A, Berekatain M, Mafi M, Berekatain M, Rafei E. 2014. The effect of chronic periodontitis on serum levels of tumor necrosis factor-alpha in Alzheimer disease. *Dent Res J (Isfahan)*. 11(5):549–552.
- Harding A, Singhrao SK. 2019. Periodontitis to dementia or converse? *Br Dent J*. 226(9):634.
- Hyman BT, Phelps CH, Beach TG, Bigio EH, Cairns NJ, Carrillo MC, Dickson DW, Duyckaerts C, Frosch MP, Masliah E, et al. 2012. National Institute on Aging–Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimers Dement*. 8(1):1–13.
- Ma KS-K, Hasturk H, Carreras I, Dedeoglu A, Veeravalli JJ, Huang J-Y, Kantarci A, Cheng-Chung Wei J. 2021. Dementia and the risk of periodontitis: a population-based cohort study. *J Dent Res*. In press.
- Merchant SN, Vovk A, Kalash D, Hovencamp N, Aukhil I, Harrison P, Zapert E, Bidwell J, Varnado P, Shaddox LM. 2014. Localized aggressive periodontitis treatment response in primary and permanent dentitions. *J Periodontol*. 85(12):1722–1729.
- Nadim R, Tang J, Dilmohamed A, Yuan S, Wu C, Bakre AT, Partridge M, Ni J, Copeland JR, Anstey KJ, et al. 2020. Influence of periodontal disease on risk of dementia: a systematic literature review and a meta-analysis. *Eur J Epidemiol*. 35(9):821–833.
- Raittio E, Farmer J. 2021. Methodological gaps in studying the oral-systemic disease connection. *J Dent Res*. 100(5):445–447.
- Selkoe DJ. 2001. Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev*. 81(2):741–766.
- Stein PS, Desrosiers M, Donegan SJ, Yepes JF, Kryscio RJ. 2007. Tooth loss, dementia and neuropathy in the Nun study. *J Am Dent Assoc*. 138(10):1314–1322.
- Sparks Stein P, Steffen MJ, Smith C, Jicha G, Ebersole JL, Abner E, Dawson D 3rd. 2012. Serum antibodies to periodontal pathogens are a risk factor for Alzheimer's disease. *Alzheimers Dement*. 8(3):196–203.