

Northumbria Research Link

Citation: Roche, Johanna, Rae, Dale E., Redman, Kirsten N., Knutson, Kristen L., von Schantz, Malcolm, Gómez-Olivé, F. Xavier and Scheuermaier, Karine (2021) Sleep disorders in low- and middle-income countries: a call for action. *Journal of Clinical Sleep Medicine*, 17 (1). pp. 2341-2342. ISSN 1550-9389

Published by: American Academy of Sleep Medicine

URL: <https://doi.org/10.5664/jcsm.9614> <<https://doi.org/10.5664/jcsm.9614>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/46970/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

The *Journal of Clinical Sleep Medicine* is dedicated to advancing the science of clinical sleep medicine. In order to provide subscribers with access to new scientific developments as early as possible, accepted papers are posted prior to their final publication in an issue.

These papers are posted as received—without copyediting or formatting by the publisher. In some instances, substantial changes are made during the copyediting and formatting processes; therefore, the final version of the paper may differ significantly from this version.

Unless indicated otherwise, all papers are copyright of the American Academy of Sleep Medicine. No paper in whole or in part may be used in any form without written permission from the American Academy of Sleep Medicine.

Media & Public Relations: Embargo Policy

A submitted paper is embargoed until it is posted as an “accepted paper” on the journal website. At that time, the paper is no longer embargoed and may be promoted before its final publication in an issue. Although the full text of an accepted paper is available to subscribers only, the abstract is available to the public. The DOI number is a permanent identifier for the article, and the DOI link below should be used in a press release to link to the study. Please contact the public relations department of the American Academy of Sleep Medicine with any questions at media@aasm.org or 630-737-9700.

<https://doi.org/10.5664/jcsm.9614>

Sleep disorders in low- and middle-income countries: a call for action

Response to Zeng L, Chen R, Hu L, et al. Concern about sleep disorders in underresourced settings is imminent. *J Clin Sleep Med*. 2021;17(XX):XXX–XXX. doi:10.5664/jcsm.9590

Johanna Roche, PhD^{1,2}; Dale E. Rae, PhD³; Kirsten N. Redman, MSc¹; Kristen L. Knutson, PhD⁴; Malcolm von Schantz, PhD^{1,5}; F. Xavier Gómez-Olivé, MD, PhD⁶; Karine Scheuermaier, MD¹

¹*Wits Sleep Laboratory, Brain Function Research Group, School of Physiology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa*

²*Institute of Mountain Emergency Medicine, Eurac Research, Bolzano, Italy*

³*Division of Physiological Sciences & Health through Physical Activity, Lifestyle and Sport Research Centre, Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa*

⁴*Feinberg School of Medicine, Northwestern University, Chicago, IL*

⁵*Department of Psychology, Northumbria University, Newcastle-upon-Tyne, United Kingdom*

⁶*MRC/Wits Rural Public Health and Health Transitions Research Unit (Agincourt), School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa*

Address correspondence to: Johanna Roche, Institute of Mountain Emergency Medicine, Eurac Research Via Ipazia 2, 39000 Bolzano, Italy; Email: johanna.roche@eurac.edu

DISCLOSURE STATEMENT

This was not an industry supported study. All authors have reported no financial conflicts of interest.

Manuscript word count: 486

We are grateful to Zeng et al for their response¹ to our recent paper², where we used portable polysomnography in the homes of randomly sampled older adults at the Agincourt Health and Demographic Surveillance System site in rural northeast South Africa. In this increasingly hypertensive and overweight/obese population, we observed an obstructive sleep apnea (OSA) prevalence of 29.3%. Importantly, apnea-hypopnea index (AHI) severity and being a woman were independently associated with a greater cardiometabolic risk score. In their letter¹, Zeng *et al* drew attention to the fact that since ± 4.5 billion people globally live in rural communities, the potential for undiagnosed and untreated sleep disorders across the developing world is huge. In fact, an earlier study of 43,935 persons across eight African and Asian rural communities found a 16.6% overall prevalence (ranging from 4% to 40%) of severe problems falling or staying asleep³.

Target 3.4 of the UN sustainable development goals (SDGs) aims to reduce mortality from non-communicable diseases (NCDs) by one-third by 2030 through prevention and treatment⁴. Primary measures addressing this target, such as pharmacological and behavioral treatments of hypertension and diabetes, are being widely implemented across the world. As our study and Zeng et al's response highlight, there is a need beyond this to address issues such as OSA, which are more complex to diagnose and treat, but nonetheless prevalent and important in combating NCDs.

Our knowledge about the prevalence and natural history of OSA and other sleep disorders in low- and middle-income countries (LMIC) is still insufficient. We agree with Zeng et al that our findings require replication at a larger scale and across LMICs, capturing differences in cultural and genetic

backgrounds. Our results provide important proof of concept for the utility of portable polysomnography in order to study the prevalence of and clinically diagnose OSA in under-resourced settings. This is important not only for addressing cardiometabolic morbidity and mortality, but also the growing health inequality both between social groups and between men and women stemming from untreated OSA, as the daytime effects of OSA affect the ability of patients to function during the daytime and to make a living⁵.

Beyond the issue of diagnosis, there is a paucity of treatment options for OSA in LMICs. In addition to the worldwide need to develop novel specific pharmacological interventions against OSA, there is a need to make CPAP therapy, the current gold-standard treatment, available in and feasible for populations in developing countries, especially in rural areas where electricity supply is expensive and unreliable. This presents a technological challenge for developing and mass-producing affordable CPAP equipment that is resilient to power supply interruptions, or even usable in homes without electricity. Once realistic treatment options are available, there will be a need to complement education about controlling cardiometabolic risk through a healthier lifestyle with specific education about the causes, symptoms, and risks associated with OSA. All of this represents an increasingly urgent world-wide research agenda.

REFERENCES

1. Zeng L, Chen R, Hu L, et al. Concern about sleep disorders in underresourced settings is imminent. *J Clin Sleep Med*. 2021;17(XX):XXX–XXX. doi:10.5664/jcsm.9590
2. Roche J, Rae DE, Redman KN, et al. Impact of obstructive sleep apnea on cardiometabolic health in a random sample of older adults in rural South Africa: building the case for the treatment of sleep disorders in underresourced settings. *J Clin Sleep Med*. 2021;17(7):1423-1434. doi:10.5664/jcsm.9214
3. Stranges S, Tigbe W, Gómez-Olivé FX, Thorogood M, Kandala N-B. Sleep problems: an emerging global epidemic? Findings from the INDEPTH WHO-SAGE study among more than 40,000 older adults from 8 countries across Africa and Asia. *Sleep*. 2012;35(8):1173-1181. doi:10.5665/sleep.2012
4. NCD Countdown 2030 collaborators. NCD Countdown 2030: worldwide trends in non-communicable disease mortality and progress towards Sustainable Development Goal target 3.4. *Lancet*. 2018;392(10152):1072-1088. doi:10.1016/S0140-6736(18)31992-5
5. Durmer JS, Dinges DF. Neurocognitive consequences of sleep deprivation. *Semin Neurol*. 2005;25(1):117-129. doi:10.1055/s-2005-867080

Accepted Article