

Online media reporting of suicidal behaviour in Ghana: Analysis of adherence to the WHO guidelines

International Journal of
Social Psychiatry
2021, Vol. 67(3) 251–259
© The Author(s) 2020



Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/0020764020919787
journals.sagepub.com/home/isp



Emmanuel Nii-Boye Quarshie^{1,2} , Johnny Andoh-Arthur^{2,3},
Kwaku Oppong Asante^{2,3,4} and Winifred Asare-Doku^{2,5}

Abstract

Background: Irresponsible media reporting of suicide is a potential risk for copycat suicide. There is a paucity of studies from sub-Saharan Africa on the quality of media reporting of suicide.

Objectives: We assessed the compliance of Ghanaian online media outlets with the World Health Organization (WHO) guidelines for media reporting of suicide.

Methods: We searched 10 local media outlets with strong online presence in Ghana, to identify suicide-related news reports from 2000 through 2019. We applied summative content analysis and chi-square (χ^2) test to the data.

Results: We included 288 news reports, of which 261 (90.6%) were completed suicides, 7 (2.4%) were attempted suicides and 20 (6.9%) were homicide suicides. Most of the news reports failed to comply with the WHO guidelines: 92.7% mentioned the specific method of the suicide act, 82.6% included 'suicide' in the headline and 55.6% included photos of the victims. The χ^2 tests indicated that privately owned media outlets were more likely than publicly owned to post a photo of the victim, $\chi^2(1) = 17.37, p < .001$, and report the incident location in the headline, $\chi^2(1) = 15.00, p < .001$. However, generally, there were no statistically significant relationships between the quality of reporting and media outlet ownership. Each of the 288 reports failed to mention any of the potentially helpful features recommended by the WHO guidelines.

Conclusion: Regardless of the ownership of the media outlet (whether private or publicly owned), mostly, the online reportage of suicidal behaviour in Ghana deviates sharply from the international recommended best practice by the WHO.

Keywords

Attempted suicide, completed suicide, Ghana, online media, sub-Saharan Africa, suicide reporting

Introduction

Suicide is newsworthy (Machlin et al., 2013; Sullivan, 2007); however, irresponsible media reporting of the phenomenon has been implicated as a risk factor for death by suicide, particularly, in vulnerable groups within the population (Niederkröthaler et al., 2009, 2010; Sinyor, Schaffer, Nishikawa, et al., 2018; Sisask & Värnik, 2012; World Health Organization [WHO], 2014). Studies have consistently identified that persons experiencing suicidal crisis could become aware of suicide methods, which they might have not thought about previously, through exposure to irresponsible media reporting of suicide, thereby increasing the chances of copying the suicidal behaviour (Cheng et al., 2017, 2018; Niederkröthaler et al., 2012; Nutt et al., 2015; Pirkis, Burgess, et al., 2006; Tsai, 2010; WHO & International Association for Suicide Prevention [IASP], 2017). This form of social learning has been found mainly among students

and young people (Cheng et al., 2018; Gould et al., 2014; O'Connor et al., 2014; Zalsman et al., 2016).

The suicide reporting guidelines of the WHO provide standards for responsible media reporting of suicide around the world (WHO, 2000; WHO & IASP, 2017), but several countries have developed context-specific

¹School of Psychology, University of Leeds, Leeds, UK

²Centre for Suicide and Violence Research (CSVR), Accra, Ghana

³Department of Psychology, University of Ghana, Accra, Ghana

⁴Department of Psychology, University of the Free State, Bloemfontein, South Africa

⁵School of Medicine and Public Health, The University of Newcastle, Callaghan, NSW, Australia

Corresponding author:

Emmanuel Nii-Boye Quarshie, School of Psychology, University of Leeds, Leeds LS2 9JT, West Yorkshire, UK.

Email: enquarshie@gmail.com

standard guidelines for professional media practitioners (Mishara & Dargis, 2019; Pirkis, Blood, et al., 2006; Samaritans, 2013; Sinyor, Schaffer, Heisel, et al., 2018; Utterson et al., 2017). The WHO reporting guidelines outline certain potentially helpful features, which may help prevent suicide, and potentially harmful characteristics, which may provoke suicide, thereby providing a source of education on best practice for media practitioners (WHO, 2000; WHO & IASP, 2017). Evidence from high-income countries suggests that guidelines on media reporting of suicide have been useful, although reliable evaluative evidence on the impact of the guidelines on professional practice of media practitioners and on suicide and suicidal behaviours remains limited (Bolzern et al., 2019; Marzano et al., 2018; Niederkrotenthaler et al., 2010; Pirkis, Blood, et al., 2006; Pirkis et al., 2017; Stack, 2020; Utterson et al., 2017; Williams & Witte, 2018).

In most low- and middle-income countries (LAMICs), the WHO reporting guidelines remain the main reference document guiding media practitioners in reporting suicide. Similar to research findings from many high-income countries (Bolzern et al., 2019; Marzano et al., 2018; McTernan et al., 2018; Stack, 2020; Utterson et al., 2017), recent evidence shows that adherence to the WHO's recommendations and compliance with standard reporting guidelines are still lacking among journalists and media outlets in LAMICs (Arafat et al., 2019; Armstrong et al., 2018; Chandra et al., 2014; Chang & Freedman, 2018; Cheng et al., 2017; Chiang et al., 2016; Chu et al., 2018; Chun et al., 2018; Yang et al., 2013). For example, quality assessment of media reporting of suicide in India, within the lens of the WHO reporting guidelines, shows that reporting of potentially harmful features (e.g. details of suicide method) is common, while recommended helpful reporting practices are scarce – for example, providing contact details of available and accessible suicide support services (Armstrong et al., 2018; Chandra et al., 2014). In Bangladesh, Bhutan, China, India, Indonesia, Korea, Sri Lanka and Taiwan, both online and newspaper reporting of suicide have been found to fail in adhering to the WHO recommended reporting guidelines (Arafat et al., 2019; Chandra et al., 2014; Chang & Freedman, 2018; Chu et al., 2018; Chun et al., 2018; Nisa et al., 2020; Sørensen et al., 2019; Zangmo & Zangmo, 2019).

Recent evidence from Bhutan, India, Indonesia and Sri Lanka indicates that although some media reports show some level of adherence to recommended reporting guidelines, the majority of the available media reports are non-compliant (Chandra et al., 2014; Nisa et al., 2020; Sørensen et al., 2019; Zangmo & Zangmo, 2019). Frequently, they reported the method of self-harm and suicide in the headline, included identifiable information (e.g. picture) of the victim, used sensational and insensitive language and attributed a single-factor cause to the self-harm or suicide, without providing any information on help-seeking and available support services.

The majority of news reports on suicide in Korea mentioned the method used and provided specific details about the location where the suicidal act took place, and almost half of the reports that were analysed mentioned the contents of suicide notes, while less than 3% of the news reports provided information about helplines, research evidence about suicide and suicide prevention (Chun et al., 2018). Similarly, in Bangladesh, most of the news reports mentioned the names of victims, victims' occupations and provided details of the suicide method; others reported life events and made monocausal attributions of suicide, the term 'suicide' appeared in the headlines of the majority of the reports, some of the reports posted pictures of victims, but no news report provided any potentially helpful recommended information – for example, expert opinion or information about suicide prevention (Arafat et al., 2019, 2020a, 2020b; Arafat, Khan, et al., 2020). Besides these breaches of the recommended reporting practices being associated with increased chances of copycat suicide and imitative suicidal behaviours among vulnerable groups (Niederkrotenthaler et al., 2010, 2012; Pirkis, Blood, et al., 2006), loved ones and families bereaved by suicide and the general public have reported being upset by the media providing details of the death, posting photos of the deceased and the general careless and insensitive nature of media reporting of suicide (Chang & Freedman, 2018; Chapple et al., 2013; Gregory et al., 2020).

Furthermore, the WHO and the IASP caution against the use of the 'c-word' ('committing', 'commits' or 'committed' suicide), as the phrase 'committed suicide' connotes criminality and a moral sin and reinforces stigma (WHO & IASP, 2017). In place of c-words, neutral terms such as 'ended/took her or his life' are suggested (Beaton et al., 2013). Despite this caution, there are reports that the use of the c-word with suicide is still common, particularly, in media reports of suicide (Beaton et al., 2013; Nielsen et al., 2016; WHO & IASP, 2017).

There is a paucity of studies from sub-Saharan African countries on the quality of media reporting of suicide, while no published study from Ghana has specifically examined the quality and compliance of media reporting of suicidal behaviours with any known standard reporting guidelines (Akotia et al., 2019; Quarshie et al., 2015). In the present study, we assessed the compliance of Ghanaian online media outlets with the WHO guidelines for media reporting of suicide, by performing a summative content analysis of the frequency and patterns of both breaches and recommended contents of news reports on suicide, attempted suicide and homicide suicide.

Methods

Design and data source

We reviewed the contents of news reports on suicide and suicidal behaviours in Ghana from 2000 through 2019.

Two authors (E.N.-B.Q. and W.A.-D.) performed keyword searches in January 2020 of the portals of 10 local media outlets with strong online presence in Ghana. Three of the media outlets were publicly owned (*Daily Graphic*, *Ghana News Agency* and *Ghanaian Times*), while seven were privately owned (Adom FM, Citi Newsroom, Daily Guide Africa, GhanaWeb, MyJoyOnline, Peace FM and Starr FM). Previous studies have found the news portals of these media houses as having wider readership and listenership among Ghanaians (Quarshie et al., 2015, 2018). We retrieved eligible news reports about completed suicide, attempted suicide and homicide suicide; we excluded news reports about euthanasia, and suicide bombing, as were also editorials and opinion pieces. English is the official language of Ghana and the language for all written media contents in the country. Across the review period, we identified a total of 513 potentially eligible news reports. We independently assessed the eligibility of the total hits and identified 288 news reports included in the final analysis of this study.

Data extraction, coding and analysis

Consistent with the WHO guidelines for media reporting of suicide (WHO, 2000; WHO & IASP, 2017), we followed the summative content analysis procedure recommended by Hsieh and Shannon (2005) to identify potentially harmful characteristics (breaches) and potentially helpful characteristics (recommended features) in each of the included 288 news reports. Potentially harmful characteristics identified in the news reports were related to breaches of the WHO reporting guidelines, for example, mention of the term 'suicide' in the headline, provision of detailed account of suicide method used, giving monocausal explanation for suicidality, romanticising or glamorising suicide, mention of the age, name, school/workplace of victims, posting a photo of the victim or a photo symbolic of self-harm/suicide (e.g. hangman's noose) and inclusion of additional problematic online features (e.g. links to other articles/websites about suicide, reader-generated comment threads).

We also examined each news story for potentially helpful characteristics related to providing mental health literacy (e.g. dispelling common myths about suicide or providing public education about the facts of suicide), drawing on health experts' views, research and data to inform public and promoting help-seeking behaviour (e.g. raising awareness of crisis support or prevention services available). In accordance with the WHO reporting guidelines, we included each of the identified breaches and recommended features as the variables of the analysis, and coded absent (0) or present (1).

Using the Statistical Package for Social Sciences (SPSS, Version 26.0, for Windows), we analysed the data for the frequencies and patterns of both breaches and recommended contents of the included news reports. We

applied chi-square (χ^2) test to compare the compliance of suicide reporting between public and privately owned online media; we used Fisher's exact test where a cell expected frequency was <5 (Fleiss et al., 2003). The conventional statistical threshold of $p < .05$ was used to determine statistically significant results.

Ethics

The protocol guiding this study was not submitted to an Institutional Review Board for ethical approval, as the study did not include the recruitment and involvement of human participants. However, in developing this article, we have preserved the ethical position of this study by excluding identifying information of media outlets, and suicidal persons and their families mentioned in the included news reports (e.g. names, address, place of work or school).

Results

A total of 288 news reports drawn from public ($n=62$ (21.5%)) and privately owned ($n=226$ (78.5%)) online media outlets met the inclusion criteria for this study. Of the 288 news reports, 261 (90.6%) were completed suicides, 7 (2.4%) were attempted suicides and 20 (6.9%) were homicide suicides. Of the 261 completed suicides, 13 (5%) were identified as celebrity suicides.

Table 1 presents the quality assessment of reporting compliance with WHO guidelines for media reporting of suicide. Generally, most of the included news reports failed to comply with the WHO guidelines: 82.6% included the term 'suicide' in the headline, as was a 'c-word' included in the headlines of 79.9%. While 92.7% of the news reports mentioned the specific method of the suicide act, 66.3% provided detailed descriptions of the methods used for the act. Also, 92.7% of the news reports mentioned the specific place or location where the suicidal act took place. Furthermore, 83.7% reported the names of the victims, 55.6% included photos of the victims and 50.3% provided reader-generated comment threads.

Surprisingly, each of the 288 news reports analysed failed to mention any of the potentially helpful features recommended by the WHO reporting guidelines (e.g. providing public education about the facts of suicide, mentioning a suicide prevention programme or support service, or providing contact details for a suicide support service). The χ^2 tests indicated that privately owned online media outlets were more likely than those publicly owned to report the incident location in the news headline, $\chi^2(1) = 15.00, p < .001$; include a photo of the victim, $\chi^2(1) = 17.37, p < .001$; and provide a detailed account of the method used for the suicidal act, $\chi^2(1) = 4.36, p < .05$. However, generally, across the range of variables analysed, the χ^2 results did not show statistically significant relationships between the quality of reporting and media outlet ownership.

Table 1. Quality assessment of reporting compliance with WHO suicide reporting guidelines.

Potentially harmful breaches	Absent				Present				χ^2 test	p value
	Total		Private media		Total		Private media			
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)		
Headlines										
'Suicide' in the headline	50 (17.4)	8 (16.0)	42 (84.0)	238 (82.6)	54 (22.7)	184 (77.3)	1.094	.296		
Suicide method in the headline	187 (64.9)	57 (30.5)	130 (69.5)	101 (35.1)	5 (5.0)	96 (95.0)	25.304	.000		
Life event(s) in the headline	216 (75.0)	50 (23.1)	166 (76.9)	72 (25.0)	12 (16.7)	60 (83.3)	1.343	.247		
'C-word' in the headline ^a	58 (20.1)	8 (13.8)	50 (86.2)	230 (79.9)	54 (23.5)	176 (76.5)	2.572	.109		
Incident location	151 (52.4)	46 (30.5)	105 (69.5)	137 (47.6)	16 (11.7)	121 (88.3)	15.004	.000		
Suicidal act (as presented in the body of story)										
Suicide method reported	21 (7.3)	3 (14.3)	18 (85.7)	267 (92.7)	59 (22.1)	208 (77.9)	0.701	.437		
Detailed account of method	97 (33.7)	14 (14.4)	83 (85.6)	191 (66.3)	48 (25.1)	143 (74.9)	4.358	.037		
Place or public site named as location of a suicide death/attempt	21 (7.3)	3 (14.3)	18 (85.7)	267 (92.7)	59 (22.1)	208 (77.9)	0.701	.437		
Time of the act (e.g. morning, afternoon, night)	142 (49.3)	33 (23.2)	109 (76.8)	146 (50.7)	29 (19.9)	117 (80.1)	0.486	.486		
Suicidal behaviour described as 'successful' or 'unsuccessful'	232 (80.6)	62 (26.7)	170 (73.3)	56 (19.4)	0	56 (100)	19.071	.000		
Causes of suicidality										
Monocausal explanation for suicidality	180 (62.5)	43 (23.9)	137 (76.1)	108 (37.5)	19 (17.6)	89 (82.4)	1.584	.208		
Narrative brushes over the complex realities of suicide	232 (80.6)	52 (22.4)	180 (77.6)	56 (19.4)	10 (17.9)	46 (82.1)	0.554	.457		
Contents/details from suicide note reported	267 (92.7)	58 (21.7)	209 (78.3)	21 (7.3)	4 (19.0)	17 (81.0)	0.082	.774		
Consideration for bereaved persons										
Romanticising or glamorising suicide or over-emphasising family/community expressions of grief	217 (75.3)	48 (22.1)	169 (77.9)	71 (24.7)	14 (19.7)	57 (80.3)	0.183	.669		
Interview with bereaved persons (comments from survivors or bereaved)	130 (45.1)	33 (25.4)	97 (74.6)	158 (54.9)	29 (18.4)	129 (81.6)	2.087	.149		

(Continued)

Table 1. (Continued)

Potentially harmful breaches	Absent		Present		χ^2 test	p value		
	Total	Public media	Private media	Total			Public media	Private media
	n (%)	n (%)	n (%)	n (%)			n (%)	n (%)
Disclosure of victim's identity								
Name of suicidal person mentioned	47 (16.3)	9 (19.1)	38 (80.9)	241 (83.7)	53 (22.0)	188 (78.0)	0.188	.661
Age of suicidal person mentioned	53 (18.4)	15 (28.3)	38 (71.7)	235 (81.6)	47 (20.0)	188 (80.0)	1.764	.184
Occupation or student status mentioned	88 (30.6)	12 (13.6)	76 (86.4)	200 (69.4)	50 (25.0)	150 (75.0)	4.671	.031
Name of school or place of work mentioned	148 (51.4)	26 (17.6)	122 (82.4)	140 (48.6)	36 (25.7)	104 (74.3)	2.826	.093
Photo of victim	128 (44.4)	42 (32.8)	86 (67.2)	160 (55.6)	20 (12.5)	140 (87.5)	17.368	.000
Celebrity status								
Victim identified as a celebrity	275 (95.5)	60 (21.8)	215 (78.2)	13 (4.5)	2 (15.4)	11 (84.6)	0.303	.741
Victim related to or has link with a celebrity	283 (98.3)	61 (21.6)	222 (78.4)	5 (1.7)	1 (20.0)	4 (80.0)	0.007	1.000
Additional potentially harmful features								
Link(s) to other articles/websites about suicide	273 (94.8)	62 (22.7)	211 (77.3)	15 (5.2)	0	15 (100)	4.326	0.47
Reader-generated comment threads	143 (49.7)	25 (17.5)	118 (82.5)	145 (50.3)	37 (25.5)	108 (74.5)	2.751	.097
Suggests or encourages a suicide cluster	268 (93.1)	58 (21.6)	210 (78.4)	20 (6.9)	4 (20.0)	16 (80.0)	0.030	1.000
Potentially helpful features (recommended features)								
Mental health literacy								
Dispels the myths about suicide or educate the public about the facts of suicide	288 (100)	62 (21.5)	226 (78.5)	–	–	–	–	–
Draws on health experts, research and data to inform public	288 (100)	62 (21.5)	226 (78.5)	–	–	–	–	–
Promoting help-seeking behaviour								
Mentions a suicide prevention programme/support service	288 (100)	62 (21.5)	226 (78.5)	–	–	–	–	–
Provides contact details for a suicide support service	288 (100)	62 (21.5)	226 (78.5)	–	–	–	–	–

Note: WHO: World Health Organization; χ^2 : chi-square.

^aThe c-word criterion refers to the mention of 'commit' suicide, 'committed' suicide or 'committing' suicide in the headline of the news report.

Discussion

While it is relevant to compare the evidence of the present study to findings of the large number of previous international studies – mainly, from high-income countries – it may be more meaningful to discuss the evidence of our study in the light of the available evidence from LAMICs and the context of Ghana. The present study shows that, generally, the media reporting of suicidal behaviour in Ghana does not comply with the WHO reporting guidelines. This evidence supports recent findings from other LAMICs in Asia, including Bangladesh, Bhutan, India, Indonesia, Sri Lanka and Taiwan (Arafat et al., 2019; Chiang et al., 2016; Nisa et al., 2020; Sørensen et al., 2019; Zangmo & Zangmo, 2019). The evidence also supports a previous anecdotal observation that media coverage of suicide in Ghana is sensational, overly simplistic and explicit (Quarshie et al., 2015). The predominant inclusion of the c-word and the term ‘suicide’ in the headlines as evident in the present study is particularly troubling, even though not entirely surprising. Recent evidence indicates that news editors in Ghana consider ‘suicide’ as a catchy term which attracts reader attention (Akotia et al., 2019), while some emerging researchers of suicide in the country, without background training in suicidology, have been found to use the c-word in scientific publications (e.g., Abdulai, 2020; Acheampong & Aziato, 2018; Der et al., 2016). As pertaining in many LAMICs (Mishara & Weisstub, 2016), suicide is highly stigmatised and morally tabooed, and attempted suicide is criminalised in Ghana (Act 29 of Ghana, 1960; Adinkrah, 2013; Osafo et al., 2015). Thus, the inclusion of the c-word and ‘suicide’ in the headlines of news reports of suicidal behaviour has implications for potential inadvertent deepening of the criminality and moral condemnation of victims and survivors of (attempted) suicide and their families (Beaton et al., 2013; Nielsen et al., 2016).

As found in other LAMICs (Chandra et al., 2014), generally, the present study found no statistically significant relationships between the quality of reporting and media outlet ownership – a plausible indication that the extent of non-compliance with recommended best practice transcends the ownership of media outlets. It is possible that today’s fiercely competitive media terrain and the extreme commercial motive of media outlets could be responsible for this sensationalistic media practice – every media house, public or privately owned, wants to be the first to break the news on suicide (Akotia et al., 2019; Yang et al., 2013). Relatedly, the disclosure of victim’s identity (mention of names, school, place of work or posting of photos) remains a critical ethical issue; some researchers have interpreted this practice to mean journalists and media outlets exploiting the suicide event for commercial gains, while remaining uncompassionate to the bereaved family (Akotia et al., 2019; Harshe et al., 2016).

Furthermore, the mention and detailed description of the method and location of suicide have been identified

as cause for concern. Narrative and systematic reviews have identified the existence of the Werther effects in vulnerable groups (Blood et al., 2007; Cheng et al., 2017; Niederkrotenthaler et al., 2010, 2012; Pirkis et al., 2009; Sisask & Värnik, 2012; Stack, 2003), while adolescents with histories of self-harm have attributed their knowledge of self-harm methods to media reporting and portrayal of suicide (O’Connor et al., 2014). In online media reporting of suicide, the additional inclusion of links to other articles/websites about suicide and reader-generated comment threads could be problematic (Utterson et al., 2017). Evidence suggests that the online discussion forums are associated with suicidal behaviours in young people (Dunlop et al., 2011).

Finally, the evidence of the present study that each of the 288 media reports analysed failed to mention any of the potentially helpful features recommended by the WHO guidelines is worthy of some comments. This finding is consistent with recent evidence from Bangladesh, India and Indonesia, where no news report analysed provided any potentially helpful information recommended by the WHO guidelines (Arafat et al., 2019, 2020a, 2020b; Chandra et al., 2014; Nisa et al., 2020; Sørensen et al., 2019). Perhaps in Ghana, this troubling evidence highlights the need for the country’s Ministry of Health to consider launching stakeholder training programmes and sustained awareness creation among media practitioners on the importance of adhering to recommended standard guidelines for media reporting of suicidal behaviour.

Limitations and recommendations

The non-inclusion of media outlets with lesser online presence in Ghana might have limited the evidence of this study, regarding the extent and pattern of compliance with or breaches of the WHO guidelines. The present study fails to assess the effects of non-compliant media reporting of suicide on the actual rates of suicide and suicidal behaviours in Ghana. Future studies may consider the use of ecological approaches and longitudinal designs to examine the copycat effects of non-compliant media reporting of suicide in Ghana.

Responsible media reporting of suicide is now a key area of focus for suicide prevention among vulnerable groups (Oexle et al., 2019; WHO, 2014; Zalsman et al., 2016). Thus, the WHO and leading researchers in the area of media and suicide prevention have recommended the development of media reporting guidelines by countries around the world (Chu et al., 2018; Mishara & Dargis, 2019; Pirkis, Blood, et al., 2006; WHO, 2014). The key evidence of the present study underscores the urgent need for mental health professionals and key media role players in Ghana (the Ghana Mental Health Authority, Ghana Psychological Council, Ghana Media Commission, National Communication Authority, the Ghana Journalists Association, training institutions of media practitioners and the Centre for Suicide and

Violence Research) to collaborate towards the formulation, adoption and implementation of guidelines for responsible (online) media reporting of suicide in the country.

Conclusion

The evidence of the present study shows that, in Ghana, online media reporting of suicidal behaviour deviates sharply from the international recommended best practice by the WHO; these reports use sensationalistic language and crude practices that are insensitive to the needs of loved ones and families bereaved by suicide and may have the inherent tendency of triggering copycat suicide in vulnerable groups.

Author contributions

E.N.-B.Q helped in the conception and design. All authors contributed to the acquisition of data, drafting and critical revision of the manuscript, and the final approval of the manuscript.

Conflict of interest

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Emmanuel Nii-Boye Quarshie  <https://orcid.org/0000-0002-8720-2355>

References

- Abdulai, T. (2020). Trends of online news media reported suicides in Ghana (1997–2019). *BMC Public Health*, *20*(1), 35. <https://doi.org/10.1186/s12889-020-8149-3>
- Acheampong, A. K., & Aziato, L. (2018). Suicidal ideations and coping strategies of mothers living with physical disabilities: A qualitative exploratory study in Ghana. *BMC Psychiatry*, *18*(1), 1–8. <https://doi.org/10.1186/s12888-018-1938-x>
- Act 29 of Ghana. (1960). *The Criminal Offences Act, 1960*. Assembly Press.
- Adinkrah, M. (2013). Criminal prosecution of suicide attempt survivors in Ghana. *International Journal of Offender Therapy and Comparative Criminology*, *57*(12), 1477–1497. <https://doi.org/10.1177/0306624X12456986>
- Akotia, C. S., Osafo, J., Asare-Doku, W., & Boakye, K. E. (2019). News editors' views about suicide and suicide stories in Ghana. *Psychological Studies*, *65*, 1–8. <https://doi.org/10.1007/s12646-019-00511-4>
- Arafat, S. M. Y., Khan, M. M., Ueda, M., Niederkrotenthaler, T., & Armstrong, G. (2020). Assessing the quality of media reporting of suicide deaths in Bangladesh against World Health Organization guidelines. *Crisis*, *41*(1), 47–53. <https://doi.org/10.1027/0227-5910/a000603>
- Arafat, S. M. Y., Mali, B., & Akter, H. (2019). Quality of online news reporting of suicidal behavior in Bangladesh against World Health Organization guidelines. *Asian Journal of Psychiatry*, *40*, 126–129. <https://doi.org/10.1016/j.ajp.2018.10.010>
- Arafat, S. M. Y., Mali, B., & Akter, H. (2020a). Do Bangladeshi newspapers educate public while reporting suicide? A year round observation from content analysis of six national newspapers. *Asian Journal of Psychiatry*, *48*, e101920. <https://doi.org/10.1016/j.ajp.2019.101920>
- Arafat, S. M. Y., Mali, B., & Akter, H. (2020b). Is suicide reporting in Bangla online news portals sensible? A year-round content analysis against World Health Organization guidelines. *Asian Journal of Psychiatry*, *49*, e101943. <https://doi.org/10.1016/j.ajp.2020.101943>
- Armstrong, G., Vijayakumar, L., Niederkrotenthaler, T., Jayaseelan, M., Kannan, R., Pirkis, J., & Jorm, A. F. (2018). Assessing the quality of media reporting of suicide news in India against World Health Organization guidelines: A content analysis study of nine major newspapers in Tamil Nadu. *Australian & New Zealand Journal of Psychiatry*, *52*(9), 856–863. <https://doi.org/10.1177/0004867418772343>
- Beaton, S., Forster, P., & Maple, M. (2013). Suicide and language: Why we shouldn't use the 'C' word. *InPsych*, *35*(1), 30–31.
- Blood, R. W., Pirkis, J., & Holland, K. (2007). Media reporting of suicide methods: An Australian perspective. *Crisis*, *28*(Suppl. 1), 64–69. <https://doi.org/10.1027/0227-5910.28.S1.64>
- Bolzern, J. E., Mnyama, N. L., & McMillan, D. (2019). Responsible journalism, imitative suicide, and transgender populations: A systematic review of UK newspapers. *Journal of Gay & Lesbian Mental Health*, *23*(3), 276–288. <https://doi.org/10.1080/19359705.2019.1620148>
- Chandra, P. S., Doraiswamy, P., Padmanabh, A., & Philip, M. (2014). Do newspaper reports of suicides comply with standard suicide reporting guidelines? A study from Bangalore, India. *International Journal of Social Psychiatry*, *60*(7), 687–694. <https://doi.org/10.1177/0020764013513438>
- Chang, K.-K., & Freedman, E. (2018). WHO's media guidelines in the press and in public perception. *Journal of Media Ethics*, *33*(1), 14–25. <https://doi.org/10.1080/23736992.2017.1401930>
- Chapple, A., Ziebland, S., Simkin, S., & Hawton, K. (2013). How people bereaved by suicide perceive newspaper reporting: Qualitative study. *The British Journal of Psychiatry*, *203*(3), 228–232. <https://doi.org/10.1192/bjp.bp.112.114116>
- Cheng, Q., Chen, F., Lee, E. S. T., & Yip, P. S. F. (2018). The role of media in preventing student suicides: A Hong Kong experience. *Journal of Affective Disorders*, *227*, 643–648. <https://doi.org/10.1016/j.jad.2017.11.007>
- Cheng, Q., Chen, F., & Yip, P. S. (2017). Media effects on suicide methods: A case study on Hong Kong 1998–2005. *PLOS ONE*, *12*(4), Article e0175580. <https://doi.org/10.1371/journal.pone.0175580>
- Chiang, Y.-C., Chung, F.-Y., Lee, C.-Y., Shih, H.-L., Lin, D.-C., & Lee, M.-B. (2016). Suicide reporting on front pages of major newspapers in Taiwan violating reporting recommendations between 2001 and 2012. *Health Communication*, *31*(11), 1395–1404. <https://doi.org/10.1080/10410236.2015.1074024>

- Chu, X., Zhang, X., Cheng, P., Schwebel, D. C., & Hu, G. (2018). Assessing the use of media reporting recommendations by the World Health Organization in suicide news published in the most influential media sources in China, 2003–2015. *International Journal of Environmental Research and Public Health*, *15*(3), 451. <https://doi.org/10.3390/ijerph15030451>
- Chun, J., Kim, J., & Lee, S. (2018). Fidelity assessment of the suicide reporting guidelines in Korean newspapers. *BMC Public Health*, *18*(1), Article 1115. <https://doi.org/10.1186/s12889-018-6014-4>
- Der, E. M., Dakwah, I. A., Derkyi-Kwarteng, L., & Badu, A. A. (2016). Hanging as a method of suicide in Ghana: A 10 year autopsy study. *Pathology Discovery*, *4*, Article 2. <https://doi.org/10.7243/2052-7896-4-2>
- Dunlop, S. M., More, E., & Romer, D. (2011). Where do youth learn about suicides on the Internet, and what influence does this have on suicidal ideation? *Journal of Child Psychology and Psychiatry*, *52*(10), 1073–1080. <https://doi.org/10.1111/j.1469-7610.2011.02416.x>
- Fleiss, J. L., Levin, B., & Paik, M. C. (2003). *Statistical methods for rates and proportions* (3rd ed.). John Wiley.
- Gould, M. S., Kleinman, M. H., Lake, A. M., Forman, J., & Middle, J. B. (2014). Newspaper coverage of suicide and initiation of suicide clusters in teenagers in the USA, 1988–96: A retrospective, population-based, case-control study. *The Lancet Psychiatry*, *1*(1), 34–43. [https://doi.org/10.1016/S2215-0366\(14\)70225-1](https://doi.org/10.1016/S2215-0366(14)70225-1)
- Gregory, P., Stevenson, F., King, M., Osborn, D., & Pitman, A. (2020). The experiences of people bereaved by suicide regarding the press reporting of the death: Qualitative study. *BMC Public Health*, *20*(1), 1–14. <https://doi.org/10.1186/s12889-020-8211-1>
- Harshe, D., Karia, S., Harshe, S., Shah, N., Harshe, G., & De Sousa, A. (2016). Celebrity suicide and its effect on further media reporting and portrayal of suicide: An exploratory study. *Indian Journal of Psychiatry*, *58*(4), 443–447. <https://doi.org/10.4103/0019-5545.196704>
- Hsieh, H.-F., & Shannon, S. E. (2005). Three approaches to qualitative content analysis. *Qualitative Health Research*, *15*(9), 1277–1288. <https://doi.org/10.1177/1049732305276687>
- Machlin, A., Pirkis, J., & Spittal, M. J. (2013). Which suicides are reported in the media – And what makes them ‘newsworthy’? *Crisis*, *34*(5), 305–313. <https://doi.org/10.1027/0227-5910/a000177>
- Marzano, L., Fraser, L., Scally, M., Farley, S., & Hawton, K. (2018). News coverage of suicidal behavior in the United Kingdom and the Republic of Ireland. *Crisis*, *39*(5), 386–396. <https://doi.org/10.1027/0227-5910/a000533>
- McTernan, N., Spillane, A., Cully, G., Cusack, E., O’Reilly, T., & Arensman, E. (2018). Media reporting of suicide and adherence to media guidelines. *International Journal of Social Psychiatry*, *64*(6), 536–544. <https://doi.org/10.1177/0020764018784624>
- Mishara, B. L., & Dargis, L. (2019). Systematic comparison of recommendations for safe messaging about suicide in public communications. *Journal of Affective Disorders*, *244*, 124–154. <https://doi.org/10.1016/j.jad.2018.09.031>
- Mishara, B. L., & Weisstub, D. N. (2016). The legal status of suicide: A global review. *International Journal of Law and Psychiatry*, *44*, 54–74. <https://doi.org/10.1016/j.ijlp.2015.08.032>
- Niederkrotenthaler, T., Fu, K.-w., Yip, P. S., Fong, D. Y., Stack, S., Cheng, Q., & Pirkis, J. (2012). Changes in suicide rates following media reports on celebrity suicide: A meta-analysis. *Journal of Epidemiology & Community Health*, *66*(11), 1037–1042. <https://doi.org/10.1136/jech-2011-200707>
- Niederkrotenthaler, T., Till, B., Kapusta, N. D., Voracek, M., Dervic, K., & Sonneck, G. (2009). Copycat effects after media reports on suicide: A population-based ecologic study. *Social Science & Medicine*, *69*(7), 1085–1090. <https://doi.org/10.1016/j.socscimed.2009.07.041>
- Niederkrotenthaler, T., Voracek, M., Herberth, A., Till, B., Strauss, M., Etzersdorfer, E., . . . Sonneck, G. (2010). Role of media reports in completed and prevented suicide: Werther v. Papageno effects. *The British Journal of Psychiatry*, *197*, 234–243. <https://doi.org/10.1192/bjp.bp.109.074633>
- Nielsen, E., Padmanathan, P., & Knipe, D. (2016). Commit* to change? A call to end the publication of the phrase ‘commit* suicide’. *Wellcome Open Research*, *1*, 21. <https://doi.org/10.12688/wellcomeopenres.10333.1>
- Nisa, N., Arifin, M., Nur, M. F., Adella, S., & Marthoenis, M. (2020). Indonesian online newspaper reporting of suicidal behavior: Compliance with World Health Organization media guidelines. *International Journal of Social Psychiatry*, *66*(3), 259–262. <https://doi.org/10.1177/0020764020903334>
- Nutt, R., Kidd, B., & Matthews, K. (2015). Assessing the adherence to guidelines of media reporting of suicide using a novel instrument – The ‘Risk of Imitative Suicide Scale’ (RISc). *Suicide and Life-Threatening Behavior*, *45*(3), 360–375. <https://doi.org/10.1111/sltb.12137>
- O’Connor, R. C., Rasmussen, S., & Hawton, K. (2014). Adolescent self-harm: A school-based study in Northern Ireland. *Journal of Affective Disorders*, *159*, 46–52. <https://doi.org/10.1016/j.jad.2014.02.015>
- Oexle, N., Niederkrotenthaler, T., & DeLeo, D. (2019). Emerging trends in suicide prevention research. *Current Opinion in Psychiatry*, *32*(4), 336–341. <https://doi.org/10.1097/YCO.0000000000000507>
- Osafo, J., Akotia, C. S., Andoh-Arthur, J., & Quarshie, E. N.-B. (2015). Attempted suicide in Ghana: Motivation, stigma, and coping. *Death Studies*, *39*(5), 274–280. <https://doi.org/10.1080/07481187.2014.991955>
- Pirkis, J. E., Blood, R. W., Beautrais, A., Burgess, P., & Skehan, J. (2006). Media guidelines on the reporting of suicide. *Crisis*, *27*(2), 82–87. <https://doi.org/10.1027/0227-5910.27.2.82>
- Pirkis, J. E., Burgess, P. M., Francis, C., Blood, R. W., & Jolley, D. J. (2006). The relationship between media reporting of suicide and actual suicide in Australia. *Social Science & Medicine*, *62*(11), 2874–2886. <https://doi.org/10.1016/j.socscimed.2005.11.033>
- Pirkis, J. E., Dare, A., Blood, R. W., Rankin, B., Williamson, M., Burgess, P., & Jolley, D. (2009). Changes in media reporting of suicide in Australia between 2000/01 and 2006/07. *Crisis*, *30*(1), 25–33. <https://doi.org/10.1027/0227-5910.30.1.25>
- Pirkis, J. E., Rossetto, A., Nicholas, A., Ftanou, M., Robinson, J., & Reavley, N. (2017). Suicide prevention media campaigns: A systematic literature review. *Health Communication*, *34*(4), 402–414. <https://doi.org/10.1080/10410236.2017.1405484>

- Quarshie, E. N.-B., Asante, K. O., Andoh-Arthur, J., Asare-Doku, W., & Navelle, P. L. (2018). Suicide attempts and deaths in older persons in Ghana: A media surveillance approach. *Current Psychology*. Advance online publication. <https://doi.org/10.1007/s12144-018-9932-5>
- Quarshie, E. N.-B., Osafo, J., Akotia, C. S., & Peprah, J. (2015). Adolescent suicide in Ghana: A content analysis of media reports. *International Journal of Qualitative Studies on Health and Well-Being*, *10*(1), 27682. <https://doi.org/10.3402/qhw.v10.27682>
- Samaritans. (2013). *Media guidelines for reporting suicide*. Samaritans.
- Sinyor, M., Schaffer, A., Heisel, M. J., Picard, A., Adamson, G., Cheung, C. P., . . . Sareen, J. (2018). Media guidelines for reporting on suicide: 2017 update of the Canadian Psychiatric Association policy paper. *The Canadian Journal of Psychiatry*, *63*(3), 182–196. <https://doi.org/10.1177/0706743717753147>
- Sinyor, M., Schaffer, A., Nishikawa, Y., Redelmeier, D. A., Niederkrotenthaler, T., Sareen, J., . . . Pirkis, J. (2018). The association between suicide deaths and putatively harmful and protective factors in media reports. *Canadian Medical Association Journal*, *190*(30), E900–E907. <https://doi.org/10.1503/cmaj.180900>
- Sisask, M., & Värnik, A. (2012). Media roles in suicide prevention: A systematic review. *International Journal of Environmental Research and Public Health*, *9*(1), 123–138. <https://doi.org/10.3390/ijerph9010123>
- Sørensen, J. B., Pearson, M., Andersen, M. W., Weerasinghe, M., Rathnaweera, M., Rathnapala, D. C., . . . Konradsen, F. (2019). Self-harm and suicide coverage in Sri Lankan newspapers. *Crisis*, *10*(1), 54–61. <https://doi.org/10.1027/0227-5910/a000534>
- Stack, S. (2003). Media coverage as a risk factor in suicide. *Journal of Epidemiology & Community Health*, *57*(4), 238–240. <https://doi.org/10.1136/jech.57.4.238>
- Stack, S. (2020). Media guidelines and suicide: A critical review. *Social Science & Medicine*. Advance online publication. <https://doi.org/10.1016/j.socscimed.2019.112690>
- Sullivan, G. (2007). Should suicide be reported in the media? In M. Mitchell (Ed.), *Remember me: Constructing immortality-beliefs on immortality, life, and death* (pp. 149–158). Routledge; Taylor & Francis.
- Tsai, J.-F. (2010). The media and suicide: Evidence based on population data over 9 years in Taiwan. *Suicide and Life-Threatening Behavior*, *40*(1), 81–86. <https://doi.org/10.1521/suli.2010.40.1.81>
- Utterson, M., Daoud, J., & Dutta, R. (2017). Online media reporting of suicides: Analysis of adherence to existing guidelines. *BJPsych Bulletin*, *41*, 83–86. <https://doi.org/10.1192/pb.bp.115.052761>
- Williams, C. L., & Witte, T. K. (2018). Media reporting on suicide: Evaluating the effects of including preventative resources and psychoeducational information on suicide risk, attitudes, knowledge, and help-seeking behaviors. *Suicide and Life-Threatening Behavior*, *48*(3), 253–270. <https://doi.org/10.1111/sltb.12355>
- World Health Organization. (2000). *Preventing suicide. A resource for media professionals*. WHO.
- World Health Organization. (2014). *Preventing suicide, a global imperative*. WHO.
- World Health Organization, & International Association for Suicide Prevention. (2017). *Preventing suicide: A resource for media professionals (updated)*. World Health Organization.
- Yang, A. C., Tsai, S.-J., Yang, C.-H., Shia, B.-C., Fuh, J.-L., Wang, S.-J., . . . Huang, N. E. (2013). Suicide and media reporting: A longitudinal and spatial analysis. *Social Psychiatry and Psychiatric Epidemiology*, *48*(3), 427–435. <https://doi.org/10.1007/s00127-012-0562-1>
- Zalsman, G., Hawton, K., Wasserman, D., van Heeringen, K., Arensman, E., Sarchiapone, M., . . . Balazs, J. (2016). Suicide prevention strategies revisited: 10-year systematic review. *The Lancet Psychiatry*, *3*(7), 646–659. [https://doi.org/10.1016/S2215-0366\(16\)30030-X](https://doi.org/10.1016/S2215-0366(16)30030-X)
- Zangmo, T., & Zangmo, S. (2019). Media reporting of suicides in Bhutan: Analysis of adherence to WHO guidelines. *Journal of Bhutan Studies*, *40*, 100–126.