

Possible gauge theoretic origin for quark-lepton complementarity

To cite this article: Paul H. Frampton and Rabindra N. Mohapatra JHEP01(2005)025

View the [article online](#) for updates and enhancements.

Related content

- [A Modern Introduction to Neutrino Physics: Lepton flavour violation](#)
F F Deppisch
- [A Tour of the Subatomic Zoo \(Third edition\): The standard model!](#)
C Schwarz
- [Universal mass texture and quark-lepton complementarity](#)
F González Canales and A Mondragón

Recent citations

- [How low-scale trinification sheds light in the flavor hierarchies, neutrino puzzle, dark matter, and leptogenesis](#)
A. E. Cárcamo Hernández *et al*
- [Investigating the Sterile Neutrino Parameters with QLC in 3 + 1 Scenario](#)
Gazal Sharma and B. C. Chauhan
- [Importance of generalized symmetry and its CP extension on neutrino mixing and leptogenesis](#)
Rome Samanta *et al*

Possible gauge theoretic origin for quark-lepton complementarity

Paul H. Frampton^a and Rabindra N. Mohapatra^b

^a*Department of Physics and Astronomy, University of North Carolina
Chapel Hill, NC 27599-3255, U.S.A.*

^b*Department of Physics, University of Maryland, College Park
MD 20742, U.S.A.*

E-mail: frampton@physics.unc.edu, rmohapat@physics.umd.edu

ABSTRACT: Similarity between the weak interaction properties of quarks and leptons has led to suggestions that the origin of lepton mixing angles may be related to those of quarks. In this paper, we present a gauge model based on $SU(2)_L \times SU(2)_R \times SU(4)_c$ group that leads to a new form for the quark lepton complementarity which predicts the solar neutrino mixing angle in terms of the Cabibbo angle for the case of inverted mass hierarchy for neutrinos. We also indicate how these ideas can be implemented in an E_6 inspired trinification $SU(3)_C \times SU(3)_L \times SU(3)_R$ model, which is more closely allied to string theory by the AdS/CFT correspondence.

KEYWORDS: Beyond Standard Model, Neutrino Physics.

Contents

1.	Introduction	1
2.	Gauge model for quark-lepton complementarity (QLC) in the 1-2 sector	2
3.	Inclusion of next order terms and derivation of quark-lepton complementarity	5
4.	A possible trinification model example	7
5.	Discussion	8

1. Introduction

One of the major challenges for particle theory today is to understand the totally different pattern of mixings among neutrinos compared to quarks. Although many different ways have been proposed to tackle this challenge (for recent reviews, see [1]), no final consensus has emerged on what is the most promising way to proceed. Any new approach is therefore desirable.

In this paper, we will focus on understanding the solar mixing angle, $\theta_{\odot} \equiv \theta'_{12}$, which is large but known not to be maximal. Since there are leptonic symmetries that lead to maximal solar mixing, a possible approach to this problem is to start in that limit and understand the deviation using new physics. There have been many attempts in the literature to achieve this goal [2]. In this note, we pursue a recent suggestion [3, 4] according to which the deviation from maximality of the solar mixing may be related to the quark mixing angle $\theta_C \equiv \theta^q_{12}$. It is based on the observation that the mixing angle responsible for solar neutrino oscillations, $\theta_{\odot} \equiv \theta'_{12}$ satisfies an interesting complementarity relation with the corresponding angle in the quark sector $\theta_{Cabibbo} \equiv \theta^q_{12}$ i.e. $\theta'_{12} + \theta^q_{12} \simeq \pi/4$. This equation is satisfied to within a few per cent accuracy. While it is quite possible that this is purely accidental, it is interesting to pursue the possibility that there is a deep meaning behind it and see where it leads. We explore the possibility that this relation may be a reflection of a fundamental quark-lepton unification.

To see how quark lepton unification can possibly lead to a complementarity relation, first note that θ_{\odot} is the 12 entry of the PMNS matrix defined as $U_{PMNS} = U_{\ell}^{\dagger} U_{\nu}$ (where U_{ℓ} is the unitary matrix operating on the left handed charged leptons that diagonalizes the (e, μ, τ) mass matrix and U_{ν} is the corresponding one for the neutrino Majorana mass matrix) and θ_c is the corresponding entry in the CKM matrix defined as $V_{CKM} = U_u^{\dagger} U_d$ (where $U_{u,d}$ diagonalize the up and down quark mass matrices). Secondly suppose that

the structure of neutrino and quark mass matrices at the high scale are such that to a leading order, the PMNS matrix is exact bimaximal [5] whereas the CKM matrix is an identity matrix and to next leading order we have the down quark and charged lepton mass matrices are equal (or very nearly so) in the basis where the up quark and neutrino Dirac masses remain unchanged, then a complementarity relation between quarks and leptons can emerge from quark-lepton unification at high scale.

In this paper, we make an attempt to connect the deviation of solar mixing angle from its maximal value to the Cabibbo angle using the gauge group $SU(2)_L \times SU(2)_R \times SU(4)_c$ symmetry that unifies quarks and leptons [6] at high scale. We find that if we use the double seesaw formula for understanding small neutrino masses, then the departure of the solar angle from maximality is determined by the Cabibbo angle and the prediction for the solar mixing angle is consistent with experiments within present uncertainties. This model also predicts a “large” value for $\theta_{13} \simeq 0.15$.

We also indicate how these considerations may be extended to the case of a gauge group $[SU(3)]^3$ which unifies to the gauge group E_6 at high scale, that has quark-lepton unification.

2. Gauge model for quark-lepton complementarity (QLC) in the 1-2 sector

Before proceeding to the construction of the model, we first outline a set of conditions that are sufficient for obtaining a connection between the θ_{\odot} and θ_C . The full quark and lepton mass matrices should be written in a form such that, we have:

$$M_{u,d} = M_{u,d}^0 + \delta M_{u,d} \tag{2.1}$$

$$M_{\ell} = M_{\ell}^0 + \delta M_{\ell} \tag{2.2}$$

$$\mathcal{M}_{\nu} = U_{bm}^* \mathcal{M}_{\nu}^d U_{bm}^{\dagger} \tag{2.3}$$

where U_{bm} is the bimaximal PMNS matrix given by:

$$U_{bm} = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{2} & -\frac{1}{2} & \frac{1}{\sqrt{2}} \\ \frac{1}{2} & -\frac{1}{2} & -\frac{1}{\sqrt{2}} \end{pmatrix}. \tag{2.4}$$

We require that $M_u^0 = \tan\beta M_d^0$ so that to leading order $V_{CKM}^0 = \mathbf{I}$, where \mathbf{I} is the identity matrix and $M_{\ell}^0 = M_d^0$ due to quark lepton symmetry. We also choose $\delta M_u = 0$. Note also that we have chosen the $\delta M_{\ell} \approx \delta M_d$ at the scale where neutrino masses arise via seesaw mechanism i.e. exact quark-lepton symmetry to zeroth order but approximate quark-lepton symmetry upto the first order terms that generate the 12 mixing angles. The observed pattern in the V_{CKM} then arises from the matrix δM_d . Its similarity in form to δM_{ℓ} feeds this change to the lepton mixing matrix in a way such that a form of quark-lepton complementarity emerges in the end. The Cabibbo angle is small in our model due to it being related to the quark mass ratios. We also note that δM_d has to be different from δM_{ℓ} in order to fit the muon mass after correcting for extrapolation effects from the seesaw scale to the weak scale. The departure from maximality of solar mixing angle is predicted in terms of Cabibbo angle as well as muon and strange quark masses.

We first describe in detail how in a gauge theory based on the gauge group $SU(2)_L \times SU(2)_R \times SU(4)_c$ the above conditions can be satisfied. In a subsequent section, we will indicate how the same considerations can be carried over to a trinification theory.

Let us start with the gauge group $SU(2)_L \times SU(2)_R \times SU(4)_c$ with supersymmetry and an additional softly broken global symmetry on the theory given by $U(1)_X$ where $X = f_1 - f_2 - f_3$ with f_i representing the i -th family. In the lepton sector this corresponds to $L_e - L_\mu - L_\tau$ symmetry, which has been extensively discussed in the literature [7]. We make the usual assignment of fermions to the $\Psi(2, 1, 4) + \Psi^c(1, 2, 4^*)$ as follows:

$$\Psi = \begin{pmatrix} u_1 & u_2 & u_3 & \nu \\ d_1 & d_2 & d_3 & e \end{pmatrix}_L \quad (2.5)$$

We add a set of three singlet fermions S_i and S_i^c to the theory ($i=1,2,3$ stands for generation) which also have the same fermion number as the normal matter multiplet. For the Higgs sector, we choose the following multiplets: $\phi \equiv (2, 2, 0)$; $\chi \equiv (2, 1, 4)$ and $\chi^c \equiv (1, 2, 4^*)$. In a supersymmetric theory, anomaly cancellation requires that we also have the fields: $\bar{\chi} \equiv (2, 1, 4^*)$ and $\bar{\chi}^c \equiv (1, 2, 4)$.

We break the gauge symmetry down to the standard model by the vevs of the fields $\langle \chi^c \rangle = \langle \bar{\chi}^c \rangle = v_R$. The standard model symmetry is then broken down by the vev $\langle \phi \rangle = \begin{pmatrix} \kappa & 0 \\ 0 & \kappa' \end{pmatrix}$. We keep $\langle \chi \rangle = \langle \bar{\chi} \rangle = 0$, which can be a stable vacuum if we forbid the trilinear term $\chi\phi\chi^c$ by a discrete symmetry.

We can write down the Yukawa couplings of the model responsible for quark and lepton masses as follows:

$$W_Y = h_{ij}\Psi_i\phi\Psi_j^c + f_{ij}(\Psi_i\bar{\chi}S_j^c + \Psi_i^c\bar{\chi}^cS_j) + \mu_{ij}S_iS_j. \quad (2.6)$$

We choose the Majorana mass for the S^c fields to be very heavy so that they decouple at TeV energies. This also decouples the $\Psi_i\bar{\chi}S_j^c$ term from the lagrangian, which we wrote down above for completeness. Note that $U(1)_X$ symmetry restricts the form of μ_{ij} to be

$$\mu = \begin{pmatrix} 0 & \mu_1 & \mu_2 \\ \mu_1 & 0 & 0 \\ \mu_2 & 0 & 0 \end{pmatrix}. \quad (2.7)$$

After the $\langle \chi^c \rangle$ and the $\langle \phi \rangle$ vevs are turned on, we have the following zeroth order forms for the various fermion mass matrices:

$$\frac{\kappa'}{\kappa}M_u^0 \equiv \cot\beta M_u^0 = M_d^0 = M_\ell^0 \quad (2.8)$$

where

$$M_u^0 = \begin{pmatrix} m_{11} & 0 & 0 \\ 0 & m_{22} & m_{23} \\ 0 & m_{23} & m_{33} \end{pmatrix}.$$

Since the up and down quark masses are proportional, the CKM matrix is the unit matrix. The leptonic mixing matrix is however not proportional to a unit matrix as can be seen below.

The mass matrix for (ν, ν^c, S) is given by:

$$M = \begin{pmatrix} 0 & M_u^0 & 0 \\ M_u^0 & 0 & f v_R \\ 0 & f^T v_R & \mu \end{pmatrix} \quad (2.9)$$

where $M_u^0, f v_R$ are all 3×3 matrices whose forms are similar and the form of μ is given above. First point to note is that the matrix M_u^0 can be diagonalized without effecting the form of μ . In general the matrix f can be written in the form

$$\begin{pmatrix} f_{11} & 0 & 0 \\ 0 & f_{22} & f_{23} \\ 0 & f_{23} & f_{33} \end{pmatrix}$$

due to the $U(1)_X$ symmetry.

The matrix in eq. (2.9) is in the double seesaw form [8] which leads to small neutrino masses even with a multi-TeV scale for the seesaw. The scale however can also be close to the GUT scale¹. In this note, we keep the double seesaw scale v_R near 100 TeV and include its effect (via renormalization group) on the fermion masses through a parameter Δ .

Diagonalizing the full neutrino matrix in eq. 2.9 [8], one obtains the light neutrino mass matrix \mathcal{M}_ν to be

$$\mathcal{M}_\nu = \begin{pmatrix} 0 & m_1 & m_2 \\ m_1 & 0 & 0 \\ m_2 & 0 & 0 \end{pmatrix} \quad (2.10)$$

where $m_1 = \frac{m_{11} m_{22} \mu_1}{f_{11} f_{22} v_R^2}$ and $m_2 = \frac{m_{11} \mu_1 m_{33}}{f_{11} f_{23} v_R^2} + \frac{m_{11} \mu_2 m_{33}}{f_{11} f_{33} v_R^2}$. Thus for instance, if $m_{ij} \sim \text{GeV}$ and $\mu_i \sim \text{GeV}$ and $v_R \simeq 100 \text{ TeV}$, we get for neutrino masses $m_i \simeq 0.1 \text{ eV}$, which is of the same order as the square root of the atmospheric neutrino mass difference square. We could increase v_R by appropriately adjusting the scale μ_{ij} . For instance, if v_R is assumed to be at the GUT scale, so would be μ_{ij} . Clearly, if $m_1 \simeq m_2$ we also get maximal mixing needed for atmospheric neutrino oscillation.

Note that solar mixing angle is maximal and in fact the PMNS matrix at this zeroth order level is bimaximal for $m_1 = m_2$ and the mass ordering of neutrinos is the so-called inverted type. Of course the model at the zeroth order level has the unpleasant feature that it predicts vanishing solar neutrino mass difference which however is corrected as we include higher order terms.

From the relation $M_u^0 = \tan\beta M_d^0 = \tan\beta M_\ell^0$, it may appear that, we have unacceptable relations among the masses of quarks and leptons. Models with such a property have been discussed before [9] and it has been shown how such models, if supersymmetric can be phenomenologically fully viable. These mass relations without further corrections imply that $m_s^0/m_b^0 = m_c/m_t$ and $m_\mu^0(M_c) = m_s^0(M_c)$, where M_c is the $SU(4)$ scale. Note that m_s and m_μ cannot be close to each other at the multi-TeV scale since renormalization group extrapolation due to strong interactions increases m_s without changing m_μ much.

¹This possibility was pointed out by S. F. King (private communication).

Therefore, we will use the δM terms to get the right m_s/m_μ . We will see that we can get a sizable correction to the maximality of solar mixing angle dictated by the zeroth order terms.

We will assume that the δM terms do not affect the third generation fermion masses. The ratio m_c/m_t then implies that $m_b = m_\tau \sim 10$ GeV. These masses are much higher than their observed values. It has however been noted that in supersymmetric theories, there are large gluino and Wino contributions to the bottom and tau masses, which can correct for this [9] and bring them down to observed values. For large $\tan\beta$, there are two contributions to the quark masses and one for the charged lepton [10]:

$$\delta m_b = \frac{2\alpha_s m_{\tilde{g}}}{3\pi m_{\tilde{q}}^2} (m_b^0 \mu \tan\beta + A_{33}^d m_0) + \frac{Y_t^2 \mu}{16\pi^2 m_{\tilde{q}}^2} (m_b^0 \mu + A_{33}^u m_0 \tan\beta) \quad (2.11)$$

and

$$\delta m_\tau = \frac{g_1^2 m_{\tilde{B}}}{16\pi^2 m_{\tilde{l}}^2} (m_\tau^0 \mu \tan\beta + A_{33}^e m_0). \quad (2.12)$$

Clearly, there are several independent parameters which one may adjust to get the bottom and the tau lepton mass right. Note that the quark mixings still vanish at this stage. As noted before, if we choose a value for $m_s \simeq m_\mu$ somewhere around 100 MeV. For v_R in the 100 TeV range, when we come down to the weak scale, m_s will increase and become bigger than m_μ . In the next section, we show how the higher order effects can fix this problem, generate quark and lepton mixing leading to quark-lepton complementarity as well as give the first generation masses right.

3. Inclusion of next order terms and derivation of quark-lepton complementarity

In order to obtain the CKM angles and the departure from bimaximal pattern for neutrino mixings, we need to include higher dimensional non-renormalizable terms in the theory that break $U(1)_X$. We assume that they contribute only to the down quark and charged lepton mass matrix in a quark-lepton symmetric way. For this purpose, in the 422-model, we include a $SU(2)_R$ triplet Higgs field δ_R which transforms under the gauge group as $(1, 3, 15)$ and an $SU(2)_L \times SU(2)_R$ singlet $\sigma(1, 1, 15)$, both of which have $X = +2$. We will also include a δ_L with zero vev so that it has no effect on the masses. This allows us to include two terms which can lead to violation of X charge after symmetry breaking:

$$\delta \mathcal{L} = \frac{1}{M} [f'_{ij} \psi_i^T \phi \vec{\tau} \cdot \vec{\delta}_R \psi_j^c + f''_{ij} \psi_i^T \phi \sigma \psi_j^c] + \frac{1}{M^2} (\psi_1^{T,\alpha} \tau_2 \phi \bar{\chi}^{c\beta} \chi_\alpha^{cT} \tau_2 \psi_{1,\beta}^c). \quad (3.1)$$

In the first term, we assume that f' and f'' are symmetric matrices and choose, $f'_{11} = f''_{11} = 0$; f_{12} , f_{22} and f_{13} as well as f''_{12} , f''_{22} and f''_{13} are chosen to be nonzero. As just noted, once $\vec{\delta}_R$ and σ acquire vevs, we can arrange their values and the couplings f' and f'' so that the contribution of these terms to the fermion mass appears predominantly in the down sector. Secondly due to the fact that the multiplets transform as **15** dim. representation of $SU(4)_c$, they will make different contribution to δM_d and δM_ℓ .

In the second term, we have explicitly given the $SU(4)_c$ indices α and β . The $SU(2)_{L,R}$ indices have been suppressed. The Pauli matrix τ_2 operates in the $SU(2)_{L,R}$ space. Note that due since $\langle \chi_{\alpha=4}^c \rangle$ and $\langle \chi^c = 4 \rangle$ are the only components that acquire vevs, it contributes to the charged lepton masses and not to quark masses. There are of course other invariants to this order but we assume that they are absent.

We choose $\langle \delta_R \rangle, \langle \sigma \rangle \simeq \langle \chi^c \rangle$ and $M \sim 10^2 (\langle \delta_R \rangle, \langle \sigma \rangle)$ so that the magnitude of these terms of the right order of magnitude (i.e. 10^{-2} GeV for the first two and few $\times 10^{-3}$ GeV for the last term) so that we generate the right value for the Cabibbo angle. Clearly, as noted the last term in eq. (3.1) makes a small contribution (due to two powers of M in the denominators) to the electron mass but does not contribute to quark masses. Collecting all these together and ignoring the third generation fermions we can write the mass matrices for the first two generation of quarks and leptons in the presence of the higher dimensional terms at the weak scale:

$$\begin{aligned} M_\ell &= \begin{pmatrix} m_0 & -3\delta' \\ -3\delta' & m_\mu^0 - 3\delta \end{pmatrix} \\ M_d &= \Delta \begin{pmatrix} 0 & \delta' \\ \delta' & m_\mu^0 + \delta \end{pmatrix} \end{aligned} \tag{3.2}$$

where m_μ^0 is the zeroth order contribution; δ and δ' are the next order contributions from eq. (3.1) and Δ is renormalization effect on the quark masses from v_R down to M_Z . The m_0 term in the charged lepton mass matrix M_ℓ comes from the second term in eq. (3.1) and therefore does not have a counter part in the down quark mass matrix.

In this model $\theta_c \simeq \frac{\delta'}{m_\mu^0 + \delta} \equiv \sqrt{\frac{m_d}{m_s}}$. On the other hand, the mixing angle for charged lepton is:

$$\theta^\ell \simeq \frac{-3\theta_C m_s}{\Delta m_\mu}. \tag{3.3}$$

We choose the scale v_R to be low (in the multi-TeV range) and choose for example of $m_s \sim m_\mu/2$ at v_R scale. The value of m_s increases somewhat as we move to the weak scale but still remains below m_μ^2 . For example if $\Delta = 1.5$ (corresponding to v_R in the multi-TeV range), we estimate $\theta^\ell \simeq \theta_C$. This leads to a prediction for the solar mixing angle [4]:

$$\sin^2 \theta_\odot \simeq 0.32. \tag{3.4}$$

The present 3σ fit to the solar and KamLand data gives $\sin^2 \theta_\odot = 0.23 - 0.38$. It also leads to a prediction for the $\theta_{13} \simeq \frac{\theta_C}{\sqrt{2}} \simeq 0.15$, which can be tested in the next round of searches for this parameter.

In this model, we get $m_e \sim m_0 - \frac{9\theta_C^2 m_s^2}{\Delta^2 m_\mu}$ which for $m_s \simeq 75$ MeV is about $m_0 - 8$ MeV. We can adjust m_0 to get the right value of m_e .

The next issue we need to address is the origin of solar mass difference square. It can arise in a manner which does not affect the quark-lepton complementarity by adding a gauge singlet σ' with $X = 2$. This allows a term of the form $\lambda S_1 S_1 \sigma'$ with $\lambda \sim 10^{-2}$. This introduces a μ_{11} entry which the leads via double seesaw to the solar mass difference of the right order for $\langle \sigma' \rangle \simeq \mu$.

²Recent lattice calculations seem to prefer such low values

4. A possible trinification model example

Another gauge model which can also provide a realization of QLC is the trinification model based on the gauge group $[\text{SU}(3)]^3$ [11] which arises from the grand unification group E_6 . Here we assign the fermions to

$$\psi_i = \psi_R + \psi_L + \psi_c \equiv (3, 3^*, 1)_{Li} + (3^*, 1, 3)_{Li} + (1, 3, 3^*)_{Li}. \quad (4.1)$$

Here the subscript (C,L,R) denotes the group under which the multiplet is a singlet. We then add three singlet fermions

$$S_i = (1, 1, 1)_i. \quad (4.2)$$

For scalars, we use

$$\Phi = \phi_c + \phi_L + \phi_R \quad (4.3)$$

where $\phi_c \sim (1, 3, 3^*) + c.c.$, $\phi_R \sim (3, 3^*, 1) + c.c.$, and $\phi_L \sim (3, 1, 3^*) + c.c.$

Again we impose a $X = f_1 - f_2 - f_3$ symmetry. In addition before symmetry breaking there is an $\text{SU}(3)_C \times \text{SU}(3)_L \times \text{SU}(3)_R \times Z_3$ quasi-simple symmetry where the Z_3 interchanges the $\text{SU}(3)_i$. Here the $\text{SU}(3)_L$ plays a role similar to the generalized weak isospin of the 331-model [12]. However, it is necessary to respect the fully unified simple E_6 symmetry to obtain the required relation between quark and lepton couplings.

There are two types of Yukawa couplings possible in this theory: for quarks the Yukawa responsible giving mass are of the form $\psi_L \psi_R \phi_c$ whereas for leptons it has the form $\epsilon^{abc} \epsilon_{pqr} \psi_{c,a}^p \psi_{c,b}^q \phi_{c,c}^r$. At the $[\text{SU}(3)]^3$ level, these two couplings are independent and therefore there is no connection between quarks and leptons. However, if the theory is assumed to emerge from an E_6 GUT model, the two couplings arise from the same [27]³ coupling and become equal. In order to derive QLC relations, we need to assume this. The leptonic part of the Yukawa interaction can then be written as (i, j, k are generation indices):

$$\mathcal{L}_Y = h_{ij} \psi_{c,i} \psi_{c,j} \phi_c + f_{ij} \psi_{c,i} S_{Lj} \phi_c^* + \mu_{ij} S_{Li} S_{Lj}. \quad (4.4)$$

In this 333-model, the VEVs break the Z_3 symmetry. The $\langle \phi_c \rangle$ has the form

$$\langle \phi_c \rangle = \begin{pmatrix} v_u & 0 & 0 \\ 0 & v_d & v_R \\ 0 & v_L & M_3 \end{pmatrix}. \quad (4.5)$$

There are two kinds of vev's in $\langle \phi \rangle$: the $v_{u,d,L}$ are of order of the weak scale whereas v_R and M_3 are in the multi-TeV range and they break $\text{SU}(3)_R$. These mass matrices lead to the double seesaw neutrino mass matrix given above. The rest of the considerations are similar to the previous model and we do not go into the details here.

This $\text{SU}(3)^3 \times Z_3$ model has the advantage that it is obtainable as a quiver theory [13] from superstring theory by using the AdS/CFT correspondence [14].

5. Discussion

In this paper, we have presented two possible gauge models that realize the quark-lepton complementarity in the 12-sector. An essential ingredient in our first approach is the Pati-Salam $SU(4)_c$ gauge symmetry which has been suspected in ref. [4] to be a possible way to obtain the QLC relation. For the first time, we present explicit models that realize this relation. As we have noted in the body of the paper, the constraints required to obtain the QLC relation poses highly nontrivial challenges for model building. What we show is that it is possible to have a natural realization under a plausible set of assumptions. The double seesaw appears to be crucial in our discussion as is the inverted mass spectrum for neutrinos that is a consequence of $L_e - L_\mu - L_\tau$ symmetry. The model predicts a value of θ_{13} not far below its present upper limit.

Acknowledgments

The work of P.H.F. is supported by the US Department of Energy Grant No DE-FG02-97ER-41036 and that of R.N.M. by the National Science Foundation Grant No. PHY-0354401. One of us (R.N.M.) would like to thank S.F. King, H. Minakata and A. Smirnov for comments and discussions.

References

- [1] For a recent review and references see G. Altarelli and F. Feruglio, *Models of neutrino masses and mixings*, *New J. Phys.* **6** (2004) 106 [[hep-ph/0405048](#)];
R.N. Mohapatra, *Ictp lectures on theoretical aspects of neutrino masses and mixings*, [hep-ph/0211252](#), to appear in *New J. Phys.*
- [2] K.S. Babu and R.N. Mohapatra, *Predictive schemes for bimaximal neutrino mixings*, *Phys. Lett.* **B 532** (2002) 77 [[hep-ph/0201176](#)];
C. Giunti and M. Tanimoto, *Deviation of neutrino mixing from bi-maximal*, *Phys. Rev.* **D 66** (2002) 053013 [[hep-ph/0207096](#)];
P.H. Frampton, S.T. Petcov and W. Rodejohann, *On deviations from bimaximal neutrino mixing*, *Nucl. Phys.* **B 687** (2004) 31 [[hep-ph/0401206](#)];
W. Rodejohann, *A parametrization for the neutrino mixing matrix*, *Phys. Rev.* **D 69** (2004) 033005 [[hep-ph/0309249](#)]; *Type-II see-Saw mechanism, deviations from bimaximal neutrino mixing and leptogenesis*, *Phys. Rev.* **D 70** (2004) 073010 [[hep-ph/0403236](#)];
A. Romanino, *Charged lepton contributions to the solar neutrino mixing and θ_{13}* , *Phys. Rev.* **D 70** (2004) 013003 [[hep-ph/0402258](#)];
G. Altarelli, F. Feruglio and I. Masina, *Can neutrino mixings arise from the charged lepton sector?*, *Nucl. Phys.* **B 689** (2004) 157 [[hep-ph/0402155](#)];
C.A. de S. Pires, *More on deviation from bi-maximal neutrino mixing*, *J. Phys.* **G30** (2004) B29–B34 [[hep-ph/0404146](#)].
- [3] M. Raidal, *Relation between the neutrino and quark mixing angles and grand unification*, *Phys. Rev. Lett.* **93** (2004) 161801 [[hep-ph/0404046](#)].
- [4] H. Minakata and A.Y. Smirnov, *Neutrino mixing and quark lepton complementarity*, *Phys. Rev.* **D 70** (2004) 073009 [[hep-ph/0405088](#)].

- [5] V.D. Barger, S. Pakvasa, T.J. Weiler and K. Whisnant, *Bi-maximal mixing of three neutrinos*, *Phys. Lett.* **B 437** (1998) 107 [[hep-ph/9806387](#)];
A.J. Baltz, A.S. Goldhaber and M. Goldhaber, *The solar neutrino puzzle: an oscillation solution with maximal neutrino mixing*, *Phys. Rev. Lett.* **81** (1998) 5730 [[hep-ph/9806540](#)];
F. Vissani, *Signal of neutrinoless double beta decay, neutrino spectrum and oscillation scenarios*, *J. High Energy Phys.* **06** (1999) 022 [[hep-ph/9906525](#)];
M. Jezabek and Y. Sumino, *Neutrino masses and bimaximal mixing*, *Phys. Lett.* **B 457** (1999) 139 [[hep-ph/9904382](#)].
- [6] J.C. Pati and A. Salam, *Lepton number as fourth color*, *Phys. Rev.* **D 10** (1974) 270.
- [7] R. Barbieri, L.J. Hall, D.R. Smith, A. Strumia and N. Weiner, *Oscillations of solar and atmospheric neutrinos*, *J. High Energy Phys.* **12** (1998) 017 [[hep-ph/9807235](#)];
A.S. Joshipura and S.D. Rindani, *Vacuum solutions of neutrino anomalies through a softly broken U(1) symmetry*, *Eur. Phys. J.* **C 14** (2000) 85 [[hep-ph/9811252](#)];
R.N. Mohapatra, A. Perez-Lorenzana and C.A. de Sousa Pires, *Type-II seesaw and a gauge model for the bimaximal mixing explanation of neutrino puzzles*, *Phys. Lett.* **B 474** (2000) 355 [[hep-ph/9911395](#)];
L. Lavoura, *New model for the neutrino mass matrix*, *Phys. Rev.* **D 62** (2000) 093011 [[hep-ph/0005321](#)];
W. Grimus and L. Lavoura, *A neutrino mass matrix with seesaw mechanism and two-loop mass splitting*, *Phys. Rev.* **D 62** (2000) 093012 [[hep-ph/0007011](#)];
L. Lavoura and W. Grimus, *Seesaw model with softly broken $L_e - L_\mu - L_\tau$* , *J. High Energy Phys.* **09** (2000) 007 [[hep-ph/0008020](#)];
W. Grimus and L. Lavoura, *Softly broken lepton numbers and maximal neutrino mixing*, *J. High Energy Phys.* **07** (2001) 045 [[hep-ph/0105212](#)];
Q. Shafi and Z. Tavartkiladze, *Anomalous flavor U(1): predictive texture for bi-maximal neutrino mixing*, *Phys. Lett.* **B 482** (2000) 1451;
T. Kitabayashi and M. Yasue, *Radiatively induced neutrino masses and oscillations in an $SU(3)_L \times U(1)_N$ gauge model*, *Phys. Rev.* **D 63** (2001) 095002 [[hep-ph/0010087](#)]; *Large mixing angle msw solution in an $SU(3)_L \times U(1)_N$ gauge model with two-loop radiative mechanism*, *Phys. Lett.* **B 508** (2001) 85 [[hep-ph/0102228](#)];
T. Kitabayashi and M. Yasue, *Large solar neutrino mixing and radiative neutrino mechanism*, *Phys. Lett.* **B 524** (2002) 308 [[hep-ph/0110303](#)];
K.S. Babu and R.N. Mohapatra, *Predictive schemes for bimaximal neutrino mixings*, *Phys. Lett.* **B 532** (2002) 77 [[hep-ph/0201176](#)];
H.S. Goh, R.N. Mohapatra and S.-P. Ng, *Testing neutrino mass matrices with approximate $L_e - L_\mu - L_\tau$ symmetry*, *Phys. Lett.* **B 542** (2002) 116 [[hep-ph/0205131](#)]; *Testing neutrino mass matrices with approximate $L_e - L_\mu - L_\tau$ symmetry*, *Phys. Lett.* **B 542** (2002) 116 [[hep-ph/0205131](#)];
H.-J. He, D.A. Dicus and J.N. Ng, *Minimal schemes for large neutrino mixings with inverted hierarchy*, *Phys. Lett.* **B 536** (2002) 83 [[hep-ph/0203237](#)];
Mass matrices with $L_e - L_\mu - L_\tau$ symmetry were discussed in S. Petcov, *On pseudodirac neutrinos, neutrino oscillations and neutrinoless double beta decay*, *Phys. Lett.* **B 110** (1982) 245.
- [8] R.N. Mohapatra, *Mechanism for understanding small neutrino mass in superstring theories*, *Phys. Rev. Lett.* **56** (1986) 561;
R.N. Mohapatra and J.W.F. Valle, *Neutrino mass and Baryon-number nonconservation in superstring models*, *Phys. Rev.* **D 34** (1986) 1642.

- [9] K.S. Babu, B. Dutta and R.N. Mohapatra, *Partial yukawa unification and a supersymmetric origin of flavor mixing*, *Phys. Rev. D* **60** (1999) 095004 [[hep-ph/9812421](#)].
- [10] L. Hall, R. Rattazi and U. Sarid, *The top quark mass in supersymmetric SO(10) unification*, *Phys. Rev. D* **50** (1994) 7048 [[hep-ph/9306309](#)];
 T. Blazek, S. Raby and S. Pokorski, *Finite supersymmetric threshold corrections to ckm matrix elements in the large tan beta regime*, *Phys. Rev. D* **52** (1995) 4151 [[hep-ph/9504364](#)];
 T. Banks, *Supersymmetry and the quark mass matrix*, *Nucl. Phys. B* **303** (1988) 172;
 E. Ma, *Radiative quark and lepton masses through soft supersymmetry breaking*, *Phys. Rev. D* **39** (1989) 1922;
 R. Hempfling, *Yukawa coupling unification with supersymmetric threshold corrections*, *Phys. Rev. D* **49** (1994) 6168.
- [11] A. De Rujula, H. Georgi and S.L. Glashow, *Trinification*, in *Fifth workshop on grand unification*, P.H. Frampton, H. Fried and K. Kang eds., World Scientific, Singapore 1984.
- [12] P.H. Frampton, *Chiral dilepton model and the flavor question*, *Phys. Rev. Lett.* **69** (1992) 2889;
 F. Pisano and V. Pleitez, *An SU(3) \times U(1) model for electroweak interactions*, *Phys. Rev. D* **46** (1992) 410 [[hep-ph/9206242](#)].
- [13] P.H. Frampton, *AdS/CFT string duality and conformal gauge field theories*, *Phys. Rev. D* **60** (1999) 041901 [[hep-th/9812117](#)];
 P.H. Frampton and W.F. Shively, *Conformal N = 0 D = 4 gauge theories from AdS/CFT superstring duality?*, *Phys. Lett. B* **454** (1999) 49 [[hep-th/9902168](#)];
 P.H. Frampton, *Conformality and gauge coupling unification*, *Phys. Rev. D* **60** (1999) 085004 [[hep-th/9905042](#)];
 P.H. Frampton, *Conformality from field-string duality on abelian orbifolds*, *Phys. Rev. D* **60** (1999) 121901 [[hep-th/9907051](#)];
 P.H. Frampton, R.N. Mohapatra and S. Suh, *Prediction of $\sin^2 \Theta_W$ in a conformal approach to coupling unification*, *Phys. Lett. B* **520** (2001) 331 [[hep-ph/0104211](#)].
- [14] J.M. Maldacena, *The large-N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [[hep-th/9711200](#)];
 S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from non-critical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](#)];
 E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)].