



MIT Open Access Articles

STUDY OF ELECTRON ANTI-NEUTRINOS ASSOCIATED WITH GAMMA-RAY BURSTS USING KamLAND

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation	Asakura, K., A. Gando, Y. Gando, T. Hachiya, S. Hayashida, H. Ikeda, K. Inoue, et al. "STUDY OF ELECTRON ANTI-NEUTRINOS ASSOCIATED WITH GAMMA-RAY BURSTS USING KamLAND." The Astrophysical Journal 806, no. 1 (June 10, 2015): 87. © 2015 The American Astronomical Society
As Published	http://dx.doi.org/10.1088/0004-637x/806/1/87
Publisher	IOP Publishing
Version	Final published version
Citable link	http://hdl.handle.net/1721.1/98306
Terms of Use	Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.

STUDY OF ELECTRON ANTI-NEUTRINOS ASSOCIATED WITH GAMMA-RAY BURSTS USING KamLAND

K. ASAKURA¹, A. GANDO¹, Y. GANDO¹, T. HACHIYA¹, S. HAYASHIDA¹, H. IKEDA¹, K. INOUE^{1,2}, K. ISHIDOSHIO¹, T. ISHIKAWA¹, S. ISHIO¹, M. KOGA^{1,2}, S. MATSUDA¹, T. MITSUI¹, D. MOTOKI¹, K. NAKAMURA^{1,2}, S. OBARA¹, Y. OKI¹, T. OURA¹, I. SHIMIZU¹, Y. SHIRAHATA¹, J. SHIRAI¹, A. SUZUKI¹, H. TACHIBANA¹, K. TAMAÉ¹, K. UESHIMA¹, H. WATANABE¹, B. D. XU¹, H. YOSHIDA^{1,18}, A. KOZLOV², Y. TAKEMOTO², S. YOSHIDA³, K. FUSHIMI⁴, A. PIEPKE^{2,5}, T. I. BANKS^{6,7}, B. E. BERGER^{2,7}, B. K. FUJIKAWA^{2,7}, T. O'DONNELL^{6,7}, J. G. LEARNED⁸, J. MARICIC⁸, M. SAKAI⁸, L. A. WINSLOW⁹, Y. EFREMENKO^{2,10,11}, H. J. KARWOWSKI^{12,15}, D. M. MARKOFF^{12,14}, W. TORNOW^{2,12,13}, J. A. DETWILER¹⁶, S. ENOMOTO^{2,16}, AND M. P. DECOWSKI^{2,17}

THE KAMLAND COLLABORATION

¹ Research Center for Neutrino Science, Tohoku University, Sendai 980-8578, Japan

² Kavli Institute for the Physics and Mathematics of the Universe (WPI), University of Tokyo, Kashiwa 277-8568, Japan

³ Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

⁴ Faculty of Integrated Arts and Science, University of Tokushima, Tokushima, 770-8502, Japan

⁵ Department of Physics and Astronomy, University of Alabama, Tuscaloosa, AL 35487, USA

⁶ Physics Department, University of California, Berkeley, CA 94720, USA

⁷ Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

⁸ Department of Physics and Astronomy, University of Hawaii at Manoa, Honolulu, HI 96822, USA

⁹ Massachusetts Institute of Technology, Cambridge, MA 02139, USA

¹⁰ Department of Physics and Astronomy, University of Tennessee, Knoxville, TN 37996, USA

¹¹ National Research Nuclear University, Moscow, Russia

¹² Triangle Universities Nuclear Laboratory, Durham, NC 27708, USA

¹³ Physics Departments at Duke University, Durham, NC 27705, USA

¹⁴ North Carolina Central University, Durham, NC 27701, USA

¹⁵ The University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

¹⁶ Center for Experimental Nuclear Physics and Astrophysics, University of Washington, Seattle, WA 98195, USA

¹⁷ Nikhef and the University of Amsterdam, Science Park, Amsterdam, The Netherlands

Received 2015 March 9; accepted 2015 April 7; published 2015 June 10

ABSTRACT

We search for electron anti-neutrinos ($\bar{\nu}_e$) from long- and short-duration gamma-ray bursts (GRBs) using data taken by the Kamioka Liquid Scintillator Anti-Neutrino Detector (KamLAND) from 2002 August to 2013 June. No statistically significant excess over the background level is found. We place the tightest upper limits on $\bar{\nu}_e$ fluence from GRBs below 7 MeV and place first constraints on the relation between $\bar{\nu}_e$ luminosity and effective temperature.

Key words: gamma-ray burst: general – neutrinos

1. INTRODUCTION

Gamma-ray bursts (GRBs) are the most luminous phenomena in the universe. The duration of GRBs (Δt_{GRB}) varies from 10 ms to 1000 s, with a roughly bimodal distribution for so-called long GRBs of $\Delta t_{\text{GRB}} \gtrsim 2$ s and short GRBs of $\Delta t_{\text{GRB}} \lesssim 2$ s. The progenitors of most short GRBs are widely thought to be mergers of neutron star–neutron star or black hole–neutron star binaries (Mészáros 2006). A favored model of long GRB progenitors is a catastrophic collapse of a massive star into a black hole (Mészáros 2006). These models are supported by observations of afterglows and identification of host galaxies for short GRBs (Villasenor et al. 2005; Fox et al. 2005) and observations of supernovae associated with long GRBs (Woosley et al. 1999; Hjorth et al. 2003). Both scenarios would result in the formation of a compact rotating black hole with an accretion disk at MeV or higher temperatures, which generates collimated relativistic fireball jets leading to GRBs. Although such a fireball model is promising and attractive, the initial condition and generation mechanism of the fireball jets are still unknown, since it is difficult to observe the optically thick center region of GRBs by electromagnetic waves.

A potential scheme to directly explore the GRB center region is the use of thermal neutrinos and gravitational waves (GWs; Suwa & Murase 2009; Sekiguchi et al. 2011), since they have strong transmissivity. Thermal neutrinos are sensitive to thermodynamic profiles of the accretion disk, and GWs are sensitive to the dynamics of progenitors. Both are complementary observations to probe GRBs. Super-Kamiokande (SK) and Sudbury Neutrino Observatory (SNO) searched for MeV neutrinos related to GRBs and placed constraints on the upper fluence limits (Fukuda et al. 2002; Aharmim et al. 2014). Others placed limits on high-energy neutrinos produced in the fireball jets (Achterberg et al. 2008; Thrane et al. 2009; Abbasi et al. 2011; Avrorin et al. 2011; Vieregge et al. 2011; Adrián-Martínez et al. 2013). GWs from GRBs were studied by a GW detector network (Aasi et al. 2014).

In this paper, we present a study of electron anti-neutrinos ($\bar{\nu}_e$) of a few tens of MeV energy produced by thermal processes from the GRB center region, especially the accretion disk (Nagataki & Kohri 2002; Caballero et al. 2009; Sekiguchi & Shibata 2011), with the Kamioka Liquid Scintillator Anti-Neutrino Detector (KamLAND). We constrain the relation between the $\bar{\nu}_e$'s luminosity (L) and effective temperature (T), as well as $\bar{\nu}_e$ fluence, for the first time. The L – T relationship can be used to directly compare with theoretical predictions. These limits and constraints are established using redshift-

¹⁸ Current address: Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan




Figure 1. Schematic diagram of the KamLAND detector. The shaded region in the liquid scintillator indicates the volume for the $\bar{\nu}_e$ analysis after the IB installation.

measured GRBs. We adopt the standard Λ CDM cosmology with $\Omega_m = 0.315$, $\Omega_\Lambda = 0.685$, and $H_0 = 67.3 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (Ade et al. 2014) throughout this paper.

2. KamLAND DETECTOR

The KamLAND detector is located $\sim 1 \text{ km}$ under the peak of Mt. Ikenoyama ($36^\circ 42' \text{ N}$, $137^\circ 31' \text{ E}$) near Kamioka, Japan. The 2700 m water equivalent of vertical rock overburden reduces the cosmic-ray muon flux by almost five orders of magnitude. A schematic diagram of KamLAND is shown in Figure 1. The primary target volume consists of 1 kton of ultra-pure liquid scintillator (LS) contained in a 13 m diameter spherical balloon made of $135 \mu\text{m}$ thick transparent nylon ethylene vinyl alcohol copolymer composite film. The LS consists of 80% dodecane and 20% pseudocumene (1, 2, 4-trimethylbenzene) by volume, and $1.36 \pm 0.03 \text{ g l}^{-1}$ of the fluor PPO (2, 5-diphenyloxazole). A buffer comprising 57% isoparaffin and 43% dodecane oils by volume, which fills the region between the balloon and the surrounding 18 m diameter spherical stainless-steel outer vessel, shields the LS from external radiation. The specific gravity of the buffer oil is adjusted to be 0.04% lower than that of the LS. An array of photomultiplier tubes (PMTs)—1325 specially developed fast PMTs masked to 17-inch diameter and 554 older 20-inch diameter PMTs reused from the Kamiokande experiment (Kume et al. 1983)—are mounted on the inner surface of the outer vessel, providing 34% photocathode coverage. This inner detector is shielded by a 3.2 kton water-Cherenkov veto detector.

KamLAND uses the inverse beta-decay reaction to detect $\bar{\nu}_e$:

$$\bar{\nu}_e + p \rightarrow e^+ + n. \quad (1)$$

This process has a delayed-coincidence (DC) event-pair signature that offers powerful background suppression. The energy deposited by the positron, which generates the DC pair's prompt event, is the sum of the e^+ kinetic energy and annihilation γ energies, $E_p (\equiv T_{e^+} + 2m_e)$, and related to the incident $\bar{\nu}_e$ energy by $E_{\bar{\nu}_e} = \langle E_{e^+} \rangle + \delta + E_\nu^{\text{CM}} E_e^{\text{CM}} / m_p$, where

E_ν^{CM} and E_e^{CM} are neutrino and electron energy in the center-of-mass frame, respectively, and $\delta = (m_n^2 - m_p^2 - m_e^2) / 2m_p$ (Strumia & Vissani 2003). In the low-energy ($E_{\bar{\nu}_e} < 20 \text{ MeV}$) range, we can approximate the above relation by $E_{\bar{\nu}_e} = E_p + \delta E$, where $\delta E = 0.78 \text{ MeV}$. We use this approximation also above 20 MeV and comment on the associated uncertainty later. The delayed event in the DC pair is generated by a 2.2 (4.9) MeV γ -ray produced when the neutron captures on a proton (^{12}C). The mean neutron capture time is $207.5 \pm 2.8 \mu\text{s}$ (Abe et al. 2010). The angular distribution of the positron emission is nearly isotropic. Unlike in a water-Cherenkov detector, the scintillation light is also isotropic. As a result, the positron signal does not provide the incoming $\bar{\nu}_e$ source direction. Due to the extremely low cross section of $\bar{\nu}_e$, the Earth does not shadow MeV-energy extraterrestrial $\bar{\nu}_e$. The detector therefore has isotropic sensitivity to GRBs.

The event energy and vertex reconstruction are based on the timing and charge distributions of scintillation photons recorded by the PMTs. The reconstruction algorithms are calibrated with on-axis and off-axis radioactive sources deployed from a glove box installed at the top of the detector. The radioactive sources are ^{60}Co , ^{68}Ge , ^{203}Hg , ^{65}Zn , ^{241}Am , ^9Be , ^{137}Cs , and ^{210}Po , ^{13}C , providing energy and vertex calibration (Berger et al. 2009; Banks et al. 2015). The overall vertex reconstruction resolution is $\sim 12 \text{ cm} / \sqrt{E (\text{MeV})}$, and energy resolution is $6.4\% / \sqrt{E (\text{MeV})}$. The energy reconstruction of positrons with $E_p > 7.5 \text{ MeV}$ (i.e., $E_{\bar{\nu}_e} > 8.3 \text{ MeV}$) is verified by using tagged ^{12}B β^- -decays generated via muon spallation (Abe et al. 2010).

In 2011 September, the KamLAND-Zen double-beta ($\beta\beta$) decay search experiment was launched (Gando et al. 2012). This experiment makes use of KamLAND's extremely low background. The KamLAND detector was modified to include a $\beta\beta$ source, 13 tons of Xe-loaded LS (Xe-LS) contained in a 3.08 m diameter inner balloon (IB), at the center of the detector.

3. EVENT SELECTION

3.1. KamLAND DC Events

In this analysis, we use KamLAND data collected from 2002 August 3 to 2013 June 4. During the majority of this period, KamLAND was measuring $\bar{\nu}_e$ from nuclear power plants with a spectrum up to about 8 MeV (Gando et al. 2011a, 2013) and geological $\bar{\nu}_e$ from the Earth's deep interior (Araki et al. 2005; Gando et al. 2011b, 2013). Following the Fukushima reactor accident in 2011 March, all Japanese reactors were subject to a protracted shutdown. The data set is divided into two periods. Period I refers to data that were taken until the IB installation in 2011 September. Period II refers to the data taken after the IB installation, which mostly coincided with the low reactor $\bar{\nu}_e$ flux.

In Period I, we search only for $\bar{\nu}_e$ events with $E_{\text{low}}^{\text{I}} (= 7.5 \text{ MeV}) \leq E_p \leq 100 \text{ MeV}$, which corresponds to the energy range of interest for GRBs with almost zero contamination from the reactor $\bar{\nu}_e$ flux. During Period II, the reactor signal is minimal, allowing a reduction of the energy threshold to $E_{\text{low}}^{\text{II}} = 0.9 \text{ MeV}$.

For the DC event pair selection, we apply the following series of selection cuts: the prompt energy is required to be

$E_{\text{low}}^k \leq E_p \leq 100$ MeV in Period k , and the delayed energy to be $1.8 \text{ MeV} \leq E_d \leq 2.6$ MeV for neutron capture on protons or $4.4 \text{ MeV} \leq E_d \leq 5.6$ MeV for neutron capture on ^{12}C , a fiducial volume cut of $R \leq 6$ m from the center of the balloon on both prompt and delayed events, a spatial correlation cut of $\Delta R \leq 1.6$ m, and a time separation cut of $0.5 \mu\text{s} \leq \Delta T \leq 1.0$ ms. Spallation cuts were used to reduce backgrounds from long-lived isotopes, e.g., ^9Li ($\tau = 257$ ms and $Q = 13.6$ MeV), that are generated by cosmic muons passing through the scintillator. In Period II, we have to use an additional spatial cut for delayed events to avoid backgrounds from the IB and its support material as shown in Figure 1 (Gando et al. 2013), and a second-level cut using a likelihood discriminator to reduce accidental backgrounds in the low-energy region (Gando et al. 2011a). The selection efficiency (ϵ_s^k) is evaluated from Monte Carlo simulation separately for Period I ($k = \text{I}$) and Period II ($k = \text{II}$) due to these additional cuts. Note that ϵ_s^{II} depends on E_p because of the energy-dependent second-level cut. The number of target protons in $R \leq 6$ m is estimated to be $N_T = (5.98 \pm 0.12) \times 10^{31}$.

The total livetime during Period I was 6.91 yr, and 55 DC events were observed during this period. In Period II, KamLAND found 88 DC events with 1.2 yr livetime. The livetime is defined as the integrated period of time that the detector was sensitive to $\bar{\nu}_e$ and includes corrections for calibration periods, detector maintenance, daily run switch, etc. The event rates are 9.1×10^{-4} and 8.4×10^{-3} events per hour in Periods I and II, respectively.

3.2. GRB Events

We use GRB events observed by one or more of *SWIFT*, *HETE-2*, *Ulysses*, *INTEGRAL*, *AGILE*, *MAXI*, and *Fermi* based on the Gamma-ray Coordinates Network.¹⁹ Initial selection criteria are the requirement that the GRB be in the time period between 2002 August 3 and 2013 June 4 and the existence of redshift and GRB-duration time measurements. At this stage, 256 long GRB and 21 short GRB events are left. Subsequently, all the KamLAND runs²⁰ that include GRB events must have passed basic quality criteria (e.g., not a calibration run and stable operation). This leaves 175 long GRBs and 17 short GRBs in Period I. Period II contains 38 long GRBs and one short GRB. One can see our GRB list online.²¹

4. DATA ANALYSIS

The average number of DC events and GRB events per 3 months is shown in Figure 2. In this figure, one can see a ‘‘step’’ before and after the launch of the *SWIFT* satellite (2004 November) for GRB events. In contrast, there is no time dependence of the DC event rate during each period. We therefore decided to analyze all of the KamLAND data regardless of the GRB event rate.

4.1. Coincidence Analysis

We conduct a time-coincidence analysis between the redshift-measured GRB samples and the KamLAND DC events for long and short GRBs. The coincidence search time




Figure 2. Number of long and short GRBs per 3 months (upper panel) and DC events per 3 months (bottom panel). The number of GRBs significantly increased due to the *SWIFT* satellite after 2004 December. During Period I, the DC event rate is within the statistical uncertainty. Period II started in 2011 September and allowed for a lower energy threshold, increasing the number of DC events.

window between a GRB event and a KamLAND DC event is defined as $-t_p + T_{\text{GRB}} < T_{\text{DC}} < T_{\text{GRB}} + \Delta t_{\text{GRB}} + t_p + t_f(z)$, where T_{DC} and T_{GRB} are the absolute times of the KamLAND DC and GRB events, respectively. Δt_{GRB} is the measured GRB duration time, t_p is 150 s corresponding to a model-dependent but reasonable time difference between the thermal neutrino production and the GRB photon production (Y. Sekiguchi & Y. Suwa 2012, private communication; K. Toma 2014, private communication), and $t_f(z)$ is the relativistic flight-time delay of MeV neutrinos due to non-zero neutrino mass (Choubey & King 2003; Li et al. 2005):

$$t_f(z) = \frac{1}{2} \frac{m_{\bar{\nu}_e}^2}{E_{\bar{\nu}_e}^2} \int_0^z \frac{dz'}{(1+z')^2 H_0 \sqrt{\Omega_\Lambda + (1+z')^3 \Omega_m}}, \quad (2)$$

with the assumption of $m_{\bar{\nu}_e} = m_{\text{heaviest}} = 87.2$ meV from $\sum m_\nu \leq 0.23$ eV (Ade et al. 2014) and $E_{\bar{\nu}_e} \geq 8.3$ MeV in Period I and $E_{\bar{\nu}_e} \geq 1.8$ MeV in Period II. All parameters of the time window are fixed before the coincidence search. The total window length for the long GRBs is 25.2 hr (18.3 hr in Period I and 6.82 hr in Period II). The short GRBs sum to a total of 1.45 hr of on-time window (1.33 hr in Period I and 0.11 hr in Period II).

No coincidence DC events were found in the above time window for both long and short GRBs. We estimate the expected accidental coincidence of DC events to be 7.4×10^{-2} and 2.2×10^{-3} for long and short GRBs, respectively. For long GRBs, the background spectrum is shown in Figure 3 with several expected spectra from our 90% upper limits (see 4.3). In the absence of a signal, the Feldman–Cousins (FC) upper limits on the DC events are $N_{90} = 2.365$ and 2.435 with 90% confidence level (CL) for long and short GRBs, respectively.

If we use a much longer, exotic time window, e.g., $t_p = 6$ hr, four coincidence DC events are found for long GRBs.

¹⁹ <http://gc.n.gsf.nasa.gov/>

²⁰ KamLAND data taking is stopped and restarted every day to ensure smooth data taking. The length of a KamLAND run is typically 24 hr long.

²¹ See <http://www.awa.tohoku.ac.jp/KamLAND/GRB/2015>.




Figure 3. Combined background spectrum of Periods I and II (solid red). The integral is 7.4×10^{-2} events. The dashed curves provide our 90% upper limits under the assumption of a Fermi–Dirac distribution at temperature T of $\bar{\nu}_e$. Each dashed curve from the left to the right corresponds to $T = 5, 10, 15$ MeV.

However, the expected accidental coincidence of DC events is 3.4. There is therefore no statistical evidence for the detection of $\bar{\nu}_e$ from long GRBs.

4.2. Fluence Upper Limits

There is no established neutrino production model for GRBs. We translate our FC limits to model-independent upper limits on $\bar{\nu}_e$ fluence, $\Psi(E_{\bar{\nu}_e})$, at the detector using a Green’s function, which represents the upper limits on monoenergetic neutrinos at that specific energy. We use the same methodology to estimate $\Psi(E_{\bar{\nu}_e})$ as SK (Fukuda et al. 2002) and SNO (Aharmim et al. 2014):

$$\Psi(E_{\bar{\nu}_e}) = \frac{N_{90}}{\sum_k N_{\text{GRB}}^k I_k(E_{\bar{\nu}_e})}, \quad (3)$$

where N_{GRB}^k is the number of GRBs and $I_k(E_{\bar{\nu}_e})$ is the effective number of DC events per one GRB with a monoenergetic spectrum in the period k :

$$I_k(E_{\bar{\nu}_e}) = N_T \int_{E_{\text{low}}^k}^{100 \text{ MeV}} \epsilon_t \epsilon_s^k(E_p^{\text{vis}}) \sigma_{\text{IBD}}(E_{\bar{\nu}_e}) \delta(E_p^{\text{exp}} + \delta E - E_{\bar{\nu}_e}) R(E_p^{\text{exp}}, E_p^{\text{vis}}) dE_p^{\text{exp}} dE_p^{\text{vis}}, \quad (4)$$

and

$$R(E_p^{\text{exp}}, E_p^{\text{vis}}) = \frac{1}{\sqrt{2\pi} \sigma(E_p^{\text{exp}})} \exp\left[-\frac{(E_p^{\text{exp}} - E_p^{\text{vis}})^2}{2\sigma^2(E_p^{\text{exp}})}\right]. \quad (5)$$

ϵ_t is the mean livetime-to-runtime ratio,²² and E_p^{exp} and E_p^{vis} are the expected and measured prompt energies, respectively. $\sigma_{\text{IBD}}(E)$ is the differential cross section of the inverse beta decay. $\sigma(E)$ corresponds to the energy resolution of $6.4\%/\sqrt{E(\text{MeV})}$.

The 90% CL upper limits on $\Psi(E_{\bar{\nu}_e})$ from KamLAND are shown for both long and short GRBs together in Figure 4 with the results from SK (Fukuda et al. 2002) and SNO (Aharmim




Figure 4. Fluence upper limits on $\bar{\nu}_e$ from GRBs as a function of neutrino energy. Results from SK and SNO are also presented for comparison. SK is expected to have poorer sensitivity at low energies due to the detection threshold. Below 7 MeV, KamLAND establishes the tightest limits on neutrino fluence. The slight distortion around 3–4 MeV is from the energy dependence of the selection efficiency (ϵ_s^{II}).

et al. 2014). Note that the results from SK and SNO treated long and short GRBs as the same. Below 7 MeV, our analysis provides the best limits so far.

4.3. Constraint on Luminosity and Effective Temperature (L – T)

N_{90} can be translated to constrain the $\bar{\nu}_e$ ’s luminosity (L) and effective temperature (T) in the accretion disk using the assumption that the $\bar{\nu}_e$ flux follows the Fermi–Dirac distribution described:

$$\psi(E_{\bar{\nu}_e}, T, L) = \frac{120}{7\pi^4} \frac{L}{T^4} \frac{E_{\bar{\nu}_e}^2}{\exp(E_{\bar{\nu}_e}/T) + 1}. \quad (6)$$

The expected total flux at the detector is in Period k ,

$$\Psi^k(E_{\bar{\nu}_e}, T, L) = \sum_{i \in k} \frac{1 + z_i}{4\pi d_i^2} \psi((1 + z_i)E_{\bar{\nu}_e}, T, L), \quad (7)$$

where z_i and d_i are the redshift and luminosity distance of the i th GRB, respectively. The luminosity and effective temperature upper limits ($T_{\text{up}}, L_{\text{up}}$) are then connected to N_{90} :

$$N_{90} = \sum_k \int_{E_{\text{low}}^k}^{100 \text{ MeV}} I'_k(T_{\text{up}}, L_{\text{up}}, E_p^{\text{vis}}) dE_p^{\text{vis}}, \quad (8)$$

where I'_k is the visible spectrum of the DC events,

$$I'_k(T_{\text{up}}, L_{\text{up}}, E_p^{\text{vis}}) = \int_{E_{\text{low}}^k}^{100 \text{ MeV}} N_T \epsilon_t \epsilon_s^k(E_p^{\text{vis}}) \sigma_{\text{IBD}}(E_{\bar{\nu}_e}) \Psi^k \times (E_{\bar{\nu}_e}, T_{\text{up}}, L_{\text{up}}) R(E_p^{\text{exp}}, E_p^{\text{vis}}) dE_p^{\text{exp}}. \quad (9)$$

With the assumption of $E_{\bar{\nu}_e} = E_p + \delta E$, the results obtained from KamLAND are shown in Figure 5. The upper limit spectra ($\sum I'_k(T_{\text{up}}, L_{\text{up}}, E_p^{\text{vis}})$) with $T_{\text{up}} = 5, 10, 15$ MeV are shown in Figure 3.

The limits are six orders of magnitude higher than the supernova $\bar{\nu}_e$ luminosity and several orders of magnitude higher than theoretical studies predict. Nagataki & Kohri (2002)

²² Runtime is the total time of data taking.




Figure 5. KamLAND constraint on the luminosity (L) and effective temperature (T) relation in the GRB accretion disk, together with several theoretical calculations (Nagataki & Kohri 2002; Caballero et al. 2009; Sekiguchi & Shibata 2011).

analytically show that a collapsar emits $\bar{\nu}_e$ with $L = 10^{52}$ erg and $T = 5$ MeV in a total accretion mass of $30 M_\odot$, an initial mass of $3 M_\odot$, and a mass accretion rate of $0.1 M_\odot \text{ s}^{-1}$. Caballero et al. (2009) numerically predict $L = 3.5 \times 10^{52}$ erg for 0.15 s with $T = 7.5$ MeV for black hole–neutron star mergers. Here we assumed that the averaged $\bar{\nu}_e$ energy corresponds to $3.15 T$. Recently, Sekiguchi presented $\dot{L} = 1.5\text{--}3 \times 10^{52}$ erg s^{-1} for 2–3 s with an averaged $\bar{\nu}_e$ energy of 20–30 MeV for a merger of binary neutron stars using state-of-the-art numerical simulations (Sekiguchi & Shibata 2011).

Finally, we comment about the approximation, $E_{\bar{\nu}_e} = E_p^{\text{exp}} + \delta E$. Above 20 MeV, this approximation is not suitable. In addition, the effect of the recoiling neutron (\bar{E}_n) to E_p is no longer negligible. This effect adds a substantial energy bias, $\sim 10\%$, but the uncertainty of L_{up} is much smaller than 10%. The amount of the error has no impact on our result and discussion.

5. SUMMARY

We find no evidence for $\bar{\nu}_e$ associated with our sample of GRBs in KamLAND. We placed the lowest observational bound on the $\bar{\nu}_e$ fluence below 7 MeV. The relation of L – T , which characterizes the GRB accretion disk, is constrained. The obtained upper limits are significantly higher than several theoretical predictions (Nagataki & Kohri 2002; Sekiguchi &

Shibata 2011). However, our result is the first constraint that can be directly compared to theoretical studies.

We are indebted to the observers of GRBs for providing us with data. KamLAND is supported by MEXT KAKENHI Grant Numbers 26104002 and 26104007; the World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan; and under the U.S. Department of Energy (DOE) grants No. DE-FG03-00ER41138, DE-AC02-05CH11231, and DE-FG02-01ER41166, as well as other DOE grants to individual institutions, and Stichting Fundamenteel Onderzoek der Materie (FOM) in the Netherlands. The Kamioka Mining and Smelting Company has provided service for activities in the mine. We thank the support of NII for SINET4.

REFERENCES

- Aasi, J., Abbott, B. P., Abbott, R., et al. 2014, *PhRvD*, **89**, 122004
 Abbasi, R., Abdou, Y., Abu-Zayyad, T., et al. 2011, *PhRvL*, **106**, 141101
 Abe, S., Enomoto, S., Furuno, K., et al. 2010, *PhRvC*, **81**, 025807
 Achterberg, A., Ackermann, M., Adams, J., et al. 2008, *ApJ*, **674**, 357
 Ade, P. A. R., Aghanim, N., Armitage-Caplan, C., et al. 2014, *A&A*, **571**, A16
 Adrián-Martínez, S., Samarai, I. A., Albert, A., et al. 2013, *JCAP*, **2013**, 006
 Aharmim, B., Ahmed, S., Anthony, A., et al. 2014, *Aph*, **55**, 1
 Araki, T., Enomoto, S., Furuno, K., et al. 2005, *Natur*, **436**, 499
 Avrorin, A., Aynutdinov, V., Belolaptikov, I., et al. 2011, *AstL*, **37**, 692
 Banks, T., Freedman, S., Wallig, J., et al. 2015, *NIMPA*, **769**, 88
 Berger, B. E., Busenitz, J., Classen, T., et al. 2009, *JInst*, **4**, P04017
 Caballero, O. L., McLaughlin, G. C., & Surman, R. 2009, *PhRvD*, **80**, 123004
 Choubey, S., & King, S. F. 2003, *PhRvD*, **67**, 073005
 Fox, D. B., Frail, D. A., Price, P. A., et al. 2005, *Natur*, **437**
 Fukuda, S., Fukuda, Y., Ishitsuka, M., et al. 2002, *ApJ*, **578**, 317
 Gando, A., Gando, Y., Hanakago, H., et al. 2012, *PhRvC*, **85**, 045504
 Gando, A., Gando, Y., Hanakago, H., et al. 2013, *PhRvD*, **88**, 033001
 Gando, A., Gando, Y., Ichimura, K., et al. 2011a, *PhRvD*, **83**, 052002
 Gando, A., Gando, Y., Ichimura, K., et al. 2011b, *NatGe*, **4**, 647
 Hjorth, J., Sollerman, J., Mller, P., et al. 2003, *Natur*, **423**, 847
 Kume, H., Sawaki, S., Ito, M., et al. 1983, *NuclIM*, **205**, 443
 Li, H., Dai, Z., & Zhang, X. 2005, *PhRvD*, **71**, 113003
 Mészáros, P. 2006, *RPPh*, **69**, 2259
 Nagataki, S., & Kohri, K. 2002, *PThPh*, **108**, 789
 Sekiguchi, Y., Kiuchi, K., Kyutoku, K., & Shibata, M. 2011, *PhRvL*, **107**, 051102
 Sekiguchi, Y., & Shibata, M. 2011, *ApJ*, **737**, 6
 Strumia, A., & Vissani, F. 2003, *PhLB*, **564**, 42
 Suwa, Y., & Murase, K. 2009, *PhRvD*, **80**, 123008
 Thrane, E., Abe, K., Hayato, Y., et al. 2009, *ApJ*, **697**, 730
 Viereg, A. G., Palladino, K., Allison, P., et al. 2011, *ApJ*, **736**, 50
 Villasenor, J. S., Lamb, D. Q., Ricker, G. R., et al. 2005, *Natur*, **437**, 855
 Woosley, S. E., Eastman, R. G., & Schmidt, B. P. 1999, *ApJ*, **516**, 788