



University of Kentucky  
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII  
International Rangeland Congress

## An Investigation on Forage Quality of Three Grasses (*Aeluropus lagopoides*, *A. littoralis*, *Puccinellia distans*) in Saline and Alkaline Habitats of the Caspian Sea

Masoume Amirkhani  
*University of Agricultural Sciences, Iran*

Mansour Mesdaghi  
*University of Agricultural Sciences, Iran*

E. Bagheri Rad  
*University of Agricultural Sciences, Iran*

A. R. Mahmoodi  
*University of Agricultural Sciences, Iran*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/15-3/26>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## An investigation on forage quality of three grasses (*Aeluropus lagopoides*, *A. littoralis*, *Puccinellia distans*) in saline and alkaline habitats of the Caspian Sea

M. Amirkhani, M. Mesdaghi, E. Bagheri Rad, and A. R. Mahmoodi


Department of Range Management, College of Natural Resources, University of Agricultural Sciences, Gorgan, Iran. E-mail: maasome\_amirkhani@yahoo.com

**Key words:** *Aeluropus lagopoides*, *A. littoralis*, *Puccinellia distans*, forage quality

**Introduction** *Aeluropus lagopoides*, *A. littoralis* and *Puccinellia distans* are perennial grasses which are adapted to saline and alkaline habitats of sea shores, salt flats and marginal area of the rivers around the Caspian Sea. These species have significant roles in providing good forage quality for domestic sheep and goats on local rangelands. These grasses occur at altitude of 10 m with annual rainfall of 180 mm. The objectives of this study were to determine forage quality of these species in two phenological stages in saline and alkaline habitats.

**Material and methods** The foliage of five plants of each species was harvested at two stages, vegetative growth (VG) and seed ripening (SR), and then analyzed for nitrogen (N), acid detergent fiber (ADF), and crude fiber (CF). Crude protein (CP), dry matter digestibility (DMD), and metabolizable energy (ME) were then calculated using equations of:  $\%N=0.16CP$ ,  $\%DMD=83.58-0.824 \cdot ADF+2.626\%N$ , and  $ME=0.17\%DMD-2$  (Standing Committee on Agriculture, 1990) and analyzed statistically by using ANOVA.

**Results** The chemical compositions of the three species at two stages are shown in Figure 1.



**Figure 1** The chemical compositions of these three grasses at two phenological stages. Harvesting was accomplished at stage of vegetative growth for *A. lagopoides* and *A. littoralis* in late Feb to late May and at stage of seed ripening in early June to late July. Harvesting of *Puccinellia distans* was at stage of vegetative growth in late March to late April, and for seed ripening was at late June to late of July.

There were significant differences in chemical compositions of three species ( $P < 0.05$ ). With advances in maturity, CP decreased, ADF and CF increased. In both phenological stages for all factors *A. lagopoides* was better than *A. littoralis* ( $P < 0.05$ ). In vegetative growth, *P. distans* had the same quality as *A. lagopoides*, but in seed ripening had the same quality as *A. littoralis* ( $P > 0.05$ ). In general, *A. lagopoides* has better quality than the two other species.

**Conclusions** These species have high potential in relation to forage quality on rangelands in the study area. Further research is needed to understand the requirements for seeding these species in the areas.

### References

- Abarsaji, G. A., 2000. Identification and investigation on some of ecophysiological characteristics of *Aeluropus* spp. in saline and alkaline rangelands in the north of Gorgan. *Pajouhesh-Va-Sazandegi* 46:21-25.
- Akhani, H., M. Ghorbanli., 1993. A contribution to the halophytic vegetation and flora of Iran. Towards the rational use of high salinity tolerant plants. 1:35-44.
- Standing Committee on Agriculture., 1990. Feeding standards for Australian livestock, Ruminants, CSIRO, Australia.