



## **PHYSICAL CHEMISTRY 2014**

12<sup>th</sup> International Conference  
on Fundamental and Applied Aspects of  
Physical Chemistry

---

The Conference is dedicated to the  
25. Anniversary of the Society of Physical Chemists of Serbia

---

September 22-26, 2014  
Belgrade, Serbia

**ISBN 978-86-82475-30-9**

**Title:** PHYSICAL CHEMISTRY 2014 (Proceedings)

**Editors:** Ž. Čupić and S. Anić

**Published by:** Society of Physical Chemists of Serbia, Studenski trg 12-16, 11158, Belgrade, Serbia

**Publisher:** Society of Physical Chemists of Serbia

**For Publisher:** S. Anić, President of Society of Physical Chemists of Serbia

**Printed by:** “Jovan” Printing and Publishing Company; 200 Copies;

**Number of pages:** 6+ 441; **Format:** B5; Printing finished in September 2014.

**Text and Layout:** “Jovan”

Neither this book nor any part may be reproduced or transmitted in any form or by any means, including photocopying, or by any information storage and retrieval system, without permission in writing from the publisher.

*200 - Copy printing*

PHYSICAL CHEMISTRY 2014

*12th International Conference on  
Fundamental and Applied Aspects of  
Physical Chemistry*

*Organized by  
The Society of Physical Chemists of  
Serbia*

*in co-operation  
with\_*

*Institute of Catalysis Bulgarian Academy of Sciences*

*Boreskov Institute of Catalysis of Siberian Branch of the Russian Academy  
of Sciences*

*Faculty of Physical Chemistry, University of Belgrade, Serbia*

*Institute of Chemistry Technology and Metallurgy,  
University of Belgrade, Serbia*

*Vinča Institute, University of Belgrade, Serbia*

*Institute of General and Physical Chemistry, Serbia*

*Faculty of Pharmacy, University of Belgrade, Serbia*

### **International Organizing Committee**

*Chairman:* S. Anić (Serbia)  
*Vice-chairman:* M. Gabrovska (Bulgaria)  
V. A. Sadykov (Russia)

*Members:*

N. Cyjetičanin (Serbia), S. N. Blagojević (Serbia), M. Daković (Serbia), T. Grozdić (Serbia), D. Jovanović (Serbia), M. Kuzmanović (Serbia), D. Marković (Serbia), J. Marković-Dimitrić (Serbia), B. Milosavljević (USA), M. Mojović (Serbia), N. Ostrovski (Serbia), I. Pašti (Serbia), M. Petković (Serbia), A. Popović-Bjelić (Serbia), B. Simonović (Serbia), D. Stanisavljev (Serbia), B. Šljukić (Serbia), N. Vukelić (Serbia), V. Vukojević (Sweden)

### **International Scientific Committee**

*Chairman:* Ž. Čupić (Serbia)  
*Vice-chairmans:* V. N. Parmon (Russia)  
S. Rakovsky (Bulgaria)

*Members:*

B. Adnađević (Serbia), S. Anić (Serbia), A. Antić-Jovanović (Serbia), G. Bačić (Serbia), A. Kellarakis (Greece), R. Cervellati (Italy), V. Dondur (Serbia), Ivan Gutman (Serbia), S. D. Furrow (USA), K. Hedrih (Serbia), M. Jeremić (Serbia), A. V. Knyazev (Russia), Lj. Kolar-Anić (Serbia), V. Kuntić (Serbia), Z. Marković (Serbia), S. Mentus (Serbia), Š. Miljanić (Serbia), M. Perić (Serbia), M. Plavšić (Serbia), D. M. F. Santos (Portugal), G. Schmitz (Belgium), I. Schreiber (Czech), P. Sevcik (Slovakia), B. C. Simionescu (Romania), N. Stepanov (Russia), D. Todorović (Serbia), M. Trtica (Serbia), V. Vasić (Serbia), D. Veselinović (Serbia)

### **Local Executive Committee**

*Chairman:* S. Blagojević  
*Vice-chairmans:* A. Ivanović-Šašić

*Members:*

P. Banković, N. Bošnjaković, J. Dostanić, A. Đerić, A. Ignjatović, A. Jović, N. Jović-Jovičić, D. Lončarević, J. Krstić, J. Maksimović, V. Marković, M. Milenković, S. Maćešić, V. Marković, B. Nedić, N. Potkonjak, D. Ranković, M. Stević, M. Žunić, M. Ristić,

## SORPTION OF PERTECHNETATE ONTO ORDERED MESOPOROUS CARBON

Đ. Petrović<sup>1</sup>, A. Đukić<sup>2</sup>, K. Kumrić<sup>3</sup>, I. Milanović<sup>2</sup>, Ž. Rašković-Lovre<sup>2</sup>  
and Lj. Matović<sup>2</sup>

<sup>1</sup>Laboratory for Radioisotopes, <sup>2</sup>Laboratory of Materials Sciences,  
<sup>3</sup>Laboratory of Physics, Vinča Institute of Nuclear Sciences, University of  
Belgrade, P.O. Box 522, Belgrade, Serbia. (djpetrovic@vin.bg.ac.rs)

### ABSTRACT

Ordered mesoporous carbon (OMC) was used as an adsorbent for the removal of pertechnetate ( $\text{TcO}_4^-$ ) anion. Batch experiments were performed to study the effects of pH and ionic strength of the solution. The adsorption of  $\text{TcO}_4^-$  is almost pH-independent in very wide pH region (from 4.0 to 10.0). Maximum distribution coefficient,  $K_d$ , was obtained at pH 2.0. Adsorption of  $\text{TcO}_4^-$  was not affected by the ionic strength changes.

### INTRODUCTION

The presence of the long-lived radionuclide  $^{99}\text{Tc}$  ( $t_{1/2} = 2.1 \times 10^5$  y) in the environment comes from anthropogenic nuclear activities such as nuclear power plants, nuclear weapons testing facilities, reprocessing of spent nuclear fuel, nuclear accidents and medical application of  $^{99\text{m}}\text{Tc}$ . The predominant chemical form of the radionuclide  $^{99}\text{Tc}$  is the highly soluble and environmentally mobile  $\text{TcO}_4^-$  anion, that easily penetrates the ecosystems [1]. Because  $^{99}\text{Tc}$  is hazardous to the environment, its effective removal from contaminated groundwater and sediments is considered a primary task.

Removal of  $\text{TcO}_4^-$  by activated carbons is one of the most efficient processes, because of large specific surface area, porous structure and variety of surface functional groups of activated carbons. Ordered mesoporous carbon belongs to a family of porous materials which attracted great attention in recent years due to its unique features such as high surface area and uniform, large and tunable pore channels. These features make OMC attractive for use in the adsorption processes.

The aim of the present paper was to investigate the influence of the solution pH and ionic strength on the  $\text{TcO}_4^-$  sorption by the OMC.

## EXPERIMENTAL

The OMC was synthesized under acidic conditions as previously described by Momčilović et al [2]. Radioactive technetium (in the form of pertechnetate) was obtained from a  $^{99}\text{Mo}/^{99\text{m}}\text{Tc}$  generator and its radioactive concentration was determined by a Capintec CRC-15 dose calibrator (Capintec, Inc. Ramsey, USA). The resolution of the instrument is 0.001 MBq, i.e. 0.01  $\mu\text{Ci}$ . The measurements of  $^{99\text{m}}\text{Tc}$  activity in aliquots were made using vial geometry.


Batch experiments were performed at room temperature by mixing 0.04 g of well powdered OMC adsorbent and 10  $\text{cm}^3$  of working  $^{99\text{m}}\text{TcO}_4^-$  solution in closed glass vials. Experimental set-up is presented in Fig. 1. The total  $^{99\text{m}}\text{TcO}_4^-$  radioactive concentration in the working solution was 1.4 MBq  $\text{cm}^{-3}$  (37.8 mCi  $\text{cm}^{-3}$ ). The initial pH was adjusted by using either 0.1 M HCl or NaOH. The samples were shaken on a laboratory shaker for 60 min at a stirring speed of 200 rpm. After that, the liquid phases were separated from the solid phases by filtration through a microporous membrane filter (0.45  $\mu\text{m}$ ). The residual radioactive concentration of  $^{99\text{m}}\text{TcO}_4^-$  in each aliquot was determined with a Capintec CRC-15 dose calibrator.

The effects of solution pH (2.0-12.0) and ionic strength (0.01 and 0.1 M NaCl) were investigated with respect to the removal efficiency of  $\text{TcO}_4^-$ . All experiments were carried out in duplicate, and the data obtained were used for analysis.

The removal efficiency ( $E$ , %) and the distribution coefficients ( $K_d$ ,  $\text{cm}^3 \text{g}^{-1}$ ) of  $^{99\text{m}}\text{TcO}_4^-$  were calculated using the following equations:

$$E = \frac{(A_i - A_f)}{A_i} \times 100 \quad (1)$$

$$K_d = \frac{(A_i - A_f)}{A_f} \times \frac{V}{m} \quad (2)$$



**Figure 1.** Experimental set-up for the sorption of  $\text{TcO}_4^-$  onto the OMC.


where  $A_i$  and  $A_f$  are the initial and the final  $^{99m}\text{TcO}_4^-$  radioactive concentrations in the liquid phase ( $\text{MBq cm}^{-3}$ ),  $V$  the volume of the solution ( $\text{cm}^3$ ) and  $m$  the mass of adsorbent (g).

## RESULTS AND DISCUSSION

The effect of the solution pH and ionic strength,  $I$ , on the adsorption of  $\text{TcO}_4^-$  using the OMC was investigated at varying pH values from 2.0 to 12.0. The obtained results are presented in Fig. 2. It is evident that the adsorption of  $\text{TcO}_4^-$  was pH-dependent, with maximum removal efficiency of 99% at the pH value 2.0. Then, the removal efficiency slightly decreases and reaches the value of 93% at the pH values between 4.0 and 10.0. At the pH values beyond 10.0, the adsorption of  $\text{TcO}_4^-$  decreased sharply and the removal efficiency of  $\text{TcO}_4^-$  fell to 72%. The distribution coefficients, presented in the insert in Fig. 2, behaved in a similar manner as a function of solution pH, with the maximum  $K_d$  value of  $6.6 \times 10^3$  at the pH 2.0, plateau with the mean  $K_d$  of  $2.9 \times 10^3$  at the pH range from 4.0 to 10.0, and the  $K_d$  value less than  $1 \times 10^3$  at the pH > 10.0. Under the investigated experimental conditions,  $E$  and  $K_d$  values are insensitive to ionic strength changes, indicating that the sorption of  $\text{TcO}_4^-$  was not affected by the presence of competing  $\text{Cl}^-$  ions.

## CONCLUSION

Ordered mesoporous carbon (OMC) was investigated for its sorption capability to remove pertechnetate ( $\text{TcO}_4^-$ ) anions from aqueous solutions.



**Fig. 2** Effect of pH and ionic strength (0.01 and 0.1 M NaCl) on the sorption of  $\text{TcO}_4^-$  by the OMC. Insert: Variation of  $K_d$  values with pH and ionic strength.

The conclusions are summarized as follows:

- Maximum  $E$  and  $K_d$  values of 99% and  $6.6 \times 10^3 \text{ cm}^3 \text{ g}^{-1}$ , respectively, were obtained at the pH 2.0;
- The adsorption of  $\text{TcO}_4^-$  is almost pH-independent in very wide pH region (from 4.0 to 10.0);
- The sorption of  $\text{TcO}_4^-$  was not affected by the presence of competing  $\text{Cl}^-$  ions.

The obtained results indicate that the  $\text{TcO}_4^-$  sorption by the OMC is well classed compared to other carbon materials (adsorbents). As a material resistant to aging (to the oxidizing conditions and the presence of water), as well as its good sorption properties, the OMC appears to be an efficient adsorbent for the  $\text{TcO}_4^-$  removal from aqueous solutions.

#### ACKNOWLEDGEMENT

This work was partially supported by the Ministry of Education and Science of the Republic of Serbia through the project No. III 45012.

#### REFERENCES

- [1] K. Shi, X. Hou, P. Roos, W. Wu, *Analytica Chimica Acta*, 2012, 709, 1-20.
- [2] M. Momčilović, M. Stojmenović, N. Gavrilov, I. Pašti, S. Mentus, B. Babić, *Electrochimica Acta*, 2014, 125, 606-614.