

1 **Single paternal Dexamethasone challenge programs offspring metabolism and reveals**
2 **multiple candidates in RNA-mediated inheritance**

3

4 Katharina Gapp^{*1,2,3,4}, Guillermo Parada^{1,2,5}, Fridolin Gross³, Alberto Corcoba², Jasmine
5 Kaur³, Evelyn Grau², Martin Hemberg^{1,2,6}, Johannes Bohacek^{3,4}, Eric A. Miska^{1,2,5}

6

7 ¹ Gurdon Institute, University of Cambridge, Cambridge, CB2 1QN, UK.

8 ² Wellcome Sanger Institute, Hinxton, CB10 1SA, UK.

9 ³ Laboratory of Molecular and Behavioral Neuroscience, Institute for Neuroscience,

10 Department of Health Sciences and Technology, ETH Zürich, Zürich, 8057, Switzerland.

11 ⁴ Neuroscience Center Zurich, ETH Zurich and University of Zurich, Zürich, 8057,

12 Switzerland.

13 ⁵ Department of Genetics, University of Cambridge, Cambridge, CB2 3EH, UK.

14 ⁶ Evergrande Center for Immunologic Diseases, Harvard Medical School and Brigham and
15 Women's Hospital, Boston MA 02215, USA.

16

17

18 * corresponding & lead author: katharina.gapp@hest.ethz.ch

19

20

21

22

23

24

25

26

27

28

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

Summary

Single traumatic events that elicit an exaggerated stress response can lead to the development of neuropsychiatric conditions. Rodent studies suggested germline RNA as a mediator of effects of chronic environmental exposures to the progeny. The effects of an acute paternal stress exposure on the germline and their potential consequences on offspring remain to be seen. We find that acute administration of an agonist for the stress-sensitive Glucocorticoid receptor, using the common corticosteroid Dexamethasone, affects the RNA payload of mature sperm as soon as 3 hours post exposure. It further impacts early embryonic transcriptional trajectories, as determined by single embryo sequencing, and metabolism in the offspring.

We show persistent regulation of tRNA fragments in sperm and descendant 2-cell-embryos, suggesting transmission from sperm to embryo. Lastly, we unravel environmentally induced alterations in sperm circRNAs, and their targets in the early embryo, highlighting this class as additional candidate in RNA-mediated inheritance of disease-risk.

Introduction

Acute stress elicits a complex but well-studied cascade of neuroendocrine responses regulated by the hypothalamic pituitary adrenal axis. It involves the release of neuropeptides in the brain that induce the secretion of corticosteroid hormones from the adrenals. These hormones in turn activate two types of corticosteroid receptors, glucocorticoid receptors (GRs) and mineralocorticoid receptor (MRs). These receptors are widely expressed throughout the body and regulate gene expression, thus enabling physiological and behavioral adjustments in response to stress (de Kloet, Joëls and Holsboer, 2005). In vulnerable individuals, this response is excessive and it can lead to long lasting maladaptive changes with consequences for psychological and metabolic health (Daskalakis *et al.*, 2012).

1 It is also known that parental experiences can compromise the health of their progeny both in
2 humans (Pembrey *et al.*, 2006; Heijmans *et al.*, 2008; Bowers and Yehuda, 2016) and in
3 animal models (Benyshek, Johnston and Martin, 2006; Roth *et al.*, 2009; Jimenez-Chillaron
4 *et al.*, 2009; Carone *et al.*, 2010; Pentinat *et al.*, 2010; Shankar *et al.*, 2010; Franklin *et al.*,
5 2010, 2011; Morgan and Bale, 2011; Weiss *et al.*, 2011; Dietz *et al.*, 2011; Vassoler *et al.*,
6 2013; Fullston *et al.*, 2013; Martínez *et al.*, 2014; Gapp *et al.*, 2014; Marco *et al.*, 2014; Rodgers
7 *et al.*, 2015; Sharma *et al.*, 2015; Wu *et al.*, 2016; Chen *et al.*, 2016; Y. Y. Zhang *et al.*, 2018).
8 Research on the underlying mechanism of such transmission has found changes in germline
9 epigenetic make-up, in particular DNA methylation, histone post translational modifications
10 (PTMs), histone positioning and RNA (Gapp and Bohacek, 2017). These epigenetic regulators
11 are responsive to the environment and have been implicated in a variety of environmentally
12 induced diseases (Jirtle and Skinner, 2007). Altered modifications must circumvent epigenetic
13 reprogramming events in zygote and, depending on the timing of exposure, during germline
14 development (Bohacek and Mansuy, 2017; Gapp and Bohacek, 2017). In the male germline,
15 RNA is excluded from reprogramming and therefore a promising candidate for
16 transgenerational information delivery (Gapp and Bohacek, 2017; Bohacek and
17 Rassoulzadegan, 2019). Several studies carried out in *D. melanogaster* and *C. elegans*
18 reported on transgenerational inheritance of induced traits and provided firm evidence for the
19 involvement of small RNAs in the mechanism of transmission (Ashe *et al.*, 2012; Grentzinger
20 *et al.*, 2012; Shirayama *et al.*, 2012). In mammals, a causal implication in the transmission of
21 environmentally induced effects across generations has been demonstrated for sperm RNA
22 only (Gapp *et al.*, 2014; Grandjean *et al.*, 2015; Sharma *et al.*, 2015; Chen *et al.*, 2016). Such
23 RNA differs substantially from somatic RNA since it mainly consists of small RNA,
24 predominantly tRNA-derived small fragments (tsRNAs), but also miRNAs, piRNAs and
25 circRNAs, among others (Chen, Yan and Duan, 2016; Gapp and Bohacek, 2017; Bohacek and
26 Rassoulzadegan, 2019). CircRNAs, comprise a very stable class of RNA that has recently
27 been observed to be present in high amounts in testis but also to some extent in sperm (Barrett
28 and Salzman, 2016). Some have been shown to act as miRNA sponges, thereby competing

1 with mRNA targets, while also regulating the expression of their host genes (Barrett and
2 Salzman, 2016). Hence, circRNAs have a strong potential for amplifying an inherited signal,
3 which makes them exceptionally interesting candidates for epigenetic germline inheritance. To
4 date, the involvement of circRNAs in soma-to-germline signalling has not yet been
5 investigated.

6 tsRNAs and miRNAs are crucial regulators of early embryonic development and players in
7 non-genetic inheritance (Gapp *et al.*, 2014; Grandjean *et al.*, 2015; Rodgers *et al.*, 2015;
8 Sharma *et al.*, 2015; Chen *et al.*, 2016; Benito *et al.*, 2018; Tyebji, Hannan and Tonkin, 2020).
9 They have been reported to be acquired through exosomal uptake during epididymal transfer
10 from caput to cauda epididymis (16,44). This might explain their responsiveness to
11 environmental perturbations, despite mature sperm's presumably transcriptionally silent state
12 caused by tightly packed chromatin. Sperm RNA can indeed change in response to chronic
13 stress or by chronic treatments that mimic stress exposure, such as repeated injection of GR
14 agonists (Rodgers *et al.*, 2013; Gapp *et al.*, 2014; Short *et al.*, 2016; Wu *et al.*, 2016). In mice,
15 uptake of epididymosomal miRNA was sufficient to replicate a chronic stress induced effect on
16 stress response in the offspring (Chan *et al.*, 2020) . Surprisingly, acute stress has also
17 recently been shown to affect offspring weight and glucose metabolism in mice (Hoyer *et al.*,
18 2013) and some of these effects were germline dependent (Bohacek, von Werdt and Mansuy,
19 2016). Together these related lines of evidence led us to hypothesize that acute GR activation
20 has an intergenerational effect on offspring phenotype and that the transmission potentially
21 implicates changes in the germline. The male germline cells - including mature sperm
22 (Kaufmann *et al.*, 1992; Haeussler and Claus, 2007) and their surrounding Sertoli cells (Hazra
23 *et al.*, 2014) – as well as the epididymal epithelial cells (Silva *et al.*, 2010) express GRs that
24 mediate the effects of glucocorticoids on transcription. Dexamethasone (Dex), a specific GR
25 agonist, is known to directly activate GR in the rat epididymis (Silva *et al.*, 2014). It is unknown
26 whether acute stress affects sperm RNA, and if so, whether uptake via epididymosomes is
27 involved in establishing germline changes that are relevant for offspring phenotypic alterations.

1 Here we investigate the impact of acute GR agonist administration on the germline RNA
2 payload including circRNAs, at various time points post administration and interrogate the fate
3 of altered sperm RNA. We further test germ-line dependency of transmitted metabolic effects
4 and dissect the underlying molecular trajectories during early embryonic development using
5 single cell sequencing of in vitro fertilization (IVF) derived embryos. Identifying a readout of
6 transgenerational risk load at the level of the paternal sperm epigenome could pave the way
7 for future studies aiming at a prevention of the transmission of the effects of acute GR
8 activation to the offspring.

9

10

11 **Results**

12 *Effects of acute Dex injection on the germline small RNA payload*

13 Two reports have suggested that a single foot shock could elicit effects on offspring phenotype
14 (Hoyer *et al.*, 2013; Bohacek, von Werdt and Mansuy, 2016). To examine potential epigenetic
15 mediators of such acute stressful impacts we investigated sperm RNA of males 2 weeks after
16 a single activation of the GR (Figure 1A). This timeline was chosen to mimic the timing at which
17 breeding occurred when effects on offspring had been observed in a previous study (Bohacek,
18 von Werdt and Mansuy, 2016). We injected the specific GR agonist Dex once intraperitoneally
19 into 8 adult males. This drug is in frequent clinical use, now also as an apparently effective
20 treatment for patients suffering from lower respiratory tract infection as a consequence of
21 Covid-19 virus (EU Clinical Trials Register, no date; 'Biggest COVID-19 trial tests repurposed
22 drugs first', 2020; Horby *et al.*, 2020). A sperm population was harvested from each animal
23 and RNA was extracted for ultra-deep small RNA sequencing, resulting in 16 libraries
24 representing one injected male each (8 vehicle and 8 Dex-injected). Purity of the sperm
25 samples was confirmed by inspecting RNA size profiles generated on the bioanalyzer to be
26 absent of ribosomal RNA peaks, that would indicate contamination by somatic cells. Reaching
27 an average of 55.4 million sequencing reads while also using randomized adapters for 3'
28 ligation put us in a position to reduce PCR biases (Dard-Dascot *et al.*, 2018) and accurately

1 quantify less abundant miRNAs that are by far outnumbered in sperm by other small RNAs
2 e.g. tsRNAs (Peng *et al.*, 2012). Our data showed an average of 60% mappable reads across
3 all libraries, including 34% of multimappers. We detected an expected dominant prevalence of
4 reads mapping to tsRNAs and abundant miRNAs in all samples (Supplementary Fig. 1A).
5 Differential gene expression analysis, using DESeq2 (Love, Huber and Anders, 2014), revealed
6 that a single acute activation of GR receptors induced changes in tsRNAs, miRNAs and rRNAs
7 collected 14 days post injection (Figure 1B,C, FDR $q < 0.05$, Table 1,2), as has been observed
8 in response to chronic environmental stress previously (Rodgers *et al.*, 2013; Gapp *et al.*,
9 2014). Interestingly, tsRNA-Gly-GCC, a tsRNA previously associated with the effects of
10 nutritional challenge (Sharma *et al.*, 2015), was among the most strongly altered tsRNAs. We
11 further detected changes in ribosomal RNA-derived RNA (rsRNA) as has been observed in
12 studies investigating the impact of high fat and high sugar diet (Y. Zhang *et al.*, 2018; Nätt *et*
13 *al.*, 2019) (Figure 1D,E). It should be noted that our analysis only detects relative changes
14 within the total small RNA pool. Thus, we cannot exclude that the apparent increase in rRNA
15 is due to changes in other RNA subtypes. Accordingly, despite the suggested predictive value
16 of altered rsRNAs for fertility (Hua *et al.*, 2019) we did not observe changes in sperm count,
17 fertilization rate or litter sizes in the Dex treated sperm and resulting offspring (Supplementary
18 Fig. 6) .

19
20 Some recent publications have suggested that sperm miRNAs and tsRNAs are acquired during
21 epididymal transit from caput to cauda (Sharma *et al.*, 2015, 2018; Conine *et al.*, 2018).
22 Further, it was shown that changes in sperm tsRNAs, induced by chronic nutritional-challenge,
23 are acquired by uptake of distinct sets of tsRNAs (Sharma *et al.*, 2015). To examine whether
24 the changes observed 2 weeks post Dex injection were also apparent in caput sperm before
25 epididymal transit we decided to assess a selection of small RNAs in sperm harvested from
26 caput epididymis using q-PCR. Out of 7 selected small RNAs encompassing both tsRNAs and
27 miRNAs we found 6 unaltered (Figure 1F) in line with the assumption that epididymal transit is
28 required to allow epididysomal uptake leading to altered RNA cargo in mature sperm. Yet

1 tsArg-CCT-2 was consistently altered in caput sperm (Figure 1F) indicating that this change is
2 either induced at an earlier transcriptional level during spermatogenesis, or that certain small
3 RNAs are taken up from exosomes in caput epididymis.

4 To test these two hypotheses, we examined a small set of RNAs in serum circulating
5 exosomes. Indeed, we detected increased levels of tsArg-CCT-2, while other tsRNAs and
6 miRNAs did not show altered serum exosome payload (Figure 1G). Thus, it appears that tsArg-
7 CCT-2 is taken up by sperm from exosomes in the caput, but that other small RNA changes
8 are not necessarily reflected in the payload of circulating exosomes.

9 While the necessity of epididymal transit to acquire changes represents one explanatory
10 framework for the absence of change in all but one selected small RNA in caput sperm,
11 alternative explanations should also be considered. Our results could also indicate that
12 changes observed in mature sperm 14 days post Dex injection represent a highly specific
13 snapshot in time, that relies on the affected sperm cells to be in a specific developmental stage
14 at the time of treatment. Cells entering into more mature stages of sperm differentiation at a
15 later point after Dex administration, such as the here assessed cells sampled from caput would
16 then no longer display said changes. Therefore we cannot conclusively establish a reliance on
17 epididymosomal uptake. To further dissect the dependence on epididymosomal uptake during
18 transit from caput to cauda epididymis from a different angle, we assessed the mature sperm
19 small RNA payload at two time points, 3 hours and 7 days following injection (Figure 2A). Cells
20 collected from cauda 7 days after injection have already exited testis, and have had time to
21 pass through the entire epididymal tract before collection. In contrast, cells collected 3 hours
22 post injection have most likely not passed through the corpus epididymis, and already reside
23 in cauda epididymis at the time of injection where sperm resides up to 5 days (Meistrich, 1975;
24 Dadoune and Alfonsi, 1984). Importantly, spontaneous ejaculation regularly voids cauda
25 epididymis of sperm, even in the absence of a mating partner (Huber and Bronson, 1980),
26 excluding the retention of “old” mature sperm in cauda for prolonged periods of time. The cells
27 collected 7 days after exposure therefore represent a mixture of cells that might have already

1 resided in the cauda and those cells that indeed passed through the corpus epididymis, yet
2 the spontaneous ejaculation ensures that the sample predominantly contains the latter.

3 The collected samples were confirmed for their purity and again processed separately to
4 represent sperm from one animal per library. The resulting libraries were analysed jointly as to
5 test for (1) effects of Dex injection independent of sampling time post injection (2) effects of
6 sampling time post injection independent of Dex treatment and (3) effects depending on both
7 Dex injection and the sampling time post injection (interaction). We report an average of 64%
8 mappable reads including 46% of multimappers and observe that tsRNAs were significantly
9 affected by sampling time post injection independent of treatment. This demonstrates the
10 fluctuation of tsRNAs over time in response to external signals such as injections, or potentially
11 due to uncontrollable external inputs from the animal husbandry (Figure 2C). Interaction
12 between treatment and time was statistically significant for 27 tsRNA mapping loci including
13 Gly-GCC-6-1. All affected tsRNAs are upregulated after 7 days. 26 tsRNAs of them are
14 unchanged after 3 hours and one tsRNA (Thr-TGT1-1) is downregulated after 3 hours
15 (Supplementary Table2 sheet 3, $q < 0.05$). This finding is consistent with the dominating view
16 that tsRNAs are acquired during epididymal transit from caput to cauda epididymis. However,
17 most tsRNAs that showed a significant change in response to treatment after 7 days, but not
18 after 3 hours (interaction between treatment and time post injection, Supplementary Table 2
19 sheet 3, $q < 0.05$) were not persistently altered in the dataset of 14 days after injection (Table
20 1). This indicates that on the one hand changes in sperm RNA are dynamic and many do not
21 persist for prolonged time. On the other hand, this suggests that potentially relevant small RNA
22 changes mostly require either sperm to reside in testis at the time of exposure, or rely on a
23 prolonged residency in the exposed organism. Interestingly, we also detected 2 exceptions
24 that show a significant group effect across 3 hours and 7 days. tsRNA-Leu-CAA and tsRNA-
25 Arg-CCT (Figure 2B) were persistently affected 3 hours and 7 days post exposure, that
26 necessarily requires a mode of rapid acquisition of tsRNA-changes in cauda epididymis. While
27 the change in tsRNA-Leu-CAA was temporary and did not persist, strikingly tsRNA-Arg-CCT-
28 2 deregulation persisted until 14 days post injection (Figure 1C). To additionally validate the

1 Dex induced change of tsRNA-Arg-CCT-2 independent of epididymal transit from caput to
2 cauda we replicated the effect observed in mature sperm sampled from cauda epididymis 3
3 hours post injection using q-PCR (Figure 2D, Supplementary table 2). Additionally, we sampled
4 caput sperm 3 hours post injection and measured tsRNA-Arg-CCT-2 levels. An overall 2-way
5 ANOVA revealed a significant interaction between sperm sampling location (caput versus
6 cauda) and treatment (vehicle versus Dex). Post hoc tests confirm a significant increase in
7 tsRNA-Arg-CCT-2 levels in response to Dex in cauda but not in caput sperm and a significant
8 increase in Arg-CCT-2 levels between cauda and caput sperm independent of treatment.

9

10 The behaviour of miRNAs differed considerably from tsRNAs. As would be expected if
11 epididymal transit was required for miRNA changes to be implemented, we observe no group
12 effect of treatment on miRNAs (Supplementary Fig. 2A, Supplementary table 2) across 3 hours
13 and 7 days post injection. Further, we detected no effect of time post injection on sperm miRNA
14 payload (Supplementary Fig. 2B, Supplementary table 2) confirming the absence of an effect
15 of injection on miRNAs per se. However, we neither detected an interaction between Dex and
16 time post injection (Supplementary Fig. 2C, Supplementary table 2) in miRNAs 7 days and 3
17 hours post injection. Importantly, when inspecting those miRNAs that were significantly altered
18 14 days after injection, no alterations were apparent 3 hours or 7 days post injection
19 (Supplementary Fig. 2D), indicating that changes in miRNAs occur more slowly or require
20 sperm cells to reside in the testes at the time of injection.

21

22 *Effects of acute GR activation on in vivo offspring metabolic phenotype*

23 Based on the two reports on effects of single foot shock on offspring weight and the impact of
24 a single GR activation on germline small RNA payload, we hypothesized that this acute impact
25 on the receptor is sufficient to elicit intergenerational effects. We thus injected Dex once
26 intraperitoneally, then harvested sperm 14 days post injection, and performed *IVF* using naïve
27 oocytes to generate offspring for phenotyping (Figure 3A). Dex treatment did not affect sperm
28 count, fertility rate or resulting litter-sizes (Supplementary Fig.6).

1 The weight and size of pups was measured every 2 to 4 weeks starting at weaning (3 weeks
2 of age) until adulthood (12 weeks of age) and body mass index (BMI) was calculated as a ratio
3 of weight and squared length. Overall ANOVA of the resulting offspring showed a significant
4 effect of treatment ($(F_{1,71}) = 76.55, p < 0.0001$), time post injection ($(F_{2,087,144.7}) = 41.99,$
5 $p < 0.0001$) and sex ($F(1, 71) = 76.55, p < 0.0001$) on BMI, and a significant interaction between
6 time and sex ($F(3, 208) = 33.75, p < 0.0001$) and time and treatment ($F(3, 208) = 5.834,$
7 $p = 0.0008$) (Figure 3B, Supplementary Fig. 3C, Supplementary table 3). These results show
8 that while males had generally higher BMI, both male and female offspring of Dex injected
9 fathers had a higher BMI.

10 To further explore potential causes and consequences of altered BMI, adult animals were
11 additionally tested for their glucose tolerance following glucose injection. Overall ANOVA
12 analysis of blood glucose levels revealed a significant effect of sex ($F(1, 44) = 54.80,$
13 $p < 0.0001$) and time post injection ($F(2.593, 114.1) = 196.6, p < 0.0001$) and significant
14 interactions between sex and time post injection ($F(4, 176) = 6.115, P = 0.0001$), and sex and
15 treatment ($F(1, 44) = 15.62, P = 0.0003$) (Supplementary Fig. 3C). Follow up repeated
16 measurements ANOVA analysis separated by sex showed a significant effect of treatment,
17 time and interaction in females (treatment: $F(1, 22) = 12.35, p = 0.0020$; time: $F(4, 88) = 110.1,$
18 $p < 0.0001$; interaction: $F(4, 88) = 2.835, p = 0.0291$) and significant effects of treatment and
19 time but no interaction in males (treatment: $F(1, 22) = 6.019, p = 0.0225$; time: $F(4, 88) = 96.36,$
20 $p < 0.0001$; $F(4, 88) = 0.5401, p = 0.7067$; Figure 3C). These data hence demonstrate a sex-
21 dependent effect of paternal Dex injection on glucose tolerance, with impaired tolerance in
22 females and decreased glucose levels in males in response to glucose challenge.

23 In addition, blood glucose levels were assessed during the insulin tolerance test. Overall
24 ANOVA analysis showed significant effects of sex ($F(1, 37) = 162.6, P < 0.0001$)
25 and time ($F(3.314, 122.6) = 23.85, P < 0.0001$) and revealed a significant interaction between
26 sex and time ($F(4, 148) = 12.49, P < 0.0001$), time and treatment ($F(4, 148) = 5.380, P = 0.0005$)
27 and time and treatment and sex ($F(4, 148) = 5.392, P = 0.0004$) (Supplementary Fig. 3C).
28 Follow-up repeated measurements ANOVA separated by sex showed a significant effect of

1 time ($F(2.982, 65.60) = 44.73, p < 0.0001$), yet no significant effect of treatment ($F(1, 22) =$
2 $0.3465, p = 0.5621$) nor an interaction between time and treatment ($F(4, 88) = 0.1373,$
3 $p = 0.9681$) in females (Figure 3D). In males we observe no effect of treatment ($F(1, 15) = 1.467,$
4 $p = 0.2446$) yet detected a significant effect of time post injection time ($F(2.914, 43.71) = 4.538,$
5 $p = 0.0079$), and a significant interaction between treatment and time post injection ($F(4, 60) =$
6 $7.003, p = 0.0001$, Figure 3D). These results indicate sex and time dependent effects of
7 paternal Dex on insulin tolerance. They further show no change in insulin tolerance in female
8 descendants of fathers injected with Dex, yet impaired insulin tolerance in male progeny.
9 Lastly, we explored a potential reflection of altered BMI in tissue composition by necropsy and
10 weighing the dissected organs and fat pads. Overall ANOVA of necropsy weights revealed a
11 significant effect of sex ($F(1, 140) = 28.27, P < 0.0001$), tissue ($F(4, 140) = 232.7, P < 0.0001$)
12 and a significant interaction between sex and tissue ($F(4, 140) = 3.379, P = 0.0113$) yet no
13 effect of treatment ($F(1, 140) = 0.2587, P = 0.6118$), or interaction between treatment and sex
14 ($F(1, 140) = 0.0004794, P = 0.9826$) or treatment and tissue ($F(4, 140) = 0.1635, P = 0.9565$)
15 on tissue weight (Supplementary Fig. 3A,B,C). This confirms sex dependency, yet no effect of
16 paternal Dex injection on tissue weight in both sexes.

17

18 *Effects of acute Dex on offspring early embryonic small RNA*

19 The small quantity of paternal RNAs in the zygote relative to the large pool of maternal RNAs
20 poses serious obstacles to their accurate quantification (Chen, Yan and Duan, 2016). While
21 initial reports on small RNA transmission relied on comparative sequencing or microarray
22 analyses of unfertilized oocytes and fertilized zygotes (Ostermeier *et al.*, 2004), today we are
23 aware that such comparisons can be deceiving, as they rely heavily on both assessment
24 method (e.g. microarray restricted to a selective set versus unbiased genome-wide
25 sequencing) and sequencing depth (Dard-Dascot *et al.*, 2018; Yeri *et al.*, 2018). One such
26 example is inconsistent results regarding miRNAs that are exclusively supplied from the
27 sperm, such as miR-34c, -99a, -214 (Amanai, Brahmajosyula and Perry, 2006; Liu *et al.*, 2012).
28 Alternative approaches have used indirect measures, e.g. assessing mRNA targets of

1 paternally derived small RNAs (Amanai, Brahmajosyula and Perry, 2006; Tang *et al.*, 2007;
2 Krawetz *et al.*, 2011; Sharma *et al.*, 2015). We attempted to directly examine the relative
3 difference between the small RNA landscape in early embryos resulting from IVF of naïve
4 oocytes with sperm from either Dex or vehicle injected males (Figure 4A). We used small-RNA
5 sequencing to compare 2-cell embryos derived from Dex treated or control fathers. We
6 detected an average of 29 % mappable reads including 21% multimappers. While we only
7 detected subtle changes in miRNAs of Dex exposed progeny, we observed downregulation of
8 several tsRNAs from 6 different genomic locations ($q < 0.1$) (Figure 4B). Strikingly, two of the
9 downregulated tsRNAs (Gly-GCC at several genomic loci and Gly-CCC) were consistently
10 downregulated in sperm 14 days post Dex injection. This could either indicate a reduced
11 delivery of this sperm RNA cargo in Dex treated males to the oocytes they fertilize, or earlier
12 usage and function of respective RNA in Dex leading to a quicker elimination or shorter half-
13 life.

14
15

16 *Effects of acute Dex administration on offspring early embryonic transcriptome*

17 If sperm RNA were directly impacting the zygotic mRNA pool or if they were affecting early
18 embryonic gene expression, then this should be apparent in the 2 cell embryo's transcriptome
19 (Figure 5A). To examine the effect of paternal Dex on early embryonic RNA content we
20 subjected 2-cell embryos to the Smartseq single cell sequencing protocol (Supplementary Fig.
21 4). After performing quality control and filtering the sequenced 2-cell embryo data on criteria
22 such as minimal read count/embryo (Supplementary Fig. 4A), we carried out unsupervised
23 clustering based on their gene expression profiles using SC3 (Kiselev *et al.*, 2017). We
24 identified two distinct clusters (C1 and C2), which were composed by a balanced mixture of
25 treated and control cells. (Figure 5B). Since the resolution of single cell experiments allows
26 characterizing distinctive transcriptomic profiles within early cell division stages, we used
27 scmap (Kiselev, Yiu and Hemberg, 2018), to project each 2 cell embryo gene expression
28 profile onto a reference dataset of single cells from 2 cell embryo states previously reported

1 by Deng *et al.* (Deng *et al.*, 2014) (Figure 5C.) Most of the 2 cell embryos belonging to cluster
2 C1 projected to the late 2 cell stage, whereas embryos from C2 exclusively projected to cells
3 from the mid 2 cell stage. This shows that the two clusters identified through unsupervised
4 clustering correspond to 2-cell embryos in the mid and late 2-cell stage respectively.

5 Principal component analysis (PCA) revealed a prominent separation between C1 and C2
6 along the PC1 axis, suggesting a correlation between PC1 and developmental transitions
7 between mid and late single cell embryos (Supplementary Fig. 4B.) Interestingly, 2 cell embryo
8 offspring of males injected with Dex exhibited a significant shift of the C1 cluster across PC1
9 (two-sided Wilcox test $p < 0.03$), while the C2 clusters did not show significant differences
10 across PC1 between treatment and control groups (Figure 5D.) These results suggest that the
11 effect of paternal Dex treatment on the transcriptome only becomes apparent at the late 2 cell
12 embryo stage. To further explore this hypothesis, we calculated the silhouette coefficient
13 (Rousseeuw, 1987) on PC1, as a measure of distance between C1 and C2 clusters, for the
14 control and treatment group. We observed a significant increase of PC1 silhouette coefficient
15 between treatment and controls for both C1 (one-sided Wilcoxon test p -value < 0.005) and C2
16 (one-sided Wilcoxon test p -value $< 2 \times 10^{-5}$.) This confirms that Dex treatment affects
17 embryonic gene expression, promoting altered late 2 cell embryo stages since the divergence
18 from mid 2 cell embryos is significantly bigger in Dex offspring compared to control offspring
19 (Figure 5E.)

20 Accordingly, differential gene expression analysis using Monocle2 (Qiu *et al.*, 2017) focused
21 on late 2-cell embryos (cluster C1) revealed significant gene expression changes between
22 offspring of males injected with Dex and controls across 38 genes, some of which were already
23 apparent to a less significant extent during mid-2 cell embryos (cluster C1; e.g. *Tcl1*;
24 Supplementary Table 4 In line with a potentially altered developmental trajectory becoming
25 apparent in cluster 1, the late 2-cell stage includes several affected genes that are involved in
26 early embryonic development. For example, *Bcap31* (B-cell receptor-associated protein 31) is
27 an important element for endoplasmic reticulum and Golgi apparatus function, and *Bcap31*
28 mutations lead to developmental diseases with metabolic disturbances (Cacciagli *et al.*, 2013).

1 This is reminiscent of the metabolic phenotype observed in the adult offspring of Dex injected
2 fathers. Hprt (Hypoxanthine-guanine phosphoribosyltransferase) is crucial for cell cycle
3 division, and Tcl1 (T-cell leukemia/lymphoma) regulates cell proliferation (Kang *et al.*, 2013;
4 Miyazaki *et al.*, 2013). Hence, an upregulation of Hprt and a concomitant down-regulation of
5 Tcl1 might indicate that cell fate decisions later during development may be affected. Another
6 differentially expressed gene is Rbbp7 (RB binding protein 7), which is part of many histone
7 deacetylase complexes such as Nurd and PRC2/EED-EZH2, and thus plays an essential role
8 in chromatin mediated gene regulation (Yu *et al.*, 2018). Interestingly, several forms of PRC
9 mutations in humans lead to different kinds of overgrowth phenotypes (Deevy and Bracken,
10 2019), an abnormality reminiscent of the increased BMI observed in Dex-offspring (Figure 3B.)
11
12

13 *Effects of Dex administration on an interesting candidate for sperm RNA mediated inheritance*
14 Despite the observed changes in sperm tsRNAs following acute Dex injection, we did not find
15 an obvious causal connection to the altered 2-cell embryonic transcripts. This prompted us to
16 investigate whether other germline changes might be more crucial for the offspring's *in vivo*
17 alterations in our model. We previously showed that chronic stress exposure also led to
18 changes in sperm long RNAs that contributed functionally to the transmission of effects to the
19 offspring (Gapp *et al.*, 2018). Yet the fact that sperm RNA is stable through transmission and
20 that the minute amounts of transmitted paternal RNA can elicit major changes in the embryo
21 remains puzzling. Therefore, we evaluated the impact of Dex injection on the highly stable
22 class of circRNAs in male sperm. CircRNAs were previously detected in swine (Gòdia *et al.*,
23 2020) and human sperm (Chioccarelli *et al.*, 2019) and suggested to have functional
24 implications in epigenetic regulation. They have been attributed a critical role in the male
25 germline after cessation of transcriptional activity (Tang *et al.*, 2020). Using Circexplorer in
26 combination with EdgeR, analysis of sperm long RNA sequencing of males treated with Dex
27 and controls revealed significant upregulation of two circRNAs (Figure 6A, $q < 0.1$) and we also
28 observed several significant changes in the sperm long RNA protein coding transcripts

1 following acute Dex treatment (Figure 6B,C, Supplementary Fig. 1B, Supplementary table 5).
2 Both circRNAs are hosted in genes relevant for immune function (Taspase 1: Tasp1 and DENN
3 Domain Containing 1B Dennd1b), yet the host genes did not show differential abundance
4 of the protein coding transcript (Supplementary table 5). We then replicated the upregulation
5 of these to CircRNAs by q-PCR in cauda epididymal sperm of a separate set of animals using
6 CircRNA specific primers that span the backsplice junction. At the same time, we also
7 assessed their abundance in caput epididymal sperm to evaluate whether the observed effect
8 was also present in sperm cells under development. We observed a significant interaction
9 between treatment and tissue. Post hoc tests confirm the increased abundance of CircTasp1
10 and CircDend1 in cauda sperm detected in the sequencing analysis. No change was
11 detectable in caput sperm, arguing against an induction of the change in developing sperm
12 cells (Figure 6D,E). Given the absence of transcription, these data suggest that in control
13 conditions these CircRNAs exert their function during spermiogenesis (for example by being
14 translated) and that by contrast in Dex treated cells the CircRNA is not consumed to the same
15 extent leading to an apparent upregulation in mature cauda epididymal sperm. CircAtlas(Wu,
16 Ji and Zhao, 2020) revealed several potential miRNA sponge-targets to be captured by the
17 altered circRNAs. Some of these miRNAs are common sponge-targets of both circRNAs such
18 as mir3110-5p, mir706, mir1955 (Supplementary Fig. 5). Diamine acetyltransferase 1 (Sat1),
19 one of 3110-5p's high confidence miRNA-targets, as predicted by TargetScan(Agarwal *et al.*,
20 2015), is indeed significantly upregulated in the embryos composing cluster 1 (later
21 developmental stage). MiRNA-target-upregulation is expected if mir3110-5p was
22 downregulated through circRNA-mediated-sponging and highlights a potential effective
23 contribution of increased circRNA in sperm to embryonic pathway regulation. This is the first
24 report of a change induced by environmental exposure in this compelling class of RNA in
25 sperm.

26

27 **Discussion**

1 By generating offspring using assisted reproductive techniques (IVF), we circumvent potential
2 confounding variables such as transmission via RNA contained in seminal exosomes (Vojtech
3 *et al.*, 2014) and affected maternal care by altered mating behavior (Mashoodh *et al.*, 2012)
4 and prove germline dependence (Bohacek and Mansuy, 2017). Consistent with the significant
5 changes of miRNAs and tsRNAs in the germline 2 weeks post GR activation, previous studies
6 including our own have observed regulation of mouse sperm small RNA in a variety of contexts
7 (Rodgers *et al.*, 2013; Gapp *et al.*, 2014; Grandjean *et al.*, 2015; Sharma *et al.*, 2015; Chen *et*
8 *al.*, 2016; Wu *et al.*, 2016; Short *et al.*, 2017; Benito *et al.*, 2018; Rompala *et al.*, 2018;
9 Yeshurun and Hannan, 2018).

10 Especially relevant specifically for our analysis, sperm RNA sequencing after drinking water
11 administration of corticosterone for 4 weeks followed by mating, led to strong downregulation
12 of tsRNA-GluCTC and tsRNA CysGCA, two of our top down-regulated tsRNAs, indicating that
13 these tsRNAs are responding similarly to acute and chronic insults. At the same time this
14 chronic manipulation elicits changes of several miRNAs, e.g. 34c and 471 (Short *et al.*, 2016),
15 albeit in the opposite direction of what we find in response to acute Dex treatment. These
16 discrepancies may arise either by the Dex induced short-term suppression of internal
17 corticosteroid (Barden, 1999) or due to adaptations in response to chronic administration.

18 While four (Petropoulos, Matthews and Szyf, 2014; Short *et al.*, 2016; Wu *et al.*, 2016; Cartier
19 *et al.*, 2018) out of five (Bönisch *et al.*, 2016) previous studies did report phenotypic effects
20 following chronic paternal Dex exposure, only two assessed sperm small RNAs to associate
21 the alterations to the sperm RNA payload (Bönisch *et al.*, 2016; Cartier *et al.*, 2018) yielding
22 conflicting outcomes.

23 These differences might be due to inconsistent life stages (adulthood versus gestational),
24 sperm collection (swim up, somatic lysis, or no purification) and/or dosage of exposure.
25 Depending on the dosage and timing, the complex autoregulation of the GR can lead to GR
26 downregulation after prolonged activation (Gjerstad, Lightman and Spiga, 2018). Acute
27 exposures have the advantage of avoiding such long-term feedback regulation, and hence
28 provide an elegant approach for studying the signaling pathways leading to germline changes.

1 Mature sperm tsRNAs and miRNAs have been shown to be acquired during epididymal transit
2 (Sharma *et al.*, 2015, 2018) and miRNAs are necessary for early embryonic development
3 under certain circumstances (Liu *et al.*, 2012; Conine *et al.*, 2018)(Zhou *et al.*, 2019)(Wang *et*
4 *al.*, 2020). Furthermore, a recent publication suggests that chronic stress induced sperm
5 miRNAs are taken up primarily from epididymosomes originating from the caput epididymis or
6 the proximal epididymal tract (Chan *et al.*, 2020). Chronic nutritional manipulation with effects
7 on offspring also suggest the necessity of epididymal transit to acquire tsRNA changes in
8 sperm(Sharma *et al.*, 2015). Harvesting mature sperm 3 hours after exposure yields a
9 population enriched for cells that had been exposed while already in the cauda epididymis,
10 where spermatids reside for roughly 5 days (Meistrich, 1975). These cells have not traveled
11 through the epididymis nor have they had a chance to potentially take up small RNA from
12 caput-derived epididymosomes after Dex administration. As expected, we detect no changes
13 in miRNAs in these samples. We do however detect changes in tsRNAs 3 hours post Dex,
14 some of which even persist 14 days post injection. These results show rapid acquisition of
15 changes in vivo and corroborate previous in vitro findings that show that incubation with
16 epididymosomes can alter sperm RNA payload(Sharma *et al.*, 2015). Our acute intervention
17 assesses effects on germline payload already after a short interval, whereas chronic
18 interventions - based on their experimental design - do not assess changes in mature sperm
19 soon after the first intervention. Studies aiming at the elucidation of the origin of sperm RNA
20 changes might benefit from acute interventions to circumvent confounders such as dynamic
21 exosomal RNA supply as a result of cumulative interventive strain on animals.

22 An additional option for sperm RNA alterations in transcriptionally inert sperms was suggested
23 in a recent study that found mitochondrial tRNA cleavage in the T-loop in response to a one-
24 week high sugar diet in humans(Nätt *et al.*, 2019). In line with this observation, Dex injection
25 could trigger oxidative stress(Bera *et al.*, 2010) which provokes such cleavage to increase
26 tsRNA levels(Thompson *et al.*, 2008). A role for oxidative stress in sperm RNA dynamics is
27 further supported by a recent study in boar sperm that found seasonal differences in sperm
28 small and long RNA associated with changed abundance of transcripts mapping to oxidative

1 stress-, DNA damage- and autophagy- related genes (Gòdia *et al.*, 2019). However, such
2 potentially oxidative stress mediated mechanism does not explain a rapid decrease of tsRNAs
3 3 hours post Dex injection.

4 Importantly, we show alterations in sperm tsRNAs that persist in the oocyte concomitant with
5 changes in early embryonic gene expression and a metabolic phenotype in adulthood. tsRNAs
6 and tRNA-Gly derived fragments in particular are known to induce chromatin structure
7 mediated gene regulation and to regulate cell differentiation in various contexts (Li *et al.*, 2016;
8 Guzzi and Bellodi, 2020). Hence, we propose that the transmitted reduction in key tsRNAs
9 such as Gly-GCC and Gly-CCC explains in part the observed perturbations during late two cell
10 embryo developmental stage. This might reflect an accelerated developmental transcriptional
11 program in the preimplantation embryo of Dex injected males, ultimately resulting in aberrant
12 BMI and glucose metabolism later in life.

13 Additionally, we have discovered alterations in circRNA abundance in mature sperm that might
14 also impact the developmental program in the early embryo. CircRNAs have the potential to
15 be translated into proteins via back-splicing(Legnini *et al.*, 2017). Accordingly, they are crucial
16 contributors to spermiogenesis post transcriptional cessation, since they provide a stable
17 alternative to linear mRNA templates for protein translation(Tang *et al.*, 2020). Transmitted
18 sperm-circRNAs could likewise contribute to translation post fertilization, yet the
19 unconventional lattice-state of ribosomes preventing normal rates of translation post
20 transcription(Israel *et al.*, 2019) accompanied by a rapid increase in proteins of the
21 ubiquitine/proteasome pathway(Wang *et al.*, 2010) make this unlikely. Nevertheless, a study
22 on human sperm detected abundant levels of circRNA with predicted regulatory function of
23 early developmental genes in sperm heads, suggesting transmission and function post
24 fertilization(Chioccarelli *et al.*, 2019). By sponging miRNAs that regulate early embryonic
25 transcripts, circRNA could amplify minute signals from paternal environment, such as might be
26 the case for the gene transcript Sat1, that displays increased expression in Dex offspring in 2-
27 cell embryos from cluster 1.

1 Besides altered RNA identity, nucleic acid modifications especially of RNA but also DNA
2 methylation and chromatin accessibility might contribute further to the effects of Dex injections
3 on offspring metabolism. While detection of changes in each player should be subject of further
4 investigation and might reveal a glimpse of their potential implication, proof of the individual
5 relative causal contribution is extremely challenging since they likely require tight interaction
6 to unravel their orchestrated effects.

7 Finally, it might be useful to consider testing the translatability of our findings to humans. Here
8 we investigated the effects of a single Dex administration soon after the injection in mice,
9 mimicking a single GR activation such as elicited by treatment of an acute asthma
10 exacerbation (Cross, Paul and Goldman, 2011). The recent report that Dex can reduce the
11 number of deaths associated with the Covid-19 pandemic ('Biggest COVID-19 trial tests
12 repurposed drugs first', 2020), further prompts the re-evaluation of the impact of prolonged
13 Dex treatment on offspring phenotype. From a clinical perspective, additional consideration is
14 warranted for consequences on offspring health when extended time has elapsed between
15 treatment and time of conception. Such designs may pave the way for the extrapolation of our
16 findings.

17 We conclude that acute Dex treatment can induce germline epimodifications in the form of
18 small and long non-coding RNA, which likely are relevant in the transmission of the effects of
19 single traumatic events on offspring well-being. Our data suggest that sperm small RNAs are
20 not solely regulated via epididymosomal uptake during transition from caput to cauda
21 epididymis. This expands the interpretation from chronic dietary and stress exposures(Sharma
22 *et al.*, 2015; Chan *et al.*, 2020), where uptake of tsRNAs and miRNAs via epididymosomes
23 has been suggested to lead to differential sperm payload, yet required sperm to transit from
24 caput to cauda to bring about the changes. A persistent detection of significant fold changes
25 of the exact same sperm small RNA in the embryo suggests functional implication in the
26 information transfer from father to offspring. Together with potentially transmitted miRNA
27 sponges in the form of circRNAs, this likely contributes to a slight developmental acceleration
28 of gene expression programs in the early embryo and ultimately manifests in a metabolic

1 phenotype. Future studies may aim at testing the causal contribution of specific sperm RNAs
2 to the transmission of effects of acute impacts. Certainly, continuous methodological
3 refinement will help dissect the relative implication and the interplay of the distinct germline
4 modifications such as DNA-methylation, histone-PTMs and chromatin architecture in this
5 highly complex process.

6

7 ***Limitations of study***

8 This study identifies a highly dynamic response of sperm RNA cargo in response to a single
9 Dex administration and presents altered CircRNAs in mature sperm in response to an
10 environmental insult with consequences on the progeny's metabolism. While our study implies
11 a functional role of altered sperm RNAs in the transmission of a Dex induced phenotype to the
12 offspring a causal proof will require RNA injections into fertilized naïve oocytes. Second,
13 although we identified one tsRNA to be persistently altered in serum exosomes as well as in
14 caput and cauda sperm and at different times post Dex injection, the identification of the tissue
15 of origin of the altered tsRNAs would require technically highly challenging metabolic labelling
16 experiments.

17

18 ***Author's contributions***

19 KG performed animal exposures, collected samples and prepared sequencing libraries. JK
20 helped with sample collection and processing and performed q-PCRs. EG performed IVF and
21 embryo culture. GP analysed single cell sequencing data. MH supervised the single-embryo
22 sequencing analysis. AC and FG analyzed bulk sequencing data. KG, JB and EAM designed
23 the study, interpreted the results and wrote the manuscript with input from the other authors.

24

25

26 ***Acknowledgements***

27 We thank Wayo Matsushima and Pierre-Luc Germain for valuable advice on bulk sequencing
28 analysis and the Sanger RSF for help with animal care, especially the teams of Michael Woods

1 and Francesca Flack. We also thank Sergio Mompert Barrenechea for assistance with the q-
2 RT-PCR and Darren Kelly for CircRNA primer design.
3 KG was funded by the Swiss National Science Foundation early postdoc and advanced
4 postdoc mobility a SPARK and Novartis foundation grant. Some of this work was supported
5 by Cancer Research UK (C13474/A18583, C6946/A14492) and Wellcome (104640/Z/14/Z,
6 092096/Z/10/Z) to EAM. GP and MH were supported by a core grant from the Wellcome Trust.
7 The lab of JB is currently funded by the ETH Zurich, SNSF Project Grant 310030_172889/1,
8 ETH Research Grant ETH-20 19-1, the Kurt und Senta Herrmann-Stiftung, the Botnar
9 Research Center for Child Health and a 3R Competence Center Project Grant. JK was
10 supported by a Swiss-european mobility programme scholarship. For the purpose of Open
11 Access, the author has applied a CC BY public copyright license to any Author Accepted
12 Manuscript version arising from this submission.

13

14 ***Declaration of interests***

15 The authors declare no competing interests. E.A.M. is a founder and Director of STORM
16 Therapeutics Ltd. STORM Therapeutics had no role in the design of the study and collection,
17 analysis, and interpretation of data as well as in writing the manuscript.

18

19

20

1 **Figure legends**

2 **Figure 1** Effects of Dex on small RNA payload of sperm cells residing in testis at time of
3 administration. (A) Experimental design depicting a time window of two weeks between
4 injection of Dex and sperm collection for molecular analysis. (B) Volcano plot depicting fold
5 changes and significance level of miRNAs and tsRNAs (C) in mature sperm 14 days post
6 injection of Dex (n=8) versus vehicle (n=8) as assessed by small RNA sequencing. (D) Stacked
7 barplot showing reads of different sizes mapping to rRNAs in Dex (n=8) and vehicle (n=8). (E)
8 Volcano plot demonstrating global increased abundance of rRNA fragments. (F) Q-PCR
9 results of small RNA assays for caput sperm (tsArg-CCT-2: t(49)=3.49, q<0.001, tsGlyGCC:
10 t(49)=0.32, q>0.05; tsHls-GTG: t(49)=0.29, q>0.05; tsLeu-CAA-2: t(49)=0.62, q<0.5; mir34c-
11 5p: t(49)=0.27, q>0.05; mir6538: t(49)=0.59, q>0.05, mir677-5p t(49)=0.62 q>0.05) (G) serum
12 exosomes (tsArg-CCT-2: t(33)=3.43, q<0.01, tsGlyGCC: t(33)=0.04, q>0.05; tsHls-GTG:
13 t(33)=0.07, q>0.05; mir34c-5p: t(33)=0.51, q>0.05; mir6538: t(33)=0.02, q>0.05) days after
14 drug injection. Whiskers display minimum and maximum. **q<0.01, ***q<0.001 multiple t-tests
15 corrected for multiple testing.

16
17 **Figure 2** Effect of Dex on sperm cells at different time points post Dex administration. (A)
18 Experimental design showing the location of sperm at time of injection and timing of sperm
19 harvest. MA (log-intensity ratios (M-values) versus log-intensity averages (A-values) plots
20 depicting (B) effect of Dex (log₂ fold changes control versus dexamethasone), (C) of time post
21 injection (log₂ fold changes 7 days versus 3 hours) (7 days Dex n =4 and controls n =4, 3
22 hours Dex n =3 and controls n =4). TsRNAs are indicated by sequence identity for display only,
23 each dot represents one small RNA. MA plot depicts log₂ fold changes on the y axis and the
24 expression level on the x axis (the higher the expression the further to the right). Statistically
25 significantly changed small RNAs are highlighted in red q<0.05. (D) Relative expression of
26 ArgCCT-2 as obtained by q-RT-PCR (cauda: Dex n=4, controls n=4, caput: Dex n=4, controls
27 n=5; interaction F(1,13)=6.34, p=0.0257, treatment F(1,13)=5.97, p=0.0040, site of collection
28 (F(1,13)=12.15, p=0.0296; cauda control versus cauda Dex t(13)=3.42, p=0.0274, cauda Dex

1 verus caput Dex $t(13)=4.137$, $p=0.007$). Whiskers display minimum and maximum. $*p<0.05$,
2 mixed effect model group effect of location (REML), $**p<0.01$ multiple comparisons SIDAK
3 corrected.

4
5

6 **Figure 3** Effect of Dex on metabolic phenotype in the offspring (A) Experimental design
7 depicting timeline between injection, sperm harvest, in vitro fertilization and phenotyping. (B)
8 Impact of Dex on male and female adult offspring (B) Body mass index (males vehicle offspring
9 $n=21$, Dex offspring $n=22$, females vehicle offspring $n=17$, Dex offspring $n=17$) (C) glucose
10 tolerance (males vehicle offspring $n=12$, Dex offspring $n=12$, females vehicle offspring $n=12$,
11 Dex offspring $n=12$) and (D) insulin tolerance (males vehicle offspring $n=9$, Dex offspring $n=8$,
12 females vehicle offspring $n=12$, Dex offspring $n=12$). Error bars display SEM. Detailed
13 statistical results are depicted in Supplementary Fig. 5, raw data are provided in supplementary
14 table 3). $*p<0.05$, $**p<0.01$, $***p<0.001$, 3-way repeated measures ANOVA.

15

16 **Figure 4 Effect of paternal Dex injection on embryonic offspring small RNA** (A)
17 Experimental design depicting timeline between injection, sperm harvest, in vitro fertilization
18 and small RNA sequencing at 2-cell stage. (B) Heatmap showing effect of paternal Dex on
19 small RNA tsRNAs (vehicle embryonic offspring $n=5$, Dex embryonic offspring $n=4$). TsRNAs
20 are grouped by sequence identity for display only.

21

22

23 **Figure 5** Effect of paternal Dex injection on embryonic offspring long RNA transcriptome (A)
24 Experimental design depicting timeline between injection, sperm harvest, in vitro fertilization
25 and Smartseq2 sequencing at 2-cell stage. (B) Consensus matrix representing the similarity
26 between cells as reported by SC3. Similarity 0 indicates that a given pair of embryos were
27 never assigned to the same cluster, whereas similarity 1 means that a pair of embryos were
28 always assigned to the same cluster. (C) Sankey diagram showing projection of the obtained

1 clusters (C1 and C2) into clusters reported by Deng *et al.* for single cells obtained from 2 cell
2 embryos. (D) Principal component analysis of two-cell embryos gene expression. The top
3 panel indicates the density of 2 cell embryos along PC1 grouped by condition; control (red)
4 and Dex treatment (blue). The two bottom panels show the distribution of 2 cell embryos across
5 PC1 and PC2 for control (red) and treated (blue) groups. The cluster membership of each
6 embryo is denoted by the point shapes (C1 cycles; C2 triangles) and the centroids of each
7 cluster is indicated with a black symbol overlaid with an x. Wilcox tests were performed to
8 assess differences on PC1 values of C1 and C2 clusters between the treated and control
9 groups. NS denotes non-significant change for C2 cluster, while * indicates a significant
10 difference for C1 cluster (p -value < 0.05). (E) Silhouette coefficient comparison between
11 treatment and control, statistical significance was assessed with Wilcox test (** p -value < 0.01;
12 *** p -value < 0.005) (F) Selection of differentially expressed genes as determined by Monocle
13 within C1 corresponding to late 2-cell embryo stage (*** adjusted p -value < 0.005).

14 **Figure 6** Effects of Dex on long RNA payload of sperm cells residing in testis at time of
15 administration. (A) Experimental design depicting a time window of two weeks between
16 injection of Dex and sperm collection for molecular analysis.

17 (B) Volcano plot depicting fold changes and significance level of long RNA in mature sperm
18 14 days post injection of Dex ($n=4$) versus vehicle ($n=4$) as assessed by small RNA
19 sequencing. Statistically significant transcripts are highlighted in red (FDR < 0.05). (C)

20 Heatmap showing significantly differentially expressed long RNA transcripts of the same
21 experiment (multiple comparison corrected, $q < 0.05$). (D) Q-PCR results of CircRNA assays for

22 caput and cauda sperm for Tasp1 (interaction treatment x tissue $F(1,5)=7.53$, $p < 0.05$, caput:
23 control $n=4$, DEX $n=5$, $p > 0.05$; cauda control $n=5$, DEX $n=3$, $p < 0.01$) (* $p < 0.05$ mixed effect
24 model interaction (REML), ** $q < 0.01$ multiple comparisons SIDAK corrected) and Dennd1

25 (interaction treatment x tissue $F(1,15)=6$, $p < 0.05$, caput: control $n=4$, DEX $n=5$, $p > 0.05$; cauda
26 control $n=5$, DEX $n=5$, $p < 0.05$) (E). Whiskers display minimum and maximum. * $p < 0.05$ mixed
27 effect model interaction (REML), * $q < 0.05$ multiple comparisons SIDAK corrected.

28

1 **Table 1 Changes in miRNAs and CircRNAs across time and sample type**

Timepoint post injection	3 hours	7 days	14 days		
miRNAs	Cauda	Cauda	Cauda	Caput	Serum
3535	no ch.	no ch.	↑	not det	not det
677-5p	no ch.	no ch.	↑	no ch.	not det
677-3p	no ch.	no ch.	↑		
34c-5p	no ch.	no ch.	↓	no ch.	no ch.
7033-5p	no ch.	no ch.	↑		
703	no ch.	no ch.	↑		
5126	no ch.	no ch.	↑		
5121	no ch.	no ch.	↑		
6240	no ch.	no ch.	↑		
5114	no ch.	no ch.	↑		
1839-3p	no ch.	no ch.	↑		
1949	no ch.	no ch.	↑		
196a-5p	no ch.	no ch.	↑		
3064-5p	no ch.	no ch.	↑		
196b-5p	no ch.	no ch.	↑		
6538	no ch.	no ch.	↑	no ch.	no ch.
18a-5p	no ch.	no ch.	↓		
3963	no ch.	no ch.	↓		
471-5p	no ch.	no ch.	↓		
5099	no ch.	no ch.	↑		
1843b-3p	no ch.	no ch.	↑		
1843a-3p	no ch.	no ch.	↑		
circRNAs					
Tasp1			↑	no ch.	
Dennd1b			↑	no ch.	

2
3
4
5

Table 1: Table depicts significantly altered small RNAs and circRNAs in cauda sperm at 14 days post dexamethasone injection.

1 **Table 2** Changes in tsRNAs across time and sample type

Timepoint post injection	3 hours		7 days	14 days		
	Cauda	Caput	Cauda	Cauda	Caput	Serum
Leu-CAA	↑		↑	no ch.	no ch	
<i>Thr-TGT1</i>	↓		↑	no ch.		
<i>His-GTG-2</i>	no ch		↓	no ch	no ch	no ch.
<i>His-GTG-3</i>	no ch		↑	no ch.		
<i>Pro-AGG-1</i>	no ch		↑	no ch.		
<i>Pro-TGG-2</i>	no ch		↑	no ch.		
<i>Pro-TGG-4</i>	no ch		↑	no ch.		
<i>Glu-CTC-4</i>	no ch		↑	no ch.		
<i>Pro-CGG-1</i>	no ch		↑	no ch.		
<i>Gly-GCC-6</i>	no ch		↑	no ch		
Gly-GCC-2	no ch		no ch	↓	no ch	no ch.
Arg-CCT-2	↑	no ch.	↑	↑	↑	↑
Ala-TGC-2	no ch		no ch	↓		
Arg-ACG-3	no ch		no ch	↓		
Glu-CTC-3	no ch		no ch	↓		
Ser-TGA-1	no ch		no ch	↓		
Thr-TGT-2	no ch		no ch	↑		
Ser-TGA-2	no ch		no ch	↑		
Ser-AGA-1	no ch		no ch	↑		
Glu-TTC-3	no ch		no ch	↓		
Arg-CCT-1	no ch		no ch	↑		
Ser-AGA-2	no ch		no ch	↑		
Cys-GCA-3	no ch		no ch	↓		
Pro-AGG-3	no ch		no ch	↑		
Lys-CTT-3	no ch		no ch	↓		
Arg-CCT-3	no ch		no ch	↑		
Lys-CTT-3	no ch		no ch	↓		
Trp-CCA-5	no ch		no ch	↓		
Cys-GCA-2	no ch		no ch	↓		
Ala-TGC-5	no ch		no ch	↓		
Gly-CCC-3	no ch		no ch	↓		
Gly-CCC-4	no ch		no ch	↓		
Ala-TGC-5	no ch		no ch	↓		
Thr-AGT-5	no ch		no ch	↑		
SeC-TCA-1	no ch		no ch	↓		
Thr-AGT-7	no ch		no ch	↑		
Glu-CTC-2	no ch		no ch	↓		
His-GTG-1	no ch		no ch	↑		
Arg-CCT-4	no ch		no ch	↑		
Lys-CTT-1	no ch		no ch	↓		
Lys-CTT-2	no ch		no ch	↓		
Lys-CTT-2	no ch		no ch	↓		
Asp-GTC-4	no ch		no ch	↓		
Gly-GCC-5	no ch		no ch	↓		
Thr-TGT-3	no ch		no ch	↑		
Asn-GTT-2	no ch		no ch	↑		
Asn-GTT-4	no ch		no ch	↑		
Asn-GTT-1	no ch		no ch	↑		
Gly-GCC-4	no ch		no ch	↓		
Gly-GCC-2	no ch		no ch	↓		
Thr-CGT-4	no ch		no ch	↑		
Gly-CCC-5	no ch		no ch	↓		
Ala-TGC-6	no ch		no ch	↓		
Glu-CTC-1	no ch		no ch	↓		

1
2 **Table 2:** No highlight: Significantly altered in cauda sperm at 14 days post dexamethasone
3 injection. In bold: Persistently altered in cauda sperm at 7 days and 3 hours. Italic: Significant
4 interaction between 7 days and 3 hours.

- 1 **STAR METHODS**
- 2 Detailed methods are provided in the online version of this paper and include the following:
- 3 **KEY RESOURCES TABLE**
- 4 **RESOURCE AVAILABILITY**
 - 5 ○ Lead contact
 - 6 ○ Materials availability
 - 7 ○ Data and code availability
- 8 **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
- 9 ○ Animals
- 10 **METHOD DETAILS**
 - 11 ○ *Dex treatment and sample collection*
 - 12 ○ *In vitro fertilization and embryo culture*
 - 13 ○ *Sperm and embryo RNA extraction*
 - 14 ○ *Serum exosome RNA extraction*
 - 15 ○ *Sperm RNA sequencing (RNAseq)*
 - 16 ○ *Single embryo seq*
 - 17 ○ *Insulin and Glucose tolerance test*
 - 18 ○ *Body mass index*
 - 19 ○ *Small RNA q-RT-PCR*
 - 20 ○ *CircRNA q-RT-PCR*
- 21 **QUANTIFICATION AND STATISTICAL ANALYSIS**
 - 22 ○ Bioinformatic analysis
 - 23 ○ *Bulk RNA sequencing*
 - 24 ○ *2-cell single embryo sequencing analysis*
 - 25 ○ Remaining Statistical analysis

1 **STAR METHODS**

2 RESOURCE AVAILABILITY

3 *Lead contact*

4 Further information and requests for resources and reagents should be directed to, and will be
5 fulfilled upon reasonable request, by the Lead Contact, Katharina Gapp
6 (katharina.gapp@hest.ethz.ch).

7

8 *Materials availability*

9 The study did not generate new reagents.

10

11 *Data and code availability*

12 - The raw datasets supporting the conclusions of this article are included within the
13 article (supplementary tables).

14 - Sequencing data have been deposited at Gene omnibus and ENA and are publicly available
15 as of the date of publication. All sequencing data were deposited to Gene Omnibus and
16 ENA.

17 - All code has been deposited and is publicly available on Github.

18 Accession numbers are listed in the key resources table.

19

20 EXPERIMENTAL MODEL AND SUBJECT DETAILS

21 *Animals*

22 C57Bl/6 mice were obtained from the Sanger Research support facility in-house-breeding
23 colony. They were housed in a temperature and humidity-controlled facility in individually
24 ventilated cages under a non-reversed light-dark cycle (Sanger Research support facility) or a
25 reversed-light-dark cycle (ETH EPIC). Standard chow (LabDiet(r) 5021-3 supplied by IPS) and
26 water were provided *ad libitum after weaning* unless stated otherwise (e.g. oocyte donors).

1 Breeding colony was provided SAFE R03-10 breeding diet, supplied by SAFE diets.
2 Experimental procedures were performed during the animals' inactive cycle at Sanger. Age
3 and weight matched (margin of one week) males were used in each experimental group
4 receiving Dex injections. Animals used for Dex injection followed by sperm sequencing were
5 all sexually mature (14 and 7 days or 3 hours post treatment were 13, 11 and 9 weeks of age
6 respectively) at the time of sperm collection.

7 C57Bl/6 males used for sperm sequencing 14 days post Dex injection and q-PCR
8 experiments/validation 3 hours post Dex were obtained from the ETH's EPIC in house
9 breeding colony in Zürich and were 14-18 weeks old. These mice were fed chow #3734 by
10 Kliba/Granovit.

11 IVF oocyte donor females and embryo recipients were fed SAFE R03-10 breeding diet,
12 supplied by SAFE diets until 10 days post embryo transfer. Until this time embryo recipients
13 were housed in pairs after which they were split into single housing. IVF offspring was weaned
14 at PND21 and assigned to cages avoiding littermate cohousing. Offspring phenotyping was
15 carried out between 3.5 to 4 months and necropsy at 4.5 months of age in balanced (offspring
16 controls, offspring treatment) and age matched groups (all animals had an age spread of 3
17 days). Animals were housed in groups of 4-5 mice/cage in the Sager Institute barrier research
18 support facility (all animals apart from animals for q-RT-PCR experiments) and ETHZ's EPIC
19 facility (animals for q-RT-PCR).

20 All experiments were approved by the UK home office (project license P176396F2) and
21 Cantonal commission for animal experimentation Zürich (project license ZH222/19).

22

23 METHOD DETAILS

24 *Dex treatment and sample collection*

25 Age matched males with an age spread of 1 week were randomly assigned to control and
26 treatment groups. Males were injected with either 2mg/kg of Dex in 10 % DMSO, 0.9% saline
27 or vehicle (10%DMSO in 0.9% saline). Males used for sperm collection did not undergo any
28 metabolic testing. They were sacrificed 2 weeks, 7 days and 3 hours after Dex or vehicle

1 treatment. Cauda epididymis and vas deference were dissected and placed in M2 medium.
2 After allowing sperm to diffuse into M2 medium, cells were pelleted by short centrifugation and
3 washed with PBS. For sperm RNA sequencing and q-PCR, mature sperm cells were separated
4 from potential somatic contamination by somatic lysis, followed by 2 washes with PBS
5 (Brykczynska *et al.*, 2010). Sperm counts and fertilization rate appeared unaffected post Dex
6 injection (Supplementary Fig. 6A,B).

7

8 *In vitro fertilization and embryo culture*

9 12 randomly selected, C57BL/6 females were superovulated at 26-31 days of age with Card
10 Hyperova (Cosmo Bio, KYD-010-EX-X5), followed by 7.5 IU human chorionic gonadotrophin
11 (HCG) 48 hours later.

12 Cumulus-oocyte complexes (COCs) were released from the ampulla of the oviduct 16-17
13 hours after HCG administration, and preincubated in high calcium HTF with Glutathione
14 medium for 30-60 minutes (in CO₂ incubator at 37 deg C, 5% CO₂ in air) before insemination.
15 Frozen sperm used for insemination was pooled from 2 males that had been injected with Dex
16 or vehicle 14 days prior to cryopreservation. Thawed sperm was preincubated for 30 minutes
17 in TYH (with Methyl- β -cyclodextrin, Sigma C4555) medium at 37 deg C, 5% CO₂ in air, before
18 being added to the COC complexes for fertilisation. 4 hours after insemination, the presumptive
19 zygotes were washed through several drops of KSOM (Millipore, MR-121-D) and incubated
20 overnight in KSOM.

21 For in vivo offspring, 14-20 x 2 cell embryos from overnight culture in 6 individual IVF dishes
22 /group were implanted into 0.5 dpc pseudo-pregnant F1 females (6 females/group). Each dish
23 contained oocytes from one female with the exception of 2 dishes (out of 6) in the Dex group
24 that contained oocytes of the same female, since one female failed to super-ovulate. For
25 molecular (single) embryo gene expression analysis at the two cell stage, 2-cell-embryos from
26 overnight culture were frozen, and after thawing briefly cultured in preincubated KSOM
27 until/during plating into 96 well plates. The females used to generate these embryos were
28 superovulated with PMSG. The IVF Protocol is based on EMMA Harwell's protocol (adapted

1 from Takeo & Nakagata 2011(Nakagata, 2011)), and the Sperm Freeze Protocol is based on
2 Ostermeier G.C. et al (2008)(Ostermeier *et al.*, 2008). Resulting litter sizes did not differ
3 between Vehicle and Dex injected offspring (Supplementary Fig. 6C)

4 5 *Sperm and embryo RNA extraction*

6 Total RNA was prepared from adult mouse sperm using Trizol (Thermo Scientific 15596026)
7 and Directzol (Zymo R2080). Total RNA was prepared from zygotes using the Trizol LS
8 protocol. Quantity and purity of RNA were determined by Agilent 2100 Bioanalyser (Agilent
9 Technologies) and Qubit fluorometer (Life Technologies). Absence of prominent ribosomal
10 peaks indicated absence of somatic cell contamination.

11 12 *Serum Exosome RNA extraction*

13 Trunk blood was collected from animals following cervical dislocation and stored at room
14 temperature for 30 minutes to allow coagulation. Serum was subsequently separated by 2
15 centrifugation steps first for 10 min followed by 15 minutes at 3000. g. 100 ul of serum were
16 used as input for exosomal isolation following the manufacturers instructions (exoRNeasy
17 Qiagen). 2 ul of. 25 ul RNA eluate were used as input for cDNA conversion with the
18 miRCURY® LNA® cDNA conversion kit.

19 20 21 *Sperm RNA sequencing (RNAseq)*

22 Sequencing was done using an Illumina Genome Analyzer HiSeq 2500 (Illumina) in Rapid run
23 mode for long 100bp and small 50 bp RNA sequencing runs respectively.

24 Libraries for long RNA sequencing were prepared using the TruSeq Stranded Total RNA kit
25 according to the manufacturer's instructions with indices diluted at 1:3. 200 ng of total sperm
26 RNA was subjected to removal of rRNA using Ribozero gold kit. Approximately 100ng of sperm
27 RNA and total RNA of several 2-cell zygotes was subjected to TruSeq or Nextflex (sperm 14
28 days post injection) small RNA library preparation following the manufacturer's

1 recommendations with the following modifications: adaptors were diluted 1:4 and PCR cycles
2 were augmented to 18 and 22 (Nextflex) PCR cycles respectively. When library preparation of
3 samples was split across days groups were balanced to circumvent batch effects.

4

5 *Single embryo seq.*

6 2 cell embryos were generated using the same conditions as indicated for in vivo offspring yet
7 followed by embryo cryopreservation until processing for library preparation. They were
8 thawed and those that appeared intact (34 controls and 37 Dex) pipetted into wells of 2 96 well
9 culture plates containing lysis buffer and stored at -80°C before processing according to the
10 Smartseq 2 protocol and manufacturer's recommendations (Nextera). Libraries contained a
11 1:19 Million dilution of External RNA Controls Consortium (ERCC) spike-ins (4456740 Ambion)
12 and were amplified for 18 PCR cycles. Sequencing was performed on a HiSeq V4 under paired
13 end 75bp mode.

14

15 *Insulin and Glucose tolerance test*

16 Animals were fasted 4 hours to establish a shared baseline glucose level. They received a
17 single injection of insulin (insulin: 1mU/g body weight) (Actrapid Novo Nordisk), glucose (2mg/g
18 body weight) or vehicle (saline) intraperitoneally. Blood samples were taken from lateral tail
19 vein in adult animals to assess blood glucose level using an Accucueck aviva device.

20

21 *Body mass index*

22 Animal lengths were measured using a standard ruler and weighed for assessing body weight.
23 Body mass index was calculated using the following formula: $\text{weight (g)} / (\text{length (cm)}^2)$.

24

25 *Necropsy*

26 Organs were dissected after sacrifice and weighed immediately on a scale using "g" as a unit
27 with an accuracy of 2 decimals (accurate down to 10 mg).

28

1 *Small RNA q-RT-PCR*

2 5ng/sample RNA isolated from sperm was reverse transcribed (RT) using the miCURY LNA
3 RT kit (Qiagen #339340). Quantitative RT-PCR (qRT-PCR) was performed using SYBR green
4 based detection in a Biorad thermal cycler with MiRCURY LNA-based small RNA probes
5 designed against tRNA ArgCCT-2, Gly-GCC-2, His-GTG-2, Leu-CAA-2, with a polyA tail
6 directed reverse miRCURY primer (Qiagen # 339317). RnU6 was used as an internal control
7 in sperm samples and mir-103a-3p in serum samples (Qiagen # 339306).

8

9 *CircRNA q-RT-PCR*

10 100ng/sample RNA isolated from sperm was converted into cDNA using random hexamers.
11 Primers were designed to span the exon splice junctions. Primer sequences for CircTasp1 and
12 CircDennd1b are depicted in the resource table. *Tubulin1* was used as endogenous control in
13 the Sybr Green based quantification.

14

15 QUANTIFICATION AND STATISTICAL ANALYSIS

16 *Bioinformatic analysis*

17 *Bulk RNA sequencing*

18 Each sequencing library represented sperm harvested from a single male. Sequencing
19 quality was assessed with FastQC (Andrews, 2010) and MultiQC (Ewels *et al.*, 2016).
20 Adapters were removed from the 3' ends with cutadapt (Martin, 2011) (version 1.14) and
21 resulting sequences with 14 nucleotides of length or less were discarded. All other reads
22 were aligned end to end (no soft clipping) to the ENSEMBL *Mus musculus* genome (release
23 75) (Flicek *et al.*, 2014) with STAR (Dobin *et al.*, 2013). No mismatches were allowed.
24 Featurecounts was used to match the alignments against the miRbase (Kozomara and
25 Griffiths-Jones, 2011) annotation (version 21) and obtain a matrix of miRNA counts. We
26 applied fractional counts whenever alignment occurred at multiple genomic locations.
27 Differential expression was analyzed using DESeq2 (Love, Huber and Anders, 2014).
28 Quantification of tRNA fragments was performed as above, but all CCA-3' trinucleotides were

1 trimmed after adapter removal, sequences with 15 nucleotides or less were subsequently
2 discarded and GtRNADB (Chan and Lowe, 2016) annotation (GRCm38/mm10) was used to
3 obtain the count matrix. Quantification of rRNA fragments was performed using SPORTS(Shi
4 *et al.*, 2018) on the precompiled database included in the tool.

5

6 For the data set collected 14 days after Dex injection, library preparation included the insertion
7 of 2 random tetranucleotides between read and adapters. By including only unique sequences
8 in the analysis we removed duplicates due to PCR amplification.

9 Long RNAseq libraries were pre-processed with trimmomatic (Bolger, Lohse and Usadel,
10 2014) to remove adapters. Reads were aligned to the genome using STAR (Dobin *et al.*,
11 2013) and quantified using featurecounts (Liao, Smyth and Shi, 2014). Circular RNAs were
12 quantified using Circexplorer2 (Zhang *et al.*, 2016) based on junction reads as detected by
13 STAR. Differential expression analysis was performed on the combined set of counts for
14 circular and non-circular RNAs using edgeR (Anders *et al.*, 2013). Robust estimation of
15 dispersion was used to avoid spurious significance due to outliers.

16

17 *2-cell single embryo sequencing analysis*

18 Reads from 2 cell embryos were mapped to the mouse reference genome (mm10) and ERCC
19 spike-ins using STAR (Dobin *et al.*, 2013). Resultant alignments were processed to quantify
20 the expression of annotated genes by GENCODE (vM11) and ERCC spike-ins using
21 featureCounts (Liao, Smyth and Shi, 2014). To filter low-quality sequenced embryos we only
22 considered those which had a total read count of at least 0.5 million reads with less than 15%
23 and 10% their read counts mapping to mitochondrial genes and ERCC spike-ins respectively.
24 After these filters were applied a total of 56 embryos (29 controls and 27 treated) remained.
25 We clustered their gene expression profiles using SC3 (Kiselev *et al.*, 2017) obtaining two main

1 clusters (C1 and C2). Using scmap (Kiselev, Yiu and Hemberg, 2018), we projected the gene
2 expression profiles for the two cell embryos onto an index containing expression profiles from
3 zygotic, early/mid/late 2 cell embryos and 4 cell embryo cells reported by Deng *et al.* (Deng *et*
4 *al.*, 2014). We performed PCA analyses using scater (McCarthy *et al.*, 2017) (runPCA function)
5 and we calculated the PC1 silhouette coefficient using in-house R scripts. To perform
6 differential gene expression analyses we normalized the read counts of each embryo as FPKM
7 and we used Census (Qiu *et al.*, 2017) algorithm to convert these values into relative transcripts
8 counts. We computed the obtained 'Census counts' using Monocle (v 2.99.2), assuming a
9 negative binomial distribution and a lower detection limit of 0.5. We performed differential gene
10 expression analyses between the total treated and control embryos, and also between the
11 treated and control embryos inside of C1 and C2 clusters.

12

13 *Remaining statistical analyses.*

14 Sample size for in vivo offspring phenotyping was estimated based on previous work on similar
15 models (Hoyer *et al.*, 2013; Bohacek, von Werdt and Mansuy, 2016). 3-Way repeated
16 measures ANOVA was used to assess statistical significance for BMI, GTT and ITT
17 measurements. Necropsy data were analysed using 3-way ANOVA followed by multiple t-tests
18 and corrected for multiple comparisons using the Benjamini-Hochberg method. Normality was
19 assessed with the Kolmogorov Smirnov test and met in all necropsy data. Homogeneity of
20 variances was assessed and met in all necropsy data unless gonadal WAT. These t-tests did
21 not assume homogeneity of variances (applied Welch's correction). Q-RT-PCR results
22 comparing caput and cauda sperm RNA were analyzed by fitting a Mixed model followed by
23 posthoc tests to compare individual groups applying the Sidak correction for multiple
24 comparisons. Other q-PCRs were analysed using multiple t-tests corrected for multiple
25 comparisons applying the two stage step up method by Benjamini, Krieger and Yekutieli. All
26 reported replicates were biological replicates. Significance was set at $p < 0.05$ and where
27 applicable $q < 0.05$ for all tests. All statistics of behavioural, metabolic tests and q-RT-PCR were

1 computed with Prism. Outliers were removed from q-PCR results using Prism's inbuilt ROUT
2 method and are depicted in the supplementary tables containing raw data with a star.

3

4

5 **Supplementary Material**

6 This article contains supplementary Figures and tables. Supplementary figures and legends
7 are compiled in one document.

8

9 **Supplementary table legend**

10

11 *Supplementary Table 1 related to Figure 1: Data and analysis of sperm small RNA 14 days*
12 *post Dex*

13 Sheet 1: List of normalized miRNA read counts of sperm harvested 14 days post Dex and
14 vehicle injection.

15 Sheet 2: Deseq2 results of a comparison between miRNA from sperm harvested 14 days
16 post Dex and vehicle injection.

17 Sheet 3: List of normalized tsRNA read counts of sperm harvested 14 days post Dex and
18 vehicle injection.

19 Sheet 4: Deseq2 results of a comparison between tsRNAs from sperm harvested 14 days
20 post Dex and vehicle injection.

21 Sheet 5: List of normalized read counts for rRNA from sperm harvested 14 days post Dex
22 and vehicle injection.

23 Sheet 6: Deseq2 results for rRNA quantification from sperm harvested 14 days post Dex and
24 vehicle injection.

25 Sheet 8: Raw values of qRT-PCR analysis for serum.

26 Sheet 9: Raw values of qRT-PCR analysis of caput sperm.

27

28

1 *Supplementary table 2 related to Figure 2: Data and analysis of sperm small RNA 3 hours*
2 *and 7 days post Dex*

3 Sheet 1: Deseq2 results of the comparison of tsRNAs in sperm harvested 3 hours and 7
4 days post Dex with tsRNAs in sperm harvested 3 hours and 7 days post vehicle injection.

5 Sheet 2: Deseq2 results of the comparison between tsRNAs from sperm harvested post Dex
6 and vehicle injection at 3 hours with tsRNAs in sperm harvested 7 days post Dex and vehicle
7 injection.

8 Sheet 3: Deseq2 results of the interaction between treatment and time for tsRNAs from
9 sperm harvested 3 hours and 7 days post Dex and vehicle injection.

10 Sheet 4: Deseq2 results of the comparison of miRNAs in sperm harvested 3 hours and 7
11 days post Dex with miRNAs in sperm harvested 3 hours and 7 days post vehicle injection.

12 Sheet 5: Deseq2 results of the comparison between miRNAs from sperm harvested post Dex
13 and vehicle injection at 3 hours with miRNAs in sperm harvested 7 days post Dex and
14 vehicle injection.

15 Sheet 6: Deseq2 results of the interaction between treatment and time for miRNAs from
16 sperm harvested 3 hours and 7 days post Dex and vehicle injection.

17 Sheet 7: Raw values of qRT-PCR analysis for tsRNA-Arg-CCT-2 in caput and cauda sperm
18 sampled 3 hours post Dex and vehicle injection.

19

20 *Supplementary table 3 related to Figure 3: Raw data for Dex offspring phenotyping.*

21 Raw data for BMI (sheet 1), GTT (sheet 2) , ITT (sheet 3) and necropsy weights (sheet 4) of
22 adult offspring animals resulting from IVF of wildtype oocytes and sperm harvested 14 days
23 post Dex and vehicle injection.

24

25 *Supplementary table 4 related to Figure 5: Single embryo RNA sequencing analysis.*

26 Sheet 1: Monocle output list of significantly differentially regulated genes between 2-cell
27 embryos resulting from IVF of wildtype oocytes and sperm harvested 14 days post Dex and
28 vehicle injection.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

Supplementary table 5 related to Figure 6: Data and analysis of sperm long RNA 14 days post Dex

Sheet 1: List of normalized long RNA seq counts of sperm harvested 14 days post Dex and vehicle injection.

Sheet 2: EdgeR results of a comparison between long RNA reads from sperm harvested 14 days post Dex and vehicle injection.

Sheet 3: Raw values of qRT-PCR analysis for caput and cauda sperm circRNAs.

References

Agarwal, V., Bell, G. W., Nam, J.-W. and Bartel, D. P. (2015) 'Predicting effective microRNA target sites in mammalian mRNAs', *eLife*, 4. doi: 10.7554/eLife.05005.

Amanai, M., Brahmajosyula, M. and Perry, A. C. F. (2006) 'A restricted role for sperm-borne microRNAs in mammalian fertilization.', *Biology of reproduction*, 75(6), pp. 877–884. doi: 10.1095/biolreprod.106.056499.

Anders, S., McCarthy, D. J., Chen, Y., Okoniewski, M., Smyth, G. K., Huber, W. and Robinson, M. D. (2013) 'Count-based differential expression analysis of RNA sequencing data using R and Bioconductor', *Nature Protocols*, 8(9), pp. 1765–1786. doi: 10.1038/nprot.2013.099.

Andrews, S. (2010) 'FastQC: a quality control tool for high throughput sequence data.'

Ashe, A., Sapetschnig, A., Weick, E. M., Mitchell, J., Bagijn, M. P., Cording, A. C., Doebley, A. L., Goldstein, L. D., Lehrbach, N. J., Le Pen, J., *et al.* (2012) 'piRNAs can trigger a multigenerational epigenetic memory in the germline of *C. elegans*', *Cell*. 2012/06/29, 150(1), pp. 88–99. doi: 10.1016/j.cell.2012.06.018.

Barden, N. (1999) 'Regulation of corticosteroid receptor gene expression in depression and antidepressant action.', *Journal of psychiatry & neuroscience : JPN*, 24(1), pp. 25–39. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9987205>.

Barrett, S. P. and Salzman, J. (2016) 'Circular RNAs: analysis, expression and potential functions', *Development*, 143(11), pp. 1838–1847. doi: 10.1242/DEV.128074.

1 Benito, E., Kerimoglu, C., Ramachandran, B., Pena-Centeno, T., Jain, G., Stilling, R. M., Islam, M. R.,
2 Capece, V., Zhou, Q., Edbauer, D., *et al.* (2018) 'RNA-Dependent Intergenerational Inheritance of
3 Enhanced Synaptic Plasticity after Environmental Enrichment', *Cell Reports*, 23(2), pp. 546–554. doi:
4 10.1016/j.celrep.2018.03.059.

5 Benyshek, D. C., Johnston, C. S. and Martin, J. F. (2006) 'Glucose metabolism is altered in the
6 adequately-nourished grand-offspring (F3 generation) of rats malnourished during gestation and
7 perinatal life', *Diabetologia*. 2006/03/25, 49(5), pp. 1117–1119. doi: 10.1007/s00125-006-0196-5.

8 Bera, S., Greiner, S., Choudhury, A., Dispenzieri, A., Spitz, D. R., Russell, S. J. and Goel, A. (2010)
9 'Dexamethasone-induced oxidative stress enhances myeloma cell radiosensitization while sparing
10 normal bone marrow hematopoiesis', *Neoplasia*, 12(12), pp. 980–992. doi: 10.1593/neo.101146.

11 'Biggest COVID-19 trial tests repurposed drugs first' (2020) *Nature Biotechnology*, 38(5), p. 510. doi:
12 10.1038/s41587-020-0528-x.

13 Bohacek, J. and Mansuy, I. M. (2017) 'A guide to designing germline-dependent epigenetic
14 inheritance experiments in mammals.', *Nature methods*, 14(3), pp. 243–249. doi:
15 10.1038/nmeth.4181.

16 Bohacek, J. and Rassoulzadegan, M. (2019) 'Sperm RNA: Quo vadis?', *Seminars in Cell &*
17 *Developmental Biology*. doi: 10.1016/J.SEMCDB.2019.07.005.

18 Bohacek, J., von Werdt, S. and Mansuy, I. M. (2016) 'Probing the germline-dependence of epigenetic
19 inheritance using artificial insemination in mice', *Environmental Epigenetics*, 2(1), p. dvv015. doi:
20 10.1093/eep/dvv015.

21 Bolger, A. M., Lohse, M. and Usadel, B. (2014) 'Trimmomatic: a flexible trimmer for Illumina sequence
22 data.', *Bioinformatics (Oxford, England)*, 30(15), pp. 2114–2120. doi: 10.1093/bioinformatics/btu170.

23 Bönisch, C., Irmeler, M., Brachthäuser, L., Neff, F., Bamberger, M. T., Marschall, S., Hrabě de Angelis,
24 M. and Beckers, J. (2016) 'Dexamethasone treatment alters insulin, leptin, and adiponectin levels in
25 male mice as observed in DIO but does not lead to alterations of metabolic phenotypes in the
26 offspring', *Mammalian Genome*, 27(1–2), pp. 17–28. doi: 10.1007/s00335-015-9616-5.

27 Bowers, M. E. and Yehuda, R. (2016) 'Intergenerational transmission of stress in humans',
28 *Neuropsychopharmacology*, pp. 232–244. doi: 10.1038/npp.2015.247.

29 Brykczynska, U., Hisano, M., Erkek, S., Ramos, L., Oakeley, E. J., Roloff, T. C., Beisel, C., Schubeler,
30 D., Stadler, M. B. and Peters, A. H. F. M. (2010) 'Repressive and active histone methylation mark
31 distinct promoters in human and mouse spermatozoa', *Nat Struct Mol Biol*, 17(6), pp. 679–687. doi:

1 http://www.nature.com/nsmb/journal/v17/n6/suppinfo/nsmb.1821_S1.html.

2 Cacciagli, P., Sutera-Sardo, J., Borges-Correia, A., Roux, J.-C., Dorboz, I., Desvignes, J.-P., Badens,
3 C., Delepine, M., Lathrop, M., Cau, P., *et al.* (2013) 'Mutations in BCAP31 Cause a Severe X-Linked
4 Phenotype with Deafness, Dystonia, and Central Hypomyelination and Disorganize the Golgi
5 Apparatus', *The American Journal of Human Genetics*, 93(3), pp. 579–586. doi:
6 10.1016/j.ajhg.2013.07.023.

7 Carone, B. R., Fauquier, L., Habib, N., Shea, J. M., Hart, C. E., Li, R., Bock, C., Li, C., Gu, H.,
8 Zamore, P. D., *et al.* (2010) 'Paternaly induced transgenerational environmental reprogramming of
9 metabolic gene expression in mammals.', *Cell*, 143(7), pp. 1084–1096. doi:
10 10.1016/j.cell.2010.12.008.

11 Cartier, J., Smith, T., Thomson, J. P., Rose, C. M., Khulan, B., Heger, A., Meehan, R. R. and Drake,
12 A. J. (2018) 'Investigation into the role of the germline epigenome in the transmission of
13 glucocorticoid-programmed effects across generations.', *Genome biology*, 19(1), p. 50. doi:
14 10.1186/s13059-018-1422-4.

15 Chan, J. C., Morgan, C. P., Adrian Leu, N., Shetty, A., Cisse, Y. M., Nugent, B. M., Morrison, K. E.,
16 Jašarević, E., Huang, W., Kanyuch, N., *et al.* (2020) 'Reproductive tract extracellular vesicles are
17 sufficient to transmit intergenerational stress and program neurodevelopment', *Nature*
18 *Communications*, 11(1), p. 1499. doi: 10.1038/s41467-020-15305-w.

19 Chan, P. P. and Lowe, T. M. (2016) 'GtRNADB 2.0: an expanded database of transfer RNA genes
20 identified in complete and draft genomes.', *Nucleic acids research*, 44(D1), pp. D184–9. doi:
21 10.1093/nar/gkv1309.

22 Chen, Q., Yan, M., Cao, Z., Li, X., Zhang, Yunfang, Shi, J., Feng, G. H., Peng, H., Zhang, X., Zhang,
23 Ying, *et al.* (2016) 'Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic
24 disorder', *Science*, 351(6271), pp. 397–400. doi: 10.1126/science.aad7977.

25 Chen, Q., Yan, W. and Duan, E. (2016) 'Epigenetic inheritance of acquired traits through sperm RNAs
26 and sperm RNA modifications.', *Nature reviews. Genetics*, 17(12), pp. 733–743. doi:
27 10.1038/nrg.2016.106.

28 Chioccarelli, T., Manfredola, F., Ferraro, B., Sellitto, C., Cobellis, G., Migliaccio, M., Fasano, S.,
29 Pierantoni, R. and Chianese, R. (2019) 'Expression patterns of circular RNAs in high quality and poor
30 quality human spermatozoa', *Frontiers in Endocrinology*, 10(JUL), pp. 1–14. doi:
31 10.3389/fendo.2019.00435.

1 Conine, C. C., Sun, F., Song, L., Rivera-Pérez, J. A. and Rando, O. J. (2018) 'Small RNAs Gained
2 during Epididymal Transit of Sperm Are Essential for Embryonic Development in Mice.',
3 *Developmental cell*, 46(4), pp. 470–480.e3. doi: 10.1016/j.devcel.2018.06.024.

4 Cross, K. P., Paul, R. I. and Goldman, R. D. (2011) 'Single-dose dexamethasone for mild-to-moderate
5 asthma exacerbations: effective, easy, and acceptable.', *Canadian family physician Medecin de
6 famille canadien*, 57(10), pp. 1134–1136. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21998227>.

7 Dadoune, J. and Alfonsi, M. (1984) 'Autoradiographic investigation of sperm transit through the male
8 mouse genital tract after tritiated thymidine incorporation', *Reproduction Nutrition and Development*,
9 24(6), pp. 927–935. doi: 10.1051/rnd:19840709.

10 Dard-Dascot, C., Naquin, D., D'Aubenton-Carafa, Y., Alix, K., Thermes, C. and van Dijk, E. (2018)
11 'Systematic comparison of small RNA library preparation protocols for next-generation sequencing',
12 *BMC Genomics*, 19(1), p. 118. doi: 10.1186/s12864-018-4491-6.

13 Daskalakis, N. P., Oitzl, M. S., Schachinger, H., Champagne, D. L. and de Kloet, E. R. (2012) 'Testing
14 the cumulative stress and mismatch hypotheses of psychopathology in a rat model of early-life
15 adversity', *Physiol Behav.* 2012/02/07, 106(5), pp. 707–721. doi: 10.1016/j.physbeh.2012.01.015.

16 Deevy, O. and Bracken, A. P. (2019) 'PRC2 functions in development and congenital disorders',
17 *Development*, 146(19). doi: 10.1242/DEV.181354.

18 Deng, Q., Ramsköld, D., Reinius, B. and Sandberg, R. (2014) 'Single-Cell RNA-Seq Reveals
19 Dynamic, Random Monoallelic Gene Expression in Mammalian Cells', *Science*, 343(6167), pp. 193–
20 196. doi: 10.1126/SCIENCE.1245316.

21 Dietz, D. M., Laplant, Q., Watts, E. L., Hodes, G. E., Russo, S. J., Feng, J., Oosting, R. S., Vialou, V.
22 and Nestler, E. J. (2011) 'Paternal transmission of stress-induced pathologies.', *Biological psychiatry*,
23 70(5), pp. 408–414. doi: 10.1016/j.biopsych.2011.05.005.

24 Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M. and
25 Gingeras, T. R. (2013) 'STAR: ultrafast universal RNA-seq aligner.', *Bioinformatics (Oxford, England)*,
26 29(1), pp. 15–21. doi: 10.1093/bioinformatics/bts635.

27 EU Clinical Trials Register (no date) 'Randomised Evaluation of COVID-19 Therapy (RECOVERY)'.
28 Available at: <https://www.clinicaltrialsregister.eu/ctr-search/trial/2020-001113-21/GB> (Accessed: 7 July
29 2020).

30 Ewels, P., Magnusson, M., Lundin, S. and Käller, M. (2016) 'MultiQC: summarize analysis results for
31 multiple tools and samples in a single report', *Bioinformatics*, 32(19), pp. 3047–3048. doi:

1 10.1093/bioinformatics/btw354.

2 Flicek, P., Amode, M. R., Barrell, D., Beal, K., Billis, K., Brent, S., Carvalho-Silva, D., Clapham, P.,
3 Coates, G., Fitzgerald, S., *et al.* (2014) 'Ensembl 2014.', *Nucleic acids research*, 42(Database issue),
4 pp. D749--55. doi: 10.1093/nar/gkt1196.

5 Franklin, T. B., Linder, N., Russig, H., Thöny, B. and Mansuy, I. M. (2011) 'Influence of early stress on
6 social abilities and serotonergic functions across generations in mice.', *PloS one*. Edited by C. Combs,
7 6(7), p. e21842. doi: 10.1371/journal.pone.0021842.

8 Franklin, T. B., Russig, H., Weiss, I. C., Gräff, J., Linder, N., Michalon, A., Vizi, S. and Mansuy, I. M.
9 (2010) 'Epigenetic Transmission of the Impact of Early Stress Across Generations', *Biological*
10 *Psychiatry*, 68(5), pp. 408–415. doi: 10.1016/j.biopsych.2010.05.036.

11 Fullston, T., Teague, E. M. C. O., Palmer, N. O., Deblasio, M. J., Mitchell, M., Corbett, M., Print, C. G.,
12 Owens, J. A. and Lane, M. (2013) 'Paternal obesity initiates metabolic disturbances in two generations
13 of mice with incomplete penetrance to the F2 generation and alters the transcriptional profile of testis
14 and sperm microRNA content', *FASEB Journal*, 27(10), pp. 4226–4243. doi: 10.1096/fj.12-224048.

15 Gapp, K. and Bohacek, J. (2017) 'Epigenetic germline inheritance in mammals: looking to the past to
16 understand the future', *Genes, Brain and Behavior*, 17(3). doi: 10.1111/gbb.12407.

17 Gapp, K., Jawaid, A., Sarkies, P., Bohacek, J., Pelczar, P., Prados, J., Farinelli, L., Miska, E. and
18 Mansuy, I. M. (2014) 'Implication of sperm RNAs in transgenerational inheritance of the effects of early
19 trauma in mice', *Nature neuroscience*. 2014/04/15, 17(5), pp. 667–669. doi: 10.1038/nn.3695.

20 Gapp, K., van Steenwyk, G., Germain, P. L., Matsushima, W., Rudolph, K. L. M., Manuella, F.,
21 Roszkowski, M., Vernaz, G., Ghosh, T., Pelczar, P., *et al.* (2018) 'Alterations in sperm long RNA
22 contribute to the epigenetic inheritance of the effects of postnatal trauma', *Molecular Psychiatry*, p. 1.
23 doi: 10.1038/s41380-018-0271-6.

24 Gjerstad, J. K., Lightman, S. L. and Spiga, F. (2018) 'Role of glucocorticoid negative feedback in the
25 regulation of HPA axis pulsatility.', *Stress (Amsterdam, Netherlands)*, 21(5), pp. 403–416. doi:
26 10.1080/10253890.2018.1470238.

27 Gòdia, M., Castelló, A., Rocco, M., Cabrera, B., Rodríguez-Gil, J. E., Balasch, S., Lewis, C., Sánchez,
28 A. and Clop, A. (2020) 'Identification of circular RNAs in porcine sperm and evaluation of their relation
29 to sperm motility', *Scientific Reports*, 10(1), pp. 1–11. doi: 10.1038/s41598-020-64711-z.

30 Gòdia, M., Estill, M., Castelló, A., Balasch, S., Rodríguez-Gil, J. E., Krawetz, S. A., Sánchez, A. and
31 Clop, A. (2019) 'A RNA-seq analysis to describe the boar sperm transcriptome and its seasonal

1 changes', *Frontiers in Genetics*, 10(MAR), pp. 1–14. doi: 10.3389/fgene.2019.00299.

2 Grandjean, V., Fourre, S., De Abreu, D. A. F., Derieppe, M.-A. A., Remy, J.-J. J., Rassoulzadegan, M.,
3 Fourné, S., De Abreu, D. A. F., Derieppe, M.-A. A., Remy, J.-J. J., *et al.* (2015) 'RNA-mediated
4 paternal heredity of diet-induced obesity and metabolic disorders', *Sci Rep.* 2015/12/15, 5, p. 18193.
5 doi: 10.1038/srep18193.

6 Grentzinger, T., Armenise, C., Brun, C., Mugat, B., Serrano, V., Pelisson, A. and Chambeyron, S.
7 (2012) 'piRNA-mediated transgenerational inheritance of an acquired trait.', *Genome research*, 22(10),
8 pp. 1877–1888. doi: 10.1101/gr.136614.111.

9 Guzzi, N. and Bellodi, C. (2020) 'RNA Biology Novel insights into the emerging roles of tRNA-derived
10 fragments in mammalian development'. doi: 10.1080/15476286.2020.1732694.

11 Haeussler, S. and Claus, R. (2007) 'Expression of the glucocorticoid receptor and 11 β -
12 hydroxysteroid dehydrogenase 2 in pig testes cells along fetal development', *Reproduction, Fertility
13 and Development*, 19(5), p. 664. doi: 10.1071/RD07033.

14 Hazra, R., Upton, D., Jimenez, M., Desai, R., Handelsman, D. J. and Allan, C. M. (2014) 'In Vivo
15 Actions of the Sertoli Cell Glucocorticoid Receptor', *Endocrinology*, 155(3), pp. 1120–1130. doi:
16 10.1210/en.2013-1940.

17 Heijmans, B. T., Tobi, E. W., Stein, A. D., Putter, H., Blauw, G. J., Susser, E. S., Slagboom, P. E. and
18 Lumey, L. H. (2008) 'Persistent epigenetic differences associated with prenatal exposure to famine in
19 humans', *Proc Natl Acad Sci U S A.* 2008/10/29, 105(44), pp. 17046–17049. doi:
20 10.1073/pnas.0806560105.

21 Horby, P., Lim, W. S., Emberson, J., Mafham, M., Bell, J., Linsell, L., Staplin, N., Brightling, C.,
22 Ustianowski, A., Elmahi, E., *et al.* (2020) 'Effect of Dexamethasone in Hospitalized Patients with
23 COVID-19: Preliminary Report', *medRxiv*, p. 2020.06.22.20137273. doi:
24 10.1101/2020.06.22.20137273.

25 Hoyer, C., Richter, H., Brandwein, C., Riva, M. A. and Gass, P. (2013) 'Preconceptional paternal
26 exposure to a single traumatic event affects postnatal growth of female but not male offspring', 24(15).
27 doi: 10.1097/WNR.0000000000000007.

28 Hua, M., Liu, W., Chen, Y., Zhang, F., Xu, B., Liu, S., Chen, G., Shi, H. and Wu, L. (2019)
29 'Identification of small non-coding RNAs as sperm quality biomarkers for in vitro fertilization.', *Cell
30 discovery*, 5, p. 20. doi: 10.1038/s41421-019-0087-9.

31 Huber, M. H. R. and Bronson, F. H. (1980) 'Social modulation of spontaneous ejaculation in the

1 mouse', *Behavioral and Neural Biology*, 29(3), pp. 390–393. doi: 10.1016/S0163-1047(80)90373-8.

2 Israel, S., Ernst, M., Psathaki, O. E., Drexler, H. C. A., Casser, E., Suzuki, Y., Makalowski, W., Boiani,
3 M., Fuellen, G. and Taher, L. (2019) 'An integrated genome-wide multi-omics analysis of gene
4 expression dynamics in the preimplantation mouse embryo', *Scientific Reports*, 9(1), pp. 1–15. doi:
5 10.1038/s41598-019-49817-3.

6 Jimenez-Chillaron, J. C., Isganaitis, E., Charalambous, M., Gesta, S., Pentinat-Pelegrin, T., Faucette,
7 R. R., Otis, J. P., Chow, A., Diaz, R., Ferguson-Smith, A., *et al.* (2009) 'Intergenerational transmission
8 of glucose intolerance and obesity by in utero undernutrition in mice', *Diabetes*. 2008/11/20, 58(2), pp.
9 460–468. doi: 10.2337/db08-0490.

10 Jirtle, R. L. and Skinner, M. K. (2007) 'Environmental epigenomics and disease susceptibility', *Nature*
11 *reviews. Genetics*. 2007/03/17, 8(4), pp. 253–262. doi: 10.1038/nrg2045.

12 Kang, T. H., Park, Y., Bader, J. S. and Friedmann, T. (2013) 'The Housekeeping Gene Hypoxanthine
13 Guanine Phosphoribosyltransferase (HPRT) Regulates Multiple Developmental and Metabolic
14 Pathways of Murine Embryonic Stem Cell Neuronal Differentiation', *PLoS ONE*. Edited by A. J.
15 Cooney, 8(10), p. e74967. doi: 10.1371/journal.pone.0074967.

16 Kaufmann, S. H., Wright, W. W., Okret, S., Wikström, A. C., Gustafsson, J. A., Shaper, N. L. and
17 Shaper, J. H. (1992) 'Evidence that rodent epididymal sperm contain the Mr approximately 94,000
18 glucocorticoid receptor but lack the Mr approximately 90,000 heat shock protein.', *Endocrinology*,
19 130(5), pp. 3074–3084. doi: 10.1210/endo.130.5.1572314.

20 Kiselev, V. Y., Kirschner, K., Schaub, M. T., Andrews, T., Yiu, A., Chandra, T., Natarajan, K. N., Reik,
21 W., Barahona, M., Green, A. R., *et al.* (2017) 'SC3: consensus clustering of single-cell RNA-seq data',
22 *Nature Methods*, 14(5), pp. 483–486. doi: 10.1038/nmeth.4236.

23 Kiselev, V. Y., Yiu, A. and Hemberg, M. (2018) 'scmap: projection of single-cell RNA-seq data across
24 data sets', *Nature Methods*, 15(5), pp. 359–362. doi: 10.1038/nmeth.4644.

25 de Kloet, E. R., Joëls, M. and Holsboer, F. (2005) 'Stress and the brain: from adaptation to disease.',
26 *Nature reviews. Neuroscience*, 6(6), pp. 463–475. doi: 10.1038/nrn1683.

27 Kozomara, A. and Griffiths-Jones, S. (2011) 'miRBase: integrating microRNA annotation and deep-
28 sequencing data', *Nucleic Acids Research*, 39(Issue suppl_1), pp. D152–D157. doi:
29 10.1093/nar/gkq1027.

30 Krawetz, S. A., Kruger, A., Lalancette, C., Tagett, R., Anton, E., Draghici, S. and Diamond, M. P.
31 (2011) 'A survey of small RNAs in human sperm', *Hum Reprod*. 2011/10/13, 26(12), pp. 3401–3412.

1 doi: der329 [pii] 10.1093/humrep/der329.

2 Legnini, I., Di Timoteo, G., Rossi, F., Morlando, M., Briganti, F., Sthandier, O., Fatica, A., Santini, T.,
3 Andronache, A., Wade, M., *et al.* (2017) 'Circ-ZNF609 Is a Circular RNA that Can Be Translated and
4 Functions in Myogenesis', *Molecular Cell*, 66(1), pp. 22-37.e9. doi: 10.1016/j.molcel.2017.02.017.

5 Li, Q., Hu, B., Hu, G., Chen, C., Niu, X., Liu, J., Zhou, S., Zhang, C., Wang, Y. and Deng, Z.-F. (2016)
6 'tRNA-Derived Small Non-Coding RNAs in Response to Ischemia Inhibit Angiogenesis', *Scientific*
7 *Reports*, 6(1), p. 20850. doi: 10.1038/srep20850.

8 Liao, Y., Smyth, G. K. and Shi, W. (2014) 'featureCounts: an efficient general purpose program for
9 assigning sequence reads to genomic features', *Bioinformatics*, 30(7), pp. 923–930. doi:
10 10.1093/bioinformatics/btt656.

11 Liu, W.-M., Pang, R. T. K., Chiu, P. C. N., Wong, B. P. C., Lao, K., Lee, K.-F. and Yeung, W. S. B.
12 (2012) 'Sperm-borne microRNA-34c is required for the first cleavage division in mouse', *Proceedings*
13 *of the National Academy of Sciences of the United States of America*, 109(2), pp. 490–494. doi:
14 10.1073/pnas.1110368109.

15 Love, M. I., Huber, W. and Anders, S. (2014) 'Moderated estimation of fold change and dispersion for
16 RNA-seq data with DESeq2', *Genome Biology*, 15(12), p. 550. doi: 10.1186/s13059-014-0550-8.

17 Marco, A., Kisliouk, T., Tabachnik, T., Meiri, N. and Weller, A. (2014) 'Overweight and CpG
18 methylation of the Pomc promoter in offspring of high-fat-diet-fed dams are not "reprogrammed" by
19 regular chow diet in rats.', *FASEB journal : official publication of the Federation of American Societies*
20 *for Experimental Biology*, 28(9), pp. 4148–4157. doi: 10.1096/fj.14-255620.

21 Martin, M. (2011) 'Cutadapt removes adapter sequences from high-throughput sequencing reads',
22 *EMBnet.journal*, 17(1), p. 10. doi: 10.14806/ej.17.1.200.

23 Martínez, D., Pentinat, T., Ribó, S., Daviaud, C., Bloks, V. W., Cebrià, J., Villalmanzo, N., Kalko, S. G.,
24 Ramón-Krauel, M., Díaz, R., *et al.* (2014) 'In Utero Undernutrition in Male Mice Programs Liver Lipid
25 Metabolism in the Second-Generation Offspring Involving Altered Lxra DNA Methylation.', *Cell*
26 *metabolism*, pp. 1–11. doi: 10.1016/j.cmet.2014.03.026.

27 Mashoodh, R., Franks, B., Curley, J. P. and Champagne, F. A. (2012) 'Paternal social enrichment
28 effects on maternal behavior and offspring growth', *Proc Natl Acad Sci U S A*. 2012/10/10, 109 Suppl,
29 pp. 17232–17238. doi: 10.1073/pnas.1121083109.

30 McCarthy, D. J., Campbell, K. R., Lun, A. T. L. and Wills, Q. F. (2017) 'Scater: pre-processing, quality
31 control, normalization and visualization of single-cell RNA-seq data in R', *Bioinformatics*, p. btw777.

1 doi: 10.1093/bioinformatics/btw777.

2 Meistrich, M. L. (1975) 'Alteration of epididymal sperm transport and maturation in mice by oestrogen
3 and testosterone', *Nature*, (1).

4 Miyazaki, T., Miyazaki, S., Ashida, M., Tanaka, T., Tashiro, F. and Miyazaki, J. (2013) 'Functional
5 Analysis of Tc1 Using Tc1-Deficient Mouse Embryonic Stem Cells', *PLoS ONE*, 8(8). doi:
6 10.1371/JOURNAL.PONE.0071645.

7 Morgan, C. P. and Bale, T. L. (2011) 'Early Prenatal Stress Epigenetically Programs
8 Dymasculinization in Second-Generation Offspring via the Paternal Lineage', *The Journal of
9 Neuroscience*, 31(33), pp. 11748–11755. doi: 10.1523/jneurosci.1887-11.2011.

10 Nakagata, N. (2011) 'Cryopreservation of Mouse Spermatozoa and In Vitro Fertilization', in. Humana
11 Press, pp. 57–73. doi: 10.1007/978-1-60761-974-1_4.

12 Nätt, D., Kugelberg, U., Casas, E., Nedstrand, E., Zalavary, S., Henriksson, P., Nijm, C., Jäderquist,
13 J., Sandborg, J., Flincke, E., *et al.* (2019) 'Human sperm displays rapid responses to diet', *PLoS
14 Biology*, 17(12), pp. 1–25. doi: 10.1371/journal.pbio.3000559.

15 Ostermeier, G. C., Miller, D., Huntriss, J. D., Diamond, M. P. and Krawetz, S. A. (2004) 'Reproductive
16 biology: Delivering spermatozoan RNA to the oocyte', *Nature*, 429(6988), p. 154. doi:
17 10.1038/429154a.

18 Ostermeier, G. C., Wiles, M. V, Farley, J. S. and Taft, R. A. (2008) 'Conserving, Distributing and
19 Managing Genetically Modified Mouse Lines by Sperm Cryopreservation', *PLoS ONE*. Edited by H. A.
20 El-Shemy, 3(7), p. e2792. doi: 10.1371/journal.pone.0002792.

21 Pembrey, M. E., Bygren, L. O., Kaati, G., Edvinsson, S., Northstone, K., Sjöström, M., Golding, J.,
22 Alspac, T. and Team, S. (2006) 'Sex-specific, male-line transgenerational responses in humans',
23 *European Journal of Human Genetics*, 14, pp. 159–166. doi: 10.1038/sj.ejhg.5201538.

24 Peng, H., Shi, J., Zhang, Y., Zhang, H., Liao, S., Li, W., Lei, L., Han, C., Ning, L., Cao, Y., *et al.* (2012)
25 'A novel class of tRNA-derived small RNAs extremely enriched in mature mouse sperm', *Cell
26 Research*, 22(11), pp. 1609–1612. doi: 10.1038/cr.2012.141.

27 Pentinat, T., Ramon-Krauel, M., Cebria, J., Diaz, R. and Jimenez-Chillaron, J. C. (2010)
28 'Transgenerational inheritance of glucose intolerance in a mouse model of neonatal overnutrition',
29 *Endocrinology*. 2010/10/15, 151(12), pp. 5617–5623. doi: 10.1210/en.2010-0684.

30 Petropoulos, S., Matthews, S. G. and Szyf, M. (2014) 'Adult Glucocorticoid Exposure Leads to
31 Transcriptional and DNA Methylation Changes in Nuclear Steroid Receptors in the Hippocampus and

1 Kidney of Mouse Male Offspring¹, *Biology of Reproduction*, 90(2). doi:
2 10.1095/biolreprod.113.115899.

3 Qiu, X., Hill, A., Packer, J., Lin, D., Ma, Y.-A. and Trapnell, C. (2017) 'Single-cell mRNA quantification
4 and differential analysis with Census', *Nature Methods*, 14(3), pp. 309–315. doi: 10.1038/nmeth.4150.

5 Rodgers, A. B., Morgan, C. P., Bronson, S. L., Revello, S. and Bale, T. L. (2013) 'Paternal Stress
6 Exposure Alters Sperm MicroRNA Content and Reprograms Offspring HPA Stress Axis Regulation',
7 *Journal of Neuroscience*. 2013/05/24, 33(21), pp. 9003–9012. doi: 10.1523/JNEUROSCI.0914-
8 13.2013.

9 Rodgers, A. B., Morgan, C. P., Leu, N. A. and Bale, T. L. (2015) 'Transgenerational epigenetic
10 programming via sperm microRNA recapitulates effects of paternal stress', *Proceedings of the*
11 *National Academy of Sciences*, 112(44), p. 201508347. doi: 10.1073/pnas.1508347112.

12 Rompala, G. R., Mounier, A., Wolfe, C. M., Lin, Q., Lefterov, I. and Homanics, G. E. (2018) 'Heavy
13 Chronic Intermittent Ethanol Exposure Alters Small Noncoding RNAs in Mouse Sperm and
14 Epididymosomes.', *Frontiers in genetics*, 9, p. 32. doi: 10.3389/fgene.2018.00032.

15 Roth, T. L., Lubin, F. D., Funk, A. J. and Sweatt, J. D. (2009) 'Lasting Epigenetic Influence of Early-
16 Life Adversity on the BDNF Gene', *Biological Psychiatry*, 65(9), pp. 760–769. Available at:
17 [http://www.sciencedirect.com/science/article/B6T4S-4VCNF2J-](http://www.sciencedirect.com/science/article/B6T4S-4VCNF2J-1/2/f7007e8e39f1f645dd25d9dbf97487fe)
18 [1/2/f7007e8e39f1f645dd25d9dbf97487fe](http://www.sciencedirect.com/science/article/B6T4S-4VCNF2J-1/2/f7007e8e39f1f645dd25d9dbf97487fe).

19 Rousseeuw, P. J. (1987) 'Silhouettes: A graphical aid to the interpretation and validation of cluster
20 analysis', *Journal of Computational and Applied Mathematics*, 20, pp. 53–65. doi: 10.1016/0377-
21 0427(87)90125-7.

22 Shankar, K., Kang, P., Harrell, A., Zhong, Y., Marecki, J. C., Ronis, M. J. J. and Badger, T. M. (2010)
23 'Maternal overweight programs insulin and adiponectin signaling in the offspring.', *Endocrinology*,
24 151(6), pp. 2577–2589. doi: 10.1210/en.2010-0017.

25 Sharma, U., Conine, C. C., Shea, J. M., Boskovic, A., Derr, A. G., Bing, X. Y., Belleannee, C.,
26 Kucukural, A., Serra, R. W., Sun, F., *et al.* (2015) 'Biogenesis and function of tRNA fragments during
27 sperm maturation and fertilization in mammals', *Science*, 351(6271), pp. 391–396. doi:
28 10.1126/science.aad6780.

29 Sharma, U., Sun, F., Conine, C. C., Reichholf, B., Kukreja, S., Herzog, V. A., Ameres, S. L. and
30 Rando, O. J. (2018) 'Small RNAs Are Trafficked from the Epididymis to Developing Mammalian
31 Sperm', *Developmental Cell*, 46(4), pp. 481–494.e6. doi: 10.1016/J.DEVCEL.2018.06.023.

1 Shi, J., Ko, E.-A., Sanders, K. M., Chen, Q., Zhou, Tong and Zhou, T (2018) 'SPORTS1.0: A Tool for
2 Annotating and Profiling Non-coding RNAs Optimized for rRNA- and tRNA-derived Small RNAs',
3 *Genomics Proteomics Bioinformatics*, 16, pp. 144–151. doi: 10.1016/j.gpb.2018.04.004.

4 Shirayama, M., Seth, M., Lee, H. C., Gu, W., Ishidate, T., Conte Jr., D. and Mello, C. C. (2012)
5 'piRNAs initiate an epigenetic memory of nonself RNA in the *C. elegans* germline', *Cell*. 2012/06/29,
6 150(1), pp. 65–77. doi: 10.1016/j.cell.2012.06.015.

7 Short, A. K., Fennell, K. A., Perreau, V. M., Fox, A., O'Bryan, M. K., Kim, J. H., Bredy, T. W., Pang, T.
8 Y. and Hannan, A. J. (2016) 'Elevated paternal glucocorticoid exposure alters the small noncoding
9 RNA profile in sperm and modifies anxiety and depressive phenotypes in the offspring', *Translational*
10 *Psychiatry*, 6(6), p. e837. doi: 10.1038/tp.2016.109.

11 Short, A. K., Yeshurun, S., Powell, R., Perreau, V. M., Fox, A., Kim, J. H., Pang, T. Y. and Hannan, A.
12 J. (2017) 'Exercise alters mouse sperm small noncoding RNAs and induces a transgenerational
13 modification of male offspring conditioned fear and anxiety.', *Translational psychiatry*, 7(5), p. e1114.
14 doi: 10.1038/tp.2017.82.

15 Silva, E. J. R., Queiróz, D. B. C., Honda, L. and Avellar, M. C. W. (2010) 'Glucocorticoid receptor in
16 the rat epididymis: Expression, cellular distribution and regulation by steroid hormones', *Molecular and*
17 *Cellular Endocrinology*, 325(1–2), pp. 64–77. doi: 10.1016/j.mce.2010.05.013.

18 Silva, E. J. R., Vendramini, V., Restelli, A., Bertolla, R. P., Kempinas, W. G. and Avellar, M. C. W.
19 (2014) 'Impact of adrenalectomy and dexamethasone treatment on testicular morphology and sperm
20 parameters in rats: insights into the adrenal control of male reproduction', *Andrology*, 2(6), pp. 835–
21 846. doi: 10.1111/j.2047-2927.2014.00228.x.

22 Tang, C., Xie, Y., Yu, T., Liu, N., Wang, Z., Woolsey, R. J., Tang, Y., Zhang, X., Qin, W., Zhang, Y., et
23 al. (2020) 'm6A-dependent biogenesis of circular RNAs in male germ cells', *Cell Research*, 30(3), pp.
24 211–228. doi: 10.1038/s41422-020-0279-8.

25 Tang, F., Kaneda, M., O'Carroll, D., Hajkova, P., Barton, S. C., Sun, Y. A., Lee, C., Tarakhovsky, A.,
26 Lao, K. and Surani, M. A. (2007) 'Maternal microRNAs are essential for mouse zygotic development.',
27 *Genes & development*, 21(6), pp. 644–648. doi: 10.1101/gad.418707.

28 Thompson, D. M., Lu, C., Green, P. J. and Parker, R. (2008) 'tRNA cleavage is a conserved response
29 to oxidative stress in eukaryotes', *Rna*, 14(10), pp. 2095–2103. doi: 10.1261/rna.1232808.

30 Tyebji, S., Hannan, A. J. and Tonkin, C. J. (2020) 'Pathogenic Infection in Male Mice Changes Sperm
31 Small RNA Profiles and Transgenerationally Alters Offspring Behavior', *Cell Reports*, 31(4), p.

1 107573. doi: 10.1016/J.CELREP.2020.107573.

2 Vassoler, F. M., White, S. L., Schmidt, H. D., Sadri-Vakili, G. and Pierce, R. C. (2013) 'Epigenetic
3 inheritance of a cocaine-resistance phenotype', *Nat Neurosci.* 2012/12/18, 16(1), pp. 42–47. doi:
4 10.1038/nn.3280.

5 Vojtech, L., Woo, S., Hughes, S., Levy, C., Ballweber, L., Sauteraud, R. P., Strobl, J., Westerberg, K.,
6 Gottardo, R., Tewari, M., *et al.* (2014) 'Exosomes in human semen carry a distinctive repertoire of
7 small non-coding RNAs with potential regulatory functions', *Nucleic Acids Research*, 42(11), pp.
8 7290–7304. doi: 10.1093/nar/gku347.

9 Wang, S., Kou, Z., Jing, Z., Zhang, Y., Guo, X., Dong, M., Wilmut, I. and Gao, S. (2010) 'Proteome of
10 mouse oocytes at different developmental stages', *Proceedings of the National Academy of Sciences
11 of the United States of America*, 107(41), pp. 17639–17644. doi: 10.1073/pnas.1013185107.

12 Wang, Y., Yamauchi, Y., Wang, Z., Zheng, H., Yanagimachi, R., Ward, M. A. and Yan, W. (2020)
13 'Both Cauda and Caput Epididymal Sperm Are Capable of Supporting Full-Term Development in FVB
14 and CD-1 Mice.', *Developmental cell*, 55(6), pp. 675–676. doi: 10.1016/j.devcel.2020.11.022.

15 Weiss, I. C., Franklin, T. B., Vizi, S. and Mansuy, I. M. (2011) 'Inheritable effect of unpredictable
16 maternal separation on behavioral responses in mice', *Frontiers in Behavioral Neuroscience*, 5, p. doi:
17 10.3389/fnbeh.2011.00003. doi: 10.3389/fnbeh.2011.00003.

18 Wu, L., Lu, Y., Jiao, Y., Liu, B., Li, S., Li, Y., Xing, F., Chen, D., Liu, X., Zhao, J., *et al.* (2016) 'Paternal
19 psychological stress reprograms hepatic gluconeogenesis in offspring', *Cell Metabolism*, 23(4), pp.
20 735–743. doi: 10.1016/j.cmet.2016.01.014.

21 Wu, W., Ji, P. and Zhao, F. (2020) 'CircAtlas: an integrated resource of one million highly accurate
22 circular RNAs from 1070 vertebrate transcriptomes', *Genome Biology*, 21(1), p. 101. doi:
23 10.1186/s13059-020-02018-y.

24 Yeri, A., Courtright, A., Danielson, K., Hutchins, E., Alsop, E., Carlson, E., Hsieh, M., Ziegler, O., Das,
25 A., Shah, R. V, *et al.* (2018) 'Evaluation of commercially available small RNASeq library preparation
26 kits using low input RNA', *BMC Genomics*, 19(1), p. 331. doi: 10.1186/s12864-018-4726-6.

27 Yeshurun, S. and Hannan, A. J. (2018) 'Transgenerational epigenetic influences of paternal
28 environmental exposures on brain function and predisposition to psychiatric disorders', *Molecular
29 Psychiatry*, pp. 1–13. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/29520039>.

30 Yu, N., Zhang, P., Wang, L., He, X., Yang, S. and Lu, H. (2018) 'RBBP7 is a prognostic biomarker in
31 patients with esophageal squamous cell carcinoma', *Oncology Letters*, 16(6), pp. 7204–7211. doi:

1 10.3892/ol.2018.9543.

2 Zhang, X.-O., Dong, R., Zhang, Y., Zhang, J.-L., Luo, Z., Zhang, J., Chen, L.-L. and Yang, L. (2016)

3 'Diverse alternative back-splicing and alternative splicing landscape of circular RNAs.', *Genome*

4 *research*, 26(9), pp. 1277–1287. doi: 10.1101/gr.202895.115.

5 Zhang, Y. Y., Zhang, X., Shi, J., Tuorto, F., Li, X., Liu, Y. Y., Liebers, R., Zhang, L., Qu, Y., Qian, J., *et*

6 *al.* (2018) 'Dnmt2 mediates intergenerational transmission of paternally acquired metabolic disorders

7 through sperm small non-coding RNAs', *Nature Cell Biology*, 20(May). doi: 10.1038/s41556-018-0087-

8 2.

9 Zhang, Y., Zhang, X., Shi, J., Tuorto, F., Li, X., Liu, Yusheng, Liebers, R., Zhang, L., Qu, Y., Qian, J.,

10 *et al.* (2018) 'Dnmt2 mediates intergenerational transmission of paternally acquired metabolic

11 disorders through sperm small non-coding RNAs', *Nature Cell Biology*, 20(May). doi: 10.1038/s41556-

12 018-0087-2.

13 Zhou, D., Suzuki, T., Asami, M. and Perry, A. C. F. (2019) 'Caput Epididymidal Mouse Sperm Support

14 Full Development', *Developmental Cell*, 50(1), pp. 5–6. doi: 10.1016/j.devcel.2019.05.012.

15


16

KEY RESOURCES TABLE


REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
Dexamethasone	Sigma	Cat#D4902-25MG
DMSO	VWR	Cat#472301-100ML
Hyperova	Cosmo Bio	Cat#KYD-010-EX-X5
KSOM	Millipore	Cat#MR-121-D
Trizol	Thermo Scientific	Cat#15596026
Directzol	Zymo	Cat#R2080
M-MLV	Promega	Cat#M1701
Recombinant RNAsin Ribonuclease inhibitor	Promega	Cat#N2111
Insulin	Novo Nordisk	Cat#Actrapid
Critical commercial assays		
exoRNeasy Serum/Plasma Midi Kit	Qiagen	Cat#77044
Truseq small RNA library kit	Illumina	Cat#RS-200-0012
Truseq Total RNA library kit	Illumina	Cat#RS-122-2301
Nextflex small RNA library kit	Perkin Elmer	Cat#NOVA-5132-05
Nextera XT DNA Library Preparation Kit	Illumina	Cat#FC-131-1024
miRCURY LNA RT Kit	Qiagen	Cat#339340
miRCURY® LNA® miRNA SYBR® Green PCR	Qiagen	Cat#339345
Deposited data		
Sperm small and long RNA sequencing data	Gene Omnibus	GSE162112
2-cell embryo RNA sequencing data	ENA	ERP105660
Experimental models: Organisms/strains		
C57Bl/6Jrj mice	Janvier lab	
C57Bl/6 CBLT mice	Sanger Institute	
Oligonucleotides		
mirCURY LNA miRNA PCR Assay ArgCCT-2 5'GCCCCAGUGGCCUAAUGGAUAAGGCACUGGCC3'	Qiagen	Cat#339317
mirCURY LNA miRNA PCR Assay Gly-GCC-2 5'GCAUUGGUGGUUCAGUGGUAGAAUUCUCGCCU3'	Qiagen	Cat#339317
mirCURY LNA miRNA PCR Assay His-GTG-3 5'GCCGUGAUCGUUAUAGUGGUUAGUACUCUGCGU3'	Qiagen	Cat#339317
mirCURY LNA miRNA PCR Assay Leu-CAA-2 5'GUCAGGAUGCCGAGUGGUCUAAGGCGCCAGA3'	Qiagen	Cat#339317
mirCURY LNA miRNA PCR Assay RnU6	Qiagen	Cat#339317
mmu-miR-677-p5 mirCURY LNA miRNA PCR Assay	Qiagen	Cat#339306
mmu-miR-3535 mirCURY LNA miRNA PCR Assay	Qiagen	Cat#339306
mmu-miR-6538 mirCURY LNA miRNA PCR Assay	Qiagen	Cat#339306
CircTasp1 FW `CTT AGG AGA GAT TGA ATG TGA TGC C` RW `AAA GGG AGT CAA CCA CTC AG`	Microsynth	Cat#4059186 & 4059187

CircDennd1 FW `AGCTTTCCCAGTTTATTGATGGT` RW `GAAGCCACCCGAAGTGATCT`	Microsynth	Cat#4059182 & 4059183
Software and algorithms		
Code	Github	https://github.com/ETHZ-INS/Sperm-RNA-Dex).
GraphPad Prism	GraphPad Prism version 8	www.graphpad.com
featureCounts	Liao et al., 2014	http://subread.sourceforge.net/
DESeq2	Love et al., 2014	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
edgeR	Anders <i>et al.</i> , 2013	https://bioconductor.org/packages/release/bioc/html/edgeR.html
STAR	Dobin et al., 2013	https://github.com/alexdobin/STAR
CIRCexplorer2	Zhang et al., 2016	https://circexplorer2.readthedocs.io
scater	McCarthy et al., 2017	https://bioconductor.org/packages/release/bioc/html/scater.html
Monocle	Qiu et al., 2017	http://cole-trapnell-lab.github.io/monocle-release/
Trimmomatic	Bolger et al., 2014	www.usadellab.org
cutadapt	Martin, 2011	https://cutadapt.readthedocs.io


Gapp et al. Figure 1



Gapp et al. Figure 2




Gapp et al. Figure 3




Gapp et al. Figure 4


A




B



Gapp et al. Figure 5




Gapp et al. Figure 6




Supplementary Information

Supplementary Figure 1 related to Figure 1 and 4



Relative percentage of RNA Ensemble biotypes and tRNA matching reads (A) in small RNA sequencing libraries of 8 sperm samples from control males and 8 sperm samples 14 days post dexamethasone injection. **(B)** long RNA sequencing libraries of 4 sperm samples from control males and 4 sperm samples 14 days post dexamethasone injection.


Supplementary Figure 2 related to Figure 2




Mature sperm miRNA payload as determined by next generation sequencing. Deseq2 analysis of sperm samples collected 3 hours and 14 days post dexamethasone injection did not reveal any significant effects of (A) treatment (dexamethasone injection), (B) time (time elapsed since injection) nor an (C) interaction between the two. (D) Heat map depicting those miRNAs that are significantly affected in the data obtained from 14 days post injection, at 3 hours and 7 days post dexamethasone injection.

Supplementary Figure 3 related to Figure 3

A



B



C


Variable	Fixed effects (type III)	P value	P value summary	Statistical	F (DFn, DFd)	Geisser-Greenhouse's epsilon
BMI	time	<0.0001	****	Yes	F (2.087, 144.7) = 41.99	0.6957
BMI	sex	<0.0001	****	Yes	F (1, 71) = 76.55	
BMI	treatment	<0.0001	****	Yes	F (1, 71) = 49.86	
BMI	time x sex	<0.0001	****	Yes	F (3, 208) = 33.75	
BMI	time x treatment	0.0008	***	Yes	F (3, 208) = 5.834	
BMI	sex x treatment	0.2181	ns	No	F (1, 71) = 1.544	
BMI	time x sex x treatment	0.4323	ns	No	F (3, 208) = 0.9196	

	ANOVA table	SS	DF	MS	F (DFn, DFd)	P value
GTT	time	60.74	4	15.18	F (3.314, 122.6) = 23.85	P<0.0001
GTT	sex	732.6	1	732.6	F (1, 37) = 162.6	P<0.0001
GTT	treatment	9.901	1	9.901	F (1, 37) = 2.198	P=0.1467
GTT	time x sex	31.81	4	7.951	F (4, 148) = 12.49	P<0.0001
GTT	time x treatment	13.7	4	3.425	F (4, 148) = 5.380	P=0.0005
GTT	sex x treatment	3.936	1	3.936	F (1, 37) = 0.8737	P=0.3560
GTT	time x sex x treatment	13.73	4	3.432	F (4, 148) = 5.392	P=0.0004
GTT	Subject	166.7	37	4.504		
	Residual	94.21	148	0.6366		
ITT	time	60.74	4	15.18	F (3.314, 122.6) = 23.85	P<0.0001
ITT	sex	732.6	1	732.6	F (1, 37) = 162.6	P<0.0001
ITT	treatment	9.901	1	9.901	F (1, 37) = 2.198	P=0.1467
ITT	time x sex	31.81	4	7.951	F (4, 148) = 12.49	P<0.0001
ITT	time x treatment	13.7	4	3.425	F (4, 148) = 5.380	P=0.0005
ITT	sex x treatment	3.936	1	3.936	F (1, 37) = 0.8737	P=0.3560
ITT	time x sex x treatment	13.73	4	3.432	F (4, 148) = 5.392	P=0.0004
ITT	Subject	166.7	37	4.504		
	Residual	94.21	148	0.6366		

Tissue	P value	Difference	SE of difference	t ratio
Liver (g)	0.27665	-0.06789	0.05979	1.136
Kidney (g)	0.188702	-0.01267	0.009131	1.387
Inguinal WAT (g)	0.776367	-0.02972	0.1025	0.29
Gonadal WAT (g)	0.968101	0.01122	0.2753	0.04077
Interscapular BAT (g)	0.801259	-0.006444	0.02508	0.2569
Liver (g)	0.562456	0.06872	0.116	0.5923
Kidney (g)	0.788758	0.004292	0.01573	0.2728
Inguinal WAT (g)	0.872972	-0.014	0.08608	0.1626
Gonadal WAT (g)	0.434173	-0.1747	0.2174	0.8036
Interscapular BAT (g)	0.432084	0.01892	0.02343	0.8073

Necropsy followed by weighing of dissected tissue from offspring of control injected and Dexamethasone injected males. Comparison of the weight of liver, kidney, inguinal and gonadal white adipose tissue (WAT) as well as interscapular brown adipose tissue (BAT) did not reveal significant differences between (A) female and (B) male offspring of control and dexamethasone injected males. (C) Statistical values obtained by overall ANOVAs (BMI, GTT, ITT) and Multiple t-tests corrected for multiple comparisons (necropsy). All Data besides male gonadal WAT showed equal distribution of variances. Graphs show scattered dot plots with standard error of the mean. ITT = insulin tolerance test, GTT= Glucose tolerance test, BMI= Body mass index, WAT = white adipose tissue. BAT= brown adipose tissue.

Supplementary Figure 4 related to Figure 5



Single- 2-cell embryo sequencing using the Smartseq method. (A) Result of the quality filters implemented to select the 2 cell embryos that were used for downstream analyses. Libraries (embryos/cells) that contained a suboptimal number of mitochondrial (>15%) or ERCC mapping reads (>10%) or yielded less than 500 000 reads were excluded (displayed in Out). Most libraries showed a high number of gene mapping reads and were retained and processed for further analysis (displayed in In). (B) PCA results shown as a single panel, revealing segregation of embryos for the first two principle components (matching almost perfectly cluster 1 (red) and cluster 2(blue)) and additionally depicting read count number (size of circle) and assignment to treatments (offspring embryos of control fathers = CON, dots and fathers who were injected with dexamethasone 14 days prior sperm harvest = DEX, triangles). PC2 is attributed to technical factors such as read counts. (C) Marker genes identified by sc3 for C1 and C2 clusters.

Supplementary Figure 5 related to Figure 5 and 6


circAtlas ID	microRNA name	#binding sites of miRanda	#bingding sites of targetScan
mmu-Dennd1b_0017	mmu-miR-201-3p	1	1
mmu-Dennd1b_0017	mmu-miR-3110-5p	1	1
mmu-Dennd1b_0017	mmu-miR-1953	1	1
mmu-Dennd1b_0017	mmu-miR-1981-5p	1	1
mmu-Dennd1b_0017	mmu-miR-706	1	1
mmu-Dennd1b_0017	mmu-miR-6363	1	1
mmu-Dennd1b_0017	mmu-miR-3074-5p	1	1
mmu-Dennd1b_0017	mmu-miR-1955-5p	1	2
mmu-Dennd1b_0017	mmu-miR-675-3p	1	1
mmu-Dennd1b_0017	mmu-miR-343	1	2
mmu-Dennd1b_0017	mmu-miR-3083-5p	1	1
mmu-Dennd1b_0017	mmu-miR-3073a-5p	1	1
mmu-Dennd1b_0017	mmu-miR-9-5p	1	1
mmu-Dennd1b_0017	mmu-miR-378b	1	1
mmu-Tasp1_0039	mmu-miR-221-5p	1	1
mmu-Tasp1_0039	mmu-miR-5627-3p	1	3
mmu-Tasp1_0039	mmu-miR-298-5p	1	3
mmu-Tasp1_0039	mmu-miR-3058-5p	1	1
mmu-Tasp1_0039	mmu-miR-207	1	3
mmu-Tasp1_0039	mmu-miR-709	1	1
mmu-Tasp1_0039	mmu-miR-6537-3p	1	1
mmu-Tasp1_0039	mmu-miR-713	1	2
mmu-Tasp1_0039	mmu-miR-3094-3p	1	2
mmu-Tasp1_0039	mmu-miR-100-3p	1	2
mmu-Tasp1_0039	mmu-miR-1906	1	3
mmu-Tasp1_0039	mmu-miR-320-5p	1	3
mmu-Tasp1_0039	mmu-miR-3082-3p	1	2
mmu-Tasp1_0039	mmu-miR-706	1	1
mmu-Tasp1_0039	mmu-miR-3110-5p	1	1
mmu-Tasp1_0039	mmu-miR-6900-5p	1	1
mmu-Tasp1_0039	mmu-miR-1955-5p	1	1
mmu-Tasp1_0039	mmu-miR-674-5p	1	1
mmu-Tasp1_0039	mmu-miR-351-3p	1	1
mmu-Tasp1_0039	mmu-miR-5104	1	1
mmu-Tasp1_0039	mmu-miR-504-3p	1	1
mmu-Tasp1_0039	mmu-miR-152-5p	1	1
mmu-Tasp1_0039	mmu-miR-1897-5p	1	1
mmu-Tasp1_0039	mmu-miR-3105-5p	1	1
mmu-Tasp1_0039	mmu-miR-742-3p	1	2
mmu-Tasp1_0039	mmu-miR-1969	1	2
mmu-Tasp1_0039	mmu-miR-1960	1	2
mmu-Tasp1_0039	mmu-miR-24-3p	1	2
mmu-Tasp1_0039	mmu-miR-188-5p	1	1

CircRNA targets as predicted by CircAtlas. Each column displays results from miRanda and targetScan.


miRNAs that are targeted by both circRNAs are highlighted in purple.

Supplementary Figure 6 related to Figure 1


A



B



C



Fertility measures following Dexamethasone treatment. (A) Sperm cell count 14 days post vehicle (n=7) or Dex (n=7) injection did not reveal any difference in sperm number ($t(12)=1.221$, $p=0.25$) suggesting no impairing effect of Dex on spermatogenesis. (B) Count of fertilized oocytes (as of the appearance of the second pronucleus) over total available oocytes of 6 replicates of cryopreserved sperm from a pool of 2 dex injected versus 2 vehicle injected males did reveal no significant difference in the fertilization rate between dex and vehicle sperm. (C) Number of pups per litter was similar in offspring resulting from in vitro fertilization with sperm from fathers that were injected with Vehicle (n=6) or Dex (n=6) ($t(12)=1.122$, $p=0.25$).