

Palliative care after stroke: A review

Eileen Cowey¹, Markus Schichtel², Joshua D Cheyne³,
Lorna Tweedie⁴, Richard Lehman⁵, Rita Melifonwu⁶  and
Gillian E Mead⁷ 

International Journal of Stroke
2021, Vol. 16(6) 632–639
© 2021 World Stroke Organization



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/17474930211016603

journals.sagepub.com/home/wso



Abstract

Background: Palliative care is an integral aspect of stroke unit care. In 2016, the American Stroke Association published a policy statement on palliative care and stroke. Since then there has been an expansion in the literature on palliative care and stroke.

Aim: Our aim was to narratively review research on palliative care and stroke, published since 2015.

Results: The literature fell into three broad categories: (a) scope and scale of palliative care needs, (b) organization of palliative care for stroke, and (c) shared decision making. Most literature was observational. There was a lack of evidence about interventions that address specific palliative symptoms or improve shared decision making. Racial disparities exist in access to palliative care after stroke. There was a dearth of literature from low- and middle-income countries.

Conclusion: We recommend further research, especially in low- and middle-income countries, including research to explore why racial disparities in access to palliative care exist. Randomized trials are needed to address specific palliative care needs after stroke and to understand how best to facilitate shared decision making.

Keywords

Stroke, palliative care, shared decision making, end-of-life, stroke unit

Received: 16 February 2021; accepted: 19 April 2021

Introduction

Stroke is a leading cause of death and disability, with 5.5 million attributable deaths and 80 million stroke survivors worldwide in 2016.¹ Stroke incidence and standardized death rates are falling but population growth and ageing are likely to increase the burden of stroke.¹

The World Health Organization (WHO) defines palliative care as

An approach that improves the quality of life of patients and their families facing the problem associated with life-threatening illness, through the prevention and relief of suffering by means of early identification and impeccable assessment and treatment of pain and other problems, physical, psychosocial and spiritual.²

This definition of palliative care is highly relevant to stroke.³ Yet “palliative care” is often interpreted by stroke care professionals, stroke survivors, and families as meaning only end-of-life care or withdrawal of care.⁴ However, organized stroke unit care already includes multiple aspects of the WHO definition of palliative care (Table 1); this can be considered to be “general

palliative care”, while specialist palliative care teams provide “specialist” palliative care.³

Shared decision making (SDM) is an integral aspect of palliative care. SDM involves combining the best available evidence and patients’ values and preferences into decisions about care,⁵ including decisions about interventions that improve survival but with severe disability, and about transitions between care settings.^{4,6} Stroke has a profound psychological effect on family as well as patient. A truthful, compassionate, appraisal of

¹Nursing & Health Care School, University of Glasgow, Glasgow, UK

²Institute of Public Health and Primary Care, University of Cambridge, Cambridge, UK

³Cochrane Stroke Group, Centre for Clinical Brain Sciences (CCBS), University of Edinburgh, Edinburgh, UK

⁴Family member, Gorebridge, UK

⁵Institute of Applied Health Research, Murray Learning Centre, University of Birmingham, Birmingham, UK

⁶Life After Stroke Centre, Stroke Action Nigeria, Onitsha, Nigeria

⁷Royal Infirmary of Edinburgh, Edinburgh, UK

Corresponding author:

Eileen Cowey, Nursing & Health Care School, University of Glasgow, 59 Oakfield Avenue, Glasgow G12 8LL, UK.

Email: eileen.cowey@glasgow.ac.uk

Table 1. The key aspects of the WHO definition of palliative care, and how this type of care is already provided, to some extent, by stroke units

	Palliative care	Stroke unit care
Approach-to improve quality of life of patients and families	Aims to improve quality of life	Clinicians consider the impact of stroke and its treatment on quality of life, and make decisions about the risks and benefits of treatment and rehabilitation.
Life-threatening illness	Is appropriate for patients with life-threatening illness	Stroke is a life-threatening illness. Stroke clinicians regularly deal with the physical, psychological, and existential distress of sudden, life-threatening illness.
Early	Needs to be considered early in the trajectory of a life-threatening illness	Admission to a stroke unit generally occurs very early after stroke onset. Thus, stroke clinicians are ideally placed to consider palliative care.
Pain Other physical problems Psychosocial problems Spiritual problems	Addresses a wide range of physical, psychosocial, and spiritual problems	Rehabilitation addresses the direct neurological consequences of stroke. Stroke unit care does address pain, physical, and psychosocial problems, but spiritual care is not considered to the same extent.

likely prognosis is essential⁷⁻⁹ (see Supplementary Table 1 for a bereaved carer's perspective).

Since publication of the American Heart Association's policy statement on palliative care and stroke in 2016,³ new literature has been published in this field. Our aim was to identify and critically appraise this new literature; and make recommendations for future research based on gaps identified in the literature.

Search strategy and criteria

We included systematic reviews, observational studies, and trials that focused on palliative care and stroke.

Search methods

- On 1 June 2020, we searched multiple databases (Cochrane Stroke Group's trials register, the Cochrane Database of Systematic Reviews, Cochrane Central Register of Controlled Trials (CENTRAL) in the Cochrane Library, MEDLINE Ovid, Embase Ovid, CINAHL EBSCO and Clinicaltrials.gov) using the terms "palliative care" AND "stroke" (Supplementary Table 2), from 2015 onwards, to identify new literature published since the AHA policy statement on palliative care and stroke published in 2016.³ Two authors screened titles, obtained full texts of potentially relevant articles, and both read the full texts. We found that this literature fell into three broad categories: (a) scope and scale of palliative care needs, (b) organization of palliative care for stroke and (c) SDM.

- To identify randomized trials addressing the specific palliative care needs identified in the initial database searches, one author (GEM) then searched the Cochrane Library and Database of research in stroke (DORIS).
- All authors provided other key papers, not identified in the initial searches.

Figure 1 shows the search results. Further details are in Supplementary Information.

Scope and scale of palliative care needs


Overview of the literature

Observational studies were mostly limited to end-of-life care.^{4,10-17} These included retrospective cohorts,^{10,11} a retrospective registry study,¹² a national clinical audit,¹³ retrospective case note studies,^{14,15} a qualitative study with healthcare professionals,¹⁶ cross-sectional face-to-face survey,¹⁷ and one mixed methods longitudinal study.⁴ Four studies^{4,10,11,17} collected data on stroke survivors/decedents in their own homes, six were hospital based,^{4,12-16} and two were from nursing homes. The studies were from US,^{10,11} Europe (UK, Ireland),^{4,14} Sweden,¹² Portugal,¹⁵ Australia,¹³ and India.^{16,17} Three¹⁰⁻¹³ report large cohorts of stroke patients which were subsets of larger palliative care cohorts.

Frequency of individual palliative care needs

The most frequently recorded needs were dysphagia (96.8%),¹⁵ death rattles (31.5-60.7%), dyspnea

Figure 1. PRISMA flow diagram.



(16.3–48.4%), and pain (30.3–42.7%).^{12,14,15} Two studies^{14,15} reported anxiety (12.9–18.9%) and confusion or delirium (7.9–19.4%), while one¹⁴ reported agitation (25.9%). Other symptoms included constipation, dry mouth, and seizures.¹⁵ After hospital discharge, pain affected 50%, numbness or tingling 48%, sleep disturbance 43%, and nausea/vomiting and bladder/bowel incontinence affected 15%.¹⁷ In a large US cohort^{10,11} receiving home hospice care, stroke was a common cause of fecal¹⁰ or urinary incontinence.¹¹ Existential needs are also reported; in the UK,⁴ hopelessness or loss of meaning was partly related to having to move home or to a care home. In India,¹⁷ over 40% of survivors reported hopelessness and 8% thought about death.

Identification of palliative care needs after stroke

Palliative care symptoms were recognized in about two-thirds of patients who died on an acute stroke unit,¹⁴

although the true frequency is likely to be higher. Swedish healthcare professionals were less likely to know if end-of-life symptoms were present in stroke patients than in cancer patients, or to know if stroke patients' preferences regarding place of care had been met.¹² Cognitive impairment, aphasia, and dysarthria were barriers to pain assessment^{15,16} and accessing services.⁴ The largest studies of palliative care needs after stroke were from specialist palliative care settings^{10,11,13} or unspecified hospital or care home settings,¹² and so cannot easily be generalized to a broader stroke population.

Organization of services, referrals to specialist palliative care services, and place of death

We identified 21 observational studies (mostly reporting referrals to specialist palliative care services) and

one prospective study¹⁸ of a palliative care checklist used in a neurointensive care unit.

Very few patients referred to specialist palliative care had had a stroke (0.4%,^{19,20} 4.1%,^{10,11} 6%,²¹ 8.9%¹³), and stroke patients tended to be referred only in the terminal phase of their illness.^{22,23} Six papers from the USA addressed factors predicting referrals to specialist palliative care.^{24–29} Only 25% of variance was explained by hospital site,²⁶ and racial minorities were less likely to receive palliative care referral^{27,28,30} and more likely to receive life-sustaining interventions.²⁵

Five studies addressed place of death after stroke.^{4,31–34} Most people dying from stroke do not die at home,³⁴ only half of stroke patients died in their preferred place (i.e. home);³³ and a quarter of those dying in hospital die alone. Honest discussions with patients and families are needed to explore preferred place of death, to discuss the practicalities of a home death and if this is not possible, how to ensure a “good” hospital death.

Stroke patients, including those with mental capacity, are less likely than cancer patients to receive information about their transition to end-of-life care¹² and their families less likely to receive bereavement support. Healthcare professionals tend to focus instead on physical recovery rather than the psychosocial and spiritual needs.⁴ Potentially futile interventions are often ongoing when specialist palliative care referrals are made,¹⁵ or on day of death.¹² Inadequate prescribing of palliative medications is also described^{15,16} and few physicians saw spiritual care as a significant issue.¹⁶

Management of individual palliative care problems after stroke

Patients with major stroke were often excluded from the Cochrane reviews of interventions³⁵ and the trials did not include “palliative care” endpoints.³⁶ One ongoing trial is seeking to improve care in the longer term.³⁷ Two trials which explored transitions of care did not include people with incapacity or severe stroke.^{38,39} The table in the supplementary materials summarizes additional relevant reviews for each symptom.

Low- and middle-income countries

Our searches identified two studies from India, and six studies from Africa. Western African low- and middle-income countries (LMICs) have little established palliative care activity, because of resource limitations and lack of political, organizational, or interprofessional collaboration.⁴⁰ Pain control was identified in one study⁴¹ as the main unmet palliative care need, and in South Africa, physical, emotional, and social dysfunctions were major concerns for stroke survivors.⁴²

In Nigeria, informal stroke caregivers reported dissatisfaction with infrastructure, waiting times, and information quality;⁴³ and stroke survivors’ preferences for rehabilitation settings were influenced by age and finances.⁴⁴

In sub-Saharan Africa, some still believe that stroke arises from witchcraft or demons.⁴⁵ These beliefs affect treatment preferences for acute stroke care and rehabilitation, but any potential influence on choices about end-of-life care has not been researched. Palliative care research, education of health workers, improved palliative service provision, and policy development in this area are needed.⁴⁰ In Nigeria this is beginning to happen, as stroke is now included in the national non communicable diseases multi-sectoral action plan.

Shared decision making

All the literature we identified was observational research or reviews, except for two relevant trials; one is ongoing (“Team-based Versus Primary Care Clinician-led Advance Care Planning in Practice-based Research Networks”)⁴⁶ which includes stroke patients; and a feasibility trial of a “Decision Aid for Families of Critically Ill Stroke Patients” that has not yet started.⁴⁷

SDM, autonomy, and anticipatory care plans

The ethical principle of autonomy underpins SDM and affirms the right of patients to select their medical therapy from a number of appropriate options. Advanced care plans (ACPs) are rarely in place⁴⁸ pre-stroke; for example, of all stroke deaths in an Australian study, only 4% had a pre-stroke ACP in place.²³ After stroke, an ACP for ongoing care can enable preferences, values, beliefs, and goals to be articulated by patients (if they are able to do so) or by their surrogates and enable clinicians to tailor treatment in line with their patient and family wishes.

Patients expect clinicians to take responsibility for initiating SDM, and for identifying a patient’s requirement for information. Clinicians need to plan care consistent with patients’ values and goals.⁴⁹ ACPs need regular review and update of a patient’s care preferences in the context of ongoing communication between clinicians, patients, and carers,⁵⁰ and the development of a trusting relationship, particularly as prognosis is often uncertain early after stroke and sometimes patients “rally” unexpectedly.

Practical ways to promote SDM in stroke care

The ideal processes of SDM may be impossible to achieve after stroke (because of lack of capacity, need

Table 2. Strategies for health care professionals during shared decision making after severe stroke

<ul style="list-style-type: none"> • Acknowledge “shock” and suddenness of stroke and its profound effect on the patient and family
<ul style="list-style-type: none"> • Identify patient’s wishes early on; e.g. advanced directive, Power of Attorney, any previous conversations about views of living with severe disability, patient’s “values”
<ul style="list-style-type: none"> • Ask about, and address any guilt, e.g. “If only I’d found him sooner”
<ul style="list-style-type: none"> • “Truth telling:” be as honest as you can be about likely outcomes
<ul style="list-style-type: none"> • Showing CT brain scan may help to show extent of stroke and align family and health care professionals’ expectations about recovery and goals of care
<ul style="list-style-type: none"> • Try to avoid allowing the family to feel responsible for decisions about: <ul style="list-style-type: none"> • Cardiopulmonary resuscitation • Artificial feeding or intravenous fluids
<ul style="list-style-type: none"> • Let family know that dignity/symptom control are paramount whatever the decision
<ul style="list-style-type: none"> • Offer further meetings
<ul style="list-style-type: none"> • Document the discussion to ensure consistency of messaging.

(from Chest Heart & Stroke Scotland’s online Stroke Training and Awareness (STARS) training module: “Sensitive and Effective Conversations at the End of Life after Acute Stroke,” with kind permission of CHSS. <http://chslearning.org.uk>).

for quick decisions, or uncertain prognosis), but the basic principles of SDM still need to be applied.⁵¹ The key decision makers should be identified and brought together; there needs to be trustworthy and honest exchange of information about treatment options; information must be tailored towards individuals’ needs, and prognostic information must be as accurate as possible and given consistently by different members of the health care team. SDM also requires the best available knowledge of the patients’ personal preferences.

For patients who cannot communicate, the clinician should make sure that the people closest to patient are able to express their views. Discussions should always be recorded. Clinicians should ascertain the extent to which the patient and/or carers wish to have a say in the final decision.⁵² Since these decisions may involve heavy responsibility, e.g. for an indefinite life of long-term care versus probable death, they must be made in a way that allows for trust-building and reflection. For irrevocable decisions, the burden of post-decisional regret must be minimized.⁵²

Ideally, stroke teams should undergo training in SDM.⁵³ A UK third sector stroke organization provides interactive, online training on sensitive, and effective conversations at the end of life, informed by a survey⁵⁴ of the educational needs of UK stroke professionals. Table 2 summarizes potentially helpful approaches for clinicians.

There is no Cochrane review of SDM specifically in stroke.^{55–57} Stroke teams nevertheless should offer realistic planning with patients and carers, raising the

possibility of death or survival with disability.^{4,58} Realistic information should be given, avoiding the impression that all treatments result in a functional recovery.⁹ Physicians should avoid conveying information from a neutral position or wholly delegate decisions to patients or families.^{59,60}

Healthcare professionals should consider different cultural preference of decision makers, and be aware of the significant burden of family in making decisions.^{61,62} Substantial grief and stress reactions were identified in 30% of decision makers for severely ill neuro-ICU patients.⁶³ Therefore, psychological support for carers and patients together with personalized, tailored care, and realistic information should be offered.⁸ SDM tools that include predictions from prognostic models appear helpful⁶⁴ and information should be framed positively (e.g. independence rather than dependence) yet honestly.^{65,66}

Conclusion and future directions

Almost all of the existing literature is observational, most was from high-income countries, and most focused on end-of-life care and SDM in this context. Stroke patients experience multiple palliative care symptoms (physical, psychological, and existential), few stroke patients are referred to specialist palliative care services, referrals are made close to death, few deaths occur at home, and there are racial differences in palliative care referrals. Uncertainty of prognosis and the need to balance hope with reality is challenging.

Recommendations for clinical care

How can we ensure that people with severe stroke receive holistic care in keeping with their preferences and values? Based on our review of the literature, we suggest three actions:

- a. Clinicians should consider systematically seeking “palliative care problems” in the same way that medical and rehabilitation issues are identified and addressed through history taking, examination, and checklists.
- b. SDM needs careful consideration of patient’s values and beliefs, clinicians need to be honest, yet compassionate, about likely outcomes, and patients and families need considerable support to be involved in decision making.
- c. Palliative care specialists have extensive experience of similar palliative care problems in other patient groups, and stroke unit care already includes multiple aspects of palliative care. Closer working between stroke clinicians and palliative care clinicians might facilitate better sharing of ideas and knowledge, and help ensure that stroke survivors’ palliative care needs are met.

Recommendations for future research

Observational research is needed to identify the frequency of specific palliative care needs, to explore attitudes and beliefs about stroke and death, and to better understand why there appears to be racial disparities in referral for palliative care. Research is needed particularly in low- and middle-income countries.

Randomized controlled trials are needed to identify management strategies for palliative care problems including pain, delirium, psychological distress, and existential distress, perhaps by adapting interventions that have been tested in palliative care settings in other patient populations.

Developing and testing standardized ways to make shared decisions after severe stroke, including decision aids and information tailored towards patient’s individual needs, might help ensure that patient outcomes are more in line with their beliefs and values.

Authors’ contributions

GEM devised the review. JC designed and ran initial searches and exporting of results. LT reviewed search terms. GEM, JC, and EC ran follow-up searches. EC and GEM screened and selected papers for inclusion and checked reference lists. GEM wrote the outline manuscript draft. RL and MS drafted the section on shared decision making. RM drafted the section on palliative care in Africa. EC then wrote the paper. All authors approved the final draft of the paper.

Declaration of conflicting interests


The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: GEM reports grants from Chief Scientist Office, during the conduct of the study. EC reports contributing authorship to Chest Heart & Stroke Scotland Stroke Training and Awareness (STARS) online training. MS, JDC, LT, RL, RM declare no competing interests.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iDs

Rita Melifonwu  <https://orcid.org/0000-0002-5192-3871>

Gillian Mead  <https://orcid.org/0000-0001-7494-2023>

Supplemental material

Supplemental material for this article is available online.

References

1. Johnson CO, Nguyen M, Roth GA, et al. Global, regional, and national burden of stroke, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 2019; 18: 439–58.
2. World Health Organization Geneva. *National cancer control programs: policies and managerial guidelines*, 2nd edn. Geneva: Author, 2002.
3. Braun LT, Grady KL, Kutner JS, et al. Palliative care and cardiovascular disease and stroke: a policy statement from the American Heart Association/American Stroke Association. *Circulation* 2016; 134: e198–225.
4. Kendall M, Covey E, Mead GG, et al. Outcomes, experiences and palliative care in major stroke: a multicentre, mixed-method, longitudinal study. *CMAJ* 2018; 190(9): E238–E246.
5. Elwyn G, Laitner S, Coulter A, Walker E, Watson P and Thomson R. Implementing shared decision making in the NHS. *BMJ* 2010; 341: c5146.
6. Cameron JI, O’Connell C, Foley N, et al. Canadian stroke best practice recommendations: managing transitions of care following stroke, guidelines update 2016. *Int J Stroke* 2016; 11: 807–822.
7. Visvanathan A, Mead G, Dennis M, Whiteley W, Doubal F and Lawton J. Maintaining hope after a disabling stroke: a longitudinal qualitative study of patients’ experiences, views, information needs and approaches towards making treatment decisions. *PLoS One* 2019; 14(9): e0222500. DOI: 10.1371/journal.pone.0222500.
8. Visvanathan A, Mead GE, Dennis M, Whiteley WN, Doubal FN and Lawton J. The considerations, experiences and support needs of family members making treatment decisions for patients admitted with major stroke: a qualitative study. *BMC Med Inform Decis Mak* 2020; 20: 98 <https://doi.org/10.1186/s12911-020-01137-7>.
9. Clayton JM, Hancock K, Parker S, et al. Sustaining hope when communicating with terminally ill patients and

- their families: a systematic review. *Psychooncology* 2008; 17: 641–659.
10. Chughtai B, Thomas D, Russell D, Phongtankuel V, Bowles K and Prigerson H. Prevalence and risk factors for fecal incontinence in home hospice. *Am J Hosp Palliat Med* 2019; 36: 33–37.
 11. Chughtai B, Thomas D, Russell D, Bowles K and Prigerson H. Prevalence of and risk factors for urinary incontinence in home hospice patients. *Eur Urol* 2019; 75: 268–271.
 12. Eriksson H, Milberg A, Hjelm K and Friedrichsen M. End of life care for patients dying of stroke: a comparative registry study of stroke and cancer. *PLoS One* 2016; 11(2): e0147694. DOI:10.1371/journal.pone.0147694.
 13. Ding J, Johnson CE, Lee YC (Olivia), Gazey A and Cook A. Characteristics of people with dementia vs other conditions on admission to inpatient palliative care. *J Am Geriatr Soc* 2020; 1–9. DOI:10.1111/jgs.16458.
 14. Ntlholang O, Walsh S, Bradley D and Harbison J. Identifying palliative care issues in inpatients dying following stroke. *Ir J Med Sci* 2016; 185: 741–744.
 15. Monteiro NF, Cipriano P and Freire E. Palliative approach in acute neurological events: a five-year study. *Rev Assoc Med Bras* 2018; 64: 832–836.
 16. Lloyd J, Pinto AM, Nair S and Tarey S. A qualitative study on palliative needs of stroke patients in an indian tertiary care setting – doctors’ perspective. *Indian J Palliat Care* 2019; 25: 84–91.
 17. Sharma M, Lal M, Singh T and Deepti S. Factors associated with physical and psychosocial problems among Indian stroke survivors. *Indian J Palliat Care* 2019; 25: 18–23.
 18. Creutzfeldt CJ, Hanna MG, Cheever CS, et al. Palliative care needs assessment in the neuro-ICU: effect on family. *Neurocrit Care* 2017; 27: 163–172.
 19. Bennett MI, Ziegler L, Allsop M, Daniel S and Hurlow A. What determines duration of palliative care before death for patients with advanced disease? A retrospective cohort study of community and hospital palliative care provision in a large UK city. *BMJ Open* 2016; 6: e012576.
 20. Allsop M, Ziegler LE, Mulvey MR, Russell S, Taylor R and Bennett MI. Duration and determinants of hospice-based specialist palliative care: a national retrospective cohort study. *Palliat Med* 2018; 32: 1322–1333.
 21. Sganga F, Barillaro C, Tamburrano A, et al. The benefits of a hospital palliative care team. *Int J Palliat Nurs* 2019; 25: 345–352.
 22. Tran L, Back AL and Creutzfeldt CJ. Palliative care consultations in the neuro-ICU: a qualitative study. *Neurocrit Care* 2016; 25: 266–272.
 23. Quadri SZ, Huynh T, Cappelen-Smith C, et al. Reflection on stroke deaths and end of life stroke care. *Intern Med J* 2018; 48: 330–334.
 24. Creutzfeldt CJ, Wunsch H, Curtis JR and Hua M. Prevalence and outcomes of patients meeting palliative care consultation triggers in neurological intensive care units. *Neurocrit Care* 2015; 23: 14–21.
 25. Cruz-Flores S, Rodriguez GJ, Chaudhry MRA, et al. Racial/ethnic disparities in hospital utilization in intracerebral hemorrhage. *Int J Stroke* 2019; 14: 686–695.
 26. Faigle R and Gottesman RF. Variability in palliative care use after intracerebral hemorrhage at US hospitals: a multilevel analysis. *Neuroepidemiology* 2019; 53: 84–92.
 27. Faigle R, Ziai WC, Urrutia VC, Cooper LA and Gottesman RF. Racial differences in palliative care use after stroke in majority-white, minority-serving, and racially integrated U.S. hospitals. *Crit Care Med* 2017; 45: 2046–2054.
 28. Ormseth CH, Falcone GJ, Jasak SD, et al. Minority patients are less likely to undergo withdrawal of care after spontaneous intracerebral hemorrhage. *Neurocrit Care* 2018; 29: 419–425.
 29. Williams MT, Zimmerman E, Barry M, et al. A retrospective review of patients with acute stroke with and without palliative care consultations. *Am J Hosp Palliat Med* 2019; 36: 60–64.
 30. Singh T, Peters SR, Tirschwell DL and Creutzfeldt CJ. Palliative care for hospitalized patients with stroke: results from the 2010 to 2012 national inpatient sample. *Stroke* 2017; 48: 2534–2540.
 31. Cross SH, Kaufman BG, Mentz RJ, Kamal AH, Taylor DH and Warraich HJ. Trends in place of death for individuals with cardiovascular disease in the United States. *J Am Coll Cardiol* 2019; 74: 1943–1946.
 32. Jennings N, Chambaere K, Deliens L and Cohen J. Place of death in a small island state: a death certificate population study. *BMJ Support Palliat Care* 2019; 2020; 10: e30. DOI: 10.1136/bmjspcare-2018-001631.
 33. Raijmakers NJH, de Veer AJE, Zwaan R, Hofstede JM and Francke AL. Which patients die in their preferred place? A secondary analysis of questionnaire data from bereaved relatives. *Palliat Med* 2018; 32: 347–356.
 34. Asplund K, Lundstrom S and Stegmayr B. End of life after stroke: a nationwide study of 42,502 deaths occurring within a year after stroke. *Eur Stroke J* 2018; 3: 74–81.
 35. Allida S, Cox KL, Hsieh CF, House A and Hackett ML. Pharmacological, psychological and non-invasive brain stimulation interventions for preventing depression after stroke. *Cochrane Database Syst Rev* 2020; 5: CD003689.
 36. Visvanathan A, Dennis M and Whiteley W. Parenteral fluid regimens for improving functional outcome in people with acute stroke. *Cochrane Database Syst Rev* 2015; 9: CD011138.
 37. Meisel A. Managing aftercare for stroke (MAS): MAS-II – a longitudinal complex-interventional study in post-rehabilitation stroke patients, <https://clinicaltrials.gov/ct2/show/NCT03097146> (2017, accessed 22 June 2020).
 38. Hospices Civils de Lyon. Evaluation of the impact of an individual peer support intervention for stroke patients when returning home: a mixed methods pilot study, <https://clinicaltrials.gov/ct2/show/NCT04197258> (2019, accessed 22 June 2020).
 39. Hospices Civils de Lyon. Development and evaluation of a patient-centered transition program for stroke patients, combining case management and access to an internet information platform, <https://clinicaltrials.gov/ct2/show/NCT03956160> (2019, accessed 22 June 2020).

40. Ajayi I, Iken O, Powell RA, Soyannwo O, Namisango E and Mwangi-Powell F. Palliative care research in north-ern Africa. *Eur J Palliat Care* 2014; 21: 45–47.
41. de Villiers M, Maree JE and Van Belkum C. Palliative care needs of people living in a newly established informal settlement. *Afr J Nurs Midwifery* 2018; 20: 1–7.
42. Hossain MJ. *To Investigate the need for palliative care in cerebrovascular accident (stroke) patients at Ladysmith Regional Hospital*. Master of Philosophy in Palliative Care Medicine Thesis, University of Cape Town, Cape Town, South Africa, 2016.
43. Akinpelu AO, Olaleye OA, Odole AC and Otaiku OA. Informal stroke caregivers' satisfaction with healthcare services in a tertiary healthcare centre in Ibadan, Nigeria. *Int J Caring Sci* 2014; 7: 148–156.
44. Vincent-Onabajo G and Mohammed Z. Preferred rehabilitation setting among stroke survivors in Nigeria and associated personal factors. *African J Disabil* 2018; 7: 352. DOI: 10.4102/ajod.v7i0.352.
45. Nweke MC and Eze CK. The Place of spiritual and traditional beliefs in stroke rehabilitation in sub-saharan africa: a scoping review. *J Complement Altern Med Res* 2019; 8: 1–16.
46. Totten A. Team-based versus primary care clinician-led advance care planning in practice-based research networks. US Natl. Libr. Med, <https://clinicaltrials.gov/ct2/show/NCT03577002> (2018, accessed 21 August 2020).
47. Muehlschlegel S. Decision aid feasibility trial for families of critically ill stroke patients. US Natl. Libr. Med, <https://clinicaltrials.gov/ct2/show/NCT04143113> (2019, accessed 21 August 2020).
48. Green T, Gandhi S, Kleissen T, Simon J, Raffin-Bouchal S and Ryckborst K. Advance care planning in stroke: influence of time on engagement in the process. *Patient Prefer Adherence* 2014; 8: 119–126.
49. Hancock K, Clayton JM, Parker SM, et al. Truth-telling in discussing prognosis in advanced life-limiting illnesses: a systematic review. *Palliat Med* 2007; 21: 507–517.
50. Schichtel M, Wee B, Perera R and Onakpoya I. The effect of advance care planning on heart failure: a systematic review and meta-analysis. *J Gen Intern Med* 2020; 35: 874–884.
51. Marinho V, Pinto GR, Bandeira J, et al. Impaired decision-making and time perception in individuals with stroke: behavioral and neural correlates. *Rev Neurol* 2019; 175: 367–376.
52. Vickers AJ. Decisional conflict, regret, and the burden of rational decision making. *Med Decis Mak* 2017; 37: 3–5.
53. Armstrong MJ. Shared decision-making in stroke: an evolving approach to improved patient care. *Stroke Vasc Neurol* 2017; 2: 84–87.
54. Doubal F, Cowey E, Bailey F, et al. The key challenges of discussing end-of-life stroke care with patients and families: a mixed-methods electronic survey of hospital and community healthcare professionals. *J R Coll Physicians Edinb* 2018; 48: 217–224.
55. Forster A, Brown L, Smith J, et al. Information provision for stroke patients and their caregivers. *Cochrane Database Syst Rev* 2012; 11: CD001919.
56. Stacey D, Légaré F, Lewis K, et al. Decision aids for people facing health treatment or screening decisions. *Cochrane Database Syst Rev* 2017; 4: CD001431.
57. Sutter R, Meyer-Zehnder B, Baumann SM, Marsch S and Pargger H. Advance directives in the neurocritically ill: a systematic review. *Crit Care Med* 2020; 48(8): 1188–1195. DOI: 10.1097/CCM.0000000000004388.
58. Johnson PD, Ulrich A, Siv J, Taylor B, Tirschwell D and Creutzfeldt C. Planning after stroke survival: advance care planning in the stroke clinic. *J Am Heart Assoc* 2019; 8: e011317. DOI: 10.1161/JAHA.118.011317.
59. Murtagh MJ, Burges Watson DL, et al. Situationally-sensitive knowledge translation and relational decision making in hyperacute stroke: a qualitative study. *PLoS One* 2012; 7: e37066.
60. de Boer ME, Depla M, Wojtkowiak J, et al. Life-and-death decision-making in the acute phase after a severe stroke: interviews with relatives. *Palliat Med* 2015; 29: 451–457.
61. Wheeler NC, Murali S and Sattin JA. Ethical issues in vascular neurology. *Semin Neurol* 2018; 38: 515–521.
62. de Kort FAS, Geurts M, de Kort PLM, et al. Advance directives, proxy opinions, and treatment restrictions in patients with severe stroke. *BMC Palliat Care* 2017; 16: 52.
63. Trevick SA and Lord AS. Post-traumatic stress disorder and complicated grief are common in caregivers of neuro-ICU patients. *Neurocrit Care* 2017; 26: 436–443.
64. Creutzfeldt CJ, Holloway RG and Curtis JR. Palliative care: a core competency for stroke neurologists. *Stroke* 2015; 46: 2714–2719.
65. Flynn D, Nesbitt DJ, Ford GA, et al. Development of a computerised decision aid for thrombolysis in acute stroke care. *BMC Med Inform Decis Mak* 2015; 15: 6. DOI: 10.1186/s12911-014-0127-1.
66. Flynn D, Ford GA, Stobbart L, Rodgers H, Murtagh MJ and Thomson RG. A review of decision support, risk communication and patient information tools for thrombolytic treatment in acute stroke: lessons for tool developers. *BMC Health Serv Res* 2013; 13: 225.