

The association between lunch composition and obesity in Iranian adults

Akbarzade, Z., Djafarian, K., Saeidifard, N., Aliakbari Majd, S., Garousi, N., Samadi, F., Jebraeili, H., Chamari, M., Clark, C. C. T. & Shab-Bidar, S.

Author post-print (accepted) deposited by Coventry University's Repository

Original citation & hyperlink:

Akbarzade, Z, Djafarian, K, Saeidifard, N, Aliakbari Majd, S, Garousi, N, Samadi, F, Jebraeili, H, Chamari, M, Clark, CCT & Shab-Bidar, S 2022, 'The association between lunch composition and obesity in Iranian adults', *British Journal of Nutrition*, vol. 127, no. 10, pp. 1517-1527.

<https://dx.doi.org/10.1017/S0007114521002543>

DOI 10.1017/S0007114521002543

ISSN 0007-1145

ESSN 1475-2662

Publisher: Cambridge University Press

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.

The association between lunch composition and obesity in Iranian adults

Zahra Akbarzade¹, Kurosh Djafarian², Nasim Saeidifard¹, Shabnam Aliakbari Majd³, Nazila Garousi⁴, Fatemeh Samadi⁵, Hanieh Jebraeili¹, Maryam Chamari¹, Cain C. T. Clark⁶, Sakineh Shab-Bidar^{1*}

¹Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences (TUMS), Tehran, Iran.

²Department of Clinical Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran

³Bou Ali Sina Medical Training Center, Qazvin university of Medical Science, Qazvin, Iran

⁴Food Security Research Center; Department of Clinical Nutrition, Isfahan University of Medical Sciences, Isfahan, Iran

⁵Student Research Committee, Department of Clinical Nutrition and Dietetics, Faculty of Nutrition and Food Technology, National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁶Centre for Sport, Exercise, and Life Sciences, Coventry University, Coventry, CV15FB, U.K.

***Corresponding author:** Sakineh Shab-Bidar, Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences (TUMS), Tehran, Iran, No 44, Hojjat-dost Alley, Naderi St., Keshavarz Blvd, Tehran, Iran. Tele: +989111376516, E-mail address: s_shabbidar@tums.ac.ir



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114521002543

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Abbreviations:

BMI: Body mass index

24hDRs: 24-h dietary recalls

PCA: Principal Component Analysis

WC: Waist circumference

WHR: waist-to-hip ratio

IPAQ: International short-form Physical Activity Questionnaire

ANOVA: one-way analysis of variance

Short title: Lunch consumption patterns and obesity

Abstract

We aimed to assess the dietary composition of lunch meal using a *posteriori* derived dietary patterns and to determine the association of lunch composition with obesity in a sample of Iranian adults. This cross-sectional study was conducted on 850 men and women in Tehran (aged 20-59 y). Dietary intakes were assessed using three 24-h dietary recalls, and dietary patterns were identified via principal component factor analysis. For each identified pattern, scores were calculated for each participant and then classified into tertiles. Central obesity was defined WHO criteria. General obesity was defined as a body mass index of more than 30 kg/m². Three major dietary patterns were identified at lunch meal using 12 food groups: “Bread, grains and fat”, “Western”, and “Potato and eggs”. After adjustment for potential confounders, participants at the top tertile of the “Bread, grains and fat” dietary pattern had greater odds for a higher waist to hip ratio, compared with those in the lowest tertile (OR, 1.44; 95% CI: 1.01 to 2.07). However, we found no association between ‘Western or “potato and eggs” patterns and waist to hip ratio (OR 0.89, 95% CI: 0.62 to 1.28 and OR 1.16, 95% CI: 0.69 to 1.42, respectively). None of the identified dietary patterns was associated when defining obesity with waist circumference or body mass index. In conclusion, participants had a greater chance of central obesity defined based on waist to hip ratio following a lunchtime pattern with a higher and positive loading factor for “Bread, grains and fat”.

Keywords: Lunch pattern; Obesity; Body mass index; Waist circumference; Dietary patterns

Introduction

Obesity represents one of the most important metabolic diseases worldwide. In recent decades, the prevalence of overweight and obesity has doubled, globally, whilst it has been estimated that by the year 2030, the number of obese people will reach 1.21 billion ⁽¹⁾. In addition, in Iran, almost 21.7% of adults are affected by obesity ⁽²⁾. Body mass index (BMI) is one of the most common and simple methods used in many epidemiological studies, which has been proposed as a tool for screening and early clinical evaluation of obesity ⁽³⁾. However, BMI has some limitations since it is an indicator of total body fat and does not provide information about localized abdominal fat that is associated with metabolic disease. Some studies showed that the pattern of fat distribution in the body plays a key role in identifying chronic disease ⁽⁴⁾ and people with fat accumulation in the abdomen are at a higher risk of diabetes, hypertension, and cardiovascular disease ⁽⁵⁾. In this regard, other anthropometric indicators such as waist circumference (WC) and waist-hip ratio (WHR) are superior to BMI for revealing obesity and overweight. WC and WHR are mostly used to express how fat is distributed in the body, and WHR is widely applied to distinguish between the central and peripheral distribution of adipose tissue ^(6; 7).

Socio-demographic determinants of obesity have been of great research interest recently. It is demonstrated that certain health behaviors including smoking and physical activity are associated with obesity ^(8; 9) and it has been reported differences in obesity prevalence across various gender, age and socioeconomic groups ⁽¹⁰⁾. Socioeconomic status is usually measured by education, occupation, employment, income, and wealth. Overall, higher welfare level and higher and higher socioeconomic status were observed to be associated with a lower BMI, WC, and WHR ⁽¹¹⁾. This may be due to a healthier diet in subjects with a higher level of education, income, and socioeconomic status ⁽¹²⁾, as well as a higher degree of physical activity ⁽¹¹⁾.

Diet is the fundamental component of a healthy lifestyle which can play a significant role in the prevention of non-communicable diseases ⁽¹³⁾. Previous studies have tended to focus on individual foods or nutrients, however, given the complexity of human diets, several authors have proposed the analysis of overall dietary patterns ^(14; 15). Moreover, the relationship between dietary patterns, considering the complexity of diets and the potential interaction between food components, and obesity has been well documented in the literature ^(16; 17; 18). Findings, such as the association between the Mediterranean diet and low rates of chronic diseases, or the

successful treatment of hypertension through changes in dietary patterns, have suggested the investigation of dietary intake patterns^(19; 20). Apart from well-founded evidence regarding the benefits of vegetables, fruits, fiber, nuts, and fish, and the value of reducing or eliminating snacks; people regularly consume combinations of foods in three or more meals per day^(13; 21). The main meals are described as foods that are typically consumed in the largest volume^(22; 23). It is important to acknowledge that different nutritional compositions in main meals may have an impact on diet quality and influence diet-disease relationships. There is an association between meal patterns and energy balance and weight status^(24; 25). Results of a study in Brazilian population showed that the traditional Brazilian lunch pattern is inversely associated with obesity in insufficiently active individuals⁽²⁶⁾. In another study, breakfasts containing >25 % of total energy intake and lunches containing >35 % of total energy intake were associated with an increased likelihood of central obesity⁽²⁷⁾. Indeed, previous studies have shown that the lunch-time meal represents the highest proportion of protein, fat, and carbohydrate intake⁽²⁸⁾ and supplies about 30% of the daily energy intake⁽²⁹⁾. Therefore, this study aimed to, first, identify major dietary patterns at lunch, and, second, to determine their association with obesity among adults living in Tehran.

Subjects and methods

Participants

This cross-sectional study was conducted using a sample of 850 men and women, aged 20-59, y who have visited the health center and were been selected from the five regions of Tehran from 2018 to 2019. The following formula was used for sample size calculation: $n = (pqz^2)/E^2$. Considering the total prevalence of 65% for overweight and obesity⁽³⁰⁾, an error coefficient of $d=0.04$ and at a level of 0.05, the sample size of 546 people was obtained. With a design effect of 1.5 and to compensate for the potential exclusion of participants due to under- and over-reporting of total energy intake, or attrition due to other reasons, the final sample size of 850 participants was selected for inclusion. A two-stage cluster sampling was used to recruit participants from health care centers. First, we classified health centers into five districts of the city including North, South, East, West, and center. Next, a list of all health centers that existed in each district was provided. Then, 25 health centers (due to budget and time limits) were divided according to the number of health centers in each area. After a randomized selection of health care centers from enlisted health centers in each region, the number of eligible health centers was randomly

selected. Subsequently, the total sample (850) was divided by the number of health centers (25), yielding the required number of samples in each health center. Following this, we entered each of the health centers, and those who willing to participate in our study, those who were members of the health center and living in Tehran were included and conducted random sampling to ascertain the required number of samples needed and on the other hand, at the end of the interview with the individuals, nutritional information appropriate to each person's conditions was provided to them. Subjects were considered eligible for inclusion if the following criteria were met: a) participants within the age range of 20-59 years; b) apparently healthy individuals who did not report any previous diagnosis of chronic diseases such as diabetes, cardiovascular diseases and chronic kidney, lung and liver diseases by a physician; c) be willing to take part in the study; d) being a resident of Tehran, and e) being a member of the health center. Participants were excluded from the analysis if a) their daily energy intake was implausibly low or high (<800 Kcal/day or >4200 Kcal/day); and b) those who did not report any adherence to certain dietary patterns, any special diet or diet therapy such as vegetarian diet. The sample collection was facilitated by the coordination of the Health Bureau of the Municipality of Tehran and the cooperation of the health centers of Tehran.

Outcome measures

Weight and height were measured according to standard methods⁽³¹⁾. Weight was recorded to the nearest 100 grams while minimally clothed and unshod using digital scales. Height was measured in a standing position, unshod, using a tape stadiometer to the nearest 1 mm, while the shoulders were maintained in a normal position. Body mass index (BMI) was calculated using the following formula; weight in kg, divided height in meters squared, expressed as kg/m^2 . General obesity was defined as $\text{BMI} \geq 30 \text{ kg/m}^2$ ⁽³²⁾. WC was recorded to the nearest 0.1 cm at the umbilical level and hip circumference at the maximal point over light clothing, using a non-stretch tape meter, without putting pressure on the body surface. Central obesity was defined as $\text{WC} \geq 102$ in men and $\text{WC} \geq 88$ in women, respectively, and a WHR above 0.90 for males and above 0.85 for females, respectively⁽³³⁾. Participants rested for 15 minutes before blood pressure (BP) was measured. Then, a trained assessor measured BP twice, with the participant in a seated position, with a standard mercury sphygmomanometer, and the mean of two measurements was calculated.

Dietary assessment

Dietary intake was assessed using three 24-h dietary recalls (24hDRs). The first recall took place during the participants' first visit to the health center. The other two 24hDRs were obtained at random days, including weekends over the phone. All 24hDRs were performed by trained interviewers. Any food or beverage that the participants consumed during the denominated meal-time was considered. Finally, the standard unit size and items reported based on home weighing scales were converted into grams using the home scale guideline⁽³⁴⁾. The data from these questionnaires were entered into a purpose-built excel spreadsheet, where the gram equivalent was obtained for each item and each individual. In addition, we used Nutritionist IV software (First Databank, San Bruno, CA), modified for Iranian foods, to analyze the energy and nutrients of food items. Lunch was predefined as a large meal eaten between 12:00 and 16:00⁽³⁵⁾. Foods were grouped according to similar nutritional values, Iranian consumption habits, literary data, and experience of the research team in previous studies^(36; 37; 38). Some individual food items that consisted of separate items (e.g. eggs) or that represented special dietary habits (such as potatoes) were retained as a single food. Moreover, it should be noted that some single FFQ items (salt and potatoes) considered as a single group because in the FFQ there were no other food items to be appropriate for combination with these foods into multiple-item food groups. Finally, we created 12 pre-determined dietary groups (bread and grains^(39; 40), dairy products^(41; 42), poultry^(43; 44), eggs^(45; 46), fat^(47; 48), potatoes^(49; 50), processed meat and red meat^(51; 52), soft drinks^(53; 54), vegetables⁽⁵⁵⁾, legumes and nuts⁽⁵⁶⁾, salt⁽⁵⁷⁾ and sauces⁽⁵⁸⁾) which also had an association with obesity. Also, for some foods like potato, eating habits of our population has been considered. As white potato is a good source of carbohydrate, dietary fiber, and resistant starch, it is a favorite staple food in several cultures as well as a good source of vitamin C and potassium, especially in Iran. Moreover, due to the high content of carbohydrate, it is suspected to have a link to obesity, and most common foods with potato in Iran contain more fat calories than carbohydrate calories. Then, we decided to consider potato as a separate food group in our study.

Assessment of other variables

Physical activity information was obtained using participants' oral responses to the international physical activity questionnaire and expressed as the metabolic equivalent h/wk (MET-h/wk)⁽⁵⁹⁾. We asked the participants to think about all the intense and moderate activities that they engaged

in during the past 7 days, considering the time spent on these activities, before completing the questionnaire. Additional covariates, including age (year), BMI (kg/m²), an education level (illiterate, under diploma (Primary School, Secondary School, High School), diploma, University degree), marital status (married or other), occupation (employee or unemployed), medical condition (healthy or underlying disease), smoking status (not smoking, quit smoking, smoker), and lifestyle (living alone, with someone) was obtained using questionnaires. Furthermore, we defined underlying diseases in this study as diabetes, hypertension, dyslipidemia, cardiovascular disease, cancer, and respiratory disease. According to the self-reporting of participants in the study, if a person had one of the diseases considered in this research, he/she would receive code 1 and otherwise code 0. Finally, if the individual did not have any of the respective diseases, he/she was classified in the healthy group, and those who even had one of the diseases were assigned to the group with underlying disease.

Statistical analyses

Factor analysis (principal component analysis) was used to identify major lunch consumption patterns based on 12 pre-determined dietary groups, and two interpretable factors were retained based on the scree test ⁽⁶⁰⁾. Then an orthogonal rotation method (Varimax rotation) was applied to simplify the factor structure and present it in an interpretable manner. The number of factors retained from each dietary pattern classification method was determined by eigenvalues (>1.10), scree plots, and factor interpretability. Higher loadings (≥ 0.2) show that the food shares more variance with that factor. The derived factors (lunch patterns) were labelled based on our interpretation of the data, as well as on prior literature. The factor score for each pattern was calculated by collecting consumed food groups, weighted by factor loadings, and each participant determined the score for each identified pattern. Pearson correlation coefficients were also computed to identify the association between food groups. We used one-way analysis of variance (ANOVA), with Tukey post hoc comparisons where appropriate, for quantitative variables, and chi-square tests for qualitative variables, to determine significant differences across tertiles of lunch pattern scores. The association between major dietary patterns and general and central obesity were assessed by logistic regression analysis. Analysis of covariance (ANCOVA) was used to adjust for covariates as follows; in the first model for age, sex, education, marriage, lifestyle, and smoking, whilst the second model was adjusted for Model 1 plus physical activity and total energy intake. Logistic regression analysis for general and central

obesity, according to lunch patterns, was used to obtain the odds ratio and 95% confidence interval, which were adjusted for potential confounders, including age, sex, education, marriage, lifestyle, smoking; physical activity and total energy intake. Confounders were selected based on literature review including age (years), gender (male or female), physical activity level, smoking status (never smoke or former/current smoker), and total energy intake.

All data were analyzed using the statistical software package SPSS version 22, and statistical significance was accepted at $P < 0.05$.

Results

Of the 850 participants who enrolled in the study, 60 were excluded due to a lack of adequate information and lack of co-operation in their recall report, thus, 790 remained in the study for final analysis. In general, the average calorie intake at lunchtime was higher than other meals (**Figure 1**).

Using the factor analysis method, three major dietary patterns were identified and presented in **Table 1**. Factor 1, named the " Bread, grains and fat" dietary pattern, had high and positive factor loadings for fats, bread and grains, salt, vegetables, poultry and fish and high negative factor loadings for legumes and nuts and sauces; Factor 2, named the "Potato and eggs" dietary pattern had positive factor loadings for potato and eggs, and high negative factor loadings for intake of bread and grains, nuts and legumes; and finally Factor 3, which named the "Western" dietary pattern showed positive factor loadings for red or processed meat, sauces, soft drinks, nuts and legumes and negative factor loadings for poultry and fish and dairy products.

Table 2 provides an overview of the distribution of qualitative and quantitative variables across tertiles of major dietary patterns. No significant difference was found in the distribution of qualitative variables across the tertiles of three major dietary patterns. Adherence to the "Potato and eggs" pattern was associated with an increase in age ($p=0.04$) and blood pressure ($p=0.03$). Mean intake of vegetables ($p<0.001$), dairy products ($p=0.01$), grains ($p<0.001$), and meat ($p=0.006$) were higher at the third tertiles of the "Bread, grains and fat" dietary pattern in comparison to the first tertile. Moreover, adherence to the "Eggs and potato" dietary pattern was associated with the higher intake of vegetable ($p<0.001$) and grain ($p<0.001$). In addition, the mean intake of dairy products, grains and meat were significantly different across the tertiles of "Western" dietary pattern.

Heat map shows the Pearson correlation matrix of foods groups at lunchtime. Correlation analysis showed that there were strong positive correlations between potato and eggs ($p < 0.001$), whilst strong negative correlations were found between nuts and legumes and fish and poultry ($p < 0.001$), and between meat and processed meat and fish and poultry ($p < 0.001$) (**Figure 2**).

Multivariable-adjusted means for anthropometric measures and indexes across tertiles of dietary patterns are depicted in **Table 3**. The results showed that higher adherence to the “Bread, grains and fat” dietary pattern was associated with a higher WHR ($P = 0.04$). In addition, the mean weight significantly increased across the “Egg and potato” dietary pattern ($p = 0.02$). However, there was no significant relationship after controlling for confounding factors. Also, there was no significant difference in means and SD of other anthropometric measures across major dietary patterns at lunchtime.

Unadjusted and adjusted odds ratios (OR) for the participants’ general obesity in the tertiles of lunch patterns are presented in **Table 4**. According to our findings, after control for confounders, there was no significant associations between “Bread, grains and fat” pattern (OR: 0.89, 95% CI = 0.60, 1.33, P -value = 0.59), “Eggs and potato” pattern (OR: 1.07, 95% CI = 0.71, 1.60, P -value = 0.36) and the Western dietary pattern (OR: 1.32, 95% CI = 0.88, 1.99, P -value = 0.17) with general obesity.

Table 5 shows the unadjusted and adjusted odds ratios for central obesity across the tertiles of major dietary patterns at lunchtime. In the unadjusted model, no significant association was found between central obesity, based on WC definition, and dietary patterns which remained unchanged even after adjustment for confounders. No significant difference was also observed in the odds of central obesity, when defined based on a WHR, across the tertiles of the “Bread, grains and fat” dietary pattern (odds ratio: 1.38, 95% confidence interval: 0.98 - 1.95). However, after adjustment for confounders, the participants in the top tertile of the “Bread, grains and fat” pattern had a 1.44 times higher chance of central obesity (odds ratio: 1.44; 95% confidence interval: 1.01 - 2.07) relative to individuals in the first tertile. No significant associations were observed between adherence to the “Western” pattern and “Eggs and potato” patterns with OR of central obesity in our population

Discussion

The present study, which was conducted on 850 adults in Tehran, showed that 25.4% of subjects had general obesity. In addition, 46.6% and 54.8% of subjects had central obesity based on WC and WHR, respectively. The results of this study showed that participants in the top tertile of the “Bread, grains and fat” pattern had a 44% higher risk of increased WHR compared with the participants in the lowest tertile. There were no significant associations between the “Western” and “Eggs and potato” patterns at lunchtime and central obesity. Moreover, our findings showed that there was no relationship between major dietary patterns at lunchtime and general obesity.

Socio-demographic determinants of obesity have been of great research interest recently. It is demonstrated that certain health behaviors including smoking and physical activity are associated with obesity ^(8;9). Diet, total energy intake, physical activity, sedentary lifestyles, and other health-risk behaviors are known to act as proximate, intervening variables in the relationship between socioeconomic status and obesity ⁽⁶¹⁾. In addition, it was found that obesity prevalence varies across various gender, age and socioeconomic groups ⁽¹⁰⁾. Socioeconomic status is usually measured by education, occupation, employment, income, and wealth. For example, lower education, lower occupational status, and lower incomes have been associated with a higher prevalence of obesity ⁽⁶²⁾. It has been found that smokers are likely to be obese ⁽⁶³⁾. It has been shown that smoking increases levels of cortisol and testosterone, whereas the levels of estradiol and progesterone are decreased ⁽⁶⁴⁾. Furthermore, it has been reported that smoking changes dietary habits, leading to a lower intake of fibers, fruits, and vegetables ⁽⁶⁴⁾. Physical activity has also been established as one of the important predictors of weight gain. In that, those who have enough physical activity, especially during leisure time, are less likely to be obese ⁽⁹⁾. Given the complex combinations of nutrients involved in the human diet, identifying dietary patterns may represent the best way to highlight the effects of nutrients and specific foods on health ^(65;66), permitting insight into the synergistic outcomes of nutrients and foods. However, most previous studies have investigated diet in general, regardless of the timing of food intake, and consequently, they were unable to identify specific meal characteristics and compositions ⁽⁶⁷⁾. Bellisle et al., reported that the greatest nutrient consumption over an average day occurs around midday, corresponding to lunch ⁽²³⁾. Therefore, due to the large proportion of the daily energy intake attributed to lunch, such meals should, ideally, provide sufficient amounts of macro and micronutrients to help achieve and adhere to dietary guidelines.

Dietary patterns were specified based on factor analysis using the principal component analysis (PCA) method with varimax rotation. The resulting factors were judged based on the eigenvalues (the total squared load factor of food items in terms of their consumption in grams per day) of food groups, and any factor with an eigenvalue >1 was considered as the main dietary pattern. Load factor values ≥ 0.2 were used to describe the main food groups that make up each food pattern.

One of the possible reasons for the difference between the diet patterns of the present study with other studies is that the analysis of dietary patterns is strongly dependent upon the study population. Therefore, significant differences in dietary patterns of different populations are observed by geographical area, race and culture. In addition, factor analysis is limited by the researcher's choice of food items' grouping and the number of factors to be retained, and these decisions can somewhat affect the findings and their interpretation.

In the present study, three dominant lunch consumption patterns were identified among the participants: Dietary pattern 1 ("Bread, grains and fat") was rich in fats, bread and grains, salt, poultry and fish, Dietary pattern 2 ("Eggs and potatoes") included a high intake of potato and eggs, and a low intake of bread and grains, nuts and legumes, and Dietary pattern 3 ("Western") included a high intake of red or processed meat, sauces, soft drink, nuts and legumes and a low intake of poultry and fish and dairy products. Santos et al. previously identified five lunch patterns, among which the "Western" dietary pattern (positive factor loadings for soft drinks, alcoholic beverages, sweets, gnocchi/stuffed pasta, sauces/mayonnaise, and processed meats) was relatively similar to our "Western" dietary pattern⁽⁶⁸⁾. In addition, our "Bread, grains and fat" dietary pattern was similar to Santos' "meat" dietary pattern (positive factor loadings for eggs, poultry meat, and fish/seafood; negative for beef). However, some of the items found in Santos' "meat" dietary pattern (like eggs) were also found in our other dietary patterns. Similarly, Schwedhelm et al. also identified five lunch meal patterns⁽⁶⁹⁾, among which the "Western" pattern (high intake of potatoes, cabbage, red meat, beer, sauces and condiments and low intake of fresh fruits, milk and dairy products and tea) was comparable to our "Western" dietary pattern and their "traditional" dietary pattern (high intake of bread, processed meat, butter, sugar, confectionery, cakes and cookies and low intake of water) was similar to our "Bread, grains and fat" dietary pattern. Moreover, Esmailzadeh et al. (Iran) identified three

dietary patterns⁽⁷⁰⁾, among which the “Western” dietary pattern (high in refined grains, red meat, butter, processed meat, high-fat dairy products, sweets and desserts, pizza, potatoes, eggs, hydrogenated fats, and soft drinks and low in other vegetables and low-fat dairy products) was similar to our “Western” dietary pattern and the “Iranian” foods pattern (high in refined grains, potato, tea, whole grains, hydrogenated fats, legumes, and broth) was similar to our “Bread, grains and fat” dietary pattern.

Contradictory results among studies may be attributed to distinct differences in the culture, tradition, and eating habits of the Iranian people compared to other countries. Additionally, differences in dietary patterns may be attributed to the fact that some studies examined habitual dietary patterns, whilst in our study, these patterns were derived at the meal level. The present study revealed no association between the “Western” pattern, as an unhealthy dietary pattern, and obesity; indeed, some previous studies have reported similar findings and found no significant association between the patterns rich in fat and sugar and overweight and obesity^(68; 71; 72). However, it has been reported that adherence to a “Western” or unhealthy pattern dietary increases the risk of overweight and obesity^(73; 74). Given that the “Western” dietary pattern, is considered as an unhealthy pattern, the lack of a significant association with obesity was somewhat unexpected. However, the inclusion of some food groups, such as legumes and nuts. in this pattern might have hindered the identification of an association. In addition, no association was found between the “Egg and potato” pattern and obesity, which may be considered as a healthy diet, given that it is rich in fiber, vitamins, and high-quality protein. Although the healthy dietary patterns found in some previous studies have been inversely associated with obesity and overweight⁽⁷⁵⁾, few studies have shown a positive association between healthy dietary patterns and BMI⁽⁷⁶⁾. However, some studies have also reported no association between healthy eating patterns and weight status^(71; 72), which may conceivably be due to measurement error in the variables under investigation.

In the present study, no significant association was also found between the “Bread, grains and fat” lunch pattern and general obesity. However, participants in the highest tertile had greater odds of increased WHR compared with those in the lowest tertile. The complex nature of this dietary pattern makes interpretation difficult. This pattern contains bread and grains, fats and salt, hence, a positive association was expected between this dietary pattern and obesity. However, some healthy food groups, such as poultry and fish included in this dietary pattern may have

interacted with other foods and counteracted the effects on obesity. On the other hand, it can be argued that the positive association between this dietary pattern and increased WHR is justified by the fact that this food pattern is high in fat and carbohydrates. Low-nutrient density foods can lead to central nervous system insulin resistance, which may, in turn, result in leptin resistance and increased pleasurable responses to foods⁽⁷⁷⁾.

One of the strengths of the present study is the recruitment of a large sample size compared to other studies in Iran. In addition, all districts of Tehran were selected, so that various socioeconomic statuses, educational and welfare levels, and other variables affecting the outcome could be included in the study. The present study also had some potential limitations; for instance, no causal inference could be made due to the cross-sectional nature of the study. Another limitation of this study was that the information regarding the economic conditions of the families could not be collected. As a result, the possible relationship between the economic status of a family with dietary patterns could not be assessed. In addition, the data was collected using retrospective questions, which might have yielded information recall bias.

Conclusion

Greater adherence to the “Bread, grains and fat” dietary pattern at lunchtime was associated with greater odds of central obesity when defined based on WHR. However, no significant association was found between identified lunch patterns and obesity defined by WC and BMI.

Disclosure statements

Acknowledgements: Authors thanks all those who participated in this study. We wish to express our sincere thank Dr. Khalid Iqbal from Department of Human Nutrition, Institute of Basic Medical Sciences, Khyber Medical University, Peshawar, Pakistan, for his helpful suggestions during the study. We also thank Dr. Carolina Schwedhelm, from Department of Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke (DIfE), Nuthetal, Germany for her technical support and help in preparing data.

Financial Support: This manuscript has been granted by Tehran University of Medical Sciences (Grant No: 40186).

Conflict of interest: We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

Authorship: SSb contributed to conception/design of the research; ZA, ShAM, NG, FS, HJ, MCH contributed to acquisition of data. ZA and SS-b participated in analysis and interpretation of the data; ZA drafted the manuscript; KD, SSb and CC critically revised the manuscript; and SS-b agrees to be fully accountable for ensuring the integrity and accuracy of the work. All authors read and approved the final manuscript.

Ethical Standards Disclosure: This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the ethics committee of Tehran University of Medical Sciences. Written informed consent was obtained from all subjects/patients.

References

1. Antelo M, Magdalena P, Reboredo JC (2017) Obesity: A major problem for Spanish minors. *Econ Hum Biol* **24**, 61-73.
2. Rahmani A, Sayehmiri K, Asadollahi K *et al.* (2015) Investigation of the Prevalence of Obesity in Iran: a Systematic Review and Meta-Analysis Study. *Acta medica Iranica* **53**, 596-607.
3. Cornier M-A, Despres J-P, Davis N *et al.* (2011) Assessing adiposity: a scientific statement from the American Heart Association. *Circulation* **124**, 1996-2019.
4. Esmailzadeh A, Mirmiran P, Azizi F (2004) Waist-to-hip ratio is a better screening measure for cardiovascular risk factors than other anthropometric indicators in Tehranian adult men. *International journal of obesity* **28**, 1325-1332.
5. Azizi F, Esmailzadeh A, Mirmiran P *et al.* (2005) Is there an independent association between waist-to-hip ratio and cardiovascular risk factors in overweight and obese women? *International journal of cardiology* **101**, 39-46.
6. Lee CMY, Huxley RR, Wildman RP *et al.* (2008) Indices of abdominal obesity are better discriminators of cardiovascular risk factors than BMI: a meta-analysis. *Journal of clinical epidemiology* **61**, 646-653.
7. Taylor RW, Brooking L, Williams SM *et al.* (2010) Body mass index and waist circumference cutoffs to define obesity in indigenous New Zealanders. *The American journal of clinical nutrition* **92**, 390-397.
8. Gruber J, Frakes M (2006) Does falling smoking lead to rising obesity? *Journal of health economics* **25**, 183-197.
9. Fogelholm M, Kukkonen-Harjula K (2000) Does physical activity prevent weight gain—a systematic review. *Obesity reviews* **1**, 95-111.
10. Paeratakul S, Lovejoy JC, Ryan DH *et al.* (2002) The relation of gender, race and socioeconomic status to obesity and obesity comorbidities in a sample of US adults. *International journal of obesity* **26**, 1205-1210.
11. McCormack GR, Friedenreich C, McLaren L *et al.* (2017) Interactions between neighbourhood urban form and socioeconomic status and their associations with anthropometric measurements in Canadian adults. *Journal of environmental and public health* **2017**.

12. Ruf T, Nagel G, Altenburg H-P *et al.* (2005) Food and nutrient intake, anthropometric measurements and smoking according to alcohol consumption in the EPIC Heidelberg study. *Annals of nutrition and metabolism* **49**, 16-25.
13. Organization WH (2009) *Global health risks: mortality and burden of disease attributable to selected major risks*: Geneva: World Health Organization.
14. Hu FB, Rimm E, Smith-Warner SA *et al.* (1999) Reproducibility and validity of dietary patterns assessed with a food-frequency questionnaire. **69**, 243-249.
15. Hu FB, Rimm EB, Stampfer MJ *et al.* (2000) Prospective study of major dietary patterns and risk of coronary heart disease in men. **72**, 912-921.
16. Cunha DB, de Almeida RMVR, Sichieri R *et al.* (2010) Association of dietary patterns with BMI and waist circumference in a low-income neighbourhood in Brazil. **104**, 908-913.
17. Naja F, Hwalla N, Itani L *et al.* (2015) A Western dietary pattern is associated with overweight and obesity in a national sample of Lebanese adolescents (13–19 years): a cross-sectional study. **114**, 1909-1919.
18. Shu L, Zheng P-F, Zhang X-Y *et al.* (2015) Association between dietary patterns and the indicators of obesity among Chinese: a cross-sectional study. **7**, 7995-8009.
19. Helsing EJTAjocn (1995) Traditional diets and disease patterns of the Mediterranean, circa 1960. **61**, 1329S-1337S.
20. Sacks FM, Svetkey LP, Vollmer WM *et al.* (2001) Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. **344**, 3-10.
21. Hu FB (2002) Dietary pattern analysis: a new direction in nutritional epidemiology. *Current opinion in lipidology* **13**, 3-9.
22. Leech RM, Worsley A, Timperio A *et al.* (2015) Understanding meal patterns: definitions, methodology and impact on nutrient intake and diet quality. **28**, 1-21.
23. Bellisle F, Dalix AM, Mennen L *et al.* (2003) Contribution of snacks and meals in the diet of French adults: a diet-diary study. **79**, 183-189.
24. Mesas A, Muñoz-Pareja M, López-García E *et al.* (2012) Selected eating behaviours and excess body weight: a systematic review. *Obesity Reviews* **13**, 106-135.
25. Szajewska H, Ruszczyński M (2010) Systematic review demonstrating that breakfast consumption influences body weight outcomes in children and adolescents in Europe. *Critical reviews in food science and nutrition* **50**, 113-119.

26. de Oliveira Santos R, dos Santos Vieira DA, Miranda AAM *et al.* (2018) The traditional lunch pattern is inversely correlated with body mass index in a population-based study in Brazil. *BMC Public Health* **18**, 1-7.
27. Aparicio A, Rodríguez-Rodríguez EE, Aranceta-Bartrina J *et al.* (2017) Differences in meal patterns and timing with regard to central obesity in the ANIBES ('Anthropometric data, macronutrients and micronutrients intake, practice of physical activity, socioeconomic data and lifestyles in Spain') Study. *Public health nutrition* **20**, 2364-2373.
28. Almoosawi S, Winter J, Prynne C *et al.* (2012) Daily profiles of energy and nutrient intakes: are eating profiles changing over time? **66**, 678.
29. Benelam B, Stanner SJNb (2015) Development of a methodology to assess the nutrient profile of popular UK meals. **40**, 315-325.
30. Ahmad Kiadaliri A, Jafari M, Vaez Mahdavi MR *et al.* (2015) The prevalence of adulthood overweight and obesity in Tehran: findings from Urban HEART-2 study. *Medical journal of the Islamic Republic of Iran* **29**, 178.
31. Lohman TG, Roche AF, Martorell R (1988) *Anthropometric standardization reference manual*. vol. 177: Human kinetics books Champaign, IL.
32. (2019) Obesity. <https://www.who.int/topics/obesity/en/>
33. Organization WH (2011) Waist circumference and waist-hip ratio: report of a WHO expert consultation, Geneva, 8-11 December 2008.
34. Ghaffarpour M, Houshiar-Rad A, Kianfar HJTNOK (1999) The manual for household measures, cooking yields factors and edible portion of foods. **7**, 213.
35. Kahleova H, Lloren JI, Mashchak A *et al.* (2017) Meal frequency and timing are associated with changes in body mass index in Adventist health study 2. **147**, 1722-1728.
36. Karamati M, Jessri M, Shariati-Bafghi S-E *et al.* (2012) Dietary patterns in relation to bone mineral density among menopausal Iranian women. *Calcified tissue international* **91**, 40-49.
37. Azadbakht L, Esmailzadeh A (2012) Dietary patterns and attention deficit hyperactivity disorder among Iranian children. *Nutrition* **28**, 242-249.
38. Lockheart MS, Steffen LM, Rebnord HM *et al.* (2007) Dietary patterns, food groups and myocardial infarction: a case-control study. *British Journal of Nutrition* **98**, 380-387.
39. Serra-Majem L, Bautista-Castaño I (2015) Relationship between bread and obesity. *British Journal of Nutrition* **113**, S29-S35.

40. Liu S, Willett WC, Manson JE *et al.* (2003) Relation between changes in intakes of dietary fiber and grain products and changes in weight and development of obesity among middle-aged women. *The American journal of clinical nutrition* **78**, 920-927.
41. Wang W, Wu Y, Zhang D (2016) Association of dairy products consumption with risk of obesity in children and adults: a meta-analysis of mainly cross-sectional studies. *Annals of epidemiology* **26**, 870-882. e872.
42. Barba G, Russo P (2006) Dairy foods, dietary calcium and obesity: a short review of the evidence. *Nutrition, metabolism and cardiovascular diseases* **16**, 445-451.
43. Donma MM, Donma O (2017) Beneficial effects of poultry meat consumption on cardiovascular health and the prevention of childhood obesity. *Med One* **2**.
44. Petermann-Rocha F, Parra-Soto S, Gray S *et al.* (2021) Vegetarians, fish, poultry, and meat-eaters: who has higher risk of cardiovascular disease incidence and mortality? A prospective study from UK Biobank. *European Heart Journal* **42**, 1136-1143.
45. Liu R, Zhao Y, Li Q *et al.* (2020) Body Fat Mass, Fat Distribution and Egg Consumption: a Population-Based Study in Chinese Adults: Egg consumption and Body Fat in Rural Chinese. *Journal of the American College of Nutrition* **39**, 528-536.
46. Saande CJ, Bries AE, Pritchard SK *et al.* (2020) Whole Egg Consumption Decreases Cumulative Weight Gain in Diet-Induced Obese Rats. *The Journal of nutrition* **150**, 1818-1823.
47. Lissner L, Heitmann BL (1995) Dietary fat and obesity: evidence from epidemiology. *European journal of clinical nutrition* **49**, 79-90.
48. Gillis L, Kennedy L, Gillis A *et al.* (2002) Relationship between juvenile obesity, dietary energy and fat intake and physical activity. *International Journal of Obesity* **26**, 458-463.
49. Borch D, Juul-Hindsgaul N, Veller M *et al.* (2016) Potatoes and risk of obesity, type 2 diabetes, and cardiovascular disease in apparently healthy adults: a systematic review of clinical intervention and observational studies. *The American journal of clinical nutrition* **104**, 489-498.
50. Schwingshackl L, Schwedhelm C, Hoffmann G *et al.* (2019) Potatoes and risk of chronic disease: A systematic review and dose-response meta-analysis. *European journal of nutrition* **58**, 2243-2251.
51. Rouhani M, Salehi-Abargouei A, Surkan P *et al.* (2014) Is there a relationship between red or processed meat intake and obesity? A systematic review and meta-analysis of observational studies. *Obesity Reviews* **15**, 740-748.

52. Dabbagh-Moghadam A, Mozaffari-Khosravi H, Nasiri M *et al.* (2017) Association of white and red meat consumption with general and abdominal obesity: a cross-sectional study among a population of Iranian military families in 2016. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity* **22**, 717-724.
53. Gibson S (2008) Sugar-sweetened soft drinks and obesity: a systematic review of the evidence from observational studies and interventions. *Nutrition research reviews* **21**, 134-147.
54. James J, Kerr D (2005) Prevention of childhood obesity by reducing soft drinks. *International journal of obesity* **29**, S54-S57.
55. He K, Hu F, Colditz G *et al.* (2004) Changes in intake of fruits and vegetables in relation to risk of obesity and weight gain among middle-aged women. *International journal of obesity* **28**, 1569-1574.
56. Martínez-González M, Bes-Rastrollo M (2011) Nut consumption, weight gain and obesity: Epidemiological evidence. *Nutrition, metabolism and cardiovascular diseases* **21**, S40-S45.
57. Ma Y, He FJ, MacGregor GA (2015) High salt intake: independent risk factor for obesity? *Hypertension* **66**, 843-849.
58. Appleton KM (2009) Increases in energy, protein and fat intake following the addition of sauce to an older person's meal. *Appetite* **52**, 161-165.
59. Ainsworth BE, Haskell WL, Whitt MC *et al.* (2000) Compendium of physical activities: an update of activity codes and MET intensities. *Medicine and science in sports and exercise* **32**, S498-S504.
60. Rezazadeh A, Rashidkhani B, Omidvar N (2010) Association of major dietary patterns with socioeconomic and lifestyle factors of adult women living in Tehran, Iran. *Nutrition* **26**, 337-341.
61. Singh GK, Siahpush M, Hiatt RA *et al.* (2011) Dramatic increases in obesity and overweight prevalence and body mass index among ethnic-immigrant and social class groups in the United States, 1976–2008. *Journal of community health* **36**, 94-110.
62. McLaren L (2007) Socioeconomic status and obesity. *Epidemiologic reviews* **29**, 29-48.
63. Berlin I (2009) Endocrine and metabolic effects of smoking cessation. *Current medical research and opinion* **25**, 527-534.
64. Dušková M, Hrušková H, Šimůnková K *et al.* (2014) The effects of smoking on steroid metabolism and fetal programming. *The Journal of steroid biochemistry and molecular biology* **139**, 138-143.

65. Ocké MCJPotNS (2013) Evaluation of methodologies for assessing the overall diet: dietary quality scores and dietary pattern analysis. **72**, 191-199.
66. Hu FBJCoil (2002) Dietary pattern analysis: a new direction in nutritional epidemiology. **13**, 3-9.
67. Andersson J, Nydahl M, Gustafsson K *et al.* (2003) Meals and snacks among elderly self-managing and disabled women. **41**, 149-160.
68. de Oliveira Santos R, dos Santos Vieira DA, Miranda AAM *et al.* (2018) The traditional lunch pattern is inversely correlated with body mass index in a population-based study in Brazil. **18**, 33.
69. Schwedhelm C, Iqbal K, Knüppel S *et al.* (2018) Contribution to the understanding of how principal component analysis-derived dietary patterns emerge from habitual data on food consumption. **107**, 227-235.
70. Esmailzadeh A, Azadbakht LJTJon (2008) Major dietary patterns in relation to general obesity and central adiposity among Iranian women. **138**, 358-363.
71. Craig LC, McNeill G, Macdiarmid JI *et al.* (2010) Dietary patterns of school-age children in Scotland: association with socio-economic indicators, physical activity and obesity. **103**, 319-334.
72. McNaughton SA, Ball K, Mishra GD *et al.* (2008) Dietary patterns of adolescents and risk of obesity and hypertension. **138**, 364-370.
73. Lera LM, Olivares SC, Leyton BD *et al.* (2006) Dietary patterns and its relation with overweight and obesity in Chilean girls of medium-high socioeconomic level. **56**, 165-170.
74. Herber-Gast G-CM, Mishra GDJTAoCN (2013) Fruit, Mediterranean-style, and high-fat and-sugar diets are associated with the risk of night sweats and hot flushes in midlife: results from a prospective cohort study. **97**, 1092-1099.
75. Newby P, Muller D, Hallfrisch J *et al.* (2004) Food patterns measured by factor analysis and anthropometric changes in adults. **80**, 504-513.
76. Nouri M, Tarighat-Esfanjani A, Ghazizahedi SJJoh (2017) The Relationships between Dietary Patterns and Energy and Nutrient Intakes and Body Mass Index in Iranian Adults. **8**, 85-101.
77. Isganaitis E, Lustig RHJA, thrombosis, biology v (2005) Fast food, central nervous system insulin resistance, and obesity. **25**, 2451-2462.




Figure 1. The distribution of calories was consumed across the time of day




Figure 2. Heat map showing the Pearson correlation matrix for lunch food intake in grams ($n = 790$) by food groups. The color corresponds to the strength of correlations (red: positive correlation; white: no correlation; blue: negative correlation).

Table 1. Food groups used in the factor analysis and factor loadings for each of the identified lunch patterns ^a

| Food groups | Food items | Dietary patterns | | |
|----------------------------|--|-----------------------|----------------|---------|
| | | Bread, grains and Fat | Egg and potato | Western |
| 1. Bread and grains | White bread (lavash, baguettes), noodles, pasta, rice, toasted bread, white flour, Dark bread (e.g., barbari, sangak, taftun) | 0.403 | -.0523 | – |
| 2. Dairy products | Low-fat milk, skim milk, low-fat yoghurt, cheese, Kashk, yoghurt drink, High-fat milk, high-fat yoghurt, cream cheese, cream, dairy fat, ice cream, others | – | – | -0.487 |
| 3. Poultry | Chicken | 0.342 | – | -0.526 |
| 4. Eggs | Eggs | – | 0.77 | – |
| 5. Fat | Hydrogenated fats, animal fats, butter, oils Olive oil, vegetable oils, olives | 0.591 | – | 0.234 |
| 6. Potato | Potatoes | – | 0.725 | – |

| | | | | |
|--|--|--------|---------|-------|
| 7. Processed meat and red meats | Sausage, hamburger, Beef and veal, lamb, minced meat, other | – | – | 0.52 |
| 8. Soft drinks | Soft drinks | – | – | 0.39 |
| 9. Vegetables | Cauliflower, carrot, tomato and its products, spinach, lettuce, cucumber, eggplant, onion, greens, green bean, green pea, squash, mushroom, pepper, corn, garlic, turnip, others | 0.275 | – | – |
| 10. Legumes and nuts | Peanuts, almonds, pistachios, hazelnuts, roasted seeds, walnuts, Lentils, split pea, beans, chickpea, fava bean, soy, others | -0.628 | -0.0321 | 0.268 |
| 11. Salt | Salt | 0.326 | – | 0.297 |
| 12. Sauces | Mayonnaise, Ketchup, tomato paste | -0.229 | – | 0.452 |

^a Factor loadings of <0.2 have been removed to simplify the table

Table 2. Characteristics of the study participants by tertiles (T) of lunch pattern scores

| Tertiles of dietary patterns | | | | | | | | | | | | |
|------------------------------|-----------------------|-----------|-----------|---------|----------------|-----------|-----------|---------|-----------|-----------|-----------|---------|
| Characteristics | Bread, grains and fat | | | P value | Egg and potato | | | P value | Western | | | P value |
| | T1 | T2 | T3 | | T1 | T2 | T3 | | T1 | T2 | T3 | |
| Participants | 263 | 264 | 263 | | 263 | 264 | 263 | | 263 | 264 | 263 | |
| Sex, n (%) | | | | 0.21 | | | | 0.10 | | | | 0.24 |
| Male | 43(28.1) | 51(33.3) | 59(38.6) | | 62(40.5) | 45(29.4) | 46(30.1) | | 58(37.9) | 43(28.1) | 52(34.0) | |
| Female | 220(34.5) | 213(33.4) | 204(32.0) | | 201(31.6) | 219(34.4) | 217(34.1) | | 205(32.2) | 221(34.7) | 211(33.1) | |
| |) |) |) | |) |) |) | |) |) |) | |
| Education, n (%) | | | | 0.41 | | | | 0.05 | | | | 0.08 |
| Educated | 91(32.3) | 102(37.2) | 81(29.6) | | 104(38.0) | 87(31.8) | 83(30.3) | | 107(39.1) | 89(32.5) | 78(28.5) | |
| | |) | | |) | | | |) | | | |
| Activity score | | | | 0.94 | | | | 0.26 | | | | 0.78 |
| Low | 134(32.4) | 138(33.4) | 141(34.1) | | 140(33.9) | 135(32.7) | 138(33.4) | | 132(32.0) | 141(34.1) | 140(33.9) | |
| |) |) |) | |) |) |) | |) |) |) | |

| | | | | | | | | | |
|-------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Moderate | 104(34.6) | 101(33.6) | 96(31.9) | 108(35.9) | 96(31.9) | 97(32.2) | 108(35.9) | 94(31.2) | 99(32.9) |
| High | 23(31.5) | 24(32.9) | 26(35.6) | 15(20.5) | 31(42.5) | 27(37.0) | 23(31.5) | 28(38.4) | 22(30.1) |
| Occupatio n, n (%) | | | | 0.88 | | | 0.04 | | 0.69 |
| Employee | 96(33.2) | 94(32.5) | 99(34.3) | 112(38.8) | 90(31.1) | 87(30.1) | 100(34.6) | 91(31.5) | 98(33.9) |
| Unemploy ed | 166(32.2) | 170(34) | 164(32.8) | 151(30.2) | 174(34.8) | 175(35) | 163(32.6) | 172(34.4) | 165(33) |
| Marriage, n (%) | | | | 0.34 | | | 0.88 | | 0.31 |
| Married | 206(32.3) | 220(34.5) | 212(33.2) | 211(33.1) | 212(33.2) | 215(33.7) | 207(32.4) | 221(34.6) | 210(32.9) |
| Other | 57(37.5) | 44(28.9) | 51(33.6) | 52(34.2) | 52(34.2) | 48(31.6) | 56(36.8) | 43(28.3) | 53(34.9) |
| Life-style, n (%) | | | | 0.59 | | | 0.14 | | 0.19 |
| Living alone | 16(38.1) | 14(33.3) | 12(28.6) | 18(42.9) | 17(40.5) | 7(16.7) | 20(47.6) | 9(21.4) | 13(31.0) |
| Smoking, n (%) | | | | 0.94 | | | 0.44 | | 0.94 |
| Not smoking | 250(33.3) | 251(33.4) | 250(33.3) | 249(33.2) | 250(33.3) | 252(33.6) | 252(33.6) | 250(33.3) | 249(33.2) |

| | | | | | | | | | | | | |
|--------------------------------------|-------------|--------------|-------------|------|--------------|------------|--------------|------|-------------|-------------|-------------|------|
| Quit smoking | 5(41.7) | 3(25.0) | 4(33.3) | | 5(41.7) | 6(50.0) | 1(8.3) | | 3(25.0) | 5(41.7) | 4(33.3) | |
| Smoker | 8(29.6) | 10(37.0) | 9(33.3) | | 9(33.3) | 8(29.6) | 10(37.0) | | 8(29.6) | 9(33.3) | 10(37.0) | |
| *Medical condition, n (%) | | | | 0.86 | | | | 0.42 | | | | 0.33 |
| Underlying disease | 109(32.1) | 116(34.1) | 115(33.8) | | 111(32.3) | 107(31.5) | 122(35.9) | | 107(31.5) | 110(32.4) | 123(36.2) | |
| Apparently Healthy | 151(33.9) | 148(33.2) | 147(33) | | 150(33.6) | 155(34.8) | 141(31.6) | | 154(34.5) | 153(34.3) | 139(31.2) | |
| Obesity n (%) | 73(36.3) | 59(29.4) | 69(34.3) | 0.39 | 65(32.3) | 62(30.8) | 74(36.8) | 0.42 | 60(29.9) | 66(32.8) | 75(37.3) | 0.35 |
| Mean ±SD | | | | | | | | | | | | |
| Age(years) | 42.23±1.21 | 42.08±1.07 | 42.85±1.09 | 0.51 | 41.39±1.07 | 42.47±1.33 | 43.30±1.09 | 0.04 | 42.47±1.09 | 42.47±1.09 | 42.22±1.16 | 0.95 |
| Weight (kg) | 71.13±3.84 | 72.08±2.88 | 73.04±4.75 | 0.11 | 74.06±4.51 | 70.89±3.51 | 71.31±3.32 | 0.02 | 71.80±2.48 | 71.48±2.69 | 72.98±6.09 | 0.32 |
| Systolic blood pressure(mmHg) | 115.92±18.1 | 116.06±22.09 | 118.13±20.6 | 0.21 | 114.65±19.96 | 117±20.07 | 118.45±20.89 | 0.03 | 116.14±21.6 | 116.09±20.2 | 117.88±19.1 | 0.32 |
| Diastolic | 78.27±1 | 77.31±1 | 79.75±1 | 0.20 | 77.93±1 | 78.55±1 | 78.85±1 | 0.23 | 78.61±1 | 77.07±1 | 79.66±1 | 0.37 |

| | | | | | | | | | | | | |
|-------------------------------------|------------------|-----------------|-----------------|------------|-----------------|-----------------|-----------------|------------|-----------------|-----------------|-----------------|------------|
| blood pressure(mmHg) | 1.74 | 4.02 | 4.30 | | 1.44 | 3.05 | 5.51 | | 3.5 | 4.06 | 2.55 | |
| Dietary and nutrient intakes | | | | | | | | | | | | |
| Fruit | 1.81±7.7 6 | 1.52±3.8 3 | 2.14±8.2 0 | 0.58 | 1.85±7.8 1 | 1.80±4.5 0 | 1.81±7.8 0 | 0.94 | 1.60±7.5 7 | 1.79±4.3 2 | 2.08±8.1 4 | 0.42 |
| Vegetable | 64.86±4 3.97 | 81.44±4 4.55 | 96.59±5 6.48 | <0.0 01 | 69.96±4 2.61 | 86.34±5 1.01 | 86.57±5 4.88 | <0.0 01 | 78.18±4 7.06 | 79.77±5 2.76 | 84.95±5 0.88 | 0.12 |
| Dairy | 52.90±5 5.78 | 47.21±5 8.72 | 41.20±5 7.13 | 0.01 | 48.54±6 1.55 | 48.79±5 7.51 | 44.87±5 2.79 | 0.46 | 81.10±6 6.98 | 42.16±4 5.21 | 18.05±3 6.47 | <0.0 01 |
| Grains | 128.07± 47.02 | 154.4±4 0.5 | 168.4±4 3.5 | <0.0 01 | 183.9±3 7.5 | 140.1±3 6.2 | 126.8±4 5.5 | <0.0 01 | 142.02± 47.4 | 147.3±4 3.03 | 161.5±4 7.7 | <0.0 01 |
| Meats | 13.85±1 6.58 | 14.77±1 9.58 | 18.34±2 0.13 | 0.00 6 | 17.07±2 0.50 | 15.53±1 7.86 | 14.37±1 8.23 | 0.10 | 7.93±11. 93 | 12.09±1 4 | 26.96±2 3.16 | <0.0 01 |
| Carbohydrate | 66.69±2 0.46 | 66.95±1 9.32 | 68.04±2 0.19 | 0.43 | 66.35±2 0.64 | 69.06±1 9.94 | 66.27±1 9.28 | 0.96 | 66.87±1 8.32 | 67.44±2 0.35 | 67.37±2 1.22 | 0.77 |
| Protein | 22.95±8. 32 | 23.22±7. 93 | 23.55±8. 50 | 0.40 | 23.11±7. 52 | 23.18±8. 03 | 23.44±9. 14 | 0.64 | 23.51±8. 14 | 23.14±8. 82 | 23.07±7. 76 | 0.53 |
| Fat | 20.58±8. | 20.90±9 | 21.75±8. | 0.12 | 21.10±8. | 20.80±8. | 21.32±9. | 0.77 | 20.28±7. | 21.18±9. | 21.77±8. | 0.05 |

| | 84 | 5 | 82 | 44 | 12 | 57 | 74 | 88 | | | |
|--------------------|---------|---------|---------|------|---------|---------|---------|---------|---------|---------|------|
| Total | | | | | | | | | | | |
| energy | 558.71± | 556.19± | 576.34± | | 582.24± | 529.62± | 579.49± | 566.07± | 540.15± | 585.08± | |
| intake(kcal | 231.8 | 158.5 | 184.4 | 0.27 | 215.3 | 151.9 | 176.5 | 228.5 | 139.1 | 172.6 | 0.23 |
| /day) | | | | | | | | | | | |

Data are presented as mean ± standard deviation or n (%). P-values obtained using Chi-square test.

* Diabetes, Hypertension, Dyslipidemia, Cardiovascular disease, Cancer and Respiratory disease

Table 3. Multivariable-adjusted means for anthropometric measures and indexes across tertiles (T) of lunch pattern scores

| Characteristics | Bread, grains and fat | | | P value | Egg and potato | | | P value | Western | | | P value |
|---|-----------------------|-----------------|-----------------|----------|-----------------|-----------------|-----------------|------------------------|-----------------|-----------------|-----------------|----------|
| | T1 (n=263) | T2 (n=264) | T3 (n=263) | | T1 (n=263) | T2 (n=264) | T3 (n=263) | | T1 (n=263) | T2 (n=264) | T3 (n=263) | |
| | Mean±SD | | | | | | | | | | | |
| Weight (kg) | 71.13±1 3.84 | 72.08±1 2.88 | 73.04±1 4.75 | 0.1 1 | 74.06±1 4.51 | 70.89±1 3.51 | 71.31±1 3.32 | 0.0 2 | 71.80±1 2.48 | 71.48±1 2.69 | 72.98±1 6.09 | 0.3 2 |
| Model 1 | 71.15±0. 85 | 72.12±0. 84 | 72.99±0. 85 | 0.3 1 | 74.18±0. 84 | 70.88±0. 84 | 71.20±0. 84 | 0.0 1 | 71.79±0. 85 | 71.47±0. 85 | 73±0.85 | 0.4 0 |
| Model 2 | 71.31±0. 84 | 71.90±0. 84 | 73.16±0. 84 | 0.2 9 | 74.01±0. 84 | 71.05±0. 84 | 71.31±0. 84 | 0.0 2 | 73.11±0. 84 | 71.56±0. 84 | 71.71±0. 84 | 0.3 6 |
| Model 3 | 72.21±0. 90 | 71.76±0. 85 | 72.47±0. 89 | 0.8 4 | 72.99±0. 96 | 71.25±0. 85 | 72.19±0. 91 | 0.4 0 | 72.09±0. 91 | 71.62±0. 84 | 72.73±0. 90 | 0.6 6 |
| Body mass index (kg/m²) | 27.30±5. 24 | 27.04±4. 19 | 27.66±7. 20 | 0.4 5 | 27.46±4. 5 | 27.04±5. 13 | 27.46±7. 08 | 0.6 7 | 27.02±4. 10 | 27.11±4. 53 | 28.77±7. 69 | 0.1 7 |
| Model 1 | 27.31±0. 34 | 27.07±0. 34 | 27.61±0. 34 | 0.5 3 | 27.56±0. 34 | 27.07±0. 34 | 27.37±0. 34 | 0.5 9 | 27.01±0. 34 | 27.10±0. 34 | 27.88±0. 34 | 0.1 4 |
| Model 2 | 27.38±0. 34 | 27.03±0. 34 | 27.62±0. 34 | 0.4 8 | 27.57±0. 34 | 27.10±0. 34 | 27.36±0. 34 | 0.6 2 | 27.05±0. 34 | 27.11±0. 34 | 27.88±0. 34 | 0.1 6 |
| Model 3 | 27.37±0. 36 | 27.04±0. 34 | 27.64±0. 36 | 0.4 7 | 27.72±0. 39 | 27.01±0. 35 | 27.32±0. 35 | 0.4 3 | 27.06±0. 37 | 27.08±0. 34 | 27.91±0. 36 | 0.1 8 |
| Waist-circumference (cm) | 88.42±1 1.50 | 88.91±1 0.95 | 90.50±1 2.47 | 0.1 0 | 89.30±1 1.87 | 88.95±1 1.42 | 89.58±1 1.76 | 0.8 2 | 88.18±9. 60 | 89.07±1 1.02 | 90.58±1 3.9 | 0.0 5 |
| Model 1 | 88.16±0. 69 | 89.5±0.6 9 | 90.62±0. 69 | 0.0 4 | 89.57±0. 69 | 88.93±0. 69 | 89.33±0. 69 | 0.8 0 | 88.16±0. 69 | 89.05±0. 69 | 90.62±0. 69 | 0.0 4 |

| | | | | | | | | | | | | |
|---------------------------|----------------|----------------|----------------|------------------------|----------------|----------------|----------------|----------|----------------|----------------|----------------|----------|
| Model 2 | 88.53±0. 69 | 88.93±0. 69 | 90.37±0. 69 | 0.1 4 | 89.71±0. 69 | 88.85±0. 69 | 89.28±0. 69 | 0.6 8 | 88.25±0. 69 | 89.06±0. 69 | 90.53±0. 69 | 0.0 6 |
| Model 3 | 89.17±0. 74 | 88.86±0. 69 | 89.84±0. 73 | 0.6 1 | 89.03±0. 79 | 88.99±0. 70 | 89.85±0. 75 | 0.6 6 | 88.47±0. 75 | 89.11±0. 69 | 90.29±0. 74 | 0.2 5 |
| Waist to hip ratio | 0.85±0.0 9 | 0.87±0.1 7 | 0.86±0.0 8 | 0.0 4 | 0.86±0.0 9 | 0.87±0.1 7 | 0.86±0.0 8 | 0.6 7 | 0.86±0.0 9 | 0.85±0.0 8 | 0.87±0.1 7 | 0.1 3 |
| Model 1 | 0.85±0.0 08 | 0.87±0.0 08 | 0.86±0.0 08 | 0.0 4 | 0.86±0.0 08 | 0.87±0.0 08 | 0.86±0.0 08 | 0.6 4 | 0.86±0.0 08 | 0.85±0.0 08 | 0.87±0.0 08 | 0.1 3 |
| Model 2 | 0.85±0.0 08 | 0.87±0.0 08 | 0.86±0.0 08 | 0.0 3 | 0.86±0.0 08 | 0.87±0.0 08 | 0.86±0.0 08 | 0.6 8 | 0.86±0.0 08 | 0.85±0.0 08 | 0.87±0.0 08 | 0.1 5 |
| Model 3 | 0.85±0.0 08 | 0.87±0.0 08 | 0.86±0.0 08 | 0.0 6 | 0.85±0.0 08 | 0.87±0.0 08 | 0.86±0.0 08 | 0.5 7 | 0.86±0.0 08 | 0.85±0.0 08 | 0.87±0.0 08 | 0.1 5 |

Model 1: adjusted for age (continuous) Model 2: additionally adjusted for marital status, education, physical activity, smoking Model 3: further adjustment for dietary intake of fruits, vegetables, dairy, grains, energy intake

P-values obtained using ANCOVA test.

Table 4. Odds ratio (OR)s and 95% confidence intervals(CI) for general obesity (BMI \geq 30) across tertiles (T) of dietary patterns score

| | T1 (n=263) | T2 (n=264) | | P value | T3 (n=263) | | P value | P trend |
|------------------------------|----------------------|----------------------|-------------|----------------|----------------------|-------------|----------------|----------------|
| Dietary patterns | | OR | 95% CI | | OR | 95% CI | | |
| Bread, grains and fat | | | | | | | | |
| Model 1 | 1.00 (Ref) | 0.76 | (0.51,1.13) | 0.18 | 0.92 | (0.63,1.36) | 0.69 | 0.69 |
| Model 2 | 1.00 (Ref) | 0.74 | (0.49,1.12) | 0.16 | 0.88 | (0.59,1.30) | 0.52 | 0.40 |
| Model 3 | 1.00 (Ref) | 0.74 | (0.49,1.12) | 0.15 | 0.89 | (0.60,1.33) | 0.59 | 0.38 |
| Egg and potato | | | | | | | | |
| Model 1 | 1.00 (Ref) | 0.93 | (0.62,1.39) | 0.74 | 1.19 | (0.81,1.76) | 0.35 | 0.35 |
| Model 2 | 1.00 (Ref) | 0.85 | (0.56,1.29) | 0.46 | 1.06 | (0.71,1.58) | 0.76 | 0.58 |
| Model 3 | 1.00 (Ref) | 0.82 | (0.54,1.24) | 0.36 | 1.07 | (0.71,1.60) | 0.36 | 0.46 |
| Western | | | | | | | | |
| Model 1 | 1.00 (Ref) | 1.11 | (0.74,1.66) | 0.59 | 1.33 | (0.89,1.97) | 0.15 | 0.15 |
| Model 2 | 1.00 (Ref) | 1.06 | (0.70,1.60) | 0.77 | 1.28 | (0.86,1.92) | 0.21 | 0.42 |
| Model 3 | 1.00 (Ref) | 1.04 | (0.69,1.57) | 0.84 | 1.32 | (0.88,1.99) | 0.17 | 0.36 |

Model 1: unadjusted, Model2: Age, sex, education (categorical), Marriage, life-style, smoking; Model 3: Model 2 + physical activity, total energy intake.

Table 5. Odds ratio (OR)s and 95% confidence intervals(CI) for central obesity across tertiles (T) of dietary patterns score

| Dietary patterns | T1 | T2 | | P | T3 | | P | P |
|--|---------------|------|-------------|-------|------|-------------|-------|-------|
| | | OR | 95% CI | value | OR | 95% CI | value | trend |
| Waist circumference (cm) above 102 cm for men and above 88 cm for women | | | | | | | | |
| Bread, grains and fat | | | | | | | | |
| Model 1 | 1.00 (Ref) | 0.86 | (0.61,1.21) | 0.40 | 1.07 | (0.76,1.51) | 0.66 | 0.66 |
| Model 2 | 1.00 (Ref) | 0.84 | (0.58,1.22) | 0.37 | 1.12 | (0.77,1.64) | 0.53 | 0.40 |
| Model 3 | 1.00 (Ref) | 0.85 | (0.58,1.24) | 0.40 | 1.15 | (0.79,1.68) | 0.45 | 0.38 |
| Egg and potato | | | | | | | | |
| Model 1 | 1.00 (Ref) | 1.10 | (0.78,1.55) | 0.56 | 0.97 | (0.68,1.36) | 0.86 | 0.86 |
| Model 2 | 1.00 (Ref) | 0.92 | (0.63,1.34) | 0.67 | 0.74 | (0.51,1.09) | 0.13 | 0.26 |
| Model 3 | 1.00 (Ref) | 0.88 | (0.60,1.29) | 0.53 | 0.74 | (0.51,1.09) | 0.13 | 0.27 |
| Western | | | | | | | | |
| Model 1 | 1.00 | 1.34 | (0.95,1.90) | 0.08 | 1.31 | (0.93,1.86) | 0.11 | 0.11 |

| | | | | | | | | |
|--|-------|------|-------------|------|-------|-------------|------|------|
| | (Ref) | | | | | | | |
| Model 2 | 1.00 | 1.24 | (0.85,1.80) | 0.26 | 1.26 | (0.86,1.84) | 0.22 | 0.35 |
| | (Ref) | | | | | | | |
| Model 3 | 1.00 | 1.22 | (0.84,1.78) | 0.28 | 1.26 | (0.86,1.84) | 0.22 | 0.37 |
| | (Ref) | | | | | | | |
| Waist to hip ratio (<i>above 0.90 for males and above 0.85 for females</i>) | | | | | | | | |
| Bread, grains and fat | | | | | | | | |
| Model 1 | 1.00 | 1.44 | (1.02,2.03) | 0.03 | 1.38 | (0.98,1.95) | 0.06 | 0.03 |
| | (Ref) | | | | | | | |
| Model 2 | 1.00 | 1.34 | (0.94,1.92) | 0.10 | 1.42 | (0.99,2.03) | 0.05 | 0.11 |
| | (Ref) | | | | | | | |
| Model 3 | 1.00 | 1.35 | (0.94,1.93) | 0.09 | 1.44 | (1.01,2.07) | 0.04 | 0.09 |
| | (Ref) | | | | | | | |
| Egg and potato | | | | | | | | |
| Model 1 | 1.00 | 1.34 | (0.95,1.90) | 0.08 | 1.13 | (0.80,1.59) | 0.48 | 0.48 |
| | (Ref) | | | | | | | |
| Model 2 | 1.00 | 1.29 | (0.90,1.84) | 0.15 | 1.004 | (0.70,1.43) | 0.98 | 0.27 |
| | (Ref) | | | | | | | |
| Model 3 | 1.00 | 1.30 | (0.90,1.87) | 0.15 | 0.99 | (0.69,1.42) | 0.97 | 0.25 |
| | (Ref) | | | | | | | |
| Western | | | | | | | | |

| | | | | | | | | |
|----------------|---------------|------|-------------|------|------|-------------|------|------|
| Model 1 | 1.00 (Ref) | 0.83 | (0.59,1.18) | 0.31 | 0.95 | (0.67,1.34) | 0.79 | 0.79 |
| Model 2 | 1.00 (Ref) | 0.78 | (0.54,1.12) | 0.18 | 0.90 | (0.63,1.29) | 0.58 | 0.43 |
| Model 3 | 1.00 (Ref) | 0.79 | (0.55,1.14) | 0.21 | 0.89 | (0.62,1.28) | 0.54 | 0.48 |

Model 1: unadjusted, Model2: Age, sex, education (categorical), Marriage, life-style, smoking; Model 3: Model 2 + physical activity and total energy intake.