



Apr 21st, 9:00 AM - Apr 22nd, 4:45 PM

Measuring the effectiveness of Seattle's seawall enhancements on juvenile salmon-an acoustic perspective

Kerry Accola
University of Washington, kaccola@uw.edu

Follow this and additional works at: <https://cedar.wwu.edu/ssec>



Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Natural Resources and Conservation Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

Accola, Kerry, "Measuring the effectiveness of Seattle's seawall enhancements on juvenile salmon-an acoustic perspective" (2020). *Salish Sea Ecosystem Conference*. 56.
<https://cedar.wwu.edu/ssec/2020ssec/allsessions/56>


This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact westerncedar@wwu.edu.

Do Seattle's seawall enhancements influence juvenile salmon distribution along the urban waterfront? An acoustic perspective

Kerry Accola @salmon.underfoot (kaccola@uw.edu)
Washington Sea Grant Research Fellow


INTRODUCTION

- Seattle's waterfront is a key migration route for Puget Sound Pacific salmon. This includes 3 species of obligatory nearshore migrators, one of which is Puget Sound Chinook, listed as 'Threatened' (ESA)
- Seattle's new seawall (Phase I completed in 2018) is the largest eco-engineering project of its kind
- Habitat enhancements include: 1) light-penetrating glass blocks in overhanging sidewalk, 2) habitat bench to restore shallow water depths, 3) textured seawall and shelves
- Test of new application for acoustic camera
 - Mobile surveying of small fish
- First in-depth study of nighttime fish distributions along Seattle's waterfront







METHODS

- 1,070 25m acoustic surveys at set transects April-August 2019
- High frequency sonar camera quantifies salmon and other fish population densities at modified and original seawall sites, and reference beaches
- Analyses include generalized linear models and multivariate statistics





DIDSON (dual-frequency identification sonar) mounted to kayak hull

Seattle's seawall enhancements influence juvenile salmon distributions



- Salmon densities are higher at
 - New pier sidewalk corridor compared to old piers with no corridor
 - Deep open transects (10m from seawall) compared to shallow transects
 - Night compared to day
- Salmon are more evenly distributed between piers and open seawall at new sites



Above: Chum schooling in new corridor
Below: Herring simultaneously in GoPro and acoustic camera footage

FURTHER IMPLICATIONS

Seawall modifications may be important to salmon at night, with higher nighttime densities in new corridor compared to day

Salmon may navigate more around pier ends with no sidewalk corridor

