

Marshall University

Marshall Digital Scholar

Physical Therapy Faculty Research

Physical Therapy

2-2019

Variability of spatiotemporal gait parameters in children with and without Down syndrome during treadmill walking

Matthew Beerse

Gena Henderson

Huaqing Liang

Toyin Ajisafe

Jianhua Wu

Follow this and additional works at: https://mds.marshall.edu/physical_therapy_faculty

 Part of the [Physical Therapy Commons](#)

Variability of spatiotemporal gait parameters in children with and without Down syndrome during treadmill walking

Matthew Beerse ^{a,b†}, Gena Henderson ^{a†}, Huaqing Liang ^a, Toyin Ajisafe ^c, Jianhua Wu ^{a,d*}

^a Department of Kinesiology and Health, Georgia State University, Atlanta, GA, USA

^b Department of Health and Sport Science, University of Dayton, Dayton, OH, USA

^c Department of Kinesiology, Texas A&M University Corpus Christi, Corpus Christi, TX, USA

^d Center for Movement & Rehabilitation Research, Georgia State University, Atlanta, GA, USA

* Corresponding author:

Jianhua Wu, Ph.D.

Department of Kinesiology and Health, Center for Movement & Rehabilitation Research,
Georgia State University. 125 Decatur Street, Atlanta, GA 30302, USA.

Telephone: 1-404-413-8476; Fax: 1-404-413-8053; Email: jwu11@gsu.edu

† Contributed equally to this study.

Word count for Abstract: 284

Word count for Main Text: 2,995

Number of Tables: 2

Number of Figures: 2

This manuscript is submitted as an Original Article.

Acknowledgements

The authors are thankful to all the participants and their families for their participation in this study. This study was supported in part by the Jerome Lejeune Foundation.

Abstract

Background: Increasing walking speed and including bilateral external ankle load have been shown to improve aspects of the gait pattern of children with Down syndrome (DS). However, it is unknown if speed and ankle load improves the cycle-to-cycle variability in a similar way.

Research question: How do changes of walking speed and external ankle load impact spatiotemporal variability during treadmill walking in children with and without DS?

Methods: Thirteen children with DS (aged 7-10 years) and thirteen age- and sex-matched typically developing (TD) children participated in this study. Subjects completed two bouts of 60-second treadmill walking at two different speeds (*slow* and *fast*) and two load conditions (no load and ankle load equaling to 2% bodyweight at each side). Kinematic data was captured using a Vicon motion capture system. Mean and coefficient of variance of spatiotemporal gait variables were calculated and compared between children with and without DS.

Results and significance: Across all conditions, the DS group took shorter and wider steps than the TD group, but walked with a similar swing percentage, double support percentage, and foot rotation angle. Further, the DS group demonstrated greater variability of all spatiotemporal parameters, except for step width and foot rotation angle. Our results indicated that children with DS can modulate their spatiotemporal gait pattern accordingly like their TD peers when walking faster on a treadmill and/or with an external ankle load. Smaller step width variability in the DS group suggests that mediolateral stability may be prioritized during treadmill walking to safely navigate the treadmill and complete walking tasks. Similar temporal parameters but distinct spatial parameters in the DS group suggest that they may have developed similar rhythmic control but are confined by their spatial movement limitations.

1
2
3
4 **Introduction**
5

6
7 Individuals with Down syndrome (DS) typically present with hypotonia, joint
8
9 hypermobility, reduced strength, and cognitive impairments [1-3]. These conditions are
10
11 considered to underlie the often-termed “clumsy” movement patterns of individuals with DS
12
13 characterized by poor balance control, reduced joint and muscle coordination, abnormal gait
14
15 kinematics, delays in acquiring complex movement patterns, and abnormal variability of
16
17 movement [4-8]. These impairments produce stereotypical alterations in both the gait patterns of
18
19 individuals with DS and their ability to adapt to changing gait conditions [9-12].
20
21
22

23
24 When children with DS are challenged to walk faster on a treadmill they adapt similarly
25
26 to typically developing (TD) children by increasing step length and decreasing step width [13].
27
28 The inclusion of external ankle load increased stride length and decreased cadence in children
29
30 with DS similarly to their TD peers [10]. The combination of fast walking and ankle load
31
32 demonstrated a more advanced gait pattern with an improved power spectrum and general
33
34 muscular activation in children with DS [9, 10]. Although treadmill training is a common
35
36 intervention for infants [14] and children with DS [9, 10], it is unknown to what degree walking
37
38 speed and external ankle load affect cycle-to-cycle gait variability in children with DS.
39
40
41
42

43
44 All movement contains some degree of variability, which enables adaptation to new tasks
45
46 and/or environments [15, 16]. However, too little or too much variability can be associated with
47
48 movement dysfunction. When learning to walk, toddlers with DS display greater variability in
49
50 step length and width and reduced these variabilities at a slower rate than their TD peers [17].
51
52 During treadmill walking, adults with DS demonstrate greater variability of center-of-mass, step
53
54 width, and step length than healthy controls and these differences become greater at faster speeds
55
56 [18]. Moreover, adults with DS increase stride velocity variability and decrease step width
57
58
59
60
61
62
63
64
65

1
2
3
4 variability more than healthy controls when walking overground in more challenging conditions,
5
6 and there is correlation between amount of variability and self-reported frequency of falls in
7
8 adults with DS suggesting its clinical relevance and significance [19].
9

10
11 The purpose of this study was to investigate the acute effects of walking speed and
12
13 external ankle load on spatiotemporal gait parameters and their variability patterns between
14
15 children with and without DS. We hypothesized that children with DS will demonstrate a similar
16
17 spatiotemporal adaptation pattern to their TD peers such that at faster speeds or wearing external
18
19 ankle load, both groups will increase mean step length and swing percentage and decrease mean
20
21 step width and double support percentage. Between the two groups, children with DS will show
22
23 smaller step length and swing percentage and greater step width and double support percentage.
24
25 In terms of gait variability, we hypothesized that a similar variability pattern will emerge for both
26
27 groups such that variability of step length, swing percentage, and double support percentage will
28
29 increase with speed and ankle load, and variability of step width will decrease. Between the two
30
31 groups, children with DS will display greater variability in all the spatiotemporal variables.
32
33
34
35
36
37
38
39

40 **Methods**

41
42
43 Thirteen children with DS aged 7-10 years were recruited through local parent support
44
45 groups and the Down Syndrome Association of Atlanta. Thirteen age- and sex-matched TD
46
47 children also participated in the study. All participants were able to walk overground for 10-
48
49 meters independently. Exclusion criteria included a history of medical conditions or the inability
50
51 to walk independently. Anthropometric data were collected for each subject including body
52
53 mass, height, and leg length. The children with DS were shorter in height but had similar body
54
55 mass (Table 1). This study was approved by the institutional review board at the hosting
56
57
58
59
60
61
62
63
64
65

1
2
3
4 university. Written informed consent was obtained from the parents or guardians of all the
5
6 subjects before data collection.
7
8

9 Subjects completed treadmill walking at two different speeds and two load conditions for
10 a total of four conditions. The treadmill speed conditions were set by the subject's preferred
11 walking speed, calculated as the average speed of three 10-meter overground walking trials.
12
13 Previous studies have shown that preferred treadmill walking speeds are lower than overground
14 in children with DS, thus our treadmill speed conditions included a *slow* speed (75% of
15 preferred) and a *fast* speed (100% of preferred) [4, 20]. The two load conditions included *no load*
16 and *ankle load* equal to 2% of body weight on each side. This amount of ankle load increased the
17 moment of inertia of the leg by 39%, which was tolerable in children with DS, but resulted in a
18 modulated gait pattern [10]. We attached adjustable weight cuffs above each ankle and strapped
19 them tightly to minimize any motion during walking. Sixty seconds of data was collected after
20 the treadmill reached the condition speed and the subject visually appeared to be walking at a
21 steady-state. Subjects completed two treadmill walking trials at each condition. Presentation of
22 the conditions was randomized across subjects. Between trials the subjects were given adequate
23 rest. A custom full-body 18 marker set was attached to each subject, including the heel, ankle,
24 toe, knee, greater trochanter, shoulder, elbow, wrist, and temple bilaterally [10]. Kinematic data
25 was captured using a 7-camera Vicon motion capture system (Denver, CO, USA) at a sampling
26 rate of 100 Hz. Marker trajectories were filtered at a cutoff frequency of 6Hz. Custom MATLAB
27 programs (Natick, MA, USA) were written for data analysis.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 Gait events were determined using the anterior/posterior (AP) velocity of each heel
54 marker [21]. A velocity change of the heel from positive to negative indicated a heel-strike event
55 and a change from negative to positive indicated a toe-off event. To confirm the accuracy of this
56
57
58
59
60
61
62
63
64
65

1
2
3
4 algorithm with our subjects, ten subjects were randomly chosen and manually labeled for heel-
5 strikes and toe-offs and compared to the algorithm-generated gait events. Mean difference
6 between the manually labeled and algorithm-generated heel-strike events was about one frame
7 and thus deemed acceptable. However, the algorithm-generated toe-off event occurred, on
8 average, three frames before those manually labeled. Therefore, three frames were added to the
9 algorithm-generated toe-off events for each trial.

10
11
12
13
14
15
16
17
18
19 Step length was calculated as the AP difference between the ipsilateral heel marker at
20 heel-strike and the contralateral heel marker at minimum AP position during stance phase. Step
21 width was calculated as the medial/lateral (ML) difference between the heel markers at heel-
22 strike. Both step length and width were normalized by leg length. Temporal variables included
23 swing percentage of gait cycle, and double support percentage of gait cycle. Double support
24 percentage was calculated for the first double support phase of a gait cycle. Foot rotation angle
25 [22] was calculated as the angle between the line connecting the toe and heel markers of the
26 current foot step (i.e., line of foot placement) and the line connecting the heel of the current and
27 previous step from the same side (i.e., line of foot progression) [23]. A positive or negative FRA
28 indicated an out-toeing or in-toeing position of the foot with respect to the line of foot
29 progression, respectively. Spatiotemporal variables were calculated for each side and combined
30 after no asymmetry was observed between the two sides for either group. To investigate the
31 variability of spatiotemporal variables, coefficient of variation (CV) was calculated as the ratio
32 of standard deviation to mean value for each variable within each trial. For FRA, we used
33 absolute values when calculating the CV.

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

A three-way (2 group x 2 speed x 2 load) mixed ANOVA with repeated measures on speed and load was conducted on the spatiotemporal variables and their CV. Normality was

1
2
3
4 assessed using the Shapiro-Wilk test. Variables found to violate normality were log transformed
5
6 and reassessed using the Shapiro-Wilk test. This process sufficiently normalized our data. Post-
7
8 hoc pairwise comparisons were completed when necessary. The statistical analysis was
9
10 completed using the SAS 9.4 statistical software (Cary, NC). A significance level was set at
11
12 $\alpha=0.05$.
13
14
15
16
17
18

19 **Results**

20
21 Some children with DS could not complete all four walking conditions; 11 completed the
22
23 *fast* speed condition without ankle load, 10 completed the *slow* speed condition with ankle load,
24
25 and 7 completed the *fast* speed condition with ankle load. No significant differences were found
26
27 between those children with DS who were able to complete all conditions and those who were
28
29 not, so both subgroups were combined for analysis. All 13 TD children completed all walking
30
31 conditions.
32
33
34
35
36
37

38 *Mean spatiotemporal variables*

39
40 Both the TD and DS groups increased normalized step length and maintained normalized
41
42 step width when increasing walking speed (Table 2). With additional ankle load, both groups
43
44 maintained normalized step length; however, the TD group increased normalized step width
45
46 while the DS group did not change step width. Statistical analysis demonstrated that there was a
47
48 group by load interaction for normalized step length ($p=0.016$) and step width ($p<0.001$; see
49
50 Table 2 for the details). Post-hoc analysis indicated that the DS group had a smaller step length
51
52 and a greater step width than the TD group at each load condition, and the difference between the
53
54 two groups became greater at the ankle load condition. There was also a speed effect for
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 normalized step length across the two groups ($p < 0.001$). In addition, FRA was similar between
5
6 the two groups (between 0 and 5 degrees) and unaffected by speed and load conditions.
7
8

9 While walking faster, both groups displayed similar trends and values by increasing
10 swing percentage but decreasing double support percentage (Table 2). With additional ankle
11 load, both groups showed similar trends and values by increasing swing percentage but
12 decreasing double support percentage. Statistical results demonstrated that there was a speed
13
14 effect ($p < 0.001$) and a load effect ($p = 0.036$) for swing percentage, and a speed effect ($p < 0.001$)
15
16 and a load effect ($p = 0.006$) for double support percentage. There was neither a group effect nor a
17
18 group by speed/load interaction for both temporal variables.
19
20
21
22
23
24

25 26 27 28 *Variability of spatiotemporal variables* 29

30 Both groups decreased normalized step length variability but increased normalized step
31 width variability from *slow* to *fast* speed condition (Fig. 1a-d). With the additional ankle load,
32
33 both groups displayed no change in normalized step length variability but decreased normalized
34
35 step width variability. Regardless of speed and load conditions, the DS group had greater step
36
37 length variability and smaller step width variability than the TD group. For step length
38
39 variability, there was a group effect ($F(1,18) = 24.93$, $p < 0.001$) and speed effect ($F(1,17) = 8.44$,
40
41 $p = 0.010$). For step width variability, there was a group effect ($F(1,18) = 8.91$, $p = 0.008$), speed
42
43 effect ($F(1,17) = 6.82$, $p = 0.018$), and load effect ($F(1,18) = 36.22$, $p < 0.001$). Both groups showed
44
45 similar variability in FRA across the speed and load conditions with no significant statistical
46
47 results (Fig. 1e-f).
48
49
50
51
52
53

54 Both groups decreased variability in swing and double support percentage from the *slow*
55
56 to *fast* speed condition (Fig. 2a-d). The additional ankle load did not change variability in both
57
58 swing and double support percentages. Regardless of the speed and load conditions, the DS
59
60
61
62
63
64
65

1
2
3
4 group showed greater variability in both swing and double support percentages than the TD
5
6
7 group. Statistical analysis showed that there was a group effect ($F(1,18)=26.85, p<0.001$) and a
8
9 speed effect ($F(1,17)=41.95, p<0.001$) on swing percentage variability, and a group effect
10
11 ($F(1,18)=21.84, p<0.001$) and a speed effect ($F(1,17)=9.10, p=0.008$) on double support
12
13
14 percentage variability.
15
16
17

18 **Discussion**

21 Partially supporting out hypotheses, our results indicated that children with DS displayed
22
23 some similar trends in spatiotemporal pattern and its variability as their TD peers while adapting
24
25 to walking speed and ankle load but maintained group-level differences across all conditions.
26
27 This is consistent with previous findings of an overall lack of stability control in individuals with
28
29 DS [19]. In this study, modulating speed was found to induce similar acute adaptations in both
30
31 groups and the DS group demonstrated greater improvement of stability by reducing variability
32
33 in step length and swing percentage. Clinical interventions that involve modulating speed and
34
35 external ankle load during treadmill training have been shown to improve some aspects of gait
36
37 patterns of infants [14] and children with DS [9, 10]. Additionally, children with DS improve
38
39 their overground gait patterns after treadmill walking practice [13]. Clinicians can use these
40
41 findings to their advantage by modulating treadmill conditions to produce adaptations desired in
42
43 overground walking. However, in this study, modulating ankle load was found to induce some
44
45 unexpected compensations in both groups: strict step width control (lower variability) in the DS
46
47 group and wider steps in the TD group. While these were acute adaptations, when developing
48
49 clinical interventions, it is important to weigh positive effects with the potential motor
50
51 compensations. Further research on prolonged treadmill training with changes in speed and
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 external ankle load are required to quantify the potential long-term influence of motor
5
6 compensations reported in this study.
7
8

9 In contrast to previous work in adults [24], both groups of children responded to the *fast*
10 speed condition by decreasing the variability of step length, swing percentage, and double
11 support percentage. When walking on a rapidly moving treadmill belt, both groups took longer
12 steps, spent more time in swing phase and less time in double support. The consistent faster pace
13
14 might have necessitated this adaptation and reduced the variability of these parameters to
15
16 maintain a stable position in the middle of the treadmill belt. This suggests that a portion of gait
17
18 variability can be reduced or optimized when adapting to a faster speed in children with and
19
20 without DS. As excessive variability of stride velocity and limited variability of step width is
21
22 correlated with fall risk in adults with DS [18], interventions which normalize variability may
23
24 improve the overall stability and safety of individuals with DS. Our results suggest that treadmill
25
26 training at a fast speed, which decreases step length variability but increases step width
27
28 variability, may represent such an intervention.
29
30
31
32
33
34
35
36
37

38 Our results of unchanged mean step width and FRA are in opposition to previous studies
39
40 [13, 25-27]. However, Agiovlasitis et al. [24] noted a lack of change in step width at faster
41
42 speeds in young adults with DS during treadmill walking and proposed, as we speculated, that
43
44 they might maintain stability at faster speeds by modulating step length alone. Methodologically,
45
46 FRA is affected by both the degree of in-toeing or out-toeing of the foot relative to the line of
47
48 progression and the placement of the foot in the ML direction. Our definition is different from
49
50 previously reported angles, which were defined between the line of progression (i.e., the
51
52 regression line of all the steps) and the placement of the foot [25, 26]. Also, the treadmill might
53
54 substantially constrain the ML placement of the foot. Additionally, arch height has been shown
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 to influence FRA in children with DS [5] and we did not control for the arch height of our
5
6 subjects.
7

8
9 Interestingly, step width variability increased with the *fast* walking speed but decreased
10 with ankle load for both groups. Individuals with DS have been found to value frontal plane
11 stability, prioritizing equilibrium maintenance over forward progression during overground gait
12 [8, 19, 28]. The increased consistency of wider steps during the ankle load condition suggests
13
14 that the external ankle load might present an increased challenge to ML stability during gait,
15
16 requiring strict control. The prioritization of maintaining step width variability might illustrate a
17
18 posture-first strategy, which has been proposed to prevent falls during more complex walking
19
20 tasks [19, 29]. However, while the ankle load potentially perturbed ML stability, children with
21
22 DS increased swing percent and decreased double support percent, similar to the TD group. This
23
24 result furthers the proposal that external ankle load might aid the unloading of the leg during the
25
26 stance-to-swing transitions, and the increased moment of inertia most likely assisted the
27
28 pendulum motion of the swing leg [9]. Therefore, while external ankle load might acutely
29
30 improve temporal parameters associated with stability, there appeared a requirement to more
31
32 strictly maintain spatial base of support (BOS). The increased step width found in the TD group
33
34 was likely due to the added weight above the ankle, requiring more space during swing to safely
35
36 pass the legs by each other. It may warrant further investigation on whether other unexpected
37
38 compensations, such as circumduction, are utilized to accommodate the external load.
39
40

41
42 Previous studies have suggested that the more pronounced out-toeing demonstrated by
43
44 young adults with DS during fast overground walking is an attempt to increase BOS and improve
45
46 stability at the expense of efficiency [26]. Therefore, it is reasonable for modulation of FRA
47
48 variability to also emerge as a strategy to increase stability. As previously noted, we believe that
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 the ML constraints of the treadmill might have limited the capacity of both groups to alter FRA,
5
6 resulting in preferential modulation of step width to increase BOS and improve ML stability.

7
8
9 Further studies that examine the variability of FRA during fast walking overground are needed to
10
11 explore this possible explanation.
12

13
14 Although both groups demonstrated similar mean temporal parameters across both speed
15
16 conditions, the children with DS demonstrated increased variability of these parameters under all
17
18 conditions. This increased variability may be a result of their inherently lower muscle tone and
19
20 decreased ability to finely time movements [1]. However, the children with DS were able to
21
22 successfully maintain similar rhythmic timing as their TD peers across both speed conditions.
23
24 Black et al. [4] suggests that children with DS may recognize the inherent difficulty in
25
26 controlling the excessive laxity of their movements and choose to only constrain variability that
27
28 may negatively affect task performance. Our findings support this notion, as the increased
29
30 temporal variability exhibited by the children with DS did not affect their ability to demonstrate
31
32 similar temporal variables as their TD peers.
33
34
35
36
37

38 Some limitations of this study include our small sample size, exacerbated by the inability
39
40 of some children with DS to complete all walking conditions. Additionally, the short length of
41
42 our walking trials might have been insufficient for our subjects to accommodate to the task.
43
44

45 Although we only collected data when subjects visually appeared to have achieved steady-state
46
47 walking, more subtle compensations might have occurred. Further, subjects were not controlled
48
49 for their experience with treadmill walking. Although the tasks presented were novel to most
50
51 subjects, the actual degree of task novelty might have varied across the subjects. Finally, our
52
53 results can only be generalized to children with DS and should be investigated in individuals
54
55 with DS at different ages to determine if these adaptations are consistent across the lifespan.
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7 **References**
8
9

- 10 [1] Block ME. Motor Development in Children with Down Syndrome: A Review of the Literature.
11 Adapted Physical Activity Quarterly. 1991;8:179-209.
12
13 [2] Lott IT. Neurological phenotypes for Down syndrome across the life span. Prog Brain Res.
14 2012;197:101-21.
15
16 [3] Roizen NJ, Patterson D. Down's syndrome. Lancet. 2003;361:1281-9.
17
18 [4] Black DP, Smith BA, Wu J, Ulrich BD. Uncontrolled manifold analysis of segmental angle variability
19 during walking: preadolescents with and without Down syndrome. Exp Brain Res. 2007;183:511-21.
20
21 [5] Galli M, Cimolin V, Rigoldi C, Pau M, Costici P, Albertini G. The effects of low arched feet on foot
22 rotation during gait in children with Down syndrome. Journal of Intellectual Disability Research.
23 2014;58:758-64.
24
25 [6] Galli M, Rigoldi C, Brunner R, Virji-Babul N, Giorgio A. Joint stiffness and gait pattern evaluation in
26 children with Down syndrome. Gait Posture. 2008;28:502-6.
27
28 [7] Palisano RJ, Walter SD, Russell DJ, Rosenbaum PL, Gemus M, Galuppi BE, et al. Gross motor
29 function of children with down syndrome: creation of motor growth curves. Arch Phys Med Rehabil.
30 2001;82:494-500.
31
32 [8] Rigoldi C, Galli M, Mainardi L, Crivellini M, Albertini G. Postural control in children, teenagers and
33 adults with Down syndrome. Res Dev Disabil. 2011;32:170-5.
34
35 [9] Wu J, Ajisafe T. Kinetic patterns of treadmill walking in preadolescents with and without Down
36 syndrome. Gait & posture. 2014;39:241-6.
37
38 [10] Wu J, Ajisafe T, Beerse M. Children Display Adult-Like Kinetic Patterns in the Time Domain, But
39 Not in the Frequency Domain, While Walking With Ankle Load. J Appl Biomech. 2015;31:292-308.
40
41 [11] Rigoldi C, Galli M, Albertini G. Gait development during lifespan in subjects with Down syndrome.
42 Research in Developmental Disabilities. 2011;32:158-63.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 [12] Ganley KJ, Powers CM. Gait kinematics and kinetics of 7-year-old children: a comparison to adults
5 using age-specific anthropometric data. *Gait & Posture*. 2005;21:141-5.
6
7
8 [13] Smith BA, Kubo M, Black DP, Holt KG, Ulrich BD. Effect of Practice on a Novel Task—Walking
9 on a Treadmill: Preadolescents With and Without Down Syndrome. *Physical Therapy*. 2007;87:766-77.
10
11 [14] Wu J, Looper J, Ulrich BD, Ulrich DA, Angulo-Barroso RM. Exploring effects of different treadmill
12 interventions on walking onset and gait patterns in infants with Down syndrome. *Developmental*
13 *Medicine & Child Neurology*. 2007;49:839-945.
14
15 [15] Fethers L. Perspective on variability in the development of human action. *Phys Ther*. 2010;90:1860-
16 7.
17
18 [16] Vereijken B. The Complexity of Childhood Development: Variability in Perspective. *Physical*
19 *Therapy*. 2010;90:1850-9.
20
21 [17] Looper J, Wu J, Barroso RA, Ulrich D, Ulrich BD. Changes in Step Variability of New Walkers
22 With Typical Development and With Down Syndrome. *Journal of motor behavior*. 2006;38:367-72.
23
24 [18] Agiovlasitis S, McCubbin JA, Yun J, Pavol MJ, Widrick JJ. Economy and Preferred Speed of
25 Walking in Adults With and Without Down Syndrome. *Adapted Physical Activity Quarterly*.
26 2009;26:118-30.
27
28 [19] Smith BA, Ashton-Miller JA, Ulrich BD. Gait adaptations in response to perturbations in adults with
29 Down syndrome. *Gait & posture*. 2010;32:149-54.
30
31 [20] Ulrich BD, Haehl V, Buzzi UH, Kubo M, Holt KG. Modeling dynamic resource utilization in
32 populations with unique constraints: Preadolescents with and without Down syndrome. *Human*
33 *Movement Science*. 2004;23:133-56.
34
35 [21] Zeni JA, Richards JG, Higginson JS. Two simple methods for determining gait events during
36 treadmill and overground walking using kinematic data. *Gait & posture*. 2008;27:710-4.
37
38 [22] Franz JR, Kram R. The effects of grade and speed on leg muscle activations during walking. *Gait*
39 *Posture*. 2012;35:143-7.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 [23] Angulo-Barroso RM, Wu J, Ulrich DA. Long-term effect of different treadmill interventions on gait
5 development in new walkers with Down syndrome. *Gait Posture*. 2008;27:231-8.
6
7
8 [24] Agiovlasitis S, McCubbin JA, Yun J, Mpitsos G, Pavol MJ. Effects of Down syndrome on three-
9 dimensional motion during walking at different speeds. *Gait Posture*. 2009;30:345-50.
10
11 [25] Galli M, Cimolin V, Pau M, Costici P, Albertini G. Relationship between flat foot condition and gait
12 pattern alterations in children with Down syndrome. *Journal Of Intellectual Disability Research: JIDR*.
13 2014;58:269-76.
14
15 [26] Horvat M, Croce R, Zagrodnik J, Brooks B, Carter K. Spatial and temporal variability of movement
16 parameters in individuals with Down syndrome. *Percept Mot Skills*. 2012;114:774-82.
17
18 [27] Kim B-S, Bang DY, Kim B-O. Gait characteristics in down's syndrome. *Gait & Posture*. 1995;3:84.
19
20 [28] Rigoldi C, Galli M, Cimolin V, Camerota F, Celletti C, Tenore N, et al. Gait strategy in patients with
21 Ehlers-Danlos syndrome hypermobility type and Down syndrome. *Research in Developmental*
22 *Disabilities*. 2012;33:1437-42.
23
24 [29] Kelly VE, Eusterbrock AJ, Shumway-Cook A. Factors influencing dynamic prioritization during
25 dual-task walking in healthy young adults. *Gait & Posture*. 2013;37:131-4.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

6. Table(s)

Table 1: Mean (SD) of subject characteristics, overground walking speeds, and ankle load at each side

Group	Age (yrs)	Sex	Height (m)	Body mass (kg)	Leg length (m)	Fast speed (m/s)	Slow speed (m/s)	Ankle load (kg)
DS	9.12 (1.43)	11M/2F	1.25 (0.09)	31.67 (5.98)	0.57 (0.06)	0.99 (0.36)	0.74 (0.27)	0.60 (0.09)
TD	9.31 (1.49)	11M/2F	1.34 (0.08)*	30.41 (5.50)	0.65 (0.06)*	1.33 (0.11)*	1.00 (0.08)*	0.59 (0.12)

TD: typically developing children; DS: children with Down syndrome. Fast speed: treadmill speed that was equal to the subject's preferred overground walking speed; Slow speed: treadmill speed that was equal to 75% of the subject's preferred overground walking speed. Symbol * denotes a significant difference between the two groups at $p < 0.05$.

Table 2: Mean (SD) of normalized step length, normalized step width, swing percentage, double support percentage, and foot rotation angle.

Variable	Group	Slow Speed		Fast Speed		Statistical results
		No load	Ankle load	No load	Ankle load	
Normalized step length	TD	0.79 (0.09)	0.81 (0.10)	0.88 (0.10)	0.91 (0.10)	S: F(1,17)=112.17, p<0.001 G*L: F(1,18)=7.01, p=0.016
	DS	0.66 (0.12)	0.64 (0.13)	0.76 (0.12)	0.76 (0.15)	
Normalized step width	TD	0.19 (0.05)	0.21 (0.05)	0.18 (0.04)	0.21 (0.05)	G*L: F(1,18)=19.96, p<0.001
	DS	0.31 (0.07)	0.32 (0.07)	0.31 (0.08)	0.30 (0.05)	
Swing percentage (% gait cycle)	TD	62.43 (0.54)	62.05 (0.60)	61.53 (0.76)	60.87 (0.75)	S: F(1,17)=92.19, p<0.001 L: F(1,18)=5.16, p=0.036
	DS	64.46 (3.00)	64.49 (3.09)	62.66 (2.90)	60.29 (1.88)	
Double support percentage (% gait cycle)	TD	12.54 (0.56)	12.16 (0.64)	11.56 (0.74)	10.92 (0.67)	S: F(1,17)=104.39, p<0.001 L: F(1,18)=9.50, p=0.006
	DS	13.81 (2.21)	14.07 (2.77)	12.47 (2.64)	11.85 (1.93)	
Foot rotation angle (deg)	TD	4.13 (8.56)	4.42 (8.53)	5.15 (6.84)	4.93 (8.47)	
	DS	1.38 (10.50)	2.36 (8.58)	-0.08 (10.73)	3.74 (10.87)	

In statistical results, S: speed effect; L: load effect; G*L: group by load interaction; at $p<0.05$.

Figure captions:

Figure 1: Mean and SD of coefficient of variation of spatial gait variables at the *Fast* and *Slow* treadmill speeds. Normalized step length (i.e., step length/leg length) for TD group (a) and DS group (b), normalized step width (i.e., step width/leg length) for TD group (c) and DS group (d), and foot rotation angle for TD group (e) and DS group (f). Slow treadmill speed was equal to 75% of the subject's preferred overground walking speed; Fast treadmill speed was equal to the subject's preferred overground walking speed. No Load denotes the no-ankle load condition. Ankle Load denotes the ankle-load condition, which was equal to 2% of body weight attached above each ankle. Symbol * indicates a significant load effect at $p < 0.05$. Symbol + indicates a significant speed effect at $p < 0.05$.

Figure 2: Mean and SD of coefficient of variation of temporal gait variables at the *Fast* and *Slow* treadmill speeds. Swing percentage (in % gait cycle) for TD group (a) and DS group (b) and double support percentage (in % gait cycle) for TD group (c) and DS group (d). Slow treadmill speed was equal to 75% of the subject's preferred overground walking speed; Fast treadmill speed was equal to the subject's preferred overground walking speed. No Load denotes the no-ankle load condition. Ankle Load denotes the ankle-load condition, which was equal to 2% of body weight attached above each ankle. Symbol * indicates a significant load effect at $p < 0.05$. Symbol + indicates a significant speed effect at $p < 0.05$.

Figure 1:










Figure 2:



Research Highlights:

1. We study gait variability of treadmill walking in children with DS and TD.
2. Children with DS showed similar spatiotemporal adaptations to speed and ankle load.
3. Children with DS produced longer and wider steps but similar temporal variables.
4. Children with DS displayed greater variability in most spatiotemporal variables.
5. Children with DS had less step width variability and may prioritize ML stability.