

The Arabidopsis Polycomb Repressive Complex 1 (PRC1) Components AtBMI1A, B, and C Impact Gene Networks throughout All Stages of Plant Development

Wiam Merini², Francisco J. Romero-Campero², Angeles Gomez-Zambrano, Yue Zhou, Franziska Turck, and Myriam Calonje*

Institute of Plant Biochemistry and Photosynthesis, 41092 Seville, Spain (W.M., A.G.-Z., M.C.); Department of Computer Science and Artificial Intelligence, University of Seville, 41012 Seville, Spain (F.J.R.-C.); and Max Planck Institute for Plant Breeding Research, Department of Plant Developmental Biology, 50829 Cologne, Germany (F.T.)

ORCID IDs: 0000-0001-6606-7978 (W.M.); 0000-0001-9834-030X (F.J.R.-C.); 0000-0003-4921-0212 (A.G.-Z.); 0000-0002-4982-5949 (F.T.); 0000-0001-6351-6774 (M.C.).

Polycomb Group regulation in Arabidopsis (*Arabidopsis thaliana*) is required to maintain cell differentiation and allow developmental phase transitions. This is achieved by the activity of three PcG repressive complex 2s (PRC2s) and the participation of a yet poorly defined PRC1. Previous results showed that apparent PRC1 components perform discrete roles during plant development, suggesting the existence of PRC1 variants; however, it is not clear in how many processes these components participate. We show that AtBMI1 proteins are required to promote all developmental phase transitions and to control cell proliferation during organ growth and development, expanding their proposed range of action. While AtBMI1 function during germination is closely linked to B3 domain transcription factors VAL1/2 possibly in combination with *GT-box* binding factors, other AtBMI1 regulatory networks require participation of different factor combinations. Conversely, EMF1 and LHP1 bind many H3K27me3 positive genes up-regulated in *atbmi1a/b/c* mutants; however, loss of their function affects expression of a different subset, suggesting that even if EMF1, LHP1, and AtBMI1 exist in a common PRC1 variant, their role in repression depends on the functional context.

The evolutionarily conserved Polycomb Group (PcG) machinery plays a crucial role in maintaining repression of genes that are not required during a specific cell fate (Ringrose and Paro, 2004). PcG proteins form multiprotein complexes with different histone modifying activities, including PcG repressive complex 2 (PRC2), which possesses histone H3 Lys 27 (H3K27) trimethyltransferase activity (Müller et al., 2002), and PRC1, which has histone H2A Lys 119 E3 ubiquitin ligase activity (Cao et al., 2005) as well as other nonenzymatic functions critical for chromatin compaction (Francis et al., 2004). The combined activity of both complexes is required for stable repression of target genes.

In *Drosophila*, single-copy genes encode the four core subunits of PRC2: Suppressor of Zeste 12 [Su(z)12],

Extra sex combs (Esc), p55, and the catalytic subunit Enhancer of Zeste [E(z); Simon and Kingston, 2013]. Arabidopsis (*Arabidopsis thaliana*) has three E(z) homologs, CURLY LEAF (CLF), MEDEA (MEA), and SWINGER (SWN; Goodrich et al., 1997; Grossniklaus et al., 1998; Chanvivattana et al., 2004), and three Su(z)12 homologs, EMBRYONIC FLOWER2 (EMF2), VERNALIZATION2 (VRN2), and FERTILIZATION INDEPENDENT SEED2 (FIS2; Luo et al., 1999; Gendall et al., 2001; Yoshida et al., 2001), while MULTIPLE SUPPRESSOR OF IRA1 (MSI1), which is one of the five p55 homologs in Arabidopsis (Hennig et al., 2005), and the Esc homolog FERTILIZATION INDEPENDENT ENDOSPERM (FIE; Ohad et al., 1999) are common subunits to the different possible PRC2s (Mozgova et al., 2015).

Drosophila PRC1 contains Polycomb (Pc), Polyhomeotic (Ph), Posterior sex comb (Psc), and dRing1 (Shao et al., 1999; Peterson et al., 2004), each with multiple homologs in vertebrates (Schwartz and Pirrotta, 2013). Furthermore, vertebrate PRC1 complexes exist in canonical or noncanonical forms. Canonical variants harbor homologs to the four *Drosophila* core subunits (Schwartz and Pirrotta, 2013), while noncanonical PRC1 complexes contain RING1A or RING1B and one of the six different homologs of *Drosophila* Psc (PCGF) to form a H2A monoubiquitination module, along with additional subunits that further add specific biochemical properties and genomic localization to the different variants (Schwartz and Pirrotta, 2013). In Arabidopsis,

¹ This work was supported by Marie Curie CIG Grant 333748 and BIO2013-44078-P Grant from Ministerio de Economía y Competitividad. F.T. and Y.Z. are supported by core funding from the Max Planck Gesellschaft.

² These authors contributed equally to the article.

* Address correspondence to myriam.calonje@ibvf.csic.es.

The author responsible for distribution of materials integral to the

findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Myriam Calonje (myriam.calonje@ibvf.csic.es).

W.M., A.G.-Z., and Y.Z. performed the experiments; W.M. and F.J.R.-C. analyzed the data; M.C. conceived and designed the study; M.C. and F.T. prepared the manuscript.

several pieces of evidence suggest a similar high degree of complexity (Förderer et al., 2016). Two RING1 homologs, AtRING1A and AtRING1B, and three Psc/PCGF homologs, AtBMI1A, AtBMI1B, and AtBMI1C, have been characterized (Sanchez-Pulido et al., 2008; Xu and Shen, 2008; Bratzel et al., 2010, 2012; Chen et al., 2010; Yang et al., 2013; Calonje, 2014). Plants with mutations in these genes suggest a high degree of functional redundancy between AtRING1 or AtBMI1 proteins; thus, it is not clear whether each paralog can regulate a different subset of targets (Bratzel et al., 2010; Chen et al., 2010; Yang et al., 2013). The analysis is complicated by the observation that several mutant alleles are knock-downs rather than null alleles and that phenotypes show a wide range of stochastic variation among segregating siblings with “weak” and “strong” phenotypes (Bratzel et al., 2010; Chen et al., 2010).

Two other plant-specific proteins have been related to PRC1, EMBRYONIC FLOWER1 (EMF1) mediating chromatin compaction (Calonje et al., 2008; Beh et al., 2012) and LIKE-HETEROCHROMATIN PROTEIN1 (LHP1), which, as *Drosophila* Pc, binds H3K27me3 marks through its chromodomain (Turck et al., 2007). Although both proteins can interact with either AtRING1 or AtBMI1 (Bratzel et al., 2010; Chen et al., 2010), recent reports showed that they also copurify with PRC2 components (Derkacheva et al., 2013; Liang et al., 2015); thus, it is not clear in which context they carry out their functions. Additional proteins with chromatin-related functions have been shown to participate in PRC1-mediated repression of specific target genes, such as the VIVIPAROUS1 (VPT)/ABSCISIC ACID INSENSITIVE3 (ABI3)-Like1 and 2 proteins (VAL1/2; Yang et al., 2013), ALFIN1-like proteins (ALS; Molitor et al., 2014), and JM14 (Wang et al., 2014).

In plants, PcG repression maintains the differentiated state of the cells but also orchestrates developmental phase transitions by controlling the establishment of new cell identities. This likely requires different PRC1s, but little is known about their subunit composition. The repression of several seed maturation genes after germination requires the AtBMI1 and AtRING1 proteins (Bratzel et al., 2010; Chen et al., 2010; Yang et al., 2013), and a recent genome-wide study showed gene networks regulated by AtBMI1s and AtRING1s during the suppression of seed development in seedlings (Wang et al., 2016). As these results were derived from the analysis of *atring1a/b* and *atbmi1a/b* mutants developing a weak phenotype (Bratzel et al., 2010; Chen et al., 2010), their possible implication in other developmental processes or stages was not unveiled. Conversely, the repression of flower homeotic genes in seedlings requires EMF1 (Kim et al., 2012) and LHP1 (Gaudin et al., 2001), but their role in regulating other processes is not clear.

In this work, by analyzing the transcriptome of single, strong double, and triple *atbmi1* mutants, we have identified a more comprehensive set of candidate genes regulated by AtBMI1 proteins. Our results indicate that in addition to switching off the seed maturation

program after germination, AtBMI1s promote the transition from each developmental phase to the next throughout development and furthermore control cell proliferation during organ growth and development. By integrating transcriptomics datasets with previously published data, we show that AtBMI1 and VAL1/2 act together only in the regulation of seed maturation genes. Enrichment of cis-regulatory elements at VAL1/2-dependent and -independent genes suggests that AtBMI1-mediated gene repression requires different combinational modules always involving VAL-related B3 domain factors. Conversely, while EMF1 and LHP1 occupy a considerable number of genes up-regulated in *atbmi1a/b/c* mutants, loss of their function does not impact the expression of most but affects the expression of a different subset of genes. Together these results suggest that the different PRC1 variants may differ in subunit composition but also in the role that single components play all depending on the cis-regulatory context.

RESULTS

Genome-Wide Transcriptomic Data Analysis of *atbmi1* Mutants

Previous data have suggested that *AtBMI1A* and *AtBMI1B* are ubiquitously expressed and act mostly redundantly throughout development (Bratzel et al., 2010), whereas *AtBMI1C*, which is expressed in roots, endosperm, and stamen, may have functionally diverged since it cannot fully rescue *atbmi1a/b* defects when overexpressed (Yang et al., 2013; Merini and Calonje, 2015); nevertheless, *atbmi1a/c* and *atbmi1b/c* do not show phenotypic alterations (Yang et al., 2013), suggesting that loss of AtBMI1C function is compensated by the other two AtBMI1s. Therefore, to gain insight into the regulatory roles of AtBMI1s, we performed genome-wide transcriptome analysis using RNA sequencing (RNA-seq) of wild-type Col-0, *atbmi1a*, *atbmi1b*, *atbmi1a/b*, and *atbmi1a/b/c* mutants at 10 d after germination (DAG). Since individual *atbmi1a/b* double mutants display a wide range of phenotypes (Bratzel et al., 2010), we chose to select the strong *atbmi1a/b* mutant phenotype for the analysis, which differs from the *atbmi1a/b/c* phenotype mainly in the root (Yang et al., 2013; Supplemental Fig. S1). The Tuxedo protocol (Trapnell et al., 2012) was used for transcript assembly and differential expression analysis. All sequencing samples were of high quality (Supplemental Fig. S2; Supplemental Table S1). Differentially expressed genes were determined using stringent criteria consisting of a combination of fold change >4 and a *P* value < 0.05. The number of genes scored as present in at least one of our samples was 24,503, representing 72.96% of the entire Arabidopsis transcriptome. We found <3% to 4% of the surveyed transcriptome affected in single mutants and around 15% and 20% differentially expressed in strong *atbmi1a/b* double and *atbmi1a/b/c* triple mutants, respectively (Fig. 1A; Supplemental Fig. S3). Principal components analysis showed that the transcriptomes of wild

type, *atbmi1a*, and *atbmi1b* mutants clustered together, whereas the transcriptomes of *atbmi1a/b* and *atbmi1a/b/c* mutants constituted two distant and distinct clusters, indicating not only differences from the wild type and single mutant group but also in between (Fig. 1B). In any case, we found a considerable number of genes misregulated in the single mutants (Fig. 1C; Supplemental Table S2), of which a majority were a subset of those affected in double and triple mutants (Supplemental Fig. S4, A and B). The number of up-regulated genes for *atbmi1a*, *atbmi1b*, and *atbmi1a/b* was higher than down-regulated (Fig. 1C), which might confirm the role of AtBMI1 proteins in transcriptional repression. However, *atbmi1a/b/c* mutant showed higher number of down-regulated genes than up-regulated genes. This may be a consequence of the developmental stage of these mutants, in which all organs are stuck in a seed maturation phase. Up-regulation of some genes within this context may have a stronger negative impact on gene expression.

Globally, the upregulated genes in the strong *atbmi1a/b* and *atbmi1a/b/c* mutants (Supplemental Figs. S5A and S6A) showed overrepresentation of Gene Ontology terms associated with response to different stimuli (e.g. water stress, temperature, hormones) and lipid metabolism (e.g. transport, biosynthesis, storage), whereas the down-regulated genes were enriched for Gene Ontology terms related to photosynthesis and

metabolic processes (Supplemental Figs. S5B and S6B). This is consistent with the developmental fate of the mutants, which are trapped in the seed maturation phase (Yang et al., 2013). During this phase, seeds acquire desiccation tolerance and accumulate storage reserves, prevailing in the form of lipids (Vicente-Carbajosa and Carbonero, 2005), while chloroplast structure is disrupted (Delmas et al., 2013).

As PcG function is involved in the repression of master regulatory genes (Xiao and Wagner, 2015), misregulation in the different *atbmi1* mutants may be an indirect or direct consequence of the loss of AtBMI1 function, or a mix of both. Conversely, a considerable number of AtBMI1 direct target genes may not display altered expression in the absence of their upstream transcriptional activators, as has been reported for other PcG loss of function mutants (Bouyer et al., 2011; Kim et al., 2012; Derkacheva et al., 2013). In any case, although the interrelationship between PRC1 and PRC2 is not clear yet, the activity of both complexes is required for stable PcG-mediated repression; therefore, selecting genes up-regulated in *atbmi1* mutants and H3K27me3 marked in wild-type seedlings should enrich for a subset of candidate genes directly controlled by AtBMI1s. Accordingly, we intersected genes up-regulated in the different mutants with a set of 5360 H3K27me3 target genes previously identified in




Figure 1. Transcriptome analysis of wild type and selected *atbmi1* mutants at 10 DAG. A, Volcano plots representing differentially expressed genes in *atbmi1* mutants compared to wild type according to a 4-fold change and a *P* value of 0.05. Green color indicates significantly up-regulated genes and red color significantly down-regulated genes. B, Principal component analysis of the transcriptomes showing that wild type, *atbmi1a*, and *atbmi1b* cluster together, whereas *atbmi1a/b* and *atbmi1a/b/c* constitute two distinct clusters. C, Differentially expressed genes in the different genotypes, where the number of up- and down-regulated genes is indicated. D, Number of genes that were up-regulated in the different mutants and H3K27me3 marked in wild-type seedlings of the same age (up_K27).

two independent analyses in seedlings (Supplemental Table S3; Bouyer et al., 2011; Kim et al., 2012) to selected up-regulated H3K27me3 positive (up_K27) genes (Fig. 1D). The analysis showed significant overlaps between H3K27me3 marked genes and up-regulated genes in the different mutants except for *atbmi1b*, probably because it is a knock-down mutant (Bratzel et al., 2010). The same analysis using down-regulated genes showed nonsignificant overlaps in all cases excluding *atbmi1a/b/c* due to the high number of down-regulated genes in this mutant (Supplemental Fig. S7; Supplemental Table S3).

To determine whether there were AtBMI1A and AtBMI1B specific candidate targets, we compared up_K27 genes in the single and double mutants (Fig. 2A). Their number in the double mutant was considerably higher than in the single mutants, illustrating a high degree of functional redundancy. Also, most of the up_K27 genes in single mutants were included in the double mutants set of up_K27 genes; however, a group of genes seemed to be exclusively up-regulated in *atbmi1a* and *atbmi1a/b* or in *atbmi1b* and *atbmi1a/b* (104 and 27 genes, respectively). Up_K27 genes in *atbmi1a* and *atbmi1a/b* were expressed at very low levels in both single compared to the double mutants (Fig. 2B), indicating redundant regulation by

AtBMI1A and B. The *atbmi1b* mutant shows some remnant expression of *AtBMI1B*, possibly explaining higher expression in *atbmi1a* versus *atbmi1b* and the greater number of affected genes in the *atbmi1a* single mutant (Bratzel et al., 2010). Nevertheless, some genes were indeed specifically sensitive to AtBMI1B being more affected in *atbmi1b* than *atbmi1a* and not further increased in double mutants (Fig. 2B).

We next investigated the degree of redundancy between AtBMI1A/B and AtBMI1C by comparing the genes up_K27 in *atbmi1a/b* and *atbmi1a/b/c* (Fig. 3A). Clustering analysis showed that *atbmi1a/b* and *atbmi1a/b/c* shared two-thirds of the up_K27 genes (Cluster I, Supplemental Table S3) but the remaining one-third was genotype-specific (Cluster II, *atbmi1a/b/c* specific; and Cluster III, *atbmi1a/b* specific). The expression pattern of genes in Cluster I fell into two distinct subgroups. Cluster Ia included genes that displayed a gradual increase of expression in double and triple mutants, suggesting redundant regulation by AtBMI1A/B and AtBMI1C (Fig. 3B; Supplemental Fig. S8A). Cluster Ib contained genes whose regulation may depend exclusively on AtBMI1A/B, as the loss of AtBMI1C function did not affect significantly




Figure 2. Genes regulated by AtBMI1A and AtBMI1B. A, Venn diagram showing the number of up_K27 genes that overlap among *atbmi1a*, *atbmi1b*, and *atbmi1a/b* mutants. All overlaps are significant with $P < 2.2 \times 10^{-16}$ and odds ratios > 17 according to Fisher's exact test. B, Expression of levels of genes that were apparently specifically up-regulated in *atbmi1a* or *atbmi1b* mutants in the different genotypes.




Figure 3. Functional redundancy between AtBMI1A/B and AtBMI1C. A, Clustering analysis of genes up_K27 in *atbmi1a/b* and *atbmi1a/b/c* mutants. This is a significant overlap with $P < 2.2 \times 10^{-16}$ and an odds ratio >21 according to Fisher's exact test. B, Expression levels in wild type, *atbmi1a/b*, and *atbmi1a/b/c* of genes from the different clusters. The color code represents normalized expression values measured in FPKM.

their overall expression levels (Fig. 3B; Supplemental Fig. SA8). Cluster II (Supplemental Table S3) included genes exclusively up-regulated in *atbmi1a/b/c*, indicating that these are AtBMI1C specific targets or, alternatively, that AtBMI1C fully compensates the loss of AtBMI1A/B function in regulating these genes (Fig. 3B; Supplemental Fig. S8A). To discern between these two possibilities, we measured the levels of a subset of cluster II genes in wild type, *atbmi1c* single, and *atbmi1a/b/c* mutants in whole seedlings and roots at 10 DAG by quantitative reverse transcription PCR (qRT-PCR). As they were not misexpressed in *atbmi1c* single mutants (Supplemental Fig. S8B), we concluded that AtBMI1C compensates for the loss of AtBMI1A/B function in the regulation of these genes. Finally, genes in Cluster III (Supplemental Table S3) were exclusively up-regulated in *atbmi1a/b* mutants, but not in *atbmi1a/b/c* (Fig. 3B; Supplemental Fig. S8A). Although a priori unexpected, the result can be explained if the activation of these genes requires a developmental stage that is not reached in *atbmi1a/b/c*.

All together, these data indicated that AtBMI1A and B regulate genes predominantly redundantly, whereas AtBMI1C affects only a subset of AtBMI1A/B possible targets.

Deregulated Developmental Programs in *atbmi1* Mutants

AtBMI1 proteins were previously shown to participate in the regulation of several seed maturation- (Bratzel et al., 2010; Chen et al., 2010; Yang et al., 2013) and germination-related genes (Molitor et al., 2014). In addition, a recent transcriptome analysis of *atbmi1a/b* weak phenotype confirmed the role of AtBMI1 function in regulating seed development (Wang et al., 2016). When we compared the H3K27me3 up-regulated genes in the *atbmi1a/b* weak (fold change ≥ 2 , according to Wang et al., 2016; Supplemental

Table S1) to those in *atbmi1a/b* strong phenotype mutants, we found significantly more genes in the stronger mutant (Fig. 4A). Among the genes up-regulated in both datasets were genes previously identified as AtBMI1 target genes, like *ABI3*, and *DELAY OF GERMINATION1 (DOG1)*; however, other well-known AtBMI1 targets, such as *FUSCA3 (FUS3)* or *BABYBOOM (BBM)* (Yang et al., 2013), were included only in the *atbmi1a/b* strong dataset. A similar picture was obtained comparing *atbmi1a/b* weak and *atbmi1a/b/c* datasets (Supplemental Fig. S9). Therefore, to obtain a more comprehensive picture of the developmental processes regulated by AtBMI1s, we examined the annotated developmental functions of up_K27 genes in *atbmi1a/b/c* mutants, as they displayed the strongest developmental alterations.

Seed Maturation and Dormancy

Changes in the triple *atbmi1a/b/c* mutant uncovered additional genes involved in seed maturation and abscisic acid (ABA) response, such as *FUS3* and *ABI4*, and in seed dormancy, like *SOMNUS (SOM)*. Also, there were genes involved in regulating carbohydrate and lipid metabolism, like *WRINKLED1* (Supplemental Fig. S9; Supplemental Table S3). Most of these genes are switched off after germination in wild type; however, the *ABIs* are required for plant responses to various biotic and abiotic stresses (Cutler et al., 2010), suggesting involvement of AtBMI1s in regulating responses to environmental conditions.

Endosperm-Specific Genes

Maturation genes were not the only seed genes up-regulated in *atbmi1a/b/c* mutants. We found up-regulation of genes that are predominantly expressed




Figure 4. Different gene expression patterns of *atbmi1a/b* weak and strong mutants. A, Venn diagram showing overlap between the genes up_K27 in *atbmi1a/b* weak and strong mutants. The overlap is significant with $P < 2.2 \times 10^{-16}$ and an odds ratio > 15 according to Fisher's exact test. Some representative TFs in each dataset are indicated. TFs found in the two data sets are highlighted in red. B, Key flowering genes are down-regulated in *atbmi1a/b/c* mutants. The color code in upper panel represents normalized expression values measured in FPKM.

in endosperm but not in the seed coat and vegetative tissues (Wolff et al., 2011). Interestingly, among these were genes displaying a maternal (*FLOWERING WAGENINGEN*, *HOMEODOMAIN GLABROUS8*, and *AtBMI1C*) or paternal (*PICKLE RELATED2* [*PKR2*], *VARIANT IN METHYLATION5*, *AT2G21930*, and *AT3G49770*) preferred expression in the endosperm (Supplemental Fig. S9; Supplemental Table S3).

Meristem Maintenance and Cell Proliferation-Related Genes

The *atbmi1a/b/c* mutant also up-regulated genes involved in meristem maintenance and cell proliferation throughout plant life. Remarkably, two gene families with crucial roles in these processes were up-regulated in the mutants. The first encompassed the *PLETHORA* (*PLT*) or *AINTEGUMENTA-LIKE* (*AIL*) genes. Six of

eight members of this family were up_K27 in *atbmi1a/b/c* mutants (*PLT1/2/3/5/7* and *BBM*; Supplemental Figs. S9 and S10; Supplemental Table S3). Some of these *PLT* genes have overlapping roles in regulating embryo patterning, shoot and root apical meristem maintenance, and organ primordia initiation (Horstman et al., 2014). The second was the *WUS homeobox-containing* (*WOX*) gene family, which comprises 14 members (van der Graaff et al., 2009), among which *WUS* and *WOX2/3/4/5/8/9/11/12* were up-regulated in *atbmi1a/b/c* mutants (Supplemental Figs. S9 and S10; Supplemental Table S3). These factors promote cell division and prevent premature cell differentiation, which are crucial processes required for stem-cell maintenance and organ formation. In addition, we found up-regulation of other genes with related functions, for instance, *CUP SHAPED COTYLEDON3*, *ENHANCER OF SHOOT REGENERATION1*, and *GROWTH REGULATING FACTOR5*.

Root Development-Specific Genes

Apart from the genes involved in root meristem maintenance, we found in *atbmi1a/b/c* up-regulation of genes that play a crucial role in postembryonic root development, as *CEGENDUO*, *MAGPIE*, *INDOLE-3-ACETIC ACID INDUCIBLE30*, the *ROOT MERISTEM GROWTH FACTOR2*, and the Class IIB NAC transcription factor *SOMBRERO* (*SMB*), underpinning the importance of AtBMI1 function for root development (Supplemental Fig. S9; Supplemental Table S3).

Other Developmental Genes

Among the up_K27 genes in *atbmi1a/b/c* mutants were genes involved in regulating other developmental processes, such as gametophyte development, leaf development, and the flowering transition (e.g. *KANADI2*, *KNUCKLES*, *DEVELOPMENT-RELATED PcG TARGET IN THE APEX4*, *SEPALLATA2*, *FLOWERING LOCUS C* [*FLC*], *MADS AFFECTING FLOWERING4* [*MAF4*], *MAF5*, *FACTOR PROMOTING FLOWERING1*; Supplemental Fig. S9; Supplemental Table S3).

Secondary Metabolic Processes

In addition, *atbmi1a/b/c* mutants up-regulated genes involved in secondary metabolic processes like those involved in phenylpropanoid metabolism. Up-regulated genes involved in this pathway were *CHALCONE SYNTHASE* (*TRANSPARENT TESTA4* [*TT4*], *CHALCONE ISOMERASE* [*TT5*], *FLAVONOID 3'-HYDROXYLASE* [*TT7*], *DIHYDROFLAVONOL 4-REDUCTASE*, and transcription factors [TFs] such as *AtMYB90* [*PRODUCTION OF ANTHOCYANIN PIGMENT2*]), *AtMYB111*, and *AtMYB11* (Supplemental Fig. S9; Supplemental Table S3).

In summary, AtBMI1 function in Arabidopsis is required to regulate more developmental processes than previously thought.

Regulatory Cross Talk between Chromatin Complexes

RNA-seq data revealed up-regulation of several PcG or PcG-related genes in *atbmi1a/b/c* mutants, like *AtRING1A*, *AtRING1B*, *VAL1*, *VAL2*, and *VIN3*. Conversely, we did not find a significant change in the expression of *CLF*, *SWN*, *MEA*, *EMF2*, *VRN2*, *FIS2*, *MSI1*, *FIE*, *EMF1*, and *LHP1* (Supplemental Fig. S10). On the other hand, the Trithorax Group genes *ULTRA-PETALA1* (*ULT1*), *ULT2*, and *PKR2* that act antagonistically to PcG complexes were up-regulated in *atbmi1a/b/c* mutants (Supplemental Fig. S11). Misregulation of some of these chromatin factors could contribute to the strongly altered expression pattern of *atbmi1a/b/c* mutants.

Several Master Regulators of the Flowering Program Are Down-Regulated in *atbmi1a/b/c* Mutants Several MADS-box transcription factors required to specify floral meristem identity or involved in floral

organ development were down-regulated in *atbmi1a/b/c* mutants (Fig. 4B; Supplemental Table S2; e.g. *AGL42*, *SUPPRESSOR OF CONSTANS1* [*SOC1*], *SEP3*, *SEP4*, *AGL24*, *SHORT VEGETATIVE PHASE*), but also other key regulatory flowering genes, such as *TEMPRANILLO1* and several *SQUAMOSA PROMOTER BINDING PROTEIN-LIKE* (*SPLs*; e.g. *SPL2*, 3, 4, 8, 12). In addition, we found that some flowering factors that have basal expression levels in wild-type seedlings at 10 DAG expressed at lower levels in *atbmi1a/b/c* (e.g. *AGAMOUS* [*AG*], *APETALA3*, *FLOWERING LOCUS T* [*FT*]; Fig. 4B). The fact that the flowering program seems to be more repressed in *atbmi1a/b/c* mutants than in wild-type seedlings points to a requirement of AtBMI1 function for proper regulation of flower development.

VAL1/2 and the AtBMI1s Coregulate a Subset of Potential AtBMI1 Targets

VAL1/2 and AtBMI1 proteins are required for the initial repression of several seed maturation genes after germination, such as *FUS3*, *LEC1*, and *ABI3*. Furthermore, we previously showed that the VAL1/2 recruit AtBMI1 proteins to these genes; accordingly, *val1/2* and *atbmi1a/b/c* mutants display a very similar phenotype (Yang et al., 2013). However, *WUS* is an AtBMI1 but not a VAL1/2 regulated gene, indicating that there are also differences between those mutants (Yang et al., 2013). To determine to which extent the VAL1/2 and AtBMI1 proteins act together in regulating gene expression, we compared genes up-regulated in *val1/2* (Suzuki et al., 2007; Supplemental Table S2) and H3K27me3 marked in wild type according to our dataset (Supplemental Table S3) with up_K27 genes in *atbmi1a/b/c* (Fig. 5A). We found that 70% of *val1/2* up_K27 genes were included in the up_K27 *atbmi1a/b/c* dataset; these genes represented one-third of the genes up_K27 in *atbmi1a/b/c*, indicating that, despite the fact that they coregulate a considerable number of genes, AtBMI1 proteins clearly perform functions independently of VAL1/2.

The VAL proteins (VAL1, 2, and 3) belong to a subfamily of plant-specific B3 domain-containing proteins (Swaminathan et al., 2008) that is predicted to bind to *LEC2/ABI3/VP1* elements (also known as *RY* elements [*CATGCA*]; Suzuki et al., 2007); in fact, a recent report showed that a point mutation in a *LEC2/ABI3/VP1* element located at the first intron of *FLC* prevents the epigenetic silencing of the gene during vernalization (Quèsta et al., 2016). *FLC* is up-regulated in *val1/2* and *atbmi1* mutants (Supplemental Tables S2 and S3). Therefore, we investigated whether this or other cis-regulatory motifs were enriched at the promoter of AtBMI1/VAL1/2 coregulated genes. Indeed, we found enrichment of *LEC2/ABI3/VP1* motifs but also of ABA responsive elements (*ABRE*; ACGT or *G-box*; Choi et al., 2000; Fig. 5A). *ABRE/G-box* elements are recognized by bZIP transcription factors such as *ABI5* (Carles et al., 2002). *LEC2/ABI3/VP1* and *ABRE* elements are clustered in the 5' upstream regions of genes regulated by *ABI3/VP1* factors and ABA (Suzuki et al., 2005) and are




Figure 5. Interplay of AtBMI1 proteins with VAL1/2 proteins. A, Venn diagram showing overlap between the genes up_K27 in *atbmi1a/b/c* and *val1/2* mutants. Sequence LOGOs of cis-regulatory elements enriched only in up_K27 *atbmi1a/b/c* and in *atbmi1a/b/c* and *val1/2* overlapping genes. B, Co-occurrence and overlapping of ABRE/G-box and GT-box at the promoter of AtBMI1/VAL1/2 coregulated genes. P values and percentage in targets and background are indicated.

required for the correct expression of seed maturation genes (Santos-Mendoza et al., 2008). On the other hand, the plant-specific trihelix DNA binding protein ARABIDOPSIS 6B-INTERACTING PROTEIN 1-LIKE1 (ASIL1) that is involved in the repression of seed maturation genes after germination binds *GT-box* elements (GTGATT and variations of this; Gao et al., 2009). These elements are closely associated with ABRE/G-box and LEC2/ABI3/VP1 elements at the promoter of several seed maturation genes. Furthermore, *GT-box* elements frequently overlap with ABRE/G-box elements, leading to the proposal that ASIL1 represses embryonic genes by competing with the binding of transcriptional activators (Gao et al., 2009). Therefore, we looked for co-occurrence of both elements at the promoter of AtBMI1/VAL1/2 coregulated genes. Co-occurrence was indeed significant (Fig. 5B); moreover, both elements significantly overlapped at the promoter of these genes (Fig. 5B). Therefore, the combination of LEC2/ABI3/VP1 and *GT-box* co-occurring with ABRE/G-box elements represents a landmark for the subset of AtBMI1/VAL1/2 coregulated genes.

Surprisingly, the LEC2/ABI3/VP1 elements were as highly overrepresented at promoter regions of genes exclusively up_K27 in *atbmi1a/b/c*, which suggests that their repression may be functionally connected to other B3 domain transcription factors. The specific combination of LEC2/ABI3/VP1 and ABRE/GT-box elements was not detected in this group. Conversely, other motifs

were enriched in the VAL1/2-independent up_K27 subset, such as SQUAMOSA BINDING PROTEIN-, ZAP1-, ALFIN1-, and MYB-binding sites and a frequent Z-box promoter motif that is bound by a new class of transcription factors, the Z-box BINDING FACTORS, whose roles in regulating plant development have just started to be unraveled (Gangappa et al., 2013; Fig. 5A). ALFIN1 elements are bound by plant-specific ALFIN1-like proteins (AL1–7; Lee et al., 2009), which mediate gene repression (Wei et al., 2015) and interact with AtRING1 and AtBMI1 (Molitor et al., 2014), supporting the existence of other combinatorial modules involving B3 domain factors and diverse partners for AtBMI1-mediated gene repression.

Regulatory Networks of AtBMI1, EMF1, and LHP1

To investigate the functional relationship between AtBMI1 proteins and EMF1, we compared direct EMF1 targets as previously determined through genome-wide ChIP-chip analysis (Kim et al., 2012) with our WT_K27 gene dataset and with genes with altered expression (up- and down-regulated) in *atbmi1a/b/c* mutants (Supplemental Fig. S1A; Supplemental Table S4). Clustering analysis showed a subgroup of 786 overlapping genes, indicating that among the misexpressed genes in *atbmi1a/b/c* there is a significant amount of EMF1 targets. Then, we determined the number of up_K27 genes in *atbmi1a/b/c* that were included in this

subgroup (Fig. 6A). We found that one-half of *atbmi1a/b/c* up_K27 genes were EMF1 targets, suggesting interplay of EMF1 and AtBMI1 proteins in the regulation of a considerable number of genes.

There was little overlap between genes up_K27 in *atbmi1a/b/c* and *emf1-2* (Fig. 6B; Supplemental Table S4; Kim et al., 2010); furthermore, the majority of EMF1 target genes up_K27 in *atbmi1a/b/c* was not up-regulated in *emf1-2* mutants, which is consistent with the previous observation that expression of only a small percentage of EMF1 target genes is increased in *emf1-2* mutants (Kim et al., 2012; Fig. 6C). LHP1 has been shown to colocalize with 85% to 90% of H3K27me3 marked sites in Arabidopsis (Turck et al., 2007; Zhang et al., 2007; Engelhorn et al., 2012); consistent with this, 92.3% of our list of H3K27me3 marked genes (4,949 of 5,360) were occupied by LHP1 according to a recently published data set of LHP1 targets (Veluchamy et al., 2016); of these genes, 1,406 significantly overlap with the genes misexpressed (up- and down-regulated) in *atbmi1a/b/c* mutants (Supplemental Table S4; Supplemental Fig. S1B). Furthermore, we found that 93.9% of *atbmi1a/b/c* up_K27 genes were LHP1 targets (Fig. 6C), suggesting that AtBMI1 and LHP1 coregulate a high number of genes. However, when we compared H3K27me3 marked genes up-regulated in *lhp1* (fold change ≥ 2 , according to Wang et al., 2016; Supplemental Table S3) with up_K27 *atbmi1a/b/c* genes (Fig. 6D), we found very little overlap, indicating that loss of LHP1 function has also little impact on the expression of AtBMI1 regulated genes. Loss of LHP1 function, as loss of EMF1 function, mostly impacts the expression of genes involved in reproductive development. These genes were not up-regulated in *atbmi1a/b/c* mutants and some were even repressed, suggesting that LHP1 and EMF1 play different roles in their regulation. In conclusion, regulation is not correlated to the codistribution of EMF1 and LHP1 and likely also AtBMI1 proteins at target genes.

DISCUSSION

PcG regulation in Arabidopsis requires the activity of three different PRC2s, which regulate different developmental stages and display partial target specificity, and PRC1, whose identity and function is not yet well defined. Although several putative subunits have been identified (Merini and Calonje, 2015) and some evidence suggested the existence of different functional PRC1 variants (Yang et al., 2013; Calonje, 2014; Wang et al., 2014; Merini and Calonje, 2015), little is known about their composition and function. In this work, we integrated genome-wide transcriptome data with H3K27me3 and protein localization data to shed some light on the role of different PRC1 components and their possible relationship throughout plant development.

Functional Redundancy among the AtBMI1s

The identification of three AtBMI1 paralogs in Arabidopsis raised the question of whether they display

functional divergence (Sanchez-Pulido et al., 2008). We found that AtBMI1A and B display mainly redundant functions throughout development, although a small number of genes were specifically sensitive to AtBMI1B. A splice variant is annotated at the *AtBMI1B* locus (the Arabidopsis Information Resource), which encodes a variant isoform without the amino-terminal RING finger domain (Supplemental Fig. S12). It is possible that alternative roles of the variant protein explain the observed differences in gene expression between *atbmi1a* and *atbmi1b* mutants. Conversely, AtBMI1C regulates a subset of AtBMI1A/B targets. The fact that ectopic expression of *AtBMI1C* in double mutants (Yang et al., 2013; Supplemental Table S2) cannot rescue *atbmi1a/b* defects in the aerial part of the seedling points to a requirement of tissue-specific factors for AtBMI1C mediated repression. Accordingly, AtBMI1C acts redundantly to AtBMI1A/B in the regulation of a considerable number of genes involved in root development. Differences in protein sequence between AtBMI1C and AtBMI1A/B (Bratzel et al., 2010, 2012; Chen et al., 2010) may have restricted the possibilities of AtBMI1C to interact with some factors and/or favored interaction with others. Likewise, MEA cannot compensate the loss of CLF and SWN function despite its ectopic expression in *clf/swn* double mutants (Farrona et al., 2011). In any case, AtBMI1A, AtBMI1B, and in part AtBMI1C display functional redundancy, indicating how important it is to ensure AtBMI1 function throughout development.

Role of AtBMI1 Function in Plant Development

Transcriptome analysis revealed that 20% of the surveyed transcriptome was misregulated in *atbmi1a/b/c* mutants, a much higher percentage than the one reported for other PcG mutants, including *clf/swn* (Bouyer et al., 2011; Kim et al., 2012; Wang et al., 2016), thereby underlining the central role of AtBMI1s in gene regulation. To determine the AtBMI1 regulatory gene network, we focused on genes that were up-regulated in *atbmi1* mutants and H3K27me3 marked in wild-type seedlings of the same age, even though these genes may represent a subset of candidate AtBMI1 targets. Our analysis supported a requirement of AtBMI1 function for the repression of the seed maturation/dormancy program after germination (Bratzel et al., 2010; Chen et al., 2010; Molitor et al., 2014; Wang et al., 2016); however, it also unveiled the crucial role of these proteins in promoting the transition from one developmental phase to the next throughout development (Fig. 7A). After embryogenesis, plants undergo the transition from seed dormancy to germination that is antagonistically regulated by two hormones, ABA and gibberellins (GA; Shu et al., 2016). During seed maturation, endogenous ABA accumulates in the seed, inducing and maintaining seed dormancy. In contrast, before the onset of germination, endogenous ABA levels in the seed are down-regulated, while the GA content is up-regulated. Among the up-regulated genes in *atbmi1a/b/c* mutants




Figure 6. AtBMI1, EMF1, and LHP1 regulatory networks. A, Comparison of genes H3K27me3 marked bound by EMF1 and misexpressed in *atbmi1a/b/c* and with genes up_K27 in *atbmi1a/b/c*. B, Venn diagram showing up_K27 genes in *atbmi1a/b/c* and *emf1-2*. C, Comparison of genes H3K27me3 marked bound by LHP1 and misexpressed in *atbmi1a/b/c* and with genes up_K27 in *atbmi1a/b/c*. D, Venn diagram showing up_K27 genes in *atbmi1a/b/c* and *lhp1*. Some overlapping and nonoverlapping representative genes are indicated. All these overlaps are significant (P values and Fisher's exact test results are indicated).

were genes involved in inducing ABA and/or inhibiting GA signaling (e.g. *ABI3*, *ABI4*, *DOG1*, *PLT5*, *SOM*; Fig. 6A), indicating that AtBMI1-mediated repression of these genes promotes this developmental transition. Following germination, plants pass through a phase of vegetative growth that can be further divided into a juvenile and an adult vegetative phase. The *microRNA 156* (*miR156*) regulates a subset of SPL transcription factors that have been shown to promote the transition from juvenile to adult phase (Wu and Poethig, 2006); therefore, to allow phase transition, *miR156* levels need to decrease. Although our transcriptome analysis could not detect mature *miRNAs*, it has been previously shown that *pri-miR156* was up-regulated in *atbmi1a/b* mutants of all phenotypic severity (Picó et al., 2015); accordingly, we found down-regulation of several *SPLs* (e.g. *SPL2/3/4/8/12*; Fig. 6A), supporting that AtBMI1 function is required to allow this transition. Eventually, plants experience the transition from vegetative to reproductive development. This transition requires the repression of several flowering repressors such as *FLC*, *MAF4/5* (Gu et al., 2013), and *AGL15* (Fernandez et al., 2014), which are up-regulated in double and triple *atbmi1* mutants (Fig. 7A). Consequently,

flowering genes like *FT*, *SOC1*, and *AGL24* were down-regulated in *atbmi1* mutants; therefore, AtBMI1 activity is also required to switch from vegetative to reproductive development.

Furthermore, our data revealed the key role of AtBMI1 activity in controlling stem cell niche specification and cell proliferation for proper organ growth and development via the repression of several master regulators (e.g. *PLT* and *WOX* genes; Fig. 7B), which is consistent with the widespread acquisition of proliferating capacity of *atbmi1* strong mutants and the alterations in root, leaf, and flower development observed in different *atbmi1* mutants (Bratzel et al., 2010; Yang et al., 2013).

Interplay of AtBMI1 with Other PcG-Related Factors

The function of AtBMI1 has been linked to the function of VAL1/2 proteins for the regulation of several seed maturation genes (Yang et al., 2013). Here, we show that VAL1/2 and AtBMI1s act together in the regulation of the seed maturation/dormancy program; however, they do not seem to collaborate in the




Figure 7. Role of AtBMI1 proteins in regulating plant development. A, AtBMI1 proteins and PRC2 promote developmental phase transitions by the repression of key regulatory genes. B, AtBMI1 and PRC2 are required to control cell proliferation and differentiation during organ growth and development through the repression of master regulators. C, PRC1 variants differing in component composition and biochemical properties may collaborate with PRC2 activity in regulating phase transitions and different developmental processes throughout plant development. VAL and ASIL1/2 or AL1 to 7 proteins may recruit AtBMI1-containing complexes to target gene promoters by binding the appropriate combination of cis-regulatory elements.

regulation of other developmental processes. We found a specific enrichment of *LEC2/ABI3/VP1* and *ABRE/G-box* overlapping with *GT-box* cis-regulatory elements at the promoters of genes coregulated by AtBMI1 and VAL1/2 proteins. An enrichment of *LEC2/ABI3/VP1* and *ABRE BINDING FACTOR1* elements has been previously reported at the promoter of genes up-regulated in *atbmi1a/b* weak phenotype (Wang et al., 2016). Genes coregulated by ABI3/VP1-like proteins and ABA contain these motifs at their promoters (Suzuki et al., 2005). Accordingly, ABI3 and ABI5 regulate gene expression synergistically. Moreover, ABI3 interacts physically with ABI5, thereby ABI3 is also recruited to the promoters of the target genes via protein-protein interaction (Nakamura et al., 2001). A similar mechanism could be assumed for repression in which the VAL1/2 proteins bind to *LEC2/ABI3/VP1* and ASIL1 to the *GT-box* element, resulting in a direct competition with the transcriptional activators. The binding of VAL1/2 and possibly ASIL1 proteins could recruit the AtBMIs and the other PcG proteins to establish chromatin modifications that maintain gene repression. Whether ASIL1-mediated repression involves in vivo interaction with VAL and/or PcG proteins

remains to be investigated; however, in support of this, it has been shown that EMF1 interacts with ASIL1 (named EIP7) in yeast two hybrid experiments (Park et al., 2011).

We also found an enrichment of *LEC2/ABI3/VP1* elements, but not *ABRE* or *GT-box* elements, at the promoter of genes exclusively up_K27 in *atbmi1a/b/c* mutants, suggesting an implication of B3 factors in the regulation of these genes as well. Interestingly, two VAL1 splice variants have been identified through RNA sequencing analysis: a full-length form and a truncated form lacking the plant homeodomain-like domain similar to VAL3, which also lacks the plant homeodomain-like domain (Schneider et al., 2016). It is possible that truncated VAL1 and VAL3 target this group of genes, explaining their lack of up-regulation in *val1/2* mutants. Alternatively, since the B3 superfamily encompasses other subfamilies, such as the AUXIN RESPONSE FACTORS, the RELATED ABI3/VP1 and REPRODUCTIVE MERISTEM subfamilies (Swaminathan et al., 2008), some uncharacterized members of these might bind the *LEC2/ABI3/VP1* element or a variation of it. In any case, the promoters of the VAL1/2-independent genes are also enriched in other cis-regulatory elements such as

ALFIN1 motifs that are recognized in Arabidopsis by the ALs. Since the AL proteins interact with AtBMI1 proteins (Molitor et al., 2014), it is likely that a combination of B3 and AL factors participates in the regulation of a subset of these genes.

The relationship between AtBMI1 and EMF1 has been controversial. On one side, mutants in both display a very different phenotype and misexpress different subsets of PRC2 targets (Kim et al., 2010; Pu et al., 2013; Yang et al., 2013), which has led to propose the existence of PRC1 variants (Calonje, 2014; Merini and Calonje, 2015); however, they also coregulated a subset of targets (e.g. *ABI3*, *ABI4*, *FLC*) and in vitro they interact. Recent reports have shown that EMF1 copurifies with PRC2 components (Liang et al., 2015), questioning its exclusive association with PRC1. However, EMF1 colocalizes with only 45% of H3K27me3 marked genes showing a more narrow distribution at target genes than H3K27me3 marks (Kim et al., 2012). Another putative PRC1 component, LHP1, which broadly distributes across H3K27me3 marked sites (Turck et al., 2007; Zhang et al., 2007; Engelhorn et al., 2012), also copurifies with PRC2 (Derkacheva et al., 2013; Liang et al., 2015) and interacts with AtBMI1 and AtRING1 proteins in vitro (Xu and Shen, 2008; Bratzel et al., 2010). However, neither EMF1 nor LHP1 seem to be PRC2 core components, since they are required for H3K27me3 marking of only a subset of PRC2 targets (Kim et al., 2012; Wang et al., 2016).

Interestingly, when we compared the H3K27me3 marked genes that were up-regulated in *atbmi1a/b/c* with K27_EMF1 direct targets, we found that 50% of the up-regulated genes in *atbmi1* mutants were also EMF1 targets, suggesting that AtBMI1 and EMF1 could be in a complex and potentially both impact the expression of these genes. Since LHP1 is at 93.9% of genes up_K27 in *atbmi1a/b/c* mutants, the same holds true also for this PRC1 component. However, the little overlap between the genes up-regulated in *atbmi1a/b/c* and *emf1-2* or *lhp1* suggests a decisive role of AtBMI1 function in maintaining their repression. There were also genes exclusively up-regulated in *emf1-2* or *lhp1*, the majority of which are involved in flower development and these genes were not up-regulated in *atbmi1a/b/c* mutants. An interesting possibility could be that a PcG mechanism dependent on EMF1, LHP1, and PRC2 activities has evolved to specifically regulate the flower developmental program, which is consistent with the finding of these proteins copurifying with PRC2 (Liang et al., 2015).

CONCLUSION

In summary, our data point to different PRC1 functional networks in which genes may be regulated by AtBMI1 and/or EMF1 together with LHP1 and PRC2 and that additional proteins are required to regulate distinct subsets of genes. This is the case of VAL1/2 proteins in the seed development program, which built a network that apparently also includes *ABRE/GT-box*

binding factors (Fig. 7C). Furthermore, it seems highly likely that other B3 domain transcription factors and ALs are part of AtBMI1-repressive circuits. In contrast, there seems little or no overlap in gene regulation by AtBMI1 on the one side and EMF1 and LHP1 on the other, although these factors may physically interact and be simultaneously present at target genes.

MATERIALS AND METHODS

Plant Material and Growth Conditions

Arabidopsis (*Arabidopsis thaliana*) *atbmi1a* (N645041 line), *atbmi1b* (CS855837 line), *atbmi1a/b*, and *atbmi1a/b/c* (*atbmi1c* is a GT21221.Ds5.09.01.2006.jz07.348 line) mutants were described previously (Bratzel et al., 2010; Yang et al., 2013). Segregation of “weak” and “strong” *atbmi1a/b* phenotypes has been previously shown (Bratzel et al., 2010; Picó et al., 2015). Plants were grown under long-day conditions at 21°C on Murashige and Skoog agar plates containing 1.5% Suc and 0.8% agar. Seedling samples were collected at zeitgeber time 2.

Transcriptomic Analysis by RNA Sequencing

The experimental design in our study consisted of two replicates for each genotype (wild-type Col-0, *atbmi1a*, *atbmi1b*, *atbmi1a/b*, and *atbmi1a/b/c*). RNA extraction was performed using Qiagen RNeasy mini kit, following the manufacturer's instructions. RNA concentration and purity was tested using nanodrop-photometric quantification (Thermo Scientific). Library preparation was carried out following the manufacturer's recommendations (TruSeq RNA Sample Prep Kit v2, Illumina). Sequencing of RNA libraries was performed with the Illumina HiSeq2000 sequencer, yielding an average of approximately 15 million 100 bp long paired-end reads for each sample. The software package FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) was used for quality control. All sequencing samples were of high quality, and no pre-processing of the reads was required to remove low-quality reads or read fragments (Supplemental Fig. S2). The Arabidopsis Col-0 reference genome and annotation were downloaded from the Phytozome database (TAIR10; Goodstein et al., 2012). Mapping of reads to the reference genome, transcript assembly, and differential expression were performed with the software tools Bowtie, TopHat, and Cufflinks (Trapnell et al., 2012) using default parameters producing a high percentage of concordant pair alignment rate (Supplemental Table S1). The R package from Bioconductor CummeRbund (<http://www.bioconductor.org/>) was used for subsequent analysis and graphical representation of the results. Differentially expressed genes were selected as those exhibiting an expression fold change greater than four when compared with the wild type and a *P* value < 0.05. Venn diagrams comparing the different sets of differentially expressed genes were generated with Venny 2.0.2 (<http://bioinfogp.cnb.csic.es/tools/venny/index.html>) and the significance of their intersections with H3K27me3 marked genes was performed using Fisher's exact test. Gene ontology term enrichment was performed over the sets of differentially expressed genes with the web-based tools AgriGO and ReViGO (Supek et al., 2011; Yu et al., 2012) and the R bioconductor package ClusterProfiler (Du et al., 2010) using Singular Enrichment Analysis.

The clustering analysis was performed using the hierarchical algorithm implemented in the R package cluster over normalized expression levels measured using FPKM.

qRT-PCR

For qRT-PCR analysis, cDNAs were reverse-transcribed from total RNAs with QuantiTect reverse transcription kit (Qiagen). qRT-PCRs were performed using Sensi FAST SYBR & Fluorescein kit (BioLine) and an iQ5 Bio-Rad system. Expression was calculated relative to *ACTIN*. Primers used were as follows: *WOX9-RT-Fw* (5' ACTGTCGAGGGTTGAAGGTATC 3'); *WOX9-RT-Rev* (5' AGTGGTAGCGTAACAAATCTGAGTCT 3'); *WOX2-RT-Fw* (5' GCTTACTTCAATCGCTCCTCCACAA 3'); *WOX2-RT-Rev* (5' GTCGGTTTCTCGTAGG-CACCACTTG 3'); *SMB-RT-Fw* (5' ACGAATATCGCTTGGACGATAG 3'); *SMB-RT-Rev* (5' GCCTCTGTCTTGGTGAATCC 3'); *ACT2-RT-Fw* (5' CACTTGCACCAAGCAGCATGAAGA 3'); *ACT2-RT-Rev* (5' AATGGAACCACCGATC-CAGACACT 3').

Motif and Transcription Factor Binding Site Enrichment Analysis

Transcription Factor Binding Sites enrichment analysis was performed using HOMER (Heinz et al., 2010) and the known Transcription Factor Binding Sites sequences in plants from the databases AGRIS (Davuluri et al., 2003), JASPAR (Sandelin et al., 2004), and AthaMap (Steffens et al., 2004). The findMotifs.pl script was used with default parameters to perform known and de-novo motif over-representation analysis for DNA sequences of 6-, 7-, 8-, and 9-bp lengths. The target set consisted of all the gene promoters of interest. The background used for the overrepresentation analysis consisted of all the gene promoters annotated in the Arabidopsis TAIR10 genome. For the co-occurrence of the *ABRE* and *GT-box* motifs, we first identified the locations of the *ABRE* motif at the promoters and then extracted the DNA sequences 100 bp upstream and downstream from the center of the *ABRE* motif. We performed an enrichment analysis of the *GT-box* motif in these DNA sequences using the findMotifsGenome.pl HOMER script with default parameters. The significance of the overlapping between motifs was performed as an enrichment analysis of the DNA sequence resulting from the combination of both motifs. DNA sequences used in these analyses were downloaded using the BioMart functionality associated with Phytozome (Goodstein et al., 2012). Gene promoters were defined as the 1,000-bp DNA sequence upstream of the start codon of the corresponding gene.

Accession Numbers

The RNA-seq raw data generated in this study are publicly available from the GEO database identified with accession number GSE83568 (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?&acc=GSE83568>).

Supplemental Data

The following supplemental materials are available.

Supplemental Figure S1. Phenotypes of *atbmi1a/b* and *atbmi1a/b/c* mutants.

Supplemental Figure S2. Boxplots representing the read quality scores (Illumina 1.5 encoding) per base for the first replicate of all samples.

Supplemental Figure S3. Correlation among differentially expressed genes in wild type and the different genotypes.

Supplemental Figure S4. Altered gene expression in *atbmi1* mutants.

Supplemental Figure S5. Gene ontology (GO) enrichment analysis of up- and down-regulated genes in *atbmi1a/b* mutants.

Supplemental Figure S6. Gene ontology (GO) enrichment analysis of up- and down-regulated genes in *atbmi1a/b/c* mutants.

Supplemental Figure S7. Putative AtBMI1direct target genes.

Supplemental Figure S8. Genes differentially expressed in *atbmi1a/b* and *atbmi1a/b/c*.

Supplemental Figure S9. Different gene expression patterns of *atbmi1a/b* weak and *atbmi1a/b/c* mutants.

Supplemental Figure S10. Expression levels of different important developmental genes in wild type and *atbmi1a/b/c* mutants.

Supplemental Figure S11. AtBMI1, EMF1, and LHP1 functional relationship.

Supplemental Figure S12. *AtBMI1B* (*At1g06770*) splice variants.

Supplemental Table S1. Number of reads and concurrent pair alignment rate per sequencing sample.

Supplemental Table S2. Up- and down-regulated genes in *atbmi1* mutants.

Supplemental Table S3. Up-regulated genes in *atbmi1* and *val1/2* mutants that are marked with H3K27me3 marks in wild type, and genes in clusters I, II, and III after comparing genes up_K27 in *atbmi1a/b* and *atbmi1a/b/c*.

Supplemental Table S4. Up-regulated genes in *emf1-2* and *lhp1* mutants that are marked with H3K27me3 marks in wild type.

LITERATURE CITED

- Beh LY, Colwell LJ, Francis NJ (2012) A core subunit of Polycomb repressive complex 1 is broadly conserved in function but not primary sequence. *Proc Natl Acad Sci USA* **109**: E1063–E1071
- Bouyer D, Roudier F, Heese M, Andersen ED, Gey D, Nowack MK, Goodrich J, Renou J-P, Grini PE, Colot V, et al (2011) Polycomb repressive complex 2 controls the embryo-to-seedling phase transition. *PLoS Genet* **7**: e1002014
- Bratzel F, López-Torrejón G, Koch M, Del Pozo JC, Calonje M (2010) Keeping cell identity in Arabidopsis requires PRC1 RING-finger homologs that catalyze H2A monoubiquitination. *Curr Biol* **20**: 1853–1859
- Bratzel F, Yang C, Angelova A, López-Torrejón G, Koch M, del Pozo JC, Calonje M (2012) Regulation of the new Arabidopsis imprinted gene AtBMI1C requires the interplay of different epigenetic mechanisms. *Mol Plant* **5**: 260–269
- Calonje M (2014) PRC1 marks the difference in plant PcG repression. *Mol Plant* **7**: 459–471
- Calonje M, Sanchez R, Chen L, Sung ZR (2008) EMBRYONIC FLOWER1 participates in polycomb group-mediated AG gene silencing in Arabidopsis. *Plant Cell* **20**: 277–291
- Cao R, Tsukada Y, Zhang Y (2005) Role of Bmi-1 and Ring1A in H2A ubiquitylation and Hox gene silencing. *Mol Cell* **20**: 845–854
- Carles C, Bies-Etheve N, Aspart L, Léon-Kloosterziel KM, Koornneef M, Echeverria M, Delseny M (2002) Regulation of Arabidopsis thaliana Em genes: role of ABI5. *Plant J* **30**: 373–383
- Chanvavattana Y, Bishopp A, Schubert D, Stock C, Moon Y-H, Sung ZR, Goodrich J (2004) Interaction of Polycomb-group proteins controlling flowering in Arabidopsis. *Development* **131**: 5263–5276
- Chen D, Molitor A, Liu C, Shen W-H (2010) The Arabidopsis PRC1-like ring-finger proteins are necessary for repression of embryonic traits during vegetative growth. *Cell Res* **20**: 1332–1344
- Choi H, Hong J, Ha J, Kang J, Kim SY (2000) ABFs, a family of ABA-responsive element binding factors. *J Biol Chem* **275**: 1723–1730
- Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR (2010) Abscisic acid: emergence of a core signaling network. *Annu Rev Plant Biol* **61**: 651–679
- Davuluri RV, Sun H, Palaniswamy SK, Matthews N, Molina C, Kurtz M, Grotewold E (2003) AGRIS: Arabidopsis gene regulatory information server, an information resource of Arabidopsis cis-regulatory elements and transcription factors. *BMC Bioinformatics* **4**: 25
- Delmas F, Sankaranarayanan S, Deb S, Widdup E, Bourmonville C, Bollier N, Northey JGB, McCourt P, Samuel MA (2013) ABI3 controls embryo degreening through Mendel's I locus. *Proc Natl Acad Sci USA* **110**: E3888–E3894
- Derkacheva M, Steinbach Y, Wildhaber T, Mozgová I, Mahrez W, Nanni P, Bischof S, Gruissem W, Hennig L (2013) Arabidopsis MSI1 connects LHP1 to PRC2 complexes. *EMBO J* **32**: 2073–2085
- Du Z, Zhou X, Ling Y, Zhang Z, Su Z (2010) agriGO: a GO analysis toolkit for the agricultural community. *Nucleic Acids Res* **38**: W64–70
- Engelhorn J, Reimer JJ, Leuz I, Göbel U, Huettel B, Farrona S, Turck F (2012) Development-related PcG target in the apex 4 controls leaf margin architecture in Arabidopsis thaliana. *Development* **139**: 2566–2575
- Farrona S, Thorpe FL, Engelhorn J, Adrian J, Dong X, Sarid-Krebs L, Goodrich J, Turck F (2011) Tissue-specific expression of FLOWERING LOCUS T in Arabidopsis is maintained independently of polycomb group protein repression. *Plant Cell* **23**: 3204–3214
- Fernandez DE, Wang C-T, Zheng Y, Adamczyk BJ, Singhal R, Hall PK, Perry SE (2014) The MADS-domain factors AGAMOUS-LIKE15 and AGAMOUS-LIKE18, along with SHORT VEGETATIVE PHASE and AGAMOUS-LIKE24, are necessary to block floral gene expression during the vegetative phase. *Plant Physiol* **165**: 1591–1603
- Förderer A, Zhou Y, Turck F (2016) The age of multiplexity: recruitment and interactions of Polycomb complexes in plants. *Curr Opin Plant Biol* **29**: 169–178
- Francis NJ, Kingston RE, Woodcock CL (2004) Chromatin compaction by a polycomb group protein complex. *Science* **306**: 1574–1577
- Gangappa SN, Srivastava AK, Maurya JP, Ram H, Chattopadhyay S (2013) Z-box binding transcription factors (ZBFs): a new class of transcription factors in Arabidopsis seedling development. *Mol Plant* **6**: 1758–1768
- Gao M-J, Lydiate DJ, Li X, Lui H, Gjetvaj B, Hegedus DD, Rozwadowski K (2009) Repression of seed maturation genes by a trihelix transcriptional repressor in Arabidopsis seedlings. *Plant Cell* **21**: 54–71

- Gaudin V, Libault M, Pouteau S, Juul T, Zhao G, Lefebvre D, Grandjean O (2001) Mutations in LIKE HETEROCHROMATIN PROTEIN 1 affect flowering time and plant architecture in Arabidopsis. *Development* **128**: 4847–4858
- Gendall AR, Levy YY, Wilson A, Dean C (2001) The VERNALIZATION 2 gene mediates the epigenetic regulation of vernalization in Arabidopsis. *Cell* **107**: 525–535
- Goodrich J, Puangsomlee P, Martin M, Long D, Meyerowitz EM, Coupland G (1997) A Polycomb-group gene regulates homeotic gene expression in Arabidopsis. *Nature* **386**: 44–51
- Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U, Putnam N, et al (2012) Phytozome: a comparative platform for green plant genomics. *Nucleic Acids Res* **40**: D1178–D1186
- Grossniklaus U, Vielle-Calzada JP, Hoepfner MA, Gagliano WB (1998) Maternal control of embryogenesis by MEDEA, a polycomb group gene in Arabidopsis. *Science* **280**: 446–450
- Gu X, Le C, Wang Y, Li Z, Jiang D, Wang Y, He Y (2013) Arabidopsis FLC clade members form flowering-repressor complexes coordinating responses to endogenous and environmental cues. *Nat Commun* **4**: 1947
- Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK (2010) Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**: 576–589
- Hennig L, Bouveret R, Gruissem W (2005) MSI1-like proteins: an escort service for chromatin assembly and remodeling complexes. *Trends Cell Biol* **15**: 295–302
- Horstman A, Willemsen V, Boutilier K, Heidstra R (2014) AINTEGUMENTA-LIKE proteins: hubs in a plethora of networks. *Trends Plant Sci* **19**: 146–157
- Kim SY, Lee J, Eshed-Williams L, Zilberman D, Sung ZR (2012) EMF1 and PRC2 cooperate to repress key regulators of Arabidopsis development. *PLoS Genet* **8**: e1002512
- Kim SY, Zhu T, Sung ZR (2010) Epigenetic regulation of gene programs by EMF1 and EMF2 in Arabidopsis. *Plant Physiol* **152**: 516–528
- Lee WY, Lee D, Chung W-I, Kwon CS (2009) Arabidopsis ING and Alfin1-like protein families localize to the nucleus and bind to H3K4me3/2 via plant homeodomain fingers. *Plant J* **58**: 511–524
- Liang SC, Hartwig B, Perera P, Mora-García S, de Leau E, Thornton H, de Lima Alves F, Rappsilber J, Yang S, James GV, et al (2015) Kicking against the PRCs - a domesticated transposase antagonises silencing mediated by polycomb group proteins and is an accessory component of polycomb repressive complex 2. *PLoS Genet* **11**: e1005660
- Luo M, Bilodeau P, Koltunow A, Dennis ES, Peacock WJ, Chaudhury AM (1999) Genes controlling fertilization-independent seed development in Arabidopsis thaliana. *Proc Natl Acad Sci USA* **96**: 296–301
- Merini W, Calonje M (2015) PRC1 is taking the lead in PcG repression. *Plant J* **83**: 110–120
- Molitor AM, Bu Z, Yu Y, Shen W-H (2014) Arabidopsis AL PHD-PRC1 complexes promote seed germination through H3K4me3-to-H3K27me3 chromatin state switch in repression of seed developmental genes. *PLoS Genet* **10**: e1004091
- Mozgova I, Köhler C, Hennig L (2015) Keeping the gate closed: functions of the polycomb repressive complex PRC2 in development. *Plant J* **83**: 121–132
- Müller J, Hart CM, Francis NJ, Vargas ML, Sengupta A, Wild B, Miller EL, O'Connor MB, Kingston RE, Simon JA (2002) Histone methyltransferase activity of a Drosophila Polycomb group repressor complex. *Cell* **111**: 197–208
- Nakamura S, Lynch TJ, Finkelstein RR (2001) Physical interactions between ABA response loci of Arabidopsis. *Plant J* **26**: 627–635
- Ohad N, Yadegari R, Margossian L, Hannon M, Michaeli D, Harada JJ, Goldberg RB, Fischer RL (1999) Mutations in FIE, a WD polycomb group gene, allow endosperm development without fertilization. *Plant Cell* **11**: 407–416
- Park H-Y, Lee S-Y, Seok H-Y, Kim S-H, Sung ZR, Moon Y-H (2011) EMF1 interacts with EIP1, EIP6 or EIP9 involved in the regulation of flowering time in Arabidopsis. *Plant Cell Physiol* **52**: 1376–1388
- Peterson AJ, Mallin DR, Francis NJ, Ketel CS, Stamm J, Voeller RK, Kingston RE, Simon JA (2004) Requirement for sex comb on midleg protein interactions in Drosophila polycomb group repression. *Genetics* **167**: 1225–1239
- Picó S, Ortiz-Marchena MI, Merini W, Calonje M (2015) Deciphering the role of Polycomb Repressive Complex 1 (PRC1) variants in regulating the acquisition of flowering competence in Arabidopsis. *Plant Physiol* **168**: 1286–1297
- Pu L, Liu M-S, Kim SY, Chen L-FO, Fletcher JC, Sung ZR (2013) EMBRYONIC FLOWER1 and ULTRAPETALA1 act antagonistically on Arabidopsis development and stress response. *Plant Physiol* **162**: 812–830
- Qüesta JI, Song J, Geraldo N, An H, Dean C (2016) Arabidopsis transcriptional repressor VAL1 triggers Polycomb silencing at FLC during vernalization. *Science* **353**: 485–488
- Ringrose L, Paro R (2004) Epigenetic regulation of cellular memory by the Polycomb and Trithorax group proteins. *Annu Rev Genet* **38**: 413–443
- Sanchez-Pulido L, Devos D, Sung ZR, Calonje M (2008) RAWUL: a new ubiquitin-like domain in PRC1 ring finger proteins that unveils putative plant and worm PRC1 orthologs. *BMC Genomics* **9**: 308
- Sandelin A, Alkema W, Engström P, Wasserman WW, Lenhard B (2004) JASPAR: an open-access database for eukaryotic transcription factor binding profiles. *Nucleic Acids Res* **32**: D91–D94
- Santos-Mendoza M, Dubreucq B, Baud S, Parcy F, Caboche M, Lepiniec L (2008) Deciphering gene regulatory networks that control seed development and maturation in Arabidopsis. *Plant J* **54**: 608–620
- Schneider A, Aghmirzaie D, Elmarakeby H, Poudel AN, Koo AJ, Heath LS, Grene R, Collakova E (2016) Potential targets of VIVIPAROUS1/ABI3-LIKE1 (VAL1) repression in developing Arabidopsis thaliana embryos. *Plant J* **85**: 305–319
- Schwartz YB, Pirrotta V (2013) A new world of Polycombs: unexpected partnerships and emerging functions. *Nat Rev Genet* **14**: 853–864
- Shao Z, Raible F, Mollaaghababa R, Guyon JR, Wu CT, Bender W, Kingston RE (1999) Stabilization of chromatin structure by PRC1, a Polycomb complex. *Cell* **98**: 37–46
- Shu K, Liu XD, Xie Q, He ZH (2016) Two faces of one seed: hormonal regulation of dormancy and germination. *Mol Plant* **9**: 34–45
- Simon JA, Kingston RE (2013) Occupying chromatin: polycomb mechanisms for getting to genomic targets, stopping transcriptional traffic, and staying put. *Mol Cell* **49**: 808–824
- Steffens NO, Galuschka C, Schindler M, Bülow L, Hehl R (2004) Atha-Map: an online resource for in silico transcription factor binding sites in the Arabidopsis thaliana genome. *Nucleic Acids Res* **32**: D368–D372
- Supek F, Bošnjak M, Škunca N, Šmuc T (2011) REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One* **6**: e21800
- Suzuki M, Ketterling MG, McCarty DR (2005) Quantitative statistical analysis of cis-regulatory sequences in ABA/VP1- and CBF/DREB1-regulated genes of Arabidopsis. *Plant Physiol* **139**: 437–447
- Suzuki M, Wang HH-Y, McCarty DR (2007) Repression of the LEAFY COTYLEDON 1/B3 regulatory network in plant embryo development by VP1/ABSCISIC ACID INSENSITIVE 3-LIKE B3 genes. *Plant Physiol* **143**: 902–911
- Swaminathan K, Peterson K, Jack T (2008) The plant B3 superfamily. *Trends Plant Sci* **13**: 647–655
- Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L (2012) Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* **7**: 562–578
- Turck F, Roudier F, Farrona S, Martin-Magniette M-L, Guillaume E, Buisine N, Gagnot S, Martienssen RA, Coupland G, Colot V (2007) Arabidopsis TFL2/LHP1 specifically associates with genes marked by trimethylation of histone H3 lysine 27. *PLoS Genet* **3**: e86
- van der Graaff E, Laux T, Rensing SA (2009) The WUS homeobox-containing (WOX) protein family. *Genome Biol* **10**: 248
- Veluchamy A, Jégu T, Ariel F, Latrasse D, Mariappan KG, Kim S-K, Crespi M, Hirt H, Bergounioux C, Raynaud C, et al (2016) LHP1 regulates H3K27me3 spreading and shapes the three-dimensional conformation of the Arabidopsis genome. *PLoS One* **11**: e0158936
- Vicente-Carbajosa J, Carbonero P (2005) Seed maturation: developing an intrusive phase to accomplish a quiescent state. *Int J Dev Biol* **49**: 645–651
- Wang H, Liu C, Cheng J, Liu J, Zhang L, He C, Shen W-H, Jin H, Xu L, Zhang Y (2016) Arabidopsis flower and embryo developmental genes are repressed in seedlings by different combinations of polycomb group proteins in association with distinct sets of cis-regulatory elements. *PLoS Genet* **12**: e1005771
- Wang Y, Gu X, Yuan W, Schmitz RJ, He Y (2014) Photoperiodic control of the floral transition through a distinct polycomb repressive complex. *Dev Cell* **28**: 727–736

- Wei W, Zhang Y-Q, Tao J-J, Chen H-W, Li Q-T, Zhang W-K, Ma B, Lin Q, Zhang J-S, Chen S-Y** (2015) The Alfin-like homeodomain finger protein AL5 suppresses multiple negative factors to confer abiotic stress tolerance in Arabidopsis. *Plant J* **81**: 871–883
- Wolff P, Weinhofer I, Seguin J, Roszak P, Beisel C, Donoghue MTA, Spillane C, Nordborg M, Rehmsmeier M, Köhler C** (2011) High-resolution analysis of parent-of-origin allelic expression in the Arabidopsis Endosperm. *PLoS Genet* **7**: e1002126
- Wu G, Poethig RS** (2006) Temporal regulation of shoot development in Arabidopsis thaliana by miR156 and its target SPL3. *Development* **133**: 3539–3547
- Xiao J, Wagner D** (2015) Polycomb repression in the regulation of growth and development in Arabidopsis. *Curr Opin Plant Biol* **23**: 15–24
- Xu L, Shen W-H** (2008) Polycomb silencing of KNOX genes confines shoot stem cell niches in Arabidopsis. *Curr Biol* **18**: 1966–1971
- Yang C, Bratzel F, Hohmann N, Koch M, Turck F, Calonje M** (2013) VAL- and AtBMI1-mediated H2Aub initiate the switch from embryonic to post-germinative growth in Arabidopsis. *Curr Biol* **23**: 1324–1329
- Yoshida N, Yanai Y, Chen L, Kato Y, Hiratsuka J, Miwa T, Sung ZR, Takahashi S** (2001) EMBRYONIC FLOWER2, a novel polycomb group protein homolog, mediates shoot development and flowering in Arabidopsis. *Plant Cell* **13**: 2471–2481
- Yu G, Wang L-G, Han Y, He Q-Y** (2012) clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* **16**: 284–287
- Zhang X, Germann S, Blus BJ, Khorasanizadeh S, Gaudin V, Jacobsen SE** (2007) The Arabidopsis LHP1 protein colocalizes with histone H3 Lys27 trimethylation. *Nat Struct Mol Biol* **14**: 869–871