



Arnold, D. T., Attwood, M., Barratt, S., Morley, A., Elvers, K. T., McKernon, J., Donald, C., Oates, A., Noel, A., MacGowan, A., Maskell, N. A., & Hamilton, F. W. (2021). Predicting outcomes of COVID-19 from admission biomarkers: a prospective UK cohort study. *Emergency Medicine Journal*, 38(7), 543-548.
<https://doi.org/10.1136/emermed-2020-210380>

Peer reviewed version

License (if available):
CC BY-NC

Link to published version (if available):
[10.1136/emermed-2020-210380](https://doi.org/10.1136/emermed-2020-210380)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via BMJ Publishing Group at [10.1136/emermed-2020-210380](https://doi.org/10.1136/emermed-2020-210380). Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Predicting outcomes of COVID-19 from admission biomarkers; a prospective UK cohort study.

Arnold DT ¹, Attwood M ², Barratt S ¹, Morley A ¹, Elvers KT ³, McKernon J ⁴, Donald C ⁵, Oates A ⁵, Noel A ², MacGowan A ², Maskell NA ¹, Hamilton F ²

1. Academic Respiratory Unit, University of Bristol, Southmead Way, Bristol, BS105NB
2. Bristol Centre for Antimicrobial Research (BCARE), North Bristol NHS Trust, Southmead Way, Bristol, BS105NB
3. Medicines Discovery Institute Cardiff, Cardiff University, Park Place, Cardiff, CF10 3AT
4. Immunology, North Bristol NHS Trust, Southmead Way, Bristol, BS105NB
5. Biochemistry, North Bristol NHS Trust, Southmead Way, Bristol, BS105NB

Corresponding author: Dr David Arnold

david.arnold@nbt.nhs.uk

Academic Respiratory Unit
Learning and Research Centre,
Southmead Hospital
Southmead Way
BS10 5NB

Word Count: 2,931

Keywords: infectious diseases, SARS, respiratory, pneumonia/infections

Key messages

What is known on this subject?

Biomarkers that predict the clinical course of COVID-19 would be invaluable at the point of hospital admission. Several blood-based biomarkers have been suggested but have not been tested in a prospective UK cohort.

What this study adds?

In this prospective UK cohort, routinely performed blood biomarkers at the point of hospital admission had limited ability, in isolation, to predict poor outcomes from COVID-19. Biomarkers of immune activation had the best predictive performance alongside routinely collected clinical information.

ABSTRACT:

Introduction:

COVID-19 has an unpredictable clinical course so prognostic biomarkers would be invaluable when triaging patients on admission to hospital. Many biomarkers have been suggested using large observational datasets but sample timing is crucial to ensure prognostic relevance. The DISCOVER study prospectively recruited patients with COVID-19 admitted to a UK hospital and analysed a panel of putative prognostic biomarkers on the admission blood sample to identify markers of poor outcome.

Methods:

Consecutive patients admitted to hospital with proven or clinoradiological suspected COVID-19 were consented. Admission bloods were extracted from the clinical laboratory. A panel of biomarkers (IL-6, suPAR, KL-6, Troponin, Ferritin, LDH, BNP, Procalcitonin) were performed in addition to routinely performed markers (CRP, neutrophils, lymphocytes, neutrophil:lymphocyte ratio). Age, NEWS2, CURB-65 and radiographic severity score on initial chest radiograph were included as comparators. All biomarkers were tested in logistic regression against a composite outcome of non-invasive ventilation, intensive care admission, or death, with Area Under the Curve (AUC) figures calculated.

Results:

187 patients had 28-day outcomes at the time of analysis. CRP (AUC 0.69 ,95%CI:0.59-0.78), lymphocyte count (AUC 0.62, 95%CI:0.53-0.72), and other routine markers did not predict the primary outcome. IL-6 (AUC: 0.77,0.65-0.88) and suPAR (AUC 0.81,Ci: 0.72-0.88) showed some promise, but simple clinical features alone such as NEWS2 score (AUC: 0.70 ,0.60-0.79) or age (AUC: 0.70 ,0.62-0.77) performed nearly as well.

Discussion:

Admission blood biomarkers have only moderate predictive value for predicting COVID-19 outcomes, while simple clinical features such as age and NEWS2 score outperform many biomarkers. IL-6 and suPAR had the best performance, and further studies should focus on the additive value of these biomarkers to routine care.

INTRODUCTION:

COVID-19 causes a wide spectrum of disease, from asymptomatic to severe respiratory failure. The majority of patients who present to hospital will recover but some develop rapidly progressive respiratory compromise requiring ventilatory support. Biomarkers that might predict this deterioration would be invaluable when triaging patients on hospital admission to inform who can be safely discharged versus those who need careful respiratory monitoring.

The rapidly emerging literature base around biomarkers in COVID has mainly extracted data from electronic health records of large cohort studies.¹⁻⁴ Far fewer have prospectively recruited and followed up patients. Those that have tested novel biomarkers have often done so at later timepoints in patients limiting extrapolation to hospital admission.⁵⁻⁷ Few UK studies have assessed the additional value of biomarkers compared to the routinely recorded demographic information and clinical scales (such as National Early Warning Scores or CURB-65).

The Diagnostic and Severity markers of COVID-19 to Enable Rapid triage (DISCOVER) study prospectively recruited patients admitted with COVID-19 to a single UK hospital, with the aim of comparing the ability of various biomarkers and clinical scores to predict mortality, need for non-invasive ventilation or intensive care unit admission.

METHODS:

This study is reported in line with the STROBE guidelines for observational studies (see Supplemental Material)

Study design

This was a prospective cohort study aimed to assess prognostic clinical and blood biomarkers for COVID-19 disease based on the earliest available clinical and biochemical information.

Outcomes

The primary outcome of prediction was a composite outcome of intensive care admission, non-invasive ventilation (NIV) with Continuous Positive Airway Pressure (CPAP) or Bilevel Positive Airway Pressure (BIPAP) outside the intensive care unit, or death (defined below as “severe disease”).

The secondary outcome was a composite outcome of intensive care admission and death.

Patient recruitment

All patients were recruited via the DISCOVER study, a single centre observational study at North Bristol NHS Trust recruiting patients with COVID-19, from 30.03.2020 until 29.06.2020 (Ethics approval via South Yorks REC: 20/YH/0121, CRN approval no: 45469). Patients were recruited on the basis of a positive PCR result for SARS-CoV-2, using the established PHE assay in use at the time, or a clinicoradiological diagnosis of COVID-19 disease. During the pandemic, community testing became widely available, although the results were not available to hospital staff. As such, later patients were often recruited on the basis of a history of positive testing in the community. The only exclusion criteria was an inability to consent. For patients in intensive care, family members were able to consent on behalf of them if too unwell to consent as a personel consultee.

Clinical information

Clinical information was recorded on a REDCap (Vanderbilt University) database,⁸ by the consenting nurse or physician. Routine demographics were recorded, and presence of important comorbidities. Comorbidities were defined either by their recording in the admission notes / hospital record (for hypertension, heart disease, and chronic lung disease), the presence of a positive serological result (for HIV) or for requirement for dialysis or an estimated GFR of <30ml/min (for chronic kidney disease). Ethnicity was also recorded.

The earliest admission National Early Warning Score (NEWS2) was extracted from the clinical record. This is a numeric score (from 0-20), reflecting the degree of physiological dysfunction.⁹ Higher numbers indicate more severe physiological dysfunction.. Routine biochemistry and haematology results were extracted from the clinical record, using the earliest available figure. Outcome data recorded was in line with RECOVERY, the national RCT of therapeutic interventions for COVID-19 (<https://www.recoverytrial.net/>). This included 28-day mortality, requirement for intensive care, ventilation, renal replacement therapy, and inotropes.

Non-invasive ventilation was defined by the receipt of non-invasive ventilation by either continuous positive airway pressure (CPAP), bi-level positive airway pressure (BIPAP), or HFNO any time during the hospital stay.

The radiological severity score was calculated using the method described by Wong et al, 2020.¹⁰ A score of 0-4 was assigned to each lung depending on the extent of involvement by consolidation or ground glass opacities. 0 = no involvement, 1 = <25%, 2 = 25 - 49%, 3= 50 - 75%, 4 = >75% involvement. The scores for each lung were summed to produce a final severity score ranging from 0-8. Radiographs were scored by a respiratory and infectious diseases physician.

Biomarkers and samples

For conventional biomarkers of infection (namely C-reactive protein (CRP), components of the full blood count and routine renal function) data were prospectively recorded. The first result from that admission was taken or, for established inpatients, the result on the day of the diagnosis. These results were available to the clinician and therefore could impact on prognosis (i.e. clinicians were not blinded to these results)

For other potential predictive markers (Lactate dehydrogenase (LDH), procalcitonin (PCT) interleukin-6 (IL-6), Krebs von den Lungen 6 (KL-6), Ferritin, Troponin, B-type natriuretic peptide (NT-pro BNP), soluble urokinase plasminogen activator receptor (suPAR)), analysis was performed on frozen samples in batch analysis as described below. Laboratory staff were unaware of clinical outcome and were therefore functionally blinded.

The earliest initial sample was extracted from the blood sciences laboratory after routine testing had been performed; in admitted cases this was the initial sample taken in the Emergency Department (ED), in hospitalised cases, this was the sample from the day of diagnosis.

suPAR analysis was performed on the suPARnostic® ELISA platform, KL-6 and IL-6 were performed on the Fujerebio Lumipulse. CRP, Ferritin, LDH, Troponin, NT-pro-BNP and Procalcitonin were performed on the Roche Diagnostics cobas platform. The specific analytic platforms were spectrophotometric: c701 (CRP) and c501 (LDH), immunoassay: e501 (Ferritin, Troponin, NT-pro-BNP, Procalcitonin). The full blood count was performed on the Sysmex XN (Sysmex Diagnostics).

As the volume of blood required varied and was small, the total number of assays varied slightly as for some participants only enough blood was available to perform one or two assays, or there was technical failure and an inability to repeat due to lack of sample.

Statistical approach

In this study, we aimed to identify whether any individual biomarker (Lymphocyte count, Neutrophil count, Neutrophil:Lymphocyte ratio (NLR), CRP, IL-6, KL-6, suPAR, NT-pro BNP, LDH, PCT, Troponin T, Ferritin) had prognostic significance for the primary outcome as an individual marker, when used on the initial blood sample taken.

There was a deliberate focus on relatively simplistic (logistic regression) models: given the sample size, complex models would be at risk of overfitting, and the main target was to identify the additional value of a biomarker, all of which are reported on a linear scale.¹¹ Complete case analysis was performed for each individual biomarker.

ROC curves were generated, and AUC figures were calculated alongside sensitivity and specificity for each biomarker, at Youden's index. Confidence intervals were generated around the AUC by bootstrapping.

All analysis was performed in R (version 4.0.0), using the packages "tidyverse", "broom", "tidymodels", and "pROC". Analytic code is available at: https://github.com/gushamilton/discover_prediction/

Patient and Public Involvement

Given the difficulty of performing PPI during a pandemic, patients were not involved in the design, or conduct, or reporting, or dissemination plans of the DISCOVER study at the time this manuscript was submitted.

RESULTS:

187 participants had reached 28 days post diagnosis and were included in the analysis (prespecified timing of the primary outcome). Table 1 shows the demographics of the cohort. 101 patients (54%) were male, and the median age was 58 (IQR: 46-73). 76% of the cohort had positive PCR results for SARS-CoV-2, with the remaining clinically suspected.

Comorbidities were relatively common within the cohort with hypertension and diabetes being the most predominant. Patients with severe disease (death, intensive care admission or non-invasive ventilation) were generally older, and more comorbid than non-severe patients. 39 patients had the primary outcome. 15 patients went to the intensive care unit, of which 4 died and 11 survived. 7 patients required NIV outside the intensive care unit, 4 of whom died, see Figure 1.

Table 1: Demographics of the study cohort.

Characteristic	Survivor, N = 148	Non-survivor/ITU/NIV, N = 39	p-value ²
Age (18+)	55 (44, 71)	66 (60, 76)	<0.001
Sex			0.7
Male	79 (53%)	22 (56%)	
Female	69 (47%)	17 (44%)	
PCR proven disease			0.6
Proven	111 (75%)	31 (79%)	
Suspected	37 (25%)	8 (21%)	
Status at time of consent			0.010
Inpatient	119 (80%)	38 (97%)	
Outpatient	29 (20%)	1 (2.6%)	
Diabetes status			0.3
No	115 (78%)	34 (87%)	
T1DM	3 (2.0%)	1 (2.6%)	
T2DM	30 (20%)	4 (10%)	
Heart disease	34 (23%)	12 (31%)	0.3
Chronic Lung disease	26 (18%)	20 (51%)	<0.001
Severe Liver disease	3 (2.0%)	2 (5.1%)	0.3
Severe kidney impairment (eGFR< 30 or dialysis)	16 (11%)	6 (15%)	0.4
Hypertension	37 (25%)	15 (38%)	0.10
HIV status	1 (0.7%)	1 (2.6%)	0.4
Non-white ethnicity	20 (16%)	5 (15%)	0.9
Median (IQR); n (%) Wilcoxon rank sum test; Pearson's Chi-squared test; Fisher's exact test			

There was a significant variation in physiological state (e.g. NEWS2), with a median NEWS2 score of 4 (IQR 2-6), with the highest NEWS2 score recorded was 13. Patients that had severe disease had higher NEWS2 scores. There was also wide variation in functional status, with many patients having some degree of frailty, with frailer patients more likely to die or require enhanced care. Escalation status was recorded for most patients, with patients who died more likely to have limitations on ceiling of care, recorded in Table 2.

Table 2: Escalation status

Characteristic	Survivor, N = 148	Non-Survivor/ITU/NIV, N = 39
For full escalation to ITU including intubation and ventilation	104 (83%)	18 (55%)
Not for invasive ventilation (intubation & ventilation) but would be for non-invasive ventilation (NIV) or CPAP on ward.	8 (6.3%)	11 (33%)
Not for ITU or for NIV/CPAP	14 (11%)	3 (9.1%)
For palliative treatments only	0 (0%)	1 (3.0%)
Unknown	22	6
Statistics presented: n (%)		

Admission blood tests are recorded in Table 3. There was no significant difference for many blood tests, with lymphocyte count being slightly lower in those with severe disease, and neutrophil:lymphocyte ratio being slightly higher. Renal dysfunction was more common in the severe disease.

Table 3: Conventional blood tests

Characteristic	Survivor, N = 148	Non-Survivor/ITU/NIV, N = 39	p-value
Haemoglobin (g/L)	135 (126, 148)	134 (112, 147)	0.4
White cell count (x10 ⁹ /L)	7.6 (5.7, 9.9)	6.9 (5.0, 9.7)	0.7
Neutrophils (x10 ⁹ /L)	5.7 (3.9, 8.0)	5.8 (3.7, 8.4)	0.6
Lymphocytes (x10 ⁹ /L)	1.08 (0.68, 1.55)	0.86 (0.58, 1.16)	0.018
Platelets (x10 ⁹ /L)	239 (189, 289)	202 (149, 258)	0.009
Sodium (mmol/L)	138 (135, 140)	137 (136, 138)	0.4
Urea (mmol/L)	4.7 (3.6, 7.1)	7.0 (5.0, 9.4)	<0.001
eGFR (ml/min)	89 (65, 90)	64 (50, 86)	<0.001
Albumin (mmol/L)	33.5 (31.0, 36.8)	32.0 (27.5, 34.0)	0.009
CRP (mg/L)	44 (12, 86)	87 (54, 152)	<0.001
NLR	4.9 (2.8, 9.4)	6.3 (4.7, 12.3)	0.023
¹ Median (IQR) ² Wilcoxon rank sum test			

Logistic regression and AUC calculation

Table 4 shows biomarker performance to predict severe COVID-19 outcomes. Blood was not available for all participants for all tests, with the number included in each model listed. Most biomarkers had modest predictive value, with suPAR (AUC 0.81) and IL-6 (AUC 0.77) having the best performance. Many biomarkers had weak performance (CRP, neutrophils, lymphocytes, KL-6), with AUC figures between 0.5 and 0.65. ROC plots are available in the Supplemental Material for all biomarkers. Of note, both age and NEWS2 score performed as well as most biomarkers.

Table 4: Biomarker/Clinical indicator performance to predict severe COVID-19 outcomes:

Biomarker/Clinical indicator	AUC (95% C.I.)	Sensitivity at Youden's index	Specificity at Youden's index
CRP (n = 179)	0.69 (0.59-0.78)	0.77	0.55
Neutrophils (n = 185)	0.53 (0.43-0.64)	0.44	0.69
Lymphocytes (n = 185)	0.62 (0.53-0.72)	0.85	0.38
NLR (n = 185)	0.62 (0.53-0.71)	0.82	0.43
IL-6 (n = 125)	0.77 (0.65-0.88)	0.70	0.83
KL-6 (n = 149)	0.51 (0.38-0.65)	0.21	0.92
suPAR (n = 150)	0.81 (0.72-0.88)	0.82	0.65
Procalcitonin (n = 150)	0.72 (0.62-0.81)	0.94	0.43
Ferritin (n = 149)	0.64 (0.53-0.74)	0.59	0.69
LDH (n = 149)	0.62 (0.51-0.72)	0.55	0.69
Troponin (n=149)	0.70 (0.6-0.79)	0.88	0.45
BNP (n=149)	0.64 (0.53-0.74)	0.67	0.58
Age (n=187)	0.70 (0.62-0.77)	0.95	0.41
NEWS2 score (n=182)	0.70 (0.60-0.79)	0.41	0.88
CURB-65 (n= 180)	0.65 (0.55-0.73)	0.72	0.54
Radiographic severity score (n=182)	0.60 (0.49-0.71)	0.36	0.91

Results of routine biomarkers (CRP, neutrophils, lymphocytes and NLR) are shown in Figure 2. Best performing novel biomarkers (IL-6, suPAR, LDH and PCT) box plots are shown in Figure 3 (PCT values have been logged prior to plotting to aid visualisation as they had very wide ranges).

Figure 2 – Boxplots of routine blood biomarkers

Figure 3 – Boxplots of best performing novel blood biomarkers

Secondary outcomes

For the composite outcome of ITU and mortality, results are reported in the supplementary Material. Due to a small number of events (18 deaths, 15 ITU admissions), confidence intervals were wider, but results were similar although perhaps slightly worse. Again, suPAR and IL-6 were the best performing biomarkers (AUC 0.81 for suPAR, AUC 0.70 for IL-6), with most other biomarkers having an AUC between 0.5 and 0.65.

DISCUSSION:

COVID-19 remains a clinical challenge. The majority of patients who present to hospital will recover but some develop rapidly progressive respiratory failure. Biomarkers that might predict this deterioration would be invaluable when triaging patients on hospital admission to inform who can be safely discharged versus those who might need intensive care support in the near future. This paper presents a prospectively recruited UK cohort of patients with COVID-19 with targeted biomarker sampling at presentation.

Previous literature

A large observational study recruiting from the majority of NHS hospitals (ISARIC) estimates that, of the 34608 patients with outcome data, 7374 (17%) required admission to intensive care and 11659 (33.7%) died within follow up.¹² There have been numerous studies and meta-analyses on potential predictors of respiratory decline in COVID-19 pneumonia with the development of a 'cytokine storm' in specific patients cited as a major determinant.^{13 14} As a result there has been a focus on biomarkers that rise in other similar conditions such as ferritin, lactate dehydrogenase (LDH), IL-6, and soluble urokinase plasminogen activator receptor (suPAR).¹⁵ A rise in cardiovascular events and coagulopathies has also been seen in patients with COVID-19 pneumonia so biomarkers such as troponin, NT-pro BNP, fibrinogen and D-dimer have been studied.¹⁶⁻¹⁹ Given that mortality from COVID-19 increases with age and frailty, biomarkers of frailty have been shown to be related to worse outcomes (albumin, eGFR).^{2 12 20} Finally, given this is a respiratory infection, blood-based biomarkers that prognosticate bacterial pneumonia have been included in treatment guidelines and even entry criteria for clinical trials, including lymphopenia, neutrophilia, procalcitonin and C-reactive protein.^{6 21}

Study findings

In this study, conventionally performed blood biomarkers, in isolation, did not predict outcome when performed on admission. Neutrophilia and C-reactive protein had AUC close to 0.5, with lymphopenia and NLR having only marginally better discriminative value. Cardiac markers were on average slightly higher in patients with worse clinical outcomes but should not be relied on to make treatment decisions at baseline. Literature suggests that they may have more utility when measured serially, especially in the very unwell patient.¹⁶

Markers of immune activation had more promise at the admission timepoint with IL-6 and suPAR the best-performing within this cohort. It has been hypothesized that an exaggerated immune response or 'cytokine storm' plays a significant role in COVID-19 and several therapies immunomodulatory therapies are being trialed. IL-6 is a pro-inflammatory mediator for the induction of the acute phase response and shown to be a predictive marker of deterioration in other serious lung pathologies. Specific to COVID-19, Han and colleagues assessed the cytokine profile of 102 patients, admitted to a single hospital in Wuhan, by disease severity on admission. From a panel of 6 cytokines (TNF- α , IFN- γ , IL-2, IL-4, IL-6 and IL-10), serum IL-6 was predictive of disease severity at the time with an AUC of 0.84, although longer term outcomes were not reported.²² Another prospective study of 89 patients admitted to a German hospital demonstrated that admission IL-6 was superior to other blood-based biomarkers at predicting the need for mechanical ventilation (using a cut-off of 35pg/ml).²² Another prospective study of 89 patients admitted to a German hospital demonstrated that admission IL-6 was superior to other blood-based biomarkers at predicting the need for mechanical ventilation (using a cut-off of 35pg/ml).⁶ In our study, the admission IL-6 (at a cut-off of 74.9pg/ml) had a sensitivity of 0.70 and specificity of 0.84 (AUC 0.77). Randomised trial evidence is emerging showing treatment with the IL-6 receptor antagonists, tocilizumab and sarilumab, improves outcomes, including survival in severely unwell patients with COVID-19.²³

suPAR, a marker of immune activation and has been shown to predict deterioration in several infectious and inflammatory disorders. It forms part of the fibrinolysis cascade and increased levels have been shown to pre-dispose to clotting abnormality and renal dysfunction, both of which are important drivers of morbidity and mortality in COVID-19. A prospective study of 57 patients presenting with COVID-19 demonstrated that admission levels of suPAR were significantly greater among patients who eventually required ventilatory support.¹⁵ A cut-off of 6ng/ml had a sensitivity of 86% and specificity of 92%. A prospective study of 57 patients presenting with COVID-19 demonstrated that admission levels of suPAR were significantly greater among patients who eventually required ventilatory support.¹⁵ A cut-off of 6ng/ml had a sensitivity of 86% and specificity of 92%. In our study it performed reasonably compared to other blood biomarkers but at a cut-off of 5.2ng/dl only had a sensitivity of 82% and specificity of 65%, with lower cutoffs increasing sensitivity at relatively little cost.

Clinical markers

The focus of this study was the additional role of blood biomarkers when initially assessing patients presenting with COVID-19. The DISCOVER cohort also collected routinely recorded clinical data including demographics, baseline observations and initial radiography. It is notable that this easily accessible information outperformed many of the blood-based biomarkers tested.

The National Early Warning Score (NEWS2) is used throughout the United Kingdom. There is currently limited evidence supporting its use in COVID-19.²⁴ In China, Liao and colleagues created a non-validated COVID specific NEWS2 score (where age over 65 added 3 points).²⁵ We could not find any prospective UK-based studies assessing the utility of NEWS2 in COVID-19 admission.²⁵ In the DISCOVER cohort, NEWS2 score had an AUC of 0.70. The CURB-65 score is a severity scoring system for community acquired pneumonia and is often used to make decisions around need for hospital admission. Although previous retrospective literature from our centre has shown that CURB-65 scores are higher in patients with a worse outcome from COVID-19,²⁶ this prospective study has demonstrated that it cannot be relied upon to make treatment decisions on an individual patient level.

Strengths

The DISCOVER cohort was prospectively designed and analysed the earliest emergency department blood sample, which is a key strength. Clinical meta-data was robustly recorded, and novel biomarkers were performed in batch assays, maximising replicability and reducing bias as results could not influence patient care. Unlike other datasets (e.g. EHR extractions), we included patients who had clinically suspected COVID-19, as the currently available assays still have limited sensitivity. Technicians were employed to extract blood out of auto-analyser machines to get the earliest possible sample, enabling recruitment of patients after admission or even as outpatients. Missing data around comorbidities was rare, as patients were prospectively recruited by research nurses.

Limitations

The major limitation of this study is the limited sample size, leading to imprecise estimates of biomarker performance. A second limitation is the composite outcome of non-invasive ventilation, ITU admission and death which has been used in major interventional COVID-19 trials. Although

this is clinically useful and may aid differentiation of those who require specialist care, the provision and use of non-invasive ventilation is more clinician and hospital dependent and may be harder to extrapolate from. However, our secondary analysis of intensive care admission and death was largely similar, supporting this approach. Finally, although around 80% of patients had blood available, a proportion did not have excess sample available for additional testing, so only the routinely performed biomarkers (CRP, neutrophils, lymphocytes, neutrophil:lymphocyte ratio) are recorded for those patients. It is also important to note that these results were available to clinicians, and therefore may have biased the outcome (i.e. patients with high CRP may have received more aggressive therapy). At the time of recruitment to this study, there were however no approved therapies for COVID-19 and it is therefore unlikely this strongly biased the results.

CONCLUSION:

To our knowledge this is the largest recruited UK cohort of consecutive patients presenting with COVID-19. Blood biomarkers, when performed on admission bloods, had only moderate predictive value for COVID-19 similar to age and routine clinical scores (e.g. NEWS2 score). IL-6 and suPAR had the best performance, and further, large prospective studies should validate the additional value of these biomarkers to routinely collected clinical information.

Contributorship statement:

DTA and FJ had the idea for the study. SB, AMc and NAM has a supervisory role. AM lead the collection of samples. KE, JE, CD AO and AN were involved in the analysis of samples. FH led the data analysis. All authors were involved in the writing of the manuscript.

Funding:

The DISCOVER study was supported by grants from the Southmead Hospital Charity and Elizabeth Blackwell Institute (Grant number: not applicable/ NA).

DTA is funded by a National Institute for Health Research (NIHR) Doctoral Research Fellowship (DRF-2018-11-ST2-065). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

FH is funded via the National Institute for Health Academic Clinical Fellowship scheme.

The suPARnostic™ ELISA kits and IL-6/KL-6 assays were gifts from ViroGates (Birkerød, Denmark) and Fujirebio Europe respectively for unrestricted research activity. (Grant number: not applicable/ NA).

Competing interests:

None declared

Data sharing:

Although we are unable to share raw data, the analytic code is available at https://github.com/gushamilton/discover_prediction/

Ethics Approval Statements

Ethics approval via South Yorkshire REC: 20/YH/0121, CRN approval no: 45469

References

1. Petrilli CM, Jones SA, Yang J, et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ* 2020 doi: 10.1136/bmj.m1966
2. Galloway JB, Norton S, Barker RD, et al. A clinical risk score to identify patients with COVID-19 at high risk of critical care admission or death: an observational cohort study. *J Infect* 2020 doi: 10.1016/j.jinf.2020.05.064
3. Liang W, Liang H, Ou L, et al. Development and Validation of a Clinical Risk Score to Predict the Occurrence of Critical Illness in Hospitalized Patients With COVID-19. *JAMA Intern Med* 2020 doi: 10.1001/jamainternmed.2020.2033
4. Shi S, Qin M, Cai Y, et al. Characteristics and clinical significance of myocardial injury in patients with severe coronavirus disease 2019. *Eur Heart J* 2020 doi: 10.1093/eurheartj/ehaa408
5. Smadja DM, Guerin CL, Chocron R, et al. Angiotensin-converting enzyme 2 as a marker of endothelial activation is a good predictor factor for intensive care unit admission of COVID-19 patients. *Angiogenesis* 2020 doi: 10.1007/s10456-020-09730-0
6. Herold T, Jurinovic V, Arnreich C, et al. Elevated levels of interleukin-6 and CRP predict the need for mechanical ventilation in COVID-19. *J Allergy Clin Immunol* 2020 doi: 10.1016/j.jaci.2020.05.008
7. Rastrelli G, Di Stasi V, Inglese F, et al. Low testosterone levels predict clinical adverse outcomes in SARS-CoV-2 pneumonia patients. *Andrology* 2020 doi: 10.1111/andr.12821
8. Harris PA, Taylor R, Minor BL, et al. The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform* 2019;95:103208. doi: 10.1016/j.jbi.2019.103208 [published Online First: 2019/05/13]
9. McGinley A, Pearse RM. A national early warning score for acutely ill patients. *BMJ* 2012;345:e5310. doi: 10.1136/bmj.e5310 [published Online First: 2012/08/10]
10. Wong HYF, Lam HYS, Fong AH, et al. Frequency and Distribution of Chest Radiographic Findings in COVID-19 Positive Patients. *Radiology* 2019:201160. doi: 10.1148/radiol.2020201160 [published Online First: 2020/03/29]
11. Wynants L, Van Calster B, Bonten MMJ, et al. Prediction models for diagnosis and prognosis of covid-19 infection: systematic review and critical appraisal. *BMJ* 2020 doi: 10.1136/bmj.m1328
12. ISARIC (International Severe Acute Respiratory and Emerging Infections Consortium): COVID-19 Report: 08 June 2020.
13. Henry BM, de Oliveira MHS, Benoit S, et al. Hematologic, biochemical and immune biomarker abnormalities associated with severe illness and mortality in coronavirus disease 2019 (COVID-19): a meta-analysis. *Clin Chem Lab Med* 2020 doi: 10.1515/cclm-2020-0369
14. Kermali M, Khalsa RK, Pillai K, et al. The role of biomarkers in diagnosis of COVID-19 - A systematic review. *Life Sci* 2020 doi: 10.1016/j.lfs.2020.117788
15. Rovina N, Akinosoglou K, Eugen-Olsen J, et al. Soluble urokinase plasminogen activator receptor (suPAR) as an early predictor of severe respiratory failure in patients with COVID-19 pneumonia. *Crit Care* 2020 doi: 10.1186/s13054-020-02897-4
16. Vrsalovic M, Vrsalovic Presecki A. Cardiac troponins predict mortality in patients with COVID-19: A meta-analysis of adjusted risk estimates. *J Infect* 2020 doi: 10.1016/j.jinf.2020.05.022
17. Gao L, Jiang D, Wen X-S, et al. Prognostic value of NT-proBNP in patients with severe COVID-19. *Respir Res* 2020 doi: 10.1186/s12931-020-01352-w

18. Artifoni M, Danic G, Gautier G, et al. Systematic assessment of venous thromboembolism in COVID-19 patients receiving thromboprophylaxis: incidence and role of D-dimer as predictive factors. *J Thromb Thrombolysis* 2020 doi: 10.1007/s11239-020-02146-z
19. Zhang L, Yan X, Fan Q, et al. D-dimer levels on admission to predict in-hospital mortality in patients with Covid-19. *J Thromb Haemost* 2020 doi: 10.1111/jth.14859
20. Kato H, Shimizu H, Shibue Y, et al. Clinical course of 2019 novel coronavirus disease (COVID-19) in individuals present during the outbreak on the Diamond Princess cruise ship. *J Infect Chemother* 2020 doi: 10.1016/j.jiac.2020.05.005
21. Luo Z, Wang N, Liu P, et al. Association between chest CT features and clinical course of Coronavirus Disease 2019. *Respir Med* 2020 doi: 10.1016/j.rmed.2020.105989
22. Han H, Ma Q, Li C, et al. Profiling serum cytokines in COVID-19 patients reveals IL-6 and IL-10 are disease severity predictors. *Emerg Microbes Infect* 2020 doi: 10.1080/22221751.2020.1770129
23. <https://www.medrxiv.org/content/10.1101/2021.01.07.21249390v2>.
24. Ihle-Hansen H, Berge T, Tveita A, et al. COVID-19: Symptoms, course of illness and use of clinical scoring systems for the first 42 patients admitted to a Norwegian local hospital. *Tidsskr Nor Laegeforen* 2020 doi: 10.4045/tidsskr.20.0301
25. Liao X, Wang B, Kang Y. Novel coronavirus infection during the 2019-2020 epidemic: preparing intensive care units-the experience in Sichuan Province, China. *Intensive Care Med* 2020 doi: 10.1007/s00134-020-05954-2
26. Tomlins J, Hamilton F, Gunning S, et al. Clinical features of 95 sequential hospitalised patients with novel coronavirus 2019 disease (COVID-19), the first UK cohort. *J Infect* 2020 doi: 10.1016/j.jinf.2020.04.020