





Depression Symptoms and Sleep Quality During COVID-19 Outbreak in Iran: A National Cross-sectional Study

Samaneh Torkian ¹, Elham Akhlaghi ², Vahid Khosravi ³, Reza Etesami ⁴, Ali Ahmadi ^{5,*}, Narges Khanjani ⁶ and Vajihe Rouhi ⁷

¹Department of Epidemiology, School of Public Health, Iran University of Medical Sciences, Tehran, Iran

²Department of Medical Surgical Nursing, School of Nursing and Midwifery, Iran University of Medical Sciences, Tehran, Iran

³Department of Health Education and Promotion, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

⁴Department of Statistics, Shahid Bahonar University, Kerman, Iran

⁵Department of Epidemiology and Biostatistics, School of Health and Modeling in Health Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

⁶Neurology Research Center, Kerman University of Medical Sciences, Kerman, Iran

⁷Department of Epidemiology, School of Public Health, Kerman University of Medical Sciences, Kerman, Iran

*Corresponding author: Modeling in Health Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran. Email: aliahmadi2007@gmail.com

Received 2020 August 14; Revised 2020 December 20; Accepted 2021 January 24.

Abstract

Background: Coronavirus 2019 is a new infectious disease that has been declared by the World Health Organization as a global pandemic. This pandemic may have psychological effects on populations.

Objectives: This study aimed to investigate the quality of sleep and depression in the Iranian community during COVID-19 pandemic.

Methods: This nationwide cross-sectional web-based study was conducted from April 19 to May 12, 2020 in Iran. The data collection tools were the Pittsburgh Sleep Quality Index and Depression Center for Epidemiologic Studies' Depression Scale questionnaires. Each province in Iran was assumed as a cluster, and generalized estimating equations models were used to determine the demographic variables that were related to depression symptoms and sleep quality during COVID-19 outbreak.

Results: In this study, 73.5% and 41.4% of the participants had poor sleep quality and had depression symptoms, respectively. Perceived income, hours spent outdoors, the number of people with COVID-19 that were acquainted with, and a history of depression significantly were related to sleep quality. Also, gender, age, having a child, education level, economic status, and a history of depression were significantly associated with having depression symptoms.

Conclusions: Most participants had poor sleep quality, and less than half had depression symptoms during the COVID-19 pandemic. These problems can disturb daily routines, cut social ties, and increase worries and fears of the future. Interventions are necessary to reduce psychological problems in Iran.

Keywords: Corona Virus Disease 2019 (COVID-19), Depression, Psychological Problems, Sleep Quality, Iran

1. Background

Coronavirus disease 2019 (COVID-19) is a new infectious disease that causes severe acute respiratory symptoms. It was first identified in December 2019, in Wuhan, China, and has since spread globally (1). This is a new viral infection that does not have a treatment and currently can only be treated symptomatically (2). On January 30, 2020, the World Health Organization (WHO) declared a public health emergency of international concern (PHEIC) and, on February 11, announced that the new coronavirus would be called "COVID-19" (3). On March 11, 2020, WHO declared the coronavirus outbreak a "Global Pandemic" (4). Until April 2, 2020, COVID-19 has been reported from 216 coun-

tries with 6,194,533 confirmed cases and 376,320 deaths (5). Iran was ranked 10th globally in terms of confirmed cases of the disease on July 10, 2020 (6).

The COVID-19 pandemic has affected all aspects of people's lives and has changed their lifestyles (7). This pandemic may also affect communities psychologically by increasing fear and isolation and decreasing social and economic activities (8). Lack of knowledge about COVID-19 is one of the main causes of stress among people. When WHO declared COVID-19 a global pandemic, fear, and anxiety about COVID-19 spread as fast as the virus itself. While the whole world was trying to find a vaccine and effective drugs, the best intervention to reduce the rate of disease transmission was announced to be social distancing and

quarantine (9). The change in the term "social distance" to "physical distance" by WHO has caused people to feel lonely and socially isolated. As a result, people may experience mental disorders such as depression and sleep problems (9, 10).

Sleep is a naturally reversible state that contributes to physical and mental health, but disturbed sleep can cause a range of physical, mental, and emotional problems (11). Sleep quality is a key indicator of physical and mental health (12). Research has shown that sleep deprivation suppresses the immune system and causes depression (13, 14). Depression is the first most common mental illness, and is a natural response to awkward life situations (15). In addition to stress and anxiety, there are several factors that increase the chance of depression, including loneliness, staying away from others, misinformation on social media, and economic insecurity (9). With the progress of the COVID-19 epidemic to different stages, sleep and psychological conditions such as depression may change (16), and research on these psychological outcomes is necessary because the spread of COVID-19 in the Iranian population can have long-term psychological impacts (9).

2. Objectives

The main purpose of this study was to investigate depression and sleep quality and demographic factors related to it among Iranians during the COVID-19 pandemic.

3. Methods

This nationwide web-based cross-sectional study was conducted to evaluate depression and sleep quality in the Iranian general population during the COVID-19 pandemic in 2020. Data were collected from April 19 to May 12; and 3,446 people completed the questionnaire via the link. The questionnaire included demographic variables, the Pittsburgh sleep quality index (PSQI), and the depression center for epidemiologic studies' depression scale (CES-D 20).

Convenient sampling was done. The questionnaire was distributed as a web link through social networks, including Telegram, WhatsApp, and Instagram. According to Hyland et al. (2020) (17) in which the prevalence of depression was 23%, with a 95% confidence level 0.05 and error $d = 0.014$, the minimum sample size required was 3,436. Inclusion criteria were people over 18 years old with the ability to read and write. Questionnaires were completed without a name. Participation in the study was completely voluntary.

Demographic variables included in the questionnaire were gender (male, female), age (≤ 20 , 21 - 30, 31 - 40,

40 - 51, and > 50 years), marital status (single, married, separated, which included divorced and widowed), having a history of depression or current depressive disorder (yes, no), education (high school diploma and less, graduate diploma, bachelor, masters, and doctorate), job (unemployed, retired, student, freelance job, employee), income (good, moderate and poor), having a child (yes, no), having COVID-19 disease in the last two months (yes, no, I do not know), the number of people with COVID-19 in the past two months whom the participant had known (nobody, 1 - 2, 3 - 5, > 5 people) and hours spent outside home during a day in the past two months (no time, 1 - 3, 4 - 7, > 7 hours).

The PSQI is a useful instrument used to measure the quality and patterns of sleep. It includes seven domains, which are sleep duration, sleep disturbance, sleep latency, daytime dysfunction due to sleepiness, habitual sleep efficiency, sleep quality, and the need to use medicine for sleep over the last month. The questionnaire has nine items, but question 5 itself contains ten sub-items, so the whole questionnaire has 19 items, which are scored on a 4-point Likert scale from 0 to 3. The values of "0" and "3" indicate "better" and "worse" sleep, respectively. The value of all seven domains is summed to get the total score. The overall score range of the PSQI questionnaire is from 0 to 21. A total score of ≤ 5 represents poor sleep quality and > 5 represents good sleep quality (18).

The CES-D is a 20-item scale to measure the current level of depression symptoms in the general population during the past week. The questionnaire has four subscales, somatic complaints, depressed affect, positive affect, and interpersonal problems. The Likert scale is from "0 = rarely or none of the time" to "4 = almost or all of the time". Items 4, 8, 12, and 16 are reversely scored. Scores range from 0 to 60; higher scores reflect greater levels of the depression symptoms, while lower scores reflect fewer symptoms. The cut-off point that has been recommended for depression cases is 20. Individuals with a score ≥ 20 or more are classified as having depression during the past week (19).

The validity and reliability of the questionnaires have already been confirmed in the Iranian community (20-22). This study was approved by the Ethics Committee of Shahrekord University of Medical Science (SKUMS). The ethics code for this study is IR.SKUMS.REC.1399.027.

We used descriptive statistics to summarize demographic characteristics and questionnaire scores. The mean of sleep quality and depression was compared in the demographic sub-groups. Each province was assumed as a cluster, and generalized estimating equations (GEE) models were used to determine the demographic variables that were related to depression symptoms and sleep quality. All

independent variables were entered into the univariable model, and variables with a p-value of less than 0.2 were entered into the multivariable model. The final model was achieved using a backward selection approach. Odds ratio (OR), adjusted odds ratio (AOR), and 95% confidence intervals (95% CI) were reported. P-values of < 0.05 were considered statistically significant. All analyses were conducted with SPSS software version 26.

4. Results

In the present study, most participants were female (66.7%), were married (65%), were unemployed (34.1%), had a bachelor degree (35%), had at least one child (56.6%), perceived their income was moderate (67.6%), had no history of depression (76.3%) or COVID-19 in the last two months (84.6%). The mean (\pm SD) age of the participants was 34.2 (\pm 11.60) years and the average number of hours they spent out of home during a day in the past two months was 3.71 (\pm 4.60). The median number of people with COVID-19 whom the participants had known was 2.

The mean (\pm SD) of sleep quality and depression scores in the Iranian provinces are reported in Table 1. The provinces with the worst situation in sleep quality were Ilam, Semnan, and West Azabajejan; and the provinces with the worst situation in regard to depression were Sistan and Baluchestan, Golestan, and South Khorasan.

The mean (\pm SD) of sleep quality and depression scores in demographic subgroups are presented in Table 2. Predictors of sleep quality and depression in univariable and multivariate analysis are outlined in Tables 3 and 4, respectively.

In this study, 73.5% of participants had poor sleep quality. The mean (\pm SD) of the sleep quality scores was 7.51 (\pm 2.88). The mean scores of sleep quality in different COVID-19 incidence (high, medium, and low) provinces are shown in Figure 1A. There was no significant difference between these provinces in terms of sleep quality.

The variables in the multivariable model that were significantly associated with sleep quality were poor perceived income, not knowing about getting infected with COVID-19 in the last two months, knowing more than 5 people who got COVID-19, spending 4 - 7 hours or more than 7 hours outside the home, and having a history of depression (Table 4).

About 41.4% of the participants had depression symptoms. The mean (\pm SD) of participants' depression score was 18.89 (\pm 11.72). The mean scores of depression in different COVID-19 incidence (high, medium, and low) provinces

are shown in Figure 1B. There was no significant difference between these provinces in terms of depression.

The variables that were significantly associated with depression and stayed in the final multivariable model were female sex, age groups, including ages \leq 20, 21 - 30 and 31 - 40 years, not having children, high school diploma or less education, perceived poor or moderate-income, not knowing about getting infected with COVID-19 in the last two months, and having a history of depression (Table 4).

5. Discussion

The present study found that most participants had poor sleep quality and less than half had depression symptoms during the COVID-19 pandemic in Iran.

In a web-based cross-sectional survey in China during COVID-19 outbreak, about 18.2% of the general population had poor sleep quality (4), and among medical staff who manage patients with COVID-19 in January and February 2020, social support was significantly associated with self-efficacy and sleep quality; thus anxiety, stress, and self-efficacy were mediating variables related to sleep quality (2). Also, in China, the PSQI score was significantly higher in nursing interns during outbreak of COVID-19 than national norms, and depression positively correlated with the sleep quality score (23). Studies have shown that good quality sleep enhances psychoneuroimmunity against COVID-19 (7). The world sleep society has highlighted the importance of sleep as the basis of health, which helps in better decision making and cognitive understanding, even in big issues like COVID-19 pandemic (24). Researchers think getting enough sleep is not a luxury, it is something people need for good health, and sleep disorders can increase a person's risk of health problems (25). Mental health is also affected by disturbed sleep. Insufficient sleep, less than 5 h per day, is associated with higher rates of depression, stress, and suicide. Sleep deficiency is also associated with bad moods, poor emotional regulation (26), poor mental health, and even early death (27). Sleep deficiency and sleep latency can cause behavioral problems that include (28) internalized behaviors such as anxiety, depression, excess worry, psychosomatic symptoms, and externalized behaviors such as aggression, impulsive or disruptive behaviors, delinquency, and non-compliance (29). Many factors play a role in sleep quality, including occupation, beliefs, age, the living environment, and social factors. It is important to discover populations at high risk of sleep deprivation and provide the right type of intervention (11).




Figure 1. The individual and pooled results of A, sleep quality; and B, depression in Iranian provinces

Table 1. The Mean (\pm SD) and Prevalence of Sleep Quality and Depression in the Provinces in Iran

Variables	No. %	Sleep Quality			Depression		
		Mean (\pm SD)	Good	Poor	Mean (\pm SD)	Depression	No Depression
Esfahan	360 (10.4)	18.16 (\pm 11.02)	100 (27.8)	260 (72.2)	7.28 (\pm 2.79)	48 (43.6)	62 (56.2)
Alborz	55 (1.6)	18.54 (\pm 13.21)	9 (16.4)	46 (83.6)	7.89 (\pm 2.78)	24 (43.6)	31 (56.4)
Khuzestan	162 (4.7)	18.63 (\pm 11.65)	40 (24.7)	122 (75.3)	8.00 (\pm 3.09)	63 (38.9)	99 (61.1)
Tehran	379 (11.0)	18.21 (\pm 11.28)	98 (25.9)	281 (74.1)	7.63 (\pm 2.96)	144 (38.0)	235 (62.0)
Markazi	45 (1.3)	18.44 (\pm 12.12)	9 (20.0)	36 (80.0)	7.55 (\pm 2.46)	30 (66.7)	15 (33.3)
Bushehr	38 (1.1)	18.52 (\pm 12.17)	12 (31.6)	26 (68.4)	7.36 (\pm 2.68)	17 (44.7)	21 (55.3)
Chahar Mahal Bakhtiari	482 (14.0)	18.38 (\pm 11.41)	160 (33.2)	322 (66.8)	6.97 (\pm 2.79)	188 (39.0)	294 (61.0)
Kerman	100 (2.9)	17.40 (\pm 12.18)	19 (19)	81 (81)	7.58 (\pm 0.49)	40 (40.0)	60 (60.0)
Hormozgan	71 (2.1)	20.28 (\pm 12.07)	21 (29.6)	50 (70.4)	7.38 (\pm 2.83)	31 (43.7)	40 (56.3)
Sistan and Baluchestan	153 (4.4)	23.99 (\pm 12.08)	24 (15.7)	129 (84.3)	8.15 (\pm 2.83)	100 (65.4)	53 (34.6)
Zanjan	110 (3.2)	18.58 (\pm 12.02)	30 (27.3)	80 (72.5)	7.17 (\pm 2.78)	48 (43.6)	62 (56.2)
Ilam	7 (0.2)	13.57 (\pm 8.05)	1 (14.3)	6 (85.7)	7.28 (\pm 1.79)	1 (14.3)	6 (85.7)
Qom	101 (2.9)	21.46 (\pm 12.44)	24 (23.8)	77 (76.2)	7.91 (\pm 3.20)	49 (48.5)	52 (51.5)
Lorestan	74 (2.1)	19.51 (\pm 12.23)	21 (28.4)	53 (71.6)	7.45 (\pm 3.16)	28 (37.8)	46 (62.2)
Khorasan Razavi	277 (8.0)	22.00 (\pm 12.43)	69 (24.9)	208 (75.1)	7.68 (\pm 2.79)	145 (52.3)	132 (47.7)
Kohgiluyeh and Boyer-Ahmad	131 (3.8)	17.70 (\pm 10.99)	41 (31.3)	90 (68.7)	7.57 (\pm 3.11)	49 (37.4)	82 (62.6)
Fars	160 (4.6)	17.87 (\pm 11.85)	55 (34.4)	105 (65.6)	7.45 (\pm 3.18)	61 (38.1)	99 (61.9)
Hamedan	72 (2.1)	18.51 (\pm 11.03)	10 (13.9)	62 (86.1)	7.75 (\pm 2.56)	33 (45.8)	39 (54.2)
Yazd	87 (2.5)	17.05 (\pm 11.25)	20 (23.0)	67 (77.0)	7.83 (\pm 2.84)	35 (40.2)	52 (59.8)
Semnan	22 (0.6)	15.13 (\pm 10.43)	6 (27.3)	16 (72.7)	6.81 (\pm 2.70)	3 (13.6)	19 (86.4)
Golestan	34 (1.0)	21.00 (\pm 11.75)	4 (11.8)	30 (88.2)	9.20 (\pm 3.20)	17 (50.0)	17 (50.0)
East Azarbaijan	40 (1.2)	18.57 (\pm 11.78)	16 (26.7)	44 (73.3)	7.32 (\pm 3.18)	15 (37.5)	25 (62.5)
Kurdistan	43 (1.2)	17.13 (\pm 11.55)	10 (23.3)	33 (76.7)	7.39 (\pm 2.47)	17 (39.5)	26 (60.5)
West Azarbaijan	60 (1.7)	15.55 (\pm 10.72)	13 (32.5)	27 (67.5)	7.08 (\pm 2.27)	18 (30.0)	42 (70.0)
Qazvin	72 (2.1)	16.98 (\pm 10.65)	19 (26.4)	53 (73.6)	7.34 (\pm 3.02)	27 (37.5)	45 (62.5)
Mazandaran	68 (2.0)	17.16 (\pm 10.48)	16 (23.5)	52 (76.5)	7.55 (\pm 2.60)	24 (35.3)	44 (64.7)
Kermanshah	51 (1.5)	18.70 (\pm 11.20)	13 (25.5)	38 (74.5)	7.31 (\pm 2.33)	19 (37.3)	32 (62.7)
North Khorasan	37 (1.1)	20.67 (\pm 12.54)	11 (29.7)	26 (70.3)	7.72 (\pm 3.87)	18 (48.6)	19 (51.4)
Ardabil	18 (0.5)	22.83 (\pm 10.29)	4 (22.2)	14 (77.8)	7.83 (\pm 2.83)	11 (61.1)	7 (38.9)
South Khorasan	12 (0.34)	25.41 (\pm 12.80)	1 (8.1)	11 (91.7)	8.33 (\pm 2.83)	7 (58.3)	5 (41.7)
Gilan	125 (3.6)	18.02 (\pm 12.14)	36 (28.8)	89 (71.2)	7.52 (\pm 2.91)	44 (35.2)	81 (64.8)

In this study, less than half of the participants had symptoms of depression. In a survey conducted during the COVID-19 pandemic in Iran, the prevalence of depression was significantly higher among doctors (52%) and nurses (51.4%) compared with other occupations (30). In Guiyang, during the COVID-19 outbreak, the mean scores of depression among medical staff members were significantly higher than the general Chinese population (31). In

the present study, unemployed people and students were more depressed in the univariable analysis. Similarly, during the COVID-19 pandemic in China, college students were more depressed than the normal national population (32) and younger participants were more likely to show depression symptoms than older participants, which is similar to our study (4). In Nepal, during the COVID-19 pandemic, the prevalence of depression was 34.0%, and multivariable

Table 2. Association Between Demographic Variables and the Psychological Impact of Novel Coronavirus Disease 2019 (COVID-19)

Variables	No. (%)	Depression			Sleep Quality		
		No Depression	Depression	P-Value	Good Sleep Quality	Poor Sleep Quality	P-Value
Overall	3446 (100)	2018 (58.6)	1428 (41.4)	-	912 (26.5)	2534 (73.5)	-
Gender				< 0.001			< 0.001
Male	1146 (33.3)	733 (64.0)	413 (36.0)		354 (30.9)	792 (69.1)	
Female	2300 (66.7)	1285 (55.9)	1015 (44.1)		558 (24.3)	1742 (75.7)	
Age (y)				< 0.001			0.004
20 and under	417 (12.1)	183 (43.9)	234 (56.1)		90 (21.6)	327 (78.4)	
21 - 30	973 (28.2)	524 (53.9)	449 (46.1)		246 (25.3)	727 (74.7)	
31 - 40	1132 (32.8)	675 (59.6)	457 (40.4)		305 (26.9)	827 (73.1)	
41 - 50	603 (17.5)	399 (66.2)	204 (33.8)		180 (29.9)	423 (70.1)	
Over 50	321 (9.4)	237 (73.8)	84 (26.2)		91 (28.3)	230 (71.7)	
Marital status				< 0.001			0.029
Single	1106 (32.1)	559 (50.5)	547 (49.5)		275 (24.9)	831 (75.1)	
Divorced/widow	100 (2.9)	52 (52.0)	48 (48.0)		20 (20.0)	80 (80.0)	
Married	2240 (65.0)	1407 (62.8)	833 (37.2)		617 (27.5)	1623 (72.5)	
Having a child or children				< 0.001			0.064
Yes	1950 (56.6)	1245 (63.8)	705 (36.2)		540 (27.7)	1410 (72.3)	
No	1495 (43.4)	733 (51.7)	722 (48.3)		372 (24.9)	1123 (75.1)	
Education				< 0.001			0.064
Diploma and less	1182 (34.3)	600 (50.8)	582 (49.2)		277 (23.4)	905 (76.6)	
Graduate diploma	376 (10.9)	220 (58.5)	156 (41.5)		94 (25.0)	282 (75.0)	
Bachelor	1207 (35.0)	741 (61.4)	466 (38.6)		337 (27.9)	870 (72.1)	
Masters/doctorate	681 (19.8)	457 (67.1)	224 (32.9)		204 (30.0)	477 (70.0)	
Job				0.028			0.360
Unemployed	1176 (34.1)	581 (49.4)	595 (50.6)		268 (22.8)	908 (77.2)	
Freelance job	575 (16.7)	361 (62.8)	214 (37.2)		163 (28.3)	412 (71.7)	
Employee	962 (27.9)	661 (68.7)	301 (31.3)		305 (31.7)	657 (68.3)	
Student	553 (16.0)	281 (50.8)	272 (49.2)		121 (21.9)	432 (78.1)	
Retired	180 (5.2)	134 (74.4)	46 (25.6)		55 (30.6)	125 (69.4)	
Perceived income				< 0.001			0.001
Good	486 (14.1)	338 (69.5)	148 (30.5)		147 (30.2)	339 (69.8)	
Moderate	2328 (67.6)	1419 (61.0)	909 (39.0)		627 (26.9)	1701 (73.1)	
Poor	632 (18.3)	261 (41.3)	371 (58.7)		138 (21.8)	494 (78.2)	
Having COVID-19 disease in the last two months				0.004			< 0.001
Yes	61 (1.8)	34 (55.7)	27 (44.3)		14 (23.0)	47 (77.0)	
No	2917 (84.6)	1738 (59.6)	1179 (40.4)		808 (27.7)	2109 (72.3)	
I don't know	468 (13.6)	246 (52.6)	222 (47.4)		90 (19.2)	378 (80.8)	
The number of people with COVID-19 in the past two months whom they knew				0.995			0.009
Nobody	1113 (32.3)	637 (57.2)	476 (42.8)		311 (27.9)	802 (72.1)	
1 - 2	796 (23.1)	475 (59.7)	321 (40.3)		235 (29.5)	561 (70.5)	
3 - 5	804 (23.3)	489 (60.8)	315 (39.2)		203 (25.2)	601 (74.8)	
Over 5	584 (16.9)	325 (55.7)	259 (44.3)		132 (22.6)	452 (77.4)	
Hours spent outside home				< 0.001			< 0.001
No time	1218 (35.3)	637 (52.3)	581 (47.7)		275 (22.6)	943 (77.4)	
1 - 3	933 (27.1)	544 (58.3)	389 (41.7)		237 (25.4)	696 (74.6)	
4 - 7	365 (10.6)	236 (64.6)	129 (35.4)		104 (28.5)	261 (71.5)	
Over 7	930 (27.0)	601 (64.6)	329 (35.4)		296 (31.8)	634 (68.2)	

Table 3. Univariable Generalized Estimating Equation Results about the Association between Demographic Variables, Depression, and Sleep Quality

Variables	Depression		Sleep Quality	
	OR (95% CI)	P-Value	OR (95% CI)	P-Value
Gender				
Female	1.40 (1.22, 1.61)	< 0.001	1.39 (1.20, 1.61)	< 0.001
Male	1	1	1	1
Age (y)				
20 and under	3.60 (2.66, 4.88)	< 0.001	1.43 (1.03, 2.00)	0.032
21 - 30	2.41 (1.86, 3.13)	< 0.001	1.16 (0.89, 1.53)	0.261
31 - 40	1.91 (1.47, 2.46)	< 0.001	1.07 (0.82, 1.40)	0.605
41 - 50	1.44 (1.09, 1.89)	< 0.001	0.93 (0.69, 1.24)	0.620
Over 50	1	1	1	1
Marital status				
Single	1.65 (1.43, 1.90)	< 0.001	1.14 (0.96, 1.36)	0.118
Divorced/widow	1.55 (1.04, 2.32)	0.030	1.52 (0.92, 2.49)	0.097
Married	1	1	1	1
Having a child or children				
No	1.64 (1.42, 1.90)	< 0.001	1.15 (0.97, 1.37)	0.097
Yes	1	1	1	1
Education				
Diploma and less	1.97 (1.62, 2.40)	< 0.001	1.20 (1.11, 1.75)	0.004
Associate degree	1.44 (1.11, 1.87)	0.005	1.28 (0.96, 1.70)	0.086
Bachelor	1.28 (1.05, 1.55)	0.012	1.10 (0.88, 1.37)	0.378
Masters/doctorate	1	1	1	1
Job				
Unemployed	2.98 (2.11, 4.20)	< 0.001	1.49 (1.08, 2.05)	0.015
Freelance job	1.72 (1.23, 2.42)	0.002	1.11 (0.77, 1.60)	0.571
Employee	1.327 (0.93, 1.87)	0.112	0.94 (0.67, 1.34)	0.762
Student	2.820 (1.95, 4.06)	< 0.001	1.57 (1.06, 2.32)	0.023
Retired	1	1	1	1
Perceived income				
Poor	3.24 (2.56, 4.10)	< 0.001	1.55 (1.19, 2.01)	0.001
Moderate	1.46 (1.20, 1.78)	< 0.001	0.78 (0.95, 1.45)	0.129
Good	1	1	1	1
Having COVID-19 disease in the last two months				
Yes	1.17 (0.70, 1.95)	0.545	1.28 (0.72, 2.27)	0.388
I don't know	1.33 (1.07, 1.64)	0.008	1.60 (1.27, 2.03)	< 0.001
No	1	1	1	1
The number of people with COVID-19 in the past two months whom they knew				
1 - 2	0.90 (0.75, 1.08)	0.266	0.92 (0.77, 1.10)	0.403
3 - 5	0.86 (0.71, 1.04)	0.133	1.14 (0.93, 1.40)	0.181
Over 5	1.06 (0.86, 1.31)	0.544	1.32 (1.04, 1.68)	0.018
Nobody	1	1	1	1
Hours spent outside home				
1 - 3	1.66 (1.41, 1.96)	< 0.001	1.60 (1.31, 1.95)	< 0.001
4 - 7	1.30 (1.09, 1.56)	0.004	1.37 (1.12, 1.67)	0.002
Over 7	0.99 (0.78, 1.26)	0.990	1.17 (0.90, 1.52)	0.236
No time	1	1	1	1
Depression history				
Yes	4.74 (4.09, 5.49)	< 0.001	2.97 (2.41, 3.66)	< 0.001
No	1	1	1	1

Table 4. Multivariable Generalized Estimating Equation Results about the Association between Demographic Variables, Depression, and Sleep Quality

Variables	Depression		Sleep Quality	
	OR (95% CI)	P-Value	OR (95% CI)	P-Value
Gender				
Female	1.36 (1.15, 1.59)	< 0.001	-	-
Male	1	1	-	-
Age (y)				
≤ 20	2.51 (1.73, 3.64)	< 0.001	-	-
21- 30	1.90 (1.40, 2.58)	< 0.001	-	-
31- 40	1.66 (1.24, 2.22)	0.001	-	-
41- 50	1.28 (0.93, 1.77)	0.123	-	-
Over 50	1	1	-	-
Having a child or children				
No	1.43 (1.16, 1.75)	0.001	-	-
Yes	1	1	-	-
Education				
High school Diploma or less	1.58 (1.24, 2.01)	< 0.001	-	-
Graduate diploma	1.27 (0.95, 1.71)	0.097	-	-
Bachelor	1.19 (0.95, 1.48)	0.114	-	-
Masters/doctorate	1	1	-	-
Perceived income				
Poor	3.17 (2.45, 4.11)	< 0.001	1.43 (1.09, 1.89)	0.009
Moderate	1.46 (1.25, 1.91)	< 0.001	0.78 (0.90, 1.41)	0.282
Good	1	1	1	1
Having COVID-19 disease in the last two months				
Yes	1.14 (0.61, 2.12)	0.676	1.01 (0.56, 1.84)	0.952
I don't know	1.37 (1.09, 1.73)	0.006	1.59 (1.24, 2.03)	< 0.001
No	1	1	1	1
The number of people with COVID-19 in the past two months whom they knew				
1-2	-	-	0.94 (0.78, 1.13)	0.519
3- 5	-	-	1.17 (0.95, 1.43)	0.132
Over 5	-	-	1.41 (1.10, 1.81)	0.007
Nobody	-	-	1	1
Hours spent outside home				
1-3	-	-	0.88 (0.71, 1.09)	0.245
4-7	-	-	0.73 (0.57, 0.94)	0.014
Over 7	-	-	0.63 (0.51, 0.77)	< 0.001
No time	-	-	1	1

analysis showed that females and those living alone were significantly more likely to have depression, which is consistent with the present study (33). A review study has mentioned that depression is one of the common psychological reactions to the COVID-19 pandemic (34).

Long-term self-isolation, social distancing, and lockdowns are not well tolerated by people and can have adverse psychological effects (35). In this study, most people with poor sleep quality and depression had poor income. This is probably because, COVID-19 pandemic has stopped or slowed down many businesses and industries, borders are closed, and traveling has decreased significantly (35).

The limitations of the study are that we enrolled volunteers by an online link, and we did not do random sampling. Another limitation of this cross-sectional study is that it shows associations but cannot prove causality.

5.1. Conclusion

Less than half of the participants of this study had symptoms of depression, and almost two-thirds had poor sleep quality. These psychological problems may disrupt their daily routines, cut their social ties, and increase their worries and fears of the future. These psychological issues should be monitored and need increased attention in times of crisis. Psychological interventions may be necessary during the COVID-19 pandemic.

Acknowledgments

We would like to acknowledge Shahrekord University of Medical Sciences for financially supporting this project.

Footnotes

Authors' Contribution: Study concept and design, AA, EA, and ST; Analysis and interpretation of data, ST, AA, VK, and EA; Drafting of the manuscript, ST, VK, NK, AA, VR; Critical revision of the manuscript for important intellectual content, AA, NK, and ST; Statistical analysis, RE.

Conflict of Interests: The authors declare that they have no conflicts of interest.

Ethical Approval: This study was approved by the Ethics Committee of Shahrekord University of Medical Science (SKUMS). The ethics code for this study is IR.SKUMS.REC.1399.027.

Funding/Support: This study was supported by the Shahrekord University of Medical Sciences.

Informed Consent: Written informed consent was signed by all participants.

References

- Torkian S, Akhlaghi E, Khosravi V, Etesami R, Ahmadi A, Khanjani N, et al. Social support and adjustment during COVID-19 epidemic, a community-based study in Iran. *Iran J Psychiatry Behav Sci.* 2020;**14**(4): e108503. doi: [10.5812/ijpbs.108503](https://doi.org/10.5812/ijpbs.108503).
- Xiao H, Zhang Y, Kong D, Li S, Yang N. The effects of social support on sleep quality of medical staff treating patients with coronavirus disease 2019 (COVID-19) in January and February 2020 in China. *Med Sci Monit.* 2020;**26**: e923549. doi: [10.12659/MSM.923549](https://doi.org/10.12659/MSM.923549). [PubMed: [32132521](https://pubmed.ncbi.nlm.nih.gov/32132521/)]. [PubMed Central: [PMC7075079](https://pubmed.ncbi.nlm.nih.gov/PMC7075079/)].
- Torkian S, Kazemi S, Eslahi M, Khatooni E, Etesami R, Khanjani N, et al. The association between COVID-19 incidence and mortality with socioeconomic development-A global ecological study. *BMC Public Health.* 2020. doi: [10.21203/rs.3.rs-117325/v1](https://doi.org/10.21203/rs.3.rs-117325/v1).
- Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: A web-based cross-sectional survey. *Psychiatry Res.* 2020;**288**:112954. doi: [10.1016/j.psychres.2020.112954](https://doi.org/10.1016/j.psychres.2020.112954). [PubMed: [32325383](https://pubmed.ncbi.nlm.nih.gov/32325383/)]. [PubMed Central: [PMC7152913](https://pubmed.ncbi.nlm.nih.gov/PMC7152913/)].
- World Health Organization. *Coronavirus disease (COVID-19) outbreak situation.* Geneva, Switzerland: WHO; 2020. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>.
- Statista. *Number of coronavirus (COVID-19) cases worldwide as of May 15, 2020, by country.* USA: Statista; 2020, [updated 8 Feb 2021; cited May 15]. Available from: <https://www.statista.com/statistics/1043366/novel-coronavirus-2019ncov-cases-worldwide-by-country/>.
- Kim SW, Su KP. Using psychoneuroimmunity against COVID-19. *Brain Behav Immun.* 2020;**87**:4-5. doi: [10.1016/j.bbi.2020.03.025](https://doi.org/10.1016/j.bbi.2020.03.025). [PubMed: [32234338](https://pubmed.ncbi.nlm.nih.gov/32234338/)]. [PubMed Central: [PMC7194899](https://pubmed.ncbi.nlm.nih.gov/PMC7194899/)].
- Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet.* 2020;**395**(10227):912-20. doi: [10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8). [PubMed: [32112714](https://pubmed.ncbi.nlm.nih.gov/32112714/)]. [PubMed Central: [PMC7158942](https://pubmed.ncbi.nlm.nih.gov/PMC7158942/)].
- Sood S. Psychological effects of the Coronavirus disease-2019 pandemic. *Res Humanit Med.* 2020;**7**(11):23-6.
- Rico-Uribe LA, Caballero FF, Martin-Maria N, Cabello M, Ayuso-Mateos JL, Miret M. Association of loneliness with all-cause mortality: A meta-analysis. *PLoS One.* 2018;**13**(1): e0190033. doi: [10.1371/journal.pone.0190033](https://doi.org/10.1371/journal.pone.0190033). [PubMed: [29300743](https://pubmed.ncbi.nlm.nih.gov/29300743/)]. [PubMed Central: [PMC5754055](https://pubmed.ncbi.nlm.nih.gov/PMC5754055/)].
- Filip I, Tidman M, Saheba N, Bennett H, Wick B, Rouse N, et al. Public health burden of sleep disorders: Underreported problem. *J. Public Health.* 2016;**25**(3):243-8. doi: [10.1007/s10389-016-0781-0](https://doi.org/10.1007/s10389-016-0781-0).
- Li N, Xu G, Chen G, Zheng X. Sleep quality among Chinese elderly people: A population-based study. *Arch Gerontol Geriatr.* 2020;**87**:103968. doi: [10.1016/j.archger.2019.103968](https://doi.org/10.1016/j.archger.2019.103968). [PubMed: [31751901](https://pubmed.ncbi.nlm.nih.gov/31751901/)].
- Bryant PA, Trinder J, Curtis N. Sick and tired: Does sleep have a vital role in the immune system? *Nat Rev Immunol.* 2004;**4**(6):457-67. doi: [10.1038/nri1369](https://doi.org/10.1038/nri1369). [PubMed: [15173834](https://pubmed.ncbi.nlm.nih.gov/15173834/)].
- Taylor DJ, Lichstein KL, Durrence HH. Insomnia as a health risk factor. *Behav Sleep Med.* 2003;**1**(4):227-47. doi: [10.1207/S15402010BSM0104_5](https://doi.org/10.1207/S15402010BSM0104_5). [PubMed: [15600216](https://pubmed.ncbi.nlm.nih.gov/15600216/)].
- Sadock BJ, Sadock VA, Kaplan HI. *Kaplan & Sadock's synopsis of psychiatry: Behavioral sciences/clinical psychiatry.* 9th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2003.
- Zhang R, Hou T, Kong X, Wang G, Wang H, Xu S, et al. Effects of region, epidemic stage, and demographic characteristics on sleep quality and mental disturbances among health care workers during COVID-19 outbreak. *Acta Psychiatr Scand.* 2020;**142**(3):249-56. doi: [10.21203/rs.3.rs-23260/v1](https://doi.org/10.21203/rs.3.rs-23260/v1).

17. Hyland P, Shevlin M, McBride O, Murphy J, Karatzias T, Bentall RP, et al. Anxiety and depression in the Republic of Ireland during the COVID-19 pandemic. *Acta Psychiatr Scand*. 2020;**142**(3):249–56. doi: [10.1111/acps.13219](https://doi.org/10.1111/acps.13219). [PubMed: [32716520](https://pubmed.ncbi.nlm.nih.gov/32716520/)].
18. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;**28**(2):193–213. doi: [10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4). [PubMed: [2748771](https://pubmed.ncbi.nlm.nih.gov/2748771/)].
19. Gupta R, Yick A. Validation of CES-D scale for older Chinese immigrants. *Aging Ment Health*. 2001;**7**(2):257–72.
20. Malakouti SK, Pachana NA, Najji B, Kahani S, Saeedkhani M. Reliability, validity and factor structure of the CES-D in Iranian elderly. *Asian J Psychiatr*. 2015;**18**:86–90. doi: [10.1016/j.ajp.2015.08.007](https://doi.org/10.1016/j.ajp.2015.08.007). [PubMed: [26442988](https://pubmed.ncbi.nlm.nih.gov/26442988/)].
21. Farrahi Moghaddam J, Nakhaee N, Sheibani V, Garrusi B, Amirkaifi A. Reliability and validity of the Persian version of the Pittsburgh Sleep Quality Index (PSQI-P). *Sleep Breath*. 2012;**16**(1):79–82. doi: [10.1007/s11325-010-0478-5](https://doi.org/10.1007/s11325-010-0478-5). [PubMed: [21614577](https://pubmed.ncbi.nlm.nih.gov/21614577/)].
22. Essau CA, Olaya B, Pasha G, Gilvarry C, Bray D. Depressive symptoms among children and adolescents in Iran: a confirmatory factor analytic study of the centre for epidemiological studies depression scale for children. *Child Psychiatry Hum Dev*. 2013;**44**(1):123–36. doi: [10.1007/s10578-012-0314-1](https://doi.org/10.1007/s10578-012-0314-1). [PubMed: [22714666](https://pubmed.ncbi.nlm.nih.gov/22714666/)].
23. Sheng X, Liu F, Zhou J, Liao R. [Psychological status and sleep quality of nursing interns during the outbreak of COVID-19]. *Nan Fang Yi Ke Da Xue Xue Bao*. 2020;**40**(3):346–50. Chinese. doi: [10.12122/j.issn.1673-4254.2020.03.09](https://doi.org/10.12122/j.issn.1673-4254.2020.03.09). [PubMed: [32376580](https://pubmed.ncbi.nlm.nih.gov/32376580/)]. [PubMed Central: [PMC7167307](https://pubmed.ncbi.nlm.nih.gov/PMC7167307/)].
24. Gulia KK, Kumar VM. Importance of sleep for health and well-being amidst COVID-19 pandemic. *Sleep Vigil*. 2020;**4**:49–50. doi: [10.1007/s41782-020-00087-4](https://doi.org/10.1007/s41782-020-00087-4). [PubMed: [32368715](https://pubmed.ncbi.nlm.nih.gov/32368715/)]. [PubMed Central: [PMC7197235](https://pubmed.ncbi.nlm.nih.gov/PMC7197235/)].
25. CDC. *Sleep and sleep disorders*. USA: CDC; 2020, [updated April 15, 2020]. Available from: <https://www.cdc.gov/sleep/index.html>.
26. Ballard ED, Vande Voort JL, Bernert RA, Luckenbaugh DA, Richards EM, Niciu MJ, et al. Nocturnal wakefulness is associated with next-day suicidal ideation in major depressive disorder and bipolar disorder. *J Clin Psychiatry*. 2016;**77**(6):825–31. doi: [10.4088/JCP.15m09943](https://doi.org/10.4088/JCP.15m09943). [PubMed: [27337418](https://pubmed.ncbi.nlm.nih.gov/27337418/)]. [PubMed Central: [PMC5103284](https://pubmed.ncbi.nlm.nih.gov/PMC5103284/)].
27. CDC. *Are you getting enough sleep?* USA: CDC; 2020, [updated April 20, 2020]. Available from: <https://www.cdc.gov/sleep/features/getting-enough-sleep.html>.
28. Kelsoe JR, Greenwood TA, Akiskal HS, Akiskal KK. The genetic basis of affective temperament and the bipolar spectrum. *Int Clin Psychopharmacol*. 2012;**28**:e5–6. doi: [10.1097/01.yic.0000423231.81230.9e](https://doi.org/10.1097/01.yic.0000423231.81230.9e).
29. Astill RG, Van der Heijden KB, Van Ijzendoorn MH, Van Someren EJ. Sleep, cognition, and behavioral problems in school-age children: A century of research meta-analyzed. *Psychol Bull*. 2012;**138**(6):1109–38. doi: [10.1037/a0028204](https://doi.org/10.1037/a0028204). [PubMed: [22545685](https://pubmed.ncbi.nlm.nih.gov/22545685/)].
30. Taghizadeh F, Hassannia L, Moosazadeh M, Zarghami M, Taghizadeh H, Dooki AF, et al. Anxiety and depression in health workers and general population during COVID-19 epidemic in Iran: A web-based cross-sectional study. *medRxiv*. 2020. doi: [10.1101/2020.05.05.20089292](https://doi.org/10.1101/2020.05.05.20089292).
31. Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global health concern. *Lancet*. 2020;**395**(10223):470–3. doi: [10.1016/S0140-6736\(20\)30185-9](https://doi.org/10.1016/S0140-6736(20)30185-9). [PubMed: [31986257](https://pubmed.ncbi.nlm.nih.gov/31986257/)]. [PubMed Central: [PMC7135038](https://pubmed.ncbi.nlm.nih.gov/PMC7135038/)].
32. Liu X, Liu J, Zhong X. Psychological state of college students during COVID-19 epidemic. *SSRN*. 2020. doi: [10.2139/ssrn.3552814](https://doi.org/10.2139/ssrn.3552814).
33. Sigdel A, Bista A, Bhattarai N, Pun BC, Giri G, Marqusee H, et al. Depression, anxiety and depression-anxiety comorbidity amid COVID-19 pandemic: An online survey conducted during lockdown in Nepal. *medRxiv*. 2020. doi: [10.1101/2020.04.30.20086926](https://doi.org/10.1101/2020.04.30.20086926).
34. Rajkumar RP. COVID-19 and mental health: A review of the existing literature. *Asian J Psychiatr*. 2020;**52**:102066. doi: [10.1016/j.ajp.2020.102066](https://doi.org/10.1016/j.ajp.2020.102066). [PubMed: [32302935](https://pubmed.ncbi.nlm.nih.gov/32302935/)]. [PubMed Central: [PMC7151415](https://pubmed.ncbi.nlm.nih.gov/PMC7151415/)].
35. Tahir MB, Masood A. The COVID-19 outbreak: Other parallel problems. *SSRN Electronic Journal*. 2020. doi: [10.2139/ssrn.3572258](https://doi.org/10.2139/ssrn.3572258).