



## Research Article

## Molecular identification of insecticide degradation by gut bacteria isolated from *Helicoverpa armigera* of Cotton plants

**Madhusudan S\***

Department of Biotechnology, Maharani Cluster University, Bangalore- 560001 (Karnataka), India

**S.K. Jalali**

Division of Molecular Entomology, National Bureau of Agriculturally Important Insect Resources (ICAR), Bangalore - 560043 (Karnataka), India

**Sibi G**

Department of Biotechnology, Indian Academy Degree College-Autonomous, Bangalore- 560043 (Karnataka), India

\*Corresponding author. Email: [mailmadhubiotech@gmail.com](mailto:mailmadhubiotech@gmail.com)

**Article Info**

<https://doi.org/10.31018/jans.v13i2.2678>

Received: April 14, 2021

Revised: May 27, 2021

Accepted: May 29, 2021

**How to Cite**

Madhusudan, S. *et al.* (2021). Molecular identification of insecticide degradation by gut bacteria isolated from *Helicoverpa armigera* of Cotton plants. *Journal of Applied and Natural Science*, 13(2), 641 - 653. <https://doi.org/10.31018/jans.v13i2.2678>

**Abstract**

The cotton bollworm *Helicoverpa armigera* occurs as a major pest in many economically important crops, including cotton, pigeon pea, chickpea, pea, cowpea, sunflower, tomato, sorghum, pearl millet and other crops. Intestinal microorganisms play important role in the degradation of diet components of insects. In order to know the role of gut bacteria in insecticide resistance five insecticides Chlorpyrifos (20% EC), Cypermethrin (25% EC), Malathion (50% EC), Quinalphos (25% EC), Triazophos (40% EC), were selected for the insecticide degradation studies. All the bacterial isolates from the gut of lab and field populations of *H. armigera* were identified using 16S rRNA gene-based identification and tested for their growth on minimal salt medium (MSM) along with the selected insecticides. A total of 11 bacterial isolates were tested and among them, isolate CL4 (*Rhodococcus* sp.) was found to grow on minimal salt medium (MSM) and with chlorpyrifos and isolate CL2 (*Enterococcus casseliflavus*) was able to grow in MSM with chlorpyrifos (C<sub>22</sub>H<sub>19</sub>C<sub>12</sub>NO<sub>3</sub>) and malathion (C<sub>10</sub>H<sub>19</sub>O<sub>6</sub>PS<sub>2</sub>) and no growth was seen in MSM without insecticide (control). Gas Chromatography analysis of the positive bacterial isolate cultures in MSM showed that the isolate CL4 (*Rhodococcus* sp.) was able to utilize 43.9% of chlorpyrifos and isolate CL2 (*E. casseliflavus*) was able to utilize 26% of chlorpyrifos and 57.1% of malathion in MSM broth cultures with comparison with the respective control cultures. Findings of the current work suggested that gut bacteria in the field populations of *H. armigera* plays a role in insecticide resistance.

**Keywords:** Cotton plants, Gut bacteria, *Helicoverpa armigera*, Insecticide degradation, Molecular identification

**INTRODUCTION**

Old-World bollworm (*Helicoverpa armigera*) has attained the status of national pest owing to its devastating nature on commercial crops in India. The cotton bollworm *H. armigera* occurs as a major pest in many economically important crops, including cotton, pigeon pea, chickpea, pea, cowpea, sunflower, tomato, sorghum, pearl millet and other crops. The cotton grown in over 5% of the total cultivated area has consumed more than 55% of the total amount of pesticide used in India (Kranthi *et al.* 2001)

A robust complement of prokaryotes is harbored in in-

sects' alimentary canals, facilitating nutrient availability, utilization, and detoxification of environmental toxins. Intestinal microorganisms play important role in the degradation of diet components of insects (Hayashi *et al.*, 2007). However, the indigenous gut bacteria also play a role in withstanding and colonization of the gut by non-indigenous species including pathogens (Dillon and Dillon, 2004). Precious little is known about the microbial community inhabiting the guts of various higher organisms, though it is acknowledged that microbes play an important role in metabolism, growth, development and protection of their host. Lepidoptera species feed on plants and the biomass digestion is

aided by gut associated microbes (Rajagopal *et al.*, 2002). However, the diversity of insect gut microflora is not revealed well and understanding the role of those microflora during the insect growth and development would help in effective pest management. *H. armigera* harbors diverse bacterial communities in its gut regions (Dar *et al.*, 2021). Recently, *Bacillus* sp were isolated from gut region of *H. armigera* by culture dependent methods (Shinde *et al.*, 2019).

Detoxification of the environmental pollutants, pesticide detoxification in soil and other ecosystems by microorganisms have been reported earlier. Biological removal of chemo pollutants becomes the method of choice since microorganisms can use a variety of xenobiotic compounds including pesticide for their growth and mineralize and detoxify those (Kanekar *et al.*, 2004). Microbial degradation, metabolites of degradation, enzymatic degradation and genetic basis of degradation of widely used organo phosphorous insecticides, like parathion, methyl parathion, malathion, monocrotophos and dimethoate are studied extensively. Insect gut microbes, especially insecticide-resistant pest, might play a role in detoxifying the insecticides. Microbial degradation of organo phosphorous insecticides have been reported in soil and consortia. Pure culture studies have shown cleavage of molecules and formation of different metabolites such as p-nitrophenol, diethylthiophosphoric acid, p-aminophenol, dimethyl phosphate. Verma *et al.* (2006) reported the degradation of endosulfan by a *Rhodococcus* strain isolated from gut of earthworm. Bacteria belonging to genera *Bacilli*, *Proteus*, *Micrococcus*, *Pseudomonas* and *Klebsiella* isolated from the gut of rice hispa, *Diadisa armigera* (Olivier) were shown resistant to 1000 ppm endosulfan and to some extent chloropyrifos and quinalphos (Thakur *et al.*, 2005). Carboxylesterases isolated from *H. armigera* play a role in pyrethroids detoxification (Bai *et al.*, 2019). *H. armigera* larva fed with laccases from *Yersinia enterocolitica* showed significant expression of enzymes and proteins for xenobiotic degradation (Ahlawat *et al.*, 2020). Chemical insecticides are the most widely released xenobiotics by humankind into the environment, resulting in very high contamination of food and water resources with these compounds. Many insects and bacteria have developed resistance to these compounds by evolving enzymes to metabolize these xenobiotics. In-depth research is required to identify potential candidates with respect to gut microbes in the degradation of insecticides/xenobiotics. This study was carried out to determine the influence of gut microflora isolated from *H. armigera* in insecticidal resistance.

## MATERIALS AND METHODS

### Collection of field larvae

The culture of the *H. armigera* was initiated by collect-

ing about 100 - 150 larvae on cotton (*Gossypium hirsutum* L., Kanchana variety) from the farmers' fields in Vikarabad (Andhra Pradesh-Lat-17°19'60N; Lon-77°55'0E), where insecticides were used frequently. On cotton, efforts were made to collect the larvae from non-Bt cotton, which was sprayed both for lepidopteran and sucking pests. Fourth- and fifth-instar larvae were collected in the plastic container with an artificial diet and brought to the laboratory and were used for the study. The insecticide use pattern was recorded at each place as per the protocols of the National Bureau of Agriculturally Important Resources, ICAR, Bangalore.

### Rearing of field and laboratory populations of *H. armigera*

The collected larvae were brought to the National Bureau of Agriculturally Important Insects (NBAII) laboratory, Bangalore. The larvae were transferred onto a freshly prepared semi synthetic diet (Nagarkatti and Satyaprakash, 1974). After pupation, the pupae were collected, surface sterilized in sodium hypochlorite (0.1%) and washed in distilled water. The washed pupae were air-dried at room temperature and kept in plastic jars for the emergence of moth. From the emerged moths, 25 pairs were transferred to each oviposition cage lined with muslin cloth and covered with a black cotton cloth. Two balls of absorbent cotton, one with water and another one with 50% honey solution, were provided in the oviposition cage. The eggs were collected from the cages every 24 h were washed in sodium hypochlorite (0.1%) and formaldehyde (1%) solution, filtered through a black cloth and tagged onto the plastic container (7 cm dia., 4 cm height) for hatching at 27±2°C and 70% RH.

The newly hatched neonates were transferred to 128 well diet trays containing solidified diet at 1 larva well<sup>-1</sup> using a fine hairbrush and were covered with pull-n-peel tabs. Both diet trays and neonates transfer were carried out on a laminar airflow clean bench followed by maintenance in a BOD incubator maintained at 27±2°C and 14:10 h light-dark. Fifty pupae of the susceptible strain of the *H. armigera* were also obtained from the Mass Production Unit of NBAIR, Bangalore, where it is reared on semi-synthetic diet for 10 years without replacing or adding any field population, hence was considered true susceptible populations. These pupae were also reared till the emergence of the adults and thus, the culture was further multiplied to be used in the bioassay studies with the same method as that of the field collected *H. armigera*. A homogenous stock of fifth instar larvae (10-11 days old) was for isolating the gut bacteria.

### Isolation of gut bacteria from *H. armigera*

For the isolation of gut bacteria, ten larvae of *H. armigera*

*gera* were immobilized in chloroform and sterilized in sodium hypochlorite (0.1%) and ethanol (70%) solution for 5 seconds to remove the contaminants (Gebbari *et al.*, 2001). The sterilized larvae were dissected under aseptic conditions in a Laminar air flow hood to remove the entire gut. The gut was placed in a micro tube containing 500 µl of sterile peptone water, crushed mechanically and vortexed thoroughly. A hundred µl of the gut homogenate was placed on sterile nutrient agar and nutrient glucose agar plates in replicates and incubated at 30°C for 48 h. The colonies obtained after 48 h incubation in the plates were further screened for colony morphology.

### Bacterial DNA isolation

Bacterial genomic DNA was isolated as per the standard protocol (Hoffman and Winston 1987). A single colony from each of the bacterial cultures was inoculated in nutrient broth and grown for 48 h at 30°C. The cells were harvested by centrifugation and added with 100 µl of lysozyme followed by incubation at room temperature for 30 mins to disrupt the cell membrane. The cells were lysed by the addition of lysis buffer (70 µl) containing guanidium isothiocyanate, SDS and tris-EDTA. The contents were mixed by inverting and 700 µl of isopropanol was added from the top. The two layers were gently mixed until the DNA strands were visible and the addition of ice cold ethanol precipitated the extracted DNA. The DNA was resuspended in 100 µl of TE buffer and the quality of DNA was checked in agarose gel electrophoresis. The purified genomic DNA was used as template for 16S rRNA gene amplification.

### Oligonucleotide primers

16S rRNA gene primers were procured from Aristogene Biosciences (P) Ltd, Bangalore. The oligonucleotides were reconstituted to 100 ng/µl stocks in sterile TE buffer. The primers were used at a working concentration 100 ng/µl in sterile filtered distilled water. The sequences of the primers were as follows

Forward primer – 5'-ACTCCTACGGGAGGCAGCAG-3'

Reverse primer – 5'-ATTACCGCGGCTGCTGG-3'

### Amplification of 16S rRNA gene by PCR

Gradient PCR was used to determine the optimum annealing temperature. The reaction mixture contained 100 ng of the extracted DNA, 1X PCR assay buffer (250 mM Tris-HCl, 10 mM KCl, 1.5 mM MgCl<sub>2</sub>), 100 mM dNTP's, 100 ng/µL each of forward and reverse primers, 1 unit of *Taq* DNA polymerase (Sigma Aldrich, India). PCR performed with forward and reverse primers involved an initial denaturation for 2 mins at 94°C, followed by 30 cycles of 94°C denaturation for 1 min, 58°C for annealing for 30 s and extension at 72 °C for 1 min 30 s. Finally, the reactions were heated at 72°C for 5 min. Specific and optimum amplification of the gene

was seen at 58 °C of annealing temperature. Subsequently, the gene was amplified at 58°C and the amplified PCR product (1.5 Kb) was purified from low melting agarose gel as per the standard protocols (Sambrook *et al.* 2001) for further sequencing at Aristogene Biosciences (P) Ltd, Bangalore. The obtained sequences were analyzed through BLAST search and the bacterial species were identified.

### Media preparation for insecticide degradation studies

The bacterial colonies isolated were tested for the selected eight commercial insecticides against chlorpyrifos, cypermethrin, malathion, quinalphos and triazophos, the most commonly used insecticides against *H. armigera* at a concentration range of 0.2, 0.4, 0.6, 0.8 and 1.0 ml/l, respectively. Initial studies were carried out on the bacterial isolates for their growth on Minimal Salt Medium (MSM) with and without insecticides. The composition of MSM included KH<sub>2</sub>PO<sub>4</sub> 1.0 g, K<sub>2</sub>HPO<sub>4</sub> 1.0 g, NH<sub>4</sub>NO<sub>3</sub> 1.0 g, MgSO<sub>4</sub>.7H<sub>2</sub>O 0.2 g, CaCO<sub>3</sub> 0.02 g, FeSO<sub>4</sub> 0.01g in 1 L of distilled water (pH 6.5). The isolated bacteria from the guts of lab and field larval populations *H. armigera* were purified from nutrient agar plated in duplicates and incubated at room temperature (28 ± 1°C) for 48 h. Isolated colonies were picked and streaked onto nutrient agar plates to obtain pure cultures. All the pure bacterial isolates were tested for their growth on MSM with and without insecticides. Different concentration of selected insecticides as mentioned above was added into sterile MSM (agar-based) and poured into Petri plates in Laminar airflow. The Petri dishes were labelled according to the concentration of selected insecticides after solidification. Similarly, MSM plates were also prepared without insecticides. All the bacterial isolates were streaked onto these plates in aseptic conditions and were incubated at room temperature (28 ± 1°C) for 48 h. Growth of the bacterial isolates on all the plates were recorded after 48 h. Bacterial isolates growing in a particular concentration of insecticide was isolated and subcultured on nutrient agar to obtain a pure culture. Positive cultures for growth on MSM with insecticide (0.2 ml/l) were further used for degradation studies. A series of 250 ml flask containing 50 ml of MSM (broth) supplemented with above mentioned insecticides at 0.2 ml/l concentration was incubated separately with 1 ml overnight grown positive cultures and incubated on a shaker 200 rpm at 30°C for 10 days. Controls for each bacterium were maintained without insecticides and positive control of MSM with insecticides alone was also maintained. After 10 days incubation CFU count was recorded on nutrient agar plates. The cell mass in MSM with and without insecticides was separated by centrifugation at 8000 xg for 10 min and supernatant of 50 ml was collected separately in the fresh tubes. The super-

natant was further analyzed in GC to detect the concentration of insecticides at the GC-MS facility available at Bangalore Test House, Bangalore.

### Gas Chromatography analysis of minimal salt media with insecticides

The samples containing insecticide and without insecticide were extracted with methylene chloride solvent, initially using 100 ml solution initially. Later the volume of 50 ml twice was used to partition the sample. The pooled extract was collected in the separate round bottom flask after passing through anhydrous sodium sulphate. The content was evaporated to dryness using a vacuum rotary evaporator and a volume of 2 ml of acetone was added to the residue. One µl of this solution was injected into GC-ECD.

### GC Conditions

Instrument: Agilent 6890N GC with 975 inert MSD  
 Software: MSD chemstation D.02.00.275 with NIST Library Ver.2.0d  
 Column: DB-35 MSI, 30 m X 0.25 mm X 250 micron  
 Inlet Temperature: 250°C  
 Injection Mode: Split less  
 Injection Volume: 1 µl  
 Carrier gas flow: Helium (constant flow) 1.0 ml/min  
 Oven program: 70°C for 2 min, @ 4°C/min, 160°C for 5 min, 4°C/min  
 Final temp.: 295°C for 10 min.  
 Transfer line temp.: 280°C  
 The concentration of insecticides in MSM broth cultures was calculated using the formula

$$\frac{\text{Sample peak area} \times \text{Std. Conc. } (\mu\text{g/ml}) \times \text{dilution}}{\text{Standard peak area}} \dots\dots \text{Eq. 1}$$

Working std. conc. (µg/ml): 1.2 µg/ml  
 Dilution (ml): 13 µl of insecticide formulation was diluted to 25 ml using acetone and 1.0 µl was injected into GC-ECD instrument.

## RESULTS AND DISCUSSION

### Isolation of gut bacterial flora from the insecticide resistance larval populations collected from cotton and susceptible lab populations of *H. armigera*

Gut bacteria isolated from the gut of insecticide-resistant fourth and fifth instar larvae of *H. armigera* collected from the cotton fields of Vikarabad, Andhra Pradesh were coded as CL1, CL2, CL3, CL4, CL5, CL6 and CL7. Gram staining of the bacterial isolates showed that isolates CL1, CL3 were Gram +ve rods, isolates CL2, CL4, CL6 were Gram +ve cocci and isolates CL5, CL7 were Gram -ve rods. Gut bacteria was isolated from the gut of susceptible fourth and fifth in-

star larvae of *H. armigera* collected from the mass production unit of NBAIL, Bangalore was coded as HL1, HL2, HL3, HL4, HL5 and HL6. Gram staining of the bacterial isolates showed that the isolates HL1, HL3, HL5, HL6 were Gram - ve rods and isolates HL2, HL4 were Gram +ve rods.

### Isolation of DNA and 16s rRNA PCR amplification of gut bacteria of insecticide-resistant and susceptible larvae of *H. armigera*


DNA was isolated from eleven bacterial isolates from the gut of insecticide-resistant field larvae of *H. Armigera*. The presence of genomic DNA from all the seven isolates was confirmed on a 0.8% agarose gel stained with ethidium bromide (Fig. 1). An intense single band was seen in all seven wells along with the DNA marker. The extracted DNA was used as a template for amplification of 16S rRNA gene. The primers selected were specific. Initial standardisation by gradient PCR has facilitated the specific amplification as observed by high intense band. The optimum annealing temperature was found to be 58°C and an intense single band of size approximately 1.5 kb was visible on 1% agarose gel stained with ethidium bromide (Fig. 2) in all the seven wells. Similarly, DNA was isolated from six bacterial isolates from the gut of susceptible larvae of *H. armigera*. The presence of genomic DNA from all the 6 isolates was confirmed on a 0.8% agarose gel stained with ethidium bromide (Fig. 3). An intense single band was seen in all six wells along with the DNA marker. The extracted DNA was used as the template for amplification of 16S rRNA gene. The primers selected were specific. The optimum annealing temperature was found to be 58°C. An intense single band of size approximately 1.5 kb was visible on 1% agarose gel stained with ethidium bromide (Fig. 4) in all 6 wells.

### Sequencing and sequence analysis of PCR amplified 16S rRNA gene of the eleven gut bacterial isolates


The partial sequence obtained from all the seven isolates ranged from 677, 660, 478, 459, 642, 514. 559 bp respectively in length and were analysed in BLASTn ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)) and the bacterial genera and species were determined. The partial 16S rRNAs sequence and the determined bacterial sp. along with the accession number (CL1-CL7) are shown in Table 1. The max identity of the sequence was 99-100%. The nucleotide sequences of seven isolates are submitted to NCBI-Gen Bank and the accession numbers were obtained. The determined bacterial communities were found to be **CL1-Bacillus pumulis**, **CL2-Enterococcus casseliflavus**, **CL3-Bacillus subtilis**, **CL4-Rhodococcus sp.**, **CL5-Pseudomonas sp.**, **CL6-Staphylococcus sp.**, **CL7-Pseudomonas aeruginosa**.

The partial sequence obtained from all the 6 isolates





**Fig 1.** Agarose gel electrophoresis of genomic DNA from seven bacterial isolates from gut of *H. armigera* from cotton crop. M- Standard DNA Marker, CL1 – CL7 – Genomic DNA of the bacterial isolates.



**Fig. 2.** Agarose gel electrophoresis of 16S rRNA PCR amplicon from seven bacterial isolates from gut of *H. armigera* from cotton crop. M- 0.1-3 kb Low range marker: sizes- 100bp, 200bp, 300bp, 600bp, 1kb, 1.5kb, 2 kb, 2.5kb, 3 kb., CL1 – CL7 – 16S rRNA PCR amplicon of the bacterial isolates.


from the gut of the susceptible larvae of *H. armigera* ranged from 664, 861, 779, 623, 541, 693 bp. in length. The partial 16S rRNAs sequence and the determined bacterial sp. along with the accession number (HL1-HL6) are shown in Table 2. The max identity of the sequence was 99-100%. The nucleotide sequences of 6 isolates were submitted to NCBI-Gen Bank and the accession numbers were obtained. The determined bacterial communities were found to be **HL1-*Proteus vulgaris***, **HL2-*Cellulosimicrobium cellulans***, **HL3-*Klebsiella oxytoca***, **HL4-*Bacillus subtilis***, **HL5-*Stenotrophomonas maltophilia***, and **HL6-*Pseudomonas* sp.**

**Insecticide degradation studies by the gut microbial isolates from susceptible lab and field populations of *H. armigera***


All the five insecticides were selected for the insecticide degradation studies. Growth of the bacterial isolates recorded after 7 days of incubation at 30°C is shown in Table 3. Among a total of 11 bacterial isolates tested, isolate CL4 (*Rhodococcus* sp.) was found to grow on minimal salt medium (MSM) and with chlorpyrifos and isolate CL2 (*Enterococcus casseliflavus*) was able to grow in MSM with chlorpyrifos and malathion and no growth was seen in MSM without insecticide (control). The growth of isolates CL4 and CL2 on MSM without insecticide (control) and with insecticides is shown in

Plate 1. Bacterial isolates CL4 and CL2 were further grown in MSM broth without and with insecticides and incubated in shaker at 30°C for 7 days, after incubation 0.1 ml of the culture was plated on to sterile nutrient agar plates and were incubated at 30°C for 48 h and the number of CFU was calculated along with control and data is shown in Table 4. GC graph results have been shown for MSM with insecticides, MSM with isolate CL4 without insecticide and with chlorpyrifos and MSM with isolate CL2 without insecticides and with chlorpyrifos and malathion in Figure 5, 6, and 7 respectively. The concentration of the insecticides in the MSM broth cultures along with the control are shown in Table 5. GC graph analysis showed that there was decrease in the concentrations of the insecticides in MSM broth cultures with insecticides when compared to that of MSM broth cultures without insecticides. The isolate CL4 (*Rhodococcus* sp.) was able to utilize 43.9% of chlorpyrifos and isolate CL2 (*Enterococcus casseliflavus*) was able to utilize 26% of chlorpyrifos and 57.1% of malathion in MSM broth cultures with comparison with the respective control cultures. Findings of the current results suggested that gut bacteria in the field populations of *H. armigera* is having a role in insecticide resistance.

Broderick et al., (2004) described the isolation and



**Fig. 3.** Agarose gel electrophoresis of genomic DNA from six bacterial isolates from the gut of larvae of laboratory populations of *H. armigera*. Lane M – Marker - Lambda/HindIII digest (sizes-23130, 9416, 6557, 4361, 2322, 2027, 564). Lane 1 – 6 – Genomic DNA from 6 bacterial isolates.



**Fig. 4.** Agarose gel electrophoresis of 16S rRNA PCR amplicon from six bacterial isolates from gut of the larvae of laboratory populations of *H. armigera*. Lane M- Marker - 0.1-3 kb Low range marker: sizes- 100bp, 200bp, 300bp, 600bp, 1kb, 1.5kb, 2 kb, 2.5kb, 3 kb., Lane - HL1 – HL6 – 16S rRNA PCR amplicon of the bacterial isolates, 1=HL1, 2=HL2, 3=HL3, 4=HL4, 5=HL5, 6=HL6.

**Table 1.** Partial 16S rRNA sequence and identified bacterial species with GenBank accession number of the seven bacterial isolates from the gut of insecticide resistance field larval populations of *H. armigera* collected from cotton crop.

Isolate	Partial 16S rRNA gene sequence	Size (bp)	Identified Bacteria By BLASTn	GenBank Accession Number
CL1	GTGCATTGCGGGTGTCTATACATGCAAGTCGAGCG- GACAGAAGGGAGCTTGCTCCCGGGTGTAGCGGCGGACGGGT GAGTAACACGTGGGTAACCTGCCTGTAAGACTGGGATAACTCC GGGAAACCGGAGCTAATACCGGATAGTTCCTTGAACCGCATGG TTCAAGGATGAAAGACGGTTTCGGCTGCACTTACAGATGGACC CGCGGCGCATTAGCTAGTTGGTGAGGTAACGGCTCACCAAGGC GACGATGCGTAGCCGACCTGAGAGGGTGATCGGCCACACTGG GACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTAGG GAATCTCCGCAATGGACGAAAGTCTGACGGAGCAACGCCGCG TGAGTGATGAAGGTTTTCCGGATCGTAAAGCTCTGTTGTTAGGGA AGAACAAGTGCAAGAGTAGCTGCTTGACCTTGACGGTACCTAA CCAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGTAATA CGTAGGTGGCAAGCGTTGTCGGAATTATTGGGCGTAAAGGGC TCGCAGGCGGTTTTCTTAAGTCTGATGTGAAAGCCCCCGGCTCAA CCGGGGAGGGTCAATTGAAACTGGAACTTGATTGCAGAAGAG GAGAGTGAATTCACGTGTAGCGGTGAAATGCGAAAAAA	677	<i>Bacillus pumilis</i>	HQ651050
CL2	GATTATGGCTCAGGACGAACGCTGGCGGCGTGCG- TAATACATGCAAG TCGAACGCTTTTTCTTTCACCG- GAGCNTGCTCCANCGNAAGAAAAAGAGTGGCGAACGGGTGAGT AACACGTGGGTAACTGCCATCAGAAGGGGATAACACTTGGAA AACAGGTGCTAATACCGTATAACACTATTTTCCGCATGGAAGAA AGTTGAAAGGCGCTTTTGCCTCACTGATGGATGGACCCGCGGT GCATTAGCTAGTTGGTGAGGTAACGGCTACCAAGGCAACGAT GCATAGCCGACCTGAGAGGGTGATCGGCCACACTGGGACTGAG ACACGGCCACAGACTCCTACGGGAGGCAGCAGTAGGGAATCTTC GGCAATGGACGAAAGTCTGACCGAGCAACGCCGCGTGAGTGAA GAAGGTTTTCCGGATCGTAAACTCTGTTGTTAGAGAAGAACAAG GATGAGAGTAAATGTTTCATCCCTTGACGGTATCTAACAGAAA GCCACGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGT GGCAAGCGTTGTCGGATTTATTGGGCGTAAAGCGAGCGCAGG CGGTATCTTAAGTCTGATGTGAAAGCCCCCGGCTCAACCGGGG AGGGTCATTGAAACTGGGAGACTTGAGT	660	<i>Enterococcus cas- seliflavus</i>	HQ651051
CL3	TGCAGTCGAGCGGACAGATGGGAGCTT- GCTCCCTGATGTTAGCGGCGGACGGGTGAGTAACACGTGGGTA ACCTGCCTGTAAAGACTGGGATAACTCCGGGAAACCGGGGCTAA TACCGGATGGTTGTTTGAACCGCATGGTTCAAACATAAAAGGTG GCTTCGGCTACCCTTACAGATGGACCCGCGGCGCATTAGCTA GTTGGTGAGGTAACGGCTACCAAGGCAACGATGCGTAGCCGA CCTGAGAGGGTGATCGGCCACACTGGGACTGAGACACGGCCC AGACTCCTACGGGAGGCAGCAGTAGGGAATCTCCGCAATGGA GAAAGTCTGACGGAGCAACGCCGCGTGAGTGATGAAGGTTTT CGGATCGTAAAGCTCTGTTGTTAGGGAAGAACAAGTACCGTTCC AATAGGGCGGTACCTTGACGGTACCTAACAGAAAGCCACGGC TAACTACGTGCCAGCAGCCG	478	<i>Bacillus subtilis</i>	JF266592
CL5	GGGCCGTGGGGGAGGACTACACATGCAAGTCGAGCG- GATGAAGGGAGCTTGCTCCTGGATTACGCGGCGGACGGGTGA GTAATGCCTAGGAATCTGCCTGGTAGTGGGGGATAACGTCCGG AAACGGGCGCTAATACCGCATACGTCTGAGGGAGAAAGTGGG GGATCTTCGGACCTCACGCTATCAGATGAGCCTAGGTGCGATTA GCTAGTTGGTGGGGTAAAGGCCTACCAAGGCGACGATCCGTAA CTGGTCTGAGAGGATGATCAGTCACACTGGAAGTACGACACGG TCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGGACAAT GGGCGAAAGCCTGATCCAGCCATGCCGCGTGTGTGAAGAAGGT CTTCGGATTGTAAGCACTTTAAGTTGGGAGGAAGGGCAGTAAG TTAATACCTTGCTGTTTTGACGTTACCAACAGAATAAGCACCGG CTAACTTCGTGCCAGCAGCCGCGGTAATACGAAGGGTGCAAGC GTTAATCGGAATTAAGTGGGCGTAAAGCGCGCGTAGGTGGTTCA GCAAGTTGGATGTGAAATCCCCGGGCTCAACCTGGGAACTGCA TCCAAAATACTGAGCTAGAGTACGGTAGAGGGTGGTGAAA T	642	<i>Pseudomonas sp.</i>	HQ651053

Contd.....


**Table.1.** Contd.....

<b>CL6</b>	<p>GTCTGCGGCATGCTATACATGCAGTCGAGCGAACAGACGAGGAGCTT- GCTCCTTTGACGTTAGCGGCGGACGGGTGAGTAACACGTAAGGTAACCT ACCTATAAGACTGGGATAAATTCGGGAAACCGGAGCTAATACCGGATAA TATTTGAAACCGCATGGTTTCGATAGTAAAAGATGGCTTTGCTATCACTTA TAGATGGACCTGCGCCGTATTAGCTAGTTGGTAAGGTAACGGCTTACCA AGGCAACGATACGTAGCCGACCTGAGAGGGTATCGGCCACACTGGAA CTGAGACACGGTCCAGACTCCTACGGGAGGCAGCAGTAGGGAACTCTTC CGCAATGGGCGAAAGCCTGACGGAGCAACGCCGCGTGAAGTGAAG GTCTTCGGATCGTAAAACCTCTGTTATTAGGGAAGAACAAACGTGTAAGTA ACTGTGCACGTCTTGACGGTACCTAATCAGAAAGCCACGGCTAACTACG TGCCAGCAGCCGCGGTAATACGTAGGTGG</p>	514	<i>Staphylococcus</i> sp.	HQ651054
<b>CL7</b>	<p>CTGTTCTGTCCGCGAGCTAGACATGCAAGTCGAGCG- GATGAAGGGAGCTTGCTCCTGGATTACGCGGCGGACGGGTGAGTAATG CCTAGGAATCTGCCTGGTAGTGGGGGATAACGTCCGAAACGGGCGCT AATACCGCATACTGCTCCTGAGGGAGAAAAGTGGGGGATCTTCGGACCTCA CGCTATCAGATGAGCCTAGGTCGGATTAGCTAGTTGGTGGGGTAAAGG CCTACCAAGGCGACGATCCGTAACCTGGTCTGAGAGGATGATCAGTCACA CTGGAAGTGAACACGGTCCAGACTCCTACGGGAGGCAGCAGTGGGGA ATATTGGACAATGGGCGAAGCCTTGATCCAGCCATGCCGCGTGTGTGAA GAAGTCTTCGGATTGTAACACTTTAAGTTGGGAGAAGGCAGAAGTTAA TACCTTGGTTGTTTTGACGTACCACAGAATAAGCACCGGCTACTTCTGC CGCAGCCGCGTATACCAAGGTGCAGCGTAATTCGAATTACTGGCGTAAG CGCGGAAGGGTTCACAGGTGGAAGGGAATCCCCGGCTC</p>	559	<i>Pseudomonas aeruginosa</i>	HQ651055

characterization of gut microflora from Lepidoptera and the present study isolated the gut bacterial flora of *H. armigera* using culture dependent methods. Bacterial isolates were identified using 16S rRNA gene based molecular technique. Different gut bacteria belonging to different genera were isolated and identified from field and lab larval populations of *H. armigera*. Some of the similar gut bacterial isolates have been also isolated from *H. armigera* species in India (Mishra and Tandon, 2003; Priya et al., 2012) in China (Xiang et al., 2006). Actinobacteria, proteobacteria and firmicutes were the dominant bacteria in *H. armigera* fed on tomato plants (Ranjith et al., 2016). In another study, high prevalence of Gammaproteobacteria followed by Bacilli and Actinobacteria were detected in the *H. armigera* midgut (Paramasiva et al., 2014). Dar et al., (2018) isolated *Klebsiella* sp, cellulolytic bacteria, from gut of *H. armigera* with xylanase and  $\beta$ -glucosidase activities. Apart from bacterial population, a novel picorna-like virus, named *Helicoverpa armigera* Nora virus was found in the gut of *H. armigera* moths and larvae (Yang et al.,

2019). Similarly, Ifla virus was isolated from *H. armigera* by Yuan et al. (2017). Higher animals like humans contain gut bacterial diversity in the range of 1500-2500. Further, the number of gut bacteria is ten times higher than the total number of host cells and the bacteria have more than ten times the metabolic diversity (Dillon and Dillon, 2004). It was also reported that sexual performance, mating preferences, and oviposition are influenced by the gut microflora (Gavriel et al., 2010; Sharon et al., 2010).

In order to decipher the non-pathogenic interaction between the bacteria and the host, we need first to study the gut flora of the host. In the present study, culture dependent studies showed that the field collected and the lab reared population showed significant differences in bacterial population, which is in agreement with the previous studies (Xiang et al., 2006). Rani et al., (2009) reported that the difference in lab reared and field insects is due to the fact that the laboratory raised insects are exposed to narrower range of food and environmental factors when compared to the field collect-



**Plate 1.** Growth of bacterial isolate CL4 and CL2 on MSM with insecticides. CL4 – *Rhodococcus* sp. CL2 - *Enterococcus casseliflavus*. A – Control, B - MSM with Chlorpyrifos, C – MSM with Malathion.

**Table 2.** Partial 16S rRNA sequence and identified bacterial species with GenBank accession number of the six bacterial isolates from the gut of susceptible laboratory larval populations of *H. armigera*.

Isolate	Partial 16S rRNA gene sequence	Size (bp)	Identified Bacteria By BLASTn	GenBank Accession Number
HL1	CATGCAGTCGAGCGGTAACAGGAGAAAGCTTGCTTTCTT- GCTGACGAGCGGGACGGGTGAGTAATGTATGGGGATCTGCC CGATAGAGGGGGATAACTACTGGAACCGGTGGCTAATACCGCAT GACGTCTACGGACCAAGCAGGGCTCTTCGGACCTTGGCCTAT CGGATGAACCCATATGGGATTAGCTAGTAGGTGAGTAATGGCT CACCTAGGCAACGATCTCTAGCTGGTCTGAGAGGATGATCAGCC ACACTGGGACTGAGACACGGCCAGACTCCTACGGGAGGCAGC AGTGGGGAATATTGCACAATGGGCGCAAGCCTGATGCAGCCATG CCGCGTGTATGAAGAAGGCCTTAGGGTTGTAAGTACTTTTCAGC GGGGAGGAAGGTGATAAAGTTAATACCTTTGTCAATTGACGTTAC CCGCAGAAAGACACCGGCTAAGTCCGTCGCCAGCAGCCGCGGT AATACGGAGGGTGAAGCGTTAATCGGAATTACTGGGCGTAAAG CGCAGCAGGCGGTCAATTAAGTCAGATGTGAAAGCCCCGAGCT TAACTTGGGAATTGCATCTGAAACTGGTTGGCTAGAGTCTTGATG AGGGGGGTAGAATCCACGTGTAGCGGTGAAATGCGTAGAGATG TGAGGAATAC	664	<i>Proteus vulgaris</i>	JF266593
HL2	CATGCAGTCGAACGATGATGCCAGCTTGCTGGGCGGATTAG- TGGCGAACGGGTGAGTAACACGTGAGTAACCTGCCCTGACTTC GGGATAACTCCGGGAAACCGGGGCTAATACCGGATATGAGCCG CCTTCGCATGGGGTGGTTGAAAGTTTTTCGGTCAGGGATGGG CTCGCGCCTATCAGCTTGTGGTGGGGTATGGCCTACCAAGG CGACGACGGGTAGCCGGCCTGAGAGGGCGACCGGCCACACTG GGACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTGGG GAATATTGCACAATGGGCGAAAGCCTGATGCAGCGACGCCGCGT GAGGGATGAAGGCCTTCGGGTTGTAACCTCTTTCAGCAGGGAA GAAGCGCAAGTGACGGTACCTGCAGAAAGCGCCGGCTAACTA CGTGCCAGCAGCCGCGGTAATACGTAGGGCGCAAGCGTTGTCC GGAATTATTGGGCGTAAAGAGCTCGTAGGCGGTTTTGTCGCGCT GGTGTGAAAACCTCGAGGCTCAACCTCGAGCTTGCATCGGGTACG GGCAGACTAGAGTGCGGTAGGGGAGACTGGATTTCTGGTGTAG CGGTGGAATGCGCAGATATCAGGAGGAACACCGATGGCGAAGG CAGGTCTCTGGGCCGCAACTGACGCTGAGGAGCGAAAGCATGG GGAGCGAACAGGATTAGATACCCTGGTTAGTCCATGCCGTAAC GTTTGGCACTAGGTGTGGGGGC	861	<i>Cellulosimicrobium cellulans</i>	JF266594
HL3	ACGCTGGCGGCAGGCCCTAACACATGCAAGTCGAACGGTGAG- CACARAAGAGCTTGCTCTCGGGTGACGAGTGGCGGACGGGTGA GTAATGTCTGGGAAACTGCCTGATGGAGGGGGATAACTACTGGA AACGGTAGCTAATACCGCATAACGTCCGAAGACCAAGAGGGGG ACCTTCGGGCCTCTTGCCATCAGATGTGCCAGATGGGATTAGC TAGTAGGTGGGGTAAACGGCTCACCTAGGCGACGATCCCTAGCTG GTCTGAGAGGATGACCAGCCACACTGGAACCTGAGACCGGTCCA GACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGC GCAAGCCTGATGCAGCCATGCCGCGTGTATGAAGAAGGCCTTCG GGTTGTAAAGTACTTTTCAGCGGGGAGGAAGCCGATAAGGTTAAT AACCTTGTGCGATTGACGTTACCCGCGAGAAGAAGCACCAGGCTAAC TCCGTGCCAGCAGCCGCGGTAATACGGAGGGTGCAAGCGTAAT CGGAATTACTGGGCGTAAAGCGCACGCAGGCGGTCTGTCAAGTC GGATGTGAAATCCCCGGGCTCAACCTGGGAAGTGCATTGAAAC TGGCAGGCTGGAGTCTTGTAGAGGGGGTAGAATTCAGGTGTA GCGGTGAAATGCGTAGAGATCTGAGGAATACCGGTGGCGAAG GCGGCCCTTGACAAAGACTGACGCTCAGGTGCGAAAGCGTG GGGAGCAAACAGGATTAGATACCCTGGTAGTCCACGCT	779	<i>Klebsiella oxytoca</i>	JF266595
HL4	TACATGCAGTCGAGCGGACAGATGGGAGCTT- GCTCCCTGATGTTAGCGGGCGACGGGTGAGTAACACGTGGGTAA CCTGCCTGTAAGACTGGGATAACTCCGGGAAACCGGGGCTAATA CCGGATGGTTGTTTGAACCGCATGGTTCAAACATAAAAGGTGGCT TCGGCTACCACTTACAGATGGACCCGCGGCGCATTAGCTAGTTG GTGAGGTAAACGGCTCACCAAGGCAACGATGCGTAGCCGACCTGA GAGGGTGTATCGGCCACACTGGGACTGAGACACGGCCAGACTC CTACGGGAGGCAGCAGTAGGGAATCTTCCGCAATGGACGAAAGT CTGACGGAGCAACCGCGCTGAGTGATGAAGTTTTTCGGATCGT AAAGCTCTGTTGTAGGGAAAGAACAAGTACCGTTTCAATAGGGC GGTACCTTGACGGTACCTAACAGAAAGCCACGGCTAACTACGT GCCAGCAGCCGCGTAATACGTAGGTGGCAAGCGTTGTCCGGA ATTATTGGGCGTAAAGGGCTCGCAGGCGGTTTTCTTAAGTCTGAT GTGAAAGCCCCGGCTCAACCGGGGAGGGTCATTGAAACTGG GAACTTGAGTGCAAGAGGA	623	<i>Bacillus subtilis</i>	JF266596

Contd.....



**Table 2.** Contd.....


HL5	ACATGCAAGTCGAGCGGCAGCACAGGAGAGCTT- GCTCTCTGGGTGGCGAGTGGCGGACGGGTGAGGAATACATCGG AATCTACTCTGTCGTGGGGGATAACGTAGGGAACTTACGCTAAT ACCGCATACGACCTACGGGTGAAAGCAGGGGACCTTCGGGCCT TGCGCGATTGAATGAGCCGATGTCGGATTATCTAGTTGGCGGGG TAAAGGCCACCAAGGCGACGATCCGTAGCTGGTCTGAGAGGAT GATCAGCCACACTGGAAGTCCGACTCCAGACTCCTACGGG AGGCAGCAGTGGGGAATATTGGACAATGGGCGCAAGCCTGATC CAGCCATACCGCGTGGGTGAAGAAGGCCCTTCGGGTTGTAAGCC CTTTTGTTGGGAAAGAAATCCAGCTGGCTAATACCCGGTTGGGAT GACGGTACCCAAAGAATAAGCACCGGCTAATTCGTGCCAGCAG CCGCGTAATACGAAGGGTGAAGCGTTACTCGGAATTACTGGG CGTACAGCGTGCCTAGGTGGTCTGTT	549	<i>Stenotropho- monas maltophilla</i>	JF266597
HL6	GTCGAGCGGCAGCACGGGTAAGTGTAC- CTGGTGGCGAGCGGCGGACGGGTGAGTAATGCCTAGGAATCTG CCTGGTAGTGGGGGATAACGCTCGGAAACGGACGCTAATACCG CATACGTCCTACGGGAGAAAGCAGGGGACCTTCGGGCCTTGCG CTATCAGTAGCCTAGGTCCGATTAGCTAGTTGGTGGGTAAT GGCTCACCAAGGCGACGATCCGTAAGTGGTCTGAGAGGATGATC AGTCACACTGGAAGTCCGACTCCAGACTCCTACGGGAGGC AGCAGTGGGGAATATTGGACAATGGGCGAAAGCCTGATCCAGCC ATGCCGCGTGTGTGAAGAAGGCTTCGGATTGTAAGCACTTTAA GTTGGGAGGAAGGCGAGTTACCTAATACGTATCTGTTTTGACGTT ACCGACAGAATAAGCACCGGTAAGTCTGTGCCAGCAGCCGCG GTAATACAGAGGGTGAAGCGTTAATCGGAATTACTGGGCGTAA AGCGCGCTAGGTGTTTTGTTAAGTTGAATGTGAAATCCCGGG CTCAACCTGGGAAGTGCATCCAAACTGGCAGGCTAGAGTATGG TAGAGGGTGGTGAATTTCTGTGTAGCGGTGAAATGCGTAGAT ATAGGAAGGAACACCAAGTGGCGAAGGCGACCACTGGACTGATA CTGACACT	693	<i>Pseudomonas sp.</i>	JF266602

ed counterparts. However, a considerable difference in types of colonizing species was observed between the insects raised on artificial diet and the field collected ones.


As previously described in other lepidopterans (Broderick et al., 2004; Xiang et al., 2006; Priya et al., 2012), the current study has also found *Enterococcus* in abundance when compared to rest of the phylotypes. The presence of *Enterococcus* in the gut benefits the host by reducing the gut pH thereby providing alkaline conditions that have a role in toxin effects (Wilson and Benoit, 1993). It was also identified that some of the gut microflora are involved in the degradation of polycyclic aromatic hydrocarbons (Lei et al., 2004) and insecticides (Hao et al., 2002). Other factors that influence the microbial diversity of the insect gut are host plant type and the host plant's geographical location. Therefore, artificial diet can therefore be considered a poor diet compared to field insects' diet which ultimately influ-

ences the gut microflora diversity. As mentioned earlier, crop leaf surface influences the gut bacterial flora of *Helicoverpa* larvae. The phyllosphere bacteria offers functional resistance to its host plant and influences the gut bacterial community of the insect larva feeding on it (Rajendran et al. 2007). Further analysis of these systems may identify as to what other factors contribute to the significant variation in gut bacteria among larvae from different crops from different locations.

In the present study, eleven gut bacterial isolates from the gut of larvae of the insecticide resistant field and susceptible lab populations of *H. armigera* were tested for their growth on minimal salt media with chlorpyrifos, cypermethrin, malathion, quinalphos, and triazophos along with the control MSM without insecticides. The result analysis reveals that the bacterial isolates *Rhodococcus* sp. and *Enterococcus casseliflavus* from gut of the field populations of *H. armigera* were able to grow on an MSM medium with chlorpyrifos



**Fig. 5.** GC analysis of MSM with Chlorpyrifos and Malathion, A- Chlorpyrifos B- Malathion



**Fig. 6.** GC analysis of MSM with Chlorpyrifos and bacterial isolate HT3, A- Chlorpyrifos B- Control.

**Table 3.** Growth of bacterial isolates on MSM medium with insecticides from the gut of *Helicoverpa armigera*.

Bacteri- al Iso- lates	Isolate Name	MSM	MSM + Chlorpyriphos	MSM + Cyper- methrin	MSM + Malathion	MSM + Qui- nalphos	MSM + Triazo- phos
CL1	<i>Bacillus pumillis</i>	-	-	-	-	-	-
CL2	<i>Enterococcus casseliflavus</i>	--	++	-	+	--	-
CL3	<i>Bacillus subtilis</i>	-	-	-	-	-	-
CL4	<i>Rhodococcus</i> sp.	-	++	-	-	-	-
CL5	<i>Pseudomonas</i> sp.	-	-	-	-	-	-
CL6	<i>Staphylococcus</i> sp.	-	-	-	-	-	-
CL7	<i>Pseudomonas aeruginosa</i>	-	-	-	-	-	-
HL1	<i>Proteus vulgaris</i>	-	-	-	-	-	-
HL2	<i>Cellulosimicrobium cellulans</i>	-	-	-	-	-	-
HL3	<i>Klebsiella oxytoca</i>	-	-	-	-	-	-
HL4	<i>Bacillus subtilis</i>	-	-	-	-	-	-
HL5	<i>Stenotrophomonas maltophilia</i>	-	-	-	-	-	-
HL6	<i>Pseudomonas</i> sp.	-	-	-	-	-	-

MSM: Minimum Salt Medium, +: Good growth, ++: Very good growth, -: No growth  
 CL1-CL2: Bacterial isolates from the larval gut of insecticide resistant field larvae of *H. armigera* from cotton. HL1-HL2: Bacterial iso-  
 lates from the larval gut of lab populations of *H. armigera*.

(C<sub>22</sub>H<sub>19</sub>Cl<sub>2</sub>NO<sub>3</sub>) and malathion (C<sub>10</sub>H<sub>19</sub>O<sub>6</sub>PS<sub>2</sub>). From the analysis of the Gas Chromatography, the isolate CL4 (*Enterococcus* sp.) was able to utilize 43.9% of chlorpyriphos and isolate CL2 (*Enterococcus casseliflavus*) was able to utilize 26% of chlorpyriphos and 57.1% of malathion in MSM broth cultures with comparison with the respective control cultures. This indicates that these bacteria can utilize the insecticides as carbon source and why these two bacteria can utilize chlorpyriphos and malathion and why not other insecticides is not clear until now and further studies have to be done in this line of research. Earlier reports have indicated the role of different species of *Enterobacteriaceae* in degradation of organophosphorous insecticides like chlorpyriphos (Singh *et al.*, 2004), phosphates (Lee *et al.*, 1992) and glyphosates (Dick and Quinn, 1995). Carboxylesterases isolated from *H. armigera* play a role in pyrethroids detoxification (Bai *et al.*, 2019). *H. armigera* larva fed with laccases from *Yersinia enterocolitica* showed significant expression of enzymes and proteins for xenobiotic degradation (Ahlawat *et al.*, 2020). Thakur *et al.* (2005) showed that the bacterial isolates from the whole gut of the rice hispa, *Diadisa armigera* (Olivier), belonging to genera *Bacillus*, *Proteus*, *Micrococcus*, *Pseudomonas* and *Klebsiella* were resistant to 1000 ppm endosulfan and to some extent chlorpyriphos and quinalphos. The GLC analysis showed that the isolated gut bacteria had the potential to degrade endosulfan. This work probably can be a support to the current research to show that insect gut bacteria are playing a role in insecticide resistance. The study shows the potential of symbiotic bacteria in imparting insecticide resistance to insect hosts. As organ-


ic phosphorous compounds are used so widely in agriculture for pest control, symbiont detoxification could represent a rapid and previously unappreciated mechanism for insecticide resistance in insects (Whalon *et al.*, 2008). Given the general detoxification ability of microbes and their ability to evolve quickly, they could provide a potent means for rapid acquisition of insecticide resistance in hosts. Many insects harbor a robust complement of prokaryotes in their alimentary canals,

**Table 4.** Colony forming units of the bacterial isolates (CL4 and CL2) on nutrient agar.

MSM broth culture	CFU/ml
Control	00
MSM + CL4	3 X 10 <sup>-4</sup>
MSM + CL4 + Chlorpyriphos	7 X 10 <sup>-4</sup>
MSM + CL2	1 X 10 <sup>-4</sup>
MSM + CL2 + Chlorpyriphos	3 X 10 <sup>-4</sup>
MSM + CL2 + Malathion	2 X 10 <sup>-4</sup>

**Table 5.** Concentration of insecticides in MSM broth cultures with bacterial isolates CL4 and CL2 analysed by GC-ECD.

MSM broth culture	Chlorpyriphos Conc. (µg/ml)	Malathion Conc. (µg/ml)
Control	00	00
MSM	38.29	112.27
MSM + CL4	16.81	---
MSM + CL2	10.05	64.13



**Fig. 7.** GC analysis of MSM with chlorpyrifos, malathion with bacterial isolate CL2 A- Chlorpyrifos B- Malathion C - Control.

which facilitate nutrient availability, utilization and detoxification of environmental toxins, degradation of insect diet components (Williams and Roans, 2006; Hayashi et al., 2007). Earlier reports have also shown that the gut bacterial isolates of other insect species play an important role in insect growth and development. Jing et al., (2020) described the roles of gut bacteria as essential nutrient provisioning, digestion and detoxification. *Bacillus thuringiensis* isolated from the gut region of *H. armigera* play a major role in inducing pathogenicity of Bt toxin in the host (Regode et al., 2016). Carol et al., (2003) reported that the gut bacterial isolate *Enterobacter agglomerans* from the apple maggot fly was able to degrade and detoxify Dihydrochalcone Phloridzin, a plant derived compound toxic to *Rhagoletis pomonella* (Walsh) (Diptera: Tephritidae). Vesta et al., (2006) reported that bark beetles *Ips tygraphus* (Coleoptera: Scolytidae) fed on conifers which produce myrcene (MR), among some other defensive compounds, six bacterial isolates from the gut of the bark beetles were most resistant to the bactericidal compound. 16S rRNA analysis of these bacteria showed that they are related to *Enterobacteriaceae*. Studies conducted by Genta et al. (2006) on antibiotic-treated and non-treated larvae of *Tenebrio molitor* suggested that microbial products play subtle roles in the life of the insect, digestion of refractory food, detoxification of secondary plant compounds and modify the volatile profiles of the insect host. Visotto et al., (2009) treated the bacterial colonies isolated from the gut homogenate of velvetbean caterpillar, *Anticarsia gemmatalis* with tetracycline. They found that the antibiotic was sensitive to the bacteria and inhibited the bacterial proteases. Their results suggested that gut bacteria may significantly contribute to lipid- and mainly protein-digestion in velvetbean caterpillars. Anand et al. (2010) reported that the bacterial species *Proteus vulgaris*, *Klebsiella pneumoniae*, *Escherichia coli*, *Citrobacter freundii*, *Serratia liquefaciens*, *Enterobacter* sp., *Pseudomonas fluorescens*, *P. aeruginosa*, *Aeromonas* sp., and *Erwinia* sp. isolated from the gut of

*Bombyx mori* L. were able to produce digestive enzymes that degrade the carbohydrates present in the mulberry leaves. In the current study, gut bacteria from laboratory populations of *H. armigera* and laboratory and field population of *T. chilonis* could not utilize the insecticides. It can be justified that they are not exposed to insecticides, whereas in the field populations one of the total isolated bacteria could utilize the insecticides for their growth. This might be because field populations are under constant stress of insecticides, which might induce the bacteria to produce enzymes to degrade and detoxify them. In field populations, only one per cent of the total bacterial isolates were able to utilize the insecticides and a possible explanation may be that microorganisms need an adaptation period to produce the necessary degradative enzymes (Jilani and Khan, 2004).

## Conclusion

*Enterococcus casseliflavus* and *Rhodococcus* sp isolated from the gut region of *H. armigera* were able to grow in the presence of selected insecticides. Concluding from the current study results, it can be stated that gut bacteria in insecticide-resistant field populations of *H. armigera* play a significant role in insecticide resistance of the host. Understanding the insect gut microbial diversity, their functional roles in insect growth and development would offer crucial information for designing future pest management strategies.

## ACKNOWLEDGEMENTS

The authors thank The Director of National Bureau of Agriculturally Important Insect Resources (NBAIIIR), Indian Council of Agricultural Research, Bangalore for providing necessary facilities and guidance and laboratory facilities to carry out current research work.

## Conflict of interest

The authors declare that they have no conflict of interest.

## REFERENCES

- Anand, A.A.P., Vennison, S.J., Sankar, S.G., Prabhu, D.I.G., Vasani, P.T., Raghuraman, T., Geoffrey, C.J. & Vendan, S.E. (2010). Isolation and characterization of bacteria from the gut of *Bombyx mori* that degrade cellulose, xylan, pectin and starch and their impact on digestion. *Journal of Insect Science*. 10, 107-112. DOI: 10.1673/031.010.10701
- Ahluwat, S., Singh, D., Yadav, A., Singh, A.K., Viridi, J. S., & Sharma, K.K. (2020). Proteomic analysis reveals the damaging role of low redox laccase from *Yersinia enterocolitica* strain 8081 in the midgut of *Helicoverpa armigera*. *Biotechnology Letters*, 42, 2189–2210. DOI: 10.1007/s10529-020-02925-x
- Bai, L.S., Zhao, C.X., Xu, J.J., Feng, C., Li, Y.Q., Dong, Y. L. & Ma, Zhi-qing (2019). Identification and biochemical characterization of carboxylesterase 001G associated with insecticide detoxification in *Helicoverpa armigera*. *Pest. Biochem. Physiol.* 157, 69–79. DOI: 10.1016/j.pestbp.2019.03.009
- Broderick, N.A., Raffa, K.F., Goodman, R.M & Handelsman, J. (2004). Census of bacterial community of gypsy moth larval mid gut by using culturing and culture independent methods. *Applied Environmental Microbiology*. 70, 290-300. DOI: 10.1128/AEM.70.1.293-300.2004
- Carol, R.L., Sarah, E.P & Ronald, J.P. (2003). Degradation and detoxification of the dihydrochalcone phloridzin by *Enterobacter agglomerans*, a bacterium associated with the apple pest, *Rhagoletis pomonella* (Walsh) (Diptera: Tephritidae). *Environmental Entomology*. 32, 953-962. DOI: 10.1603/0046-225X-32.5.953
- Dar, M.A., Shaikh, A.F., Pawar, K.D., Xie, R., Sun, J., Kandasamy, S., & Pandit, R.S. (2021). Evaluation of cellulose degrading bacteria isolated from the gut-system of cotton bollworm, *Helicoverpa armigera* and their potential values in biomass conversion. *PeerJ* 9: e11254. DOI: 10.7717/peerj.11254
- Dar, M.A., Shaikh, A.A., Pawar, K.D., & Pandit, R.S. (2018). Exploring the gut of *Helicoverpa armigera* for cellulose degrading bacteria and evaluation of a potential strain for lignocellulosic biomass deconstruction. *Process Biochemistry*. DOI: 10.1016/j.procbio.2018.08.001
- Dick, R.E., Quinn, J.P. (1995). Glyphosate-degrading isolates from environmental samples: occurrence and pathways of degradation. *Applied Microbial Biocontrol*. 43 (8): 545-550. DOI: 10.1007/BF00218464
- Dillon, R.J and Dillon, V.M. (2004). The gut bacteria of insects: non-pathogenic interactions. *Annual Review of Entomology*. 49, 71-92. DOI: 10.1146/annurev.ento.49.061802.123416
- Gavriel, S., Jurkevitch, E., Gazit, Y. & Yuval, B. (2010). Bacterially enriched diet improves sexual performance of sterile male Mediterranean fruit flies. *Journal of Applied Entomology*. 10, 439-444. DOI: 10.1111/j.1439-0418.2010.01605.x
- Gebbardi, K., Schimana, J., Muller, J., Krantal, P., Zeeck, A & Vater, I. (2001). Screening for biologically active metabolites with endosymbiotic bacilli isolated from arthropods. *FEMS Microbiology*. 217, 199-205. DOI: 10.1111/j.1574-6968.2002.tb11475.x
- Genta, A.F., Dillon, R.J., Terra, R.W & Ferreira, C. (2006). Potential role for gut microbiota in cell wall digestion and glucoside detoxification in *Tenebrio molitor* larvae. *Journal of Insect Physiology*. 52, 593-601. DOI: 10.1016/j.jinsphys.2006.02.007
- Hao, O.J., Kim, M.H., Seagren, E.A & Kim, H. (2002). Kinetics of phenols and chlorophenol utilization by *Acinetobacter* species. *Chemosphere*. 46, 797-807. DOI: 10.1016/S0045-6535(01)00182-5
- Hayashi, A., Aoyagi, H., Yoshimura, T & Hideo Tanaka, H. (2007). Development of novel method for screening microorganisms using symbiotic association between insect and intestinal microorganisms. *Journal of Bioscience and Bioengineering*. 103, 358-367. DOI: 10.1263/jbb.103.358
- Hoffman, C.S & Winston, F. (1987). A ten-minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*. *Gene*. 57, 267-272. DOI: 10.1016/0378-1119(87)90131-4
- Jilani, S and Khan, A.M. (2004). Isolation, characterization and growth response of pesticide degrading bacteria. *Journal of Biological Science*. 4, 15-20. DOI: 10.3923/jbs.2004.15.20
- Jing, T.Z., Qi, F.H. & Wang, Z.Y. (2020). Most dominant roles of insect gut bacteria: digestion, detoxification, or essential nutrient provision? *Microbiome* 8, 38. DOI: 10.1186/s40168-020-00823-y
- Kanekar, P.P., Bhadbhade, B.J., Deshpande, N.M & Saranik, S.S. (2004). Biodegradation of Organophosphates. *Biological Sciences*. 70, 57-70
- Kranthi, K.R., Jadhav, D., Wanjari, R., Kranthi, S. & Russell, D. (2001). Pyrethroid resistance and mechanism of resistance in field strains of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Journal of Economic Entomology*. 94, 254-263. DOI: 10.1603/0022-0493-94.1.253
- Lee, K.S., Metcalf, W.W. & Wanner, B.L. (1992). Evidence for two phosphonate degradative pathways in *Enterobacter aerogenes*. *Journal of Bacteriology*. 174, 2501-2510. DOI: 10.1128/jb.174.8.2501-2510.1992
- Lei, P., Nie, M.Q., Wen, X.M., Ge, B.Z. & Zang, Z.J. (2004). Study of degradation characters of preponderant Flavobacterial strains in a mixture of anthracene, phenanthrene and pyrene. *Journal of Xi'an Jiaotong University*. 38, 657–658.
- Mishra, P.K. and Tandon, S.M. (2003). Gut bacterial flora of *Helicoverpa armigera* (Hub.) (Lepidoptera: Noctuidae). *Indian Journal of Microbiology*. 43, 55-56.
- Nagarkatti, S. and Satyaprakash, S. (1974). Rearing *Helicoverpa armigera* (Hubn.) on an artificial diet. *Technical Bulletin Commonwealth Institute of Biological Control*. 17, 169-173.
- Paramasiva, I., Shouche, Y., Kulkarni, G. J., Krishnayya, P. V., Akbar, S. M., & Sharma, H. C. (2014). Diversity in gut microflora of *Helicoverpa armigera* populations from different regions in relation to biological activity of *Bacillus thuringiensis*  $\delta$ -endotoxin Cry1Ac. *Archives of Insect Biochemistry and Physiology*, 87, 201–213. DOI: 10.1002/arch.21190
- Priya, N.G., Ojha, A., Kajla, M.K., Raj, A. & Rajagopal, R. (2012). Host plant induced variation in gut bacteria of *Helicoverpa armigera*. *PLoS One*. 7, e30768. DOI: 10.1371/journal.pone.0030768
- Rajagopal, R., Sivakumar, S., Agrawal, N., Malhotra, P & Bhatnagar, R.K. (2002). Silencing of midgut aminopepti-



- dase N of *Spodoptera litura* by dsRNA establishes its role as *Bacillus thuringiensis* toxin receptor. *Journal of Biological Chemistry*. 277, 46849-46851. DOI: 10.1074/jbc.C200523200
27. Rajendran, L., Samiyappan, R., Raguchander, T. & Saravanakumar, D. (2007). Endophytic bacteria mediate plant resistance against cotton bollworm. *Journal of Plant Interactions*. 2, 1–10. DOI: 10.1080/17429140701420003
  28. Rani, A., Sharma, A., Rajagopal, R., Adak, T. & Bhatnagar, R.K. (2009). Bacterial diversity analysis of larvae and adult midgut micro-flora using culture-dependent and culture-independent methods in lab-reared and field-collected *Anopheles stephensi*-an Asian malarial vector. *BMC Microbiology*. 9, 96. DOI: 10.1186/1471-2180-9-96
  29. Ranjith, M.T., Chellappan, M., Harish, E.R., Girija, D. & Nazeem, P.A. (2016). Bacterial communities associated with the gut of tomato fruit borer, *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) based on Illumina Next-Generation Sequencing. *Journal of Asia-Pacific Entomology*. 19, 333–340. DOI: 10.1016/j.aspen.2016.03.007
  30. Regode, V., Kuruba, S., Mohammad, A.S., & Sharma, H.C. (2016). Isolation and characterization of gut bacterial proteases involved in inducing pathogenicity of *Bacillus thuringiensis* toxin in cotton bollworm, *Helicoverpa armigera*. *Frontiers in Microbiology*, 7, 1567. DOI: 10.3389/fmicb.2016.01567
  31. Sambrook, J., Fritsch, E.F. & Maniatis, T. (2001). Molecular Cloning, A Laboratory Manual, 3<sup>rd</sup> edn (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY).
  32. Sharon, G., Segal, D., Ringo, J.M., Hefetz, A. & Rosenberg, I.Z. (2010). Commensal bacteria play a role in mating preference of *Drosophila melanogaster*. *Proceedings of National Academy of Sciences*. 107, 20051–20056. DOI: 10.1073/pnas.1009906107
  33. Shinde, A. A., Shaikh, F. K., Gadge, P. P., Padul, M. V., Govindwar, S. P., & Kachole, M. S. (2017). Conserved nature of *Helicoverpa armigera* gut bacterial flora on different host plants and in vitro interactions with PI proteins advocates role in host digestive physiology. *Journal of the Saudi Society of Agricultural Sciences*. 18: 141-149. DOI:10.1016/j.jssas.2017.03.004
  34. Singh, K., Brajesh, K., Walker, A., Alum, J., Morgan, W & Wright, D.J. (2004). Biodegradation of chlorpyrifos by *Enterobacter strain B-14* and its use in biodegradation of contaminated soils. *Applied Environmental Microbiology*. 70, 4855-4863. DOI: 10.1128/AEM.70.8.4855-4863.2004
  35. Thakur, D., Bhuyan, M., Majumdar, S., Yadav, A., Hazarika, L.K., Harman, N., Baruah, A.A.L.H., & Bora, T.C. (2005). Isolation, characterization, in-vitro antibiotic susceptibility and pesticide tolerance of gut bacteria from rice hispa, *Diadisa armigera* (Olivier). *Indian Journal of Microbiology*. 45, 217-221.
  36. Verma, K., Agarwal, M., Farooq, M., Misra, R.B & Hans, R.K. (2006). Endosulfan degradation by a *Rhodococcus* strain isolated from the earthworm gut. *Ecotoxicology and Environmental Safety*. 64, 377-381. DOI: 10.1016/j.ecoenv.2005.05.014
  37. Vesta, S.A., Sandra, R & Virmantas, S. (2006). Myrcene resistant bacteria isolated from the gut of phytophagous insects *Ips typographus*. *Ekologia*. 4, 1-6.
  38. Visotto, L.E., Oliveira, M.G.A., Guedes, R.N.C., Ribon, A.O.B & Good-God, P.I.V. (2009). Contribution of gut bacteria to digestion and development of the velvetbean caterpillar, *Anticarsia gemmatilis*. *Journal of Insect Physiology*. 55, 185-191. DOI: 10.1016/j.jinsphys.2008.10.017
  39. Whalon, M.E., Mota-Sanchez, D. & Hollingsworth, R. (2008). Global Pesticide Resistance in Arthropods. (CAB International, Oxfordshire, UK).
  40. Wilson, G.R. and Benoit, T.G. (1993). Alkaline pH activated *Bacillus thuringiensis* spores. *Journal of Invertebrate Pathology*. 62, 87–89. DOI: 10.1006/jipa.1993.1079
  41. Xiang, H., Wei, G. F., Jia, S., Huang, J. & Miao, X.X. (2006). Microbial communities in the larval midgut of laboratory and field populations of cotton bollworm (*Helicoverpa armigera*). *Canadian Journal of Microbiology*. 52, 1085–1092. DOI: 10.1139/w06-064
  42. Yang, X., Xu, P., Yuan, H., Graham, R.I., Wilson, K., & Wu, K. (2019). Discovery and characterization of a novel picorna-like RNA virus in the cotton bollworm *Helicoverpa armigera*. *Journal of Invertebrate Pathology*, 160, 1–7. DOI: 10.1016/j.jip.2018.11.003
  43. Yuan, H., Xu, P., Yang, X., Graham, R.I., Wilson, K., & Wu, K. (2017). Characterization of a novel member of genus Iflavirus in *Helicoverpa armigera*. *Journal of Invertebrate Pathology*, 144, 65–73. DOI:10.1016/j.jip.2017.01.011