

Repositório ISCTE-IUL

Deposited in *Repositório ISCTE-IUL*:

2021-07-22

Deposited version:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Santilli, Leonardo & Tierz, M. (2020). Complex (super)-matrix models with external sources and q -ensembles of Chern–Simons and ABJ(M) type. *Journal of Physics A: Mathematical and Theoretical*. 53 (42), 425201-425201

Further information on publisher's website:

[10.1088/1751-8121/abb6b0](https://doi.org/10.1088/1751-8121/abb6b0)

Publisher's copyright statement:

This is the peer reviewed version of the following article: Santilli, Leonardo & Tierz, M. (2020). Complex (super)-matrix models with external sources and q -ensembles of Chern–Simons and ABJ(M) type. *Journal of Physics A: Mathematical and Theoretical*. 53 (42), 425201-425201, which has been published in final form at <https://dx.doi.org/10.1088/1751-8121/abb6b0>. This article may be used for non-commercial purposes in accordance with the Publisher's Terms and Conditions for self-archiving.

Use policy

Creative Commons CC BY 4.0

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a link is made to the metadata record in the Repository
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

COMPLEX (SUPER)-MATRIX MODELS WITH EXTERNAL SOURCES AND q -ENSEMBLES OF CHERN–SIMONS AND ABJ(M) TYPE

LEONARDO SANTILLI AND MIGUEL TIERZ

ABSTRACT. The Langmann–Szabo–Zarembo (LSZ) matrix model is a complex matrix model with a quartic interaction and two external matrices. The model appears in the study of a scalar field theory on the non-commutative plane. We prove that the LSZ matrix model computes the probability of atypically large fluctuations in the Stieltjes–Wigert matrix model, which is a q -ensemble describing $U(N)$ Chern–Simons theory on the three-sphere. The correspondence holds in a generalized sense: depending on the spectra of the two external matrices, the LSZ matrix model either describes probabilities of large fluctuations in the Chern–Simons partition function, in the unknot invariant or in the two-unknot invariant. We extend the result to supermatrix models, and show that a generalized LSZ supermatrix model describes the probability of atypically large fluctuations in the ABJ(M) matrix model.

CONTENTS

1. Introduction	1
2. Noncommutative scalar theory with background field	4
3. Scalar theory on the fuzzy sphere	7
4. Solution of the matrix model	12
5. Solution of the supermatrix model	22
Appendix A. Schur polynomials	25
References	27

1. INTRODUCTION

The idea of introducing non-zero commutators among position or momentum coordinates in different directions goes back to the 1940s [1]. The consequences of applying ideas and results of noncommutative geometry in quantum field theory are far-reaching but also considerably involved with both several non-trivial results and difficulties as well.

In the late 1990s there was a boost of interest in noncommutative (NC) field theories in great part due to the fact that low energy string theory can be related to NC field theory [2,3]. However, soon it was established that the expectation that washing out the space-time points could weaken UV divergences in quantum field theory, and consequently simplify renormalization, did not work as expected. Rather the opposite turned out to hold: renormalization gets harder due to noncommutativity, because in planar diagrams of a perturbative expansion the UV divergences simply persist. Second, in the non-planar diagrams, they tend to “mix” with IR divergences [4]. We refer to [5,6] for classical reviews of the topic, and to [7] for insights into the relation between fuzzy spaces and matrix models.

Langmann, Szabo and Zarembo (LSZ) introduced and studied a scalar field theory on the Moyal plane, and showed that its partition function admits a matrix model representation [8,9]. The LSZ matrix model has the explicit form

$$(1.1) \quad \mathcal{Z}_{\text{LSZ}}(E, \tilde{E}) = \int \mathcal{D}M \mathcal{D}M^\dagger \exp\left(-N \text{Tr} \left\{ MEM^\dagger + M^\dagger \tilde{E}M + \widehat{V}(M^\dagger M) \right\}\right),$$

where M is a $N \times N$ complex matrix, E, \tilde{E} are external matrices with real eigenvalues determined by the model and \widehat{V} is a polynomial potential with quadratic and quartic interaction terms. Recently, a Hermitian matrix model with external field E and quartic interaction has been studied in the context of NC scalar field theories [10–12]. It is worthwhile to stress that, despite the similarities, the fact that the LSZ model is a complex random matrix ensemble leads to a different analysis and very different set of results. In particular, we will establish a relationship between this model and a family of matrix models that appear in Chern–Simons theory and are related to q -deformed random matrix ensembles.

We will also consider a supersymmetric extension of the LSZ model (sLSZ), which we define as the supermatrix model

$$(1.2) \quad \mathbf{Z}_{\text{sLSZ}}(E, \tilde{E}) = \int \mathcal{D}M \mathcal{D}M^\dagger \exp\left(- (N_1 + N_2) \text{STr} \left\{ MEM^\dagger + M^\dagger \tilde{E}M + \widehat{V}(M^\dagger M) \right\}\right),$$

where now M is a $(N_1 + N_2) \times (N_1 + N_2)$ complex supermatrix and STr is the supertrace. The details are given in section 5.

On the other hand, a number of different matrix models have been studied in gauge theory, more precisely in the study of certain topological and supersymmetric gauge theories in compact three-manifolds such as Seifert manifolds. The simplest case is that of \mathbb{S}^3 , where the partition function of $U(N)$ Chern–Simons theory, admits the expression [13]

$$(1.3) \quad \mathcal{Z}_{\text{CS}} = \int_{\mathbb{R}^N} \prod_{1 \leq j < k \leq N} \left(2 \sinh \left(\frac{x_j - x_k}{2} \right) \right)^2 \prod_{j=1}^N e^{-\frac{1}{2g_s} x_j^2} dx_j,$$

with g_s a coupling constant, which can be related to the level $k \in \mathbb{Z}$ of Chern–Simons theory by $g_s = \frac{2\pi i}{N+k}$. A review of early results is [14]. The matrix model description can also be obtained and further understood by using different types of localization of the path integral [15–17].

Interestingly, expressions such as (1.3), or generalizations thereof, with Schur polynomial insertions, describing then Wilson loop observables in the Chern–Simons theory, are completely solvable, even for finite N , using standard random matrix theory methods [18–20]. We will give an interpretation of the LSZ matrix model in terms of probabilities in the random matrix description of the observables of $U(N)$ Chern–Simons theory on \mathbb{S}^3 . This type of probability is a classical object in random matrix theory [21] and typically computed via a Fredholm determinant. For this family of models, such probabilities have also been studied, in the context of what is known as critical statistics [22, 23] (statistics interpolating between Poisson and Wigner–Dyson behaviour). However, we will be naturally lead to the consideration of atypically large fluctuations instead (like for the GUE model [24], only the model (1.3) is a q -deformation of the GUE) rather than any bulk spectral quantity. In addition, we shall see that different insertions of external matrices E, \tilde{E} will be related to different Chern–Simons observables.

Further examples beyond the topological $U(N)_k$ Chern–Simons theory are the ABJ(M) theories [25, 26], supersymmetric $U(N_1)_k \times U(N_2)_{-k}$ Chern–Simons-matter theories preserving twelve supercharges. The partition function of ABJ theory on \mathbb{S}^3 has the matrix model

representation [27]

$$(1.4) \quad \mathbf{Z}_{\text{ABJ}} = \int_{\mathbb{R}^{N_1}} \int_{\mathbb{R}^{N_2}} \frac{\prod_{1 \leq j < k \leq N_1} \left(2 \sinh \left(\frac{x_j - x_k}{2} \right) \right)^2 \prod_{1 \leq r < s \leq N_2} \left(2 \sinh \left(\frac{w_r - w_s}{2} \right) \right)^2}{\prod_{j=1}^{N_1} \prod_{r=1}^{N_2} \left(2 \cosh \left(\frac{x_j - w_r}{2} \right) \right)^2} \\ \times \prod_{j=1}^{N_1} e^{-\frac{1}{2g_s} x_j^2} dx_j \prod_{r=1}^{N_2} e^{-\frac{1}{2g_s} w_r^2} dw_r,$$

where the relation between the string coupling g_s and the Chern–Simons level $k \in \mathbb{Z}$ is $g_s = \frac{2\pi i}{k}$. It was shown in [28] that equation (1.4) defines a supermatrix model related to the ordinary matrix model for $U(N)$ Chern–Simons theory on the lens space $\mathbb{S}^3/\mathbb{Z}_2$ [29, 30]. The approach of [28] has been applied in [31] to evaluate exactly the ABJ partition function. We will show how the relation uncovered between LSZ and Chern–Simons matrix models extends to the supermatrix models (1.2)-(1.4).

This paper has two parts. In the first one, we will dwell on the interpretation of the LSZ model in noncommutative field theory because, since [8, 9], a closely related matrix model to (1.1) has appeared in other works, dealing with gauge theories in noncommutative \mathbb{R}^3 [32, 33]. Thus, it is interesting to study further this, more so given the new interpretations we will discuss in the second part paper of this paper.

The analysis of Abelian gauge theories in noncommutative \mathbb{R}^3 has been carried out in [32–37]. The deformation of three-dimensional Euclidean space considered is the so-called \mathbb{R}_λ^3 , modelled on the $\mathfrak{su}(2)$ algebra [38]. The discrete eigenvalues of the quadratic Casimir of $\mathfrak{su}(2)$ produce a “foliation” by fuzzy spheres of different radii. In [32, 36] it has been found the most general action satisfying three conditions: gauge invariance, stability of the vacuum, positivity. The second condition is tightly related to the choice of fundamental field for the theory; in particular, two different approaches, namely expanding a covariant gauge field around a gauge invariant flat connection [36] or around the null configuration [32] lead to dissimilar pictures. In both cases, after gauge fixing, the gauge theories reduce to a tower of scalar theories with quartic interaction, one on each fuzzy sphere foliating \mathbb{R}_λ^3 . On one hand, the first model yields a partition function which appears to be related to a string model with a background B -field enforcing string noncommutativity [39]. On the other hand, the similarity of the action in the latter case with the Langmann–Szabo–Zarembo model [8, 9] was highlighted in [32]. We will show that, from [32, 33] but with a slight change in the definition of the exterior calculus on \mathbb{R}_λ^3 and eventually taking the large radius limit, one reproduces the matrix model (1.1).

In the second part, which is the core of the work, we will establish the relationship between the two sets of matrix models described above, exactly in the manner explained. This is a random matrix result, linking two families of models: complex matrix models with external sources on one hand, q -ensembles that appear in Chern–Simons theory on the other. Importantly, the relationship is not between the same observables. On one hand, we have partition functions (albeit generalized via different choices of external matrices) and on the other hand probabilities in the q -ensembles, corresponding to different observables in the Chern–Simons theory. This result can also be appreciated independently of the gauge theoretic origin of both sets of models. The Chern–Simons matrix model for example, also plays a prominent role in the subject of non-intersecting Dyson Brownian motion [40, 41], see [42, 43].

The article is organized as follows. A field-theoretical background for the matrix model of interest is provided in section 2, where we review the derivation of the LSZ matrix model from a noncommutative scalar theory. In section 3 we discuss how the same matrix model

emerges as a limit in a different noncommutative field theory, the scalar theory obtained from reduction of a three-dimensional gauge theory in \mathbb{R}_λ^3 .

After that, we analyze the LSZ matrix model and explain its close relation with the Chern–Simons matrix model in section 4. We emphasize that the connection holds in a generalized sense, as the external matrices are not restricted to come from a kinetic operator of a scalar field theory, and different spectra of the external matrices will be interpreted in terms of different observables in Chern–Simons theory. In subsection 4.4 we show that the generalized LSZ partition function encodes certain probabilities in the random matrix description of topological invariants computed from Chern–Simons theory. Then, in section 5, we extend the analysis to the sLSZ supermatrix model, and establish an analogous relation between the sLSZ generalized partition function and observables in ABJ theory. These are the central results of the present work. Finally, the Appendix A contains technical details about Schur polynomials.

2. NONCOMMUTATIVE SCALAR THEORY WITH BACKGROUND FIELD

In this section we review the LSZ model: in the first subsection, the geometric construction of the Moyal plane in the presence of a background magnetic field is sketched, while the second subsection is dedicated to the construction of a scalar field theory with quartic interaction. This provides a motivation and a physical background for the study of the matrix model (1.1), that will be thoroughly analyzed in section 4.

The content of this section follows [9], although for the derivation of the matrix model in subsection 2.2 we use a slightly different formalism than the original work, that will allow us a more direct comparison with the results in the next section.

2.1. Moyal plane with magnetic field. The Moyal plane is defined through the commutation relations

$$[q^j, q^k] = i\theta\epsilon^{jk},$$

where θ is the essential parameter of the theory, with dimension of length squared. A Moyal plane can always be seen as a harmonic system, in the sense that passing to dimensionless complex coordinates one has $[z, \bar{z}] = 1$. The noncommutative plane needs not to arise from a modification of space-time. In fact, an example is given by a particle moving on a plane with a magnetic field of intensity B in the transverse direction; the momentum space then becomes noncommutative \mathbb{R}^2 , as the momentum operators modify according to:

$$p_j \mapsto P_j := p_j - \frac{1}{2}B\epsilon_{jk}q^k.$$

In this case the covariant momenta P_j satisfy the commutation relation $[P_j, P_k] = -iB\epsilon_{jk}$.

If the two frameworks are put together, that is, a transverse magnetic field is plugged in on a noncommutative plane, three possible harmonic oscillator pictures arise:

- (i) on the two-dimensional position space, with annihilation and creation operators given by the complex coordinates as above;
- (ii) on the two-dimensional momentum space, with annihilation and creation operator defined analogously;
- (iii) a pair of canonical harmonic oscillators, one on each phase space plane.

However, the most suitable choice for us is none of them, and we will take a mixture of all these ingredients to form two commuting copies of annihilation and creation operators, in such a way that the problem decouples into two one-dimensional harmonic systems. To do

so, define:

$$\begin{aligned} z &:= \frac{q^1 + iq^2}{\sqrt{2\theta}}, & \bar{z} &:= \frac{q^1 - iq^2}{\sqrt{2\theta}}, \\ v &:= \frac{p_1 + ip_2}{\sqrt{2\theta^{-1}}}, & \bar{v} &:= \frac{p_1 - ip_2}{\sqrt{2\theta^{-1}}}, \end{aligned}$$

and use them to introduce the operators:

$$\begin{aligned} a_1 &= \frac{z + iv}{\sqrt{2}}, & a_1^\dagger &= \frac{\bar{z} - i\bar{v}}{\sqrt{2}}, \\ a_2 &= \frac{\bar{z} + i\bar{v}}{\sqrt{2}}, & a_2^\dagger &= \frac{z - iv}{\sqrt{2}}. \end{aligned}$$

Straightforward calculations provide:

$$\begin{aligned} [a_\alpha, a_\beta^\dagger] &= \delta_{\alpha\beta}, \\ [a_\alpha, a_\beta] &= 0 = [a_\alpha^\dagger, a_\beta^\dagger], \end{aligned}$$

for $\alpha, \beta = 1, 2$, hence we got a pair of decoupled harmonic oscillators.

Remark. Lifting the obstruction θ shifts the canonical symplectic structure on the cotangent bundle. It turns out that such shifted 2-form is still symplectic. One can then rotate to Darboux coordinates so that the new symplectic structure on the phase space $T^*\mathbb{R}^2$ is block-diagonal. The calculations above are precisely the explicit change of coordinates.

Consider now the differential operator D_j associated to the covariant momenta P_j , and \tilde{D}_j analogous but carrying a reflected magnetic field $-B$. If we take the arbitrary combination $-\sigma D^2 - \tilde{\sigma} \tilde{D}^2$ and evaluate it at the symmetric point $\sigma = \tilde{\sigma} = \frac{1}{2}$, we obtain:

$$\left(-\sigma D^2 - \tilde{\sigma} \tilde{D}^2\right)_{\sigma=\tilde{\sigma}=\frac{1}{2}} = |\vec{p}|^2 + \frac{B^2}{4} |\vec{q}|^2 = \theta^{-1} \left(\frac{B^2 \theta^2}{4} \{z, \bar{z}\} + \{v, \bar{v}\} \right),$$

where the curly bracket in the right-hand side stands for anticommutation. On the other hand, in terms of the harmonic oscillators description, we have:

$$\sum_{\alpha=1}^2 a_\alpha^\dagger a_\alpha = \frac{1}{2} (\{z, \bar{z}\} + \{v, \bar{v}\} - i[v, \bar{z}] + i[z, \bar{v}]),$$

which means

$$(2.1) \quad \left(-\sigma D^2 - \tilde{\sigma} \tilde{D}^2\right)_{\sigma=\tilde{\sigma}=\frac{1}{2}} = \frac{2}{\theta} \sum_{\alpha=1}^2 \left(a_\alpha^\dagger a_\alpha + \frac{1}{2} \right)$$

at points $B^2 \theta^2 = 4$. The preferred curves $\frac{B^2 \theta^2}{4} = 1$ correspond to the self-dual points of the Langmann–Szabo symmetry [44]. The theory is independent of the actual choice of curve in parameter space we restrict to, namely $B = \pm 2\theta^{-1}$. In fact, the two theories we obtain are equivalent in the Seiberg–Witten sense, i.e., they transform into the same theory [44]. The invariance reflects the fact that the operators D_j, \tilde{D}_j only differ by a reflection $B \mapsto -B$, thus the symmetric choice $\sigma = \tilde{\sigma}$ drops the dependence on the sign of the magnetic field.

2.2. LSZ model. Given a scalar field Φ on the Moyal plane, we can expand it in terms of the Landau basis, consisting of eigenstates of both harmonic oscillators, as:

$$\Phi = \sum_{\ell_1, \ell_2=1}^{\infty} M_{\ell_1 \ell_2} |\ell_1, \ell_2\rangle.$$

This expression naturally defines an infinite matrix M associated to the field Φ . Now recall the kinetic operator in (2.1); using the property

$$a_{\alpha}^{\dagger} a_{\alpha} |\ell_1, \ell_2\rangle = (\ell_{\alpha} - 1) |\ell_1, \ell_2\rangle, \quad \ell_{\alpha} = 1, 2, \dots, \quad \alpha = 1, 2,$$

it is possible to write:

$$\begin{aligned} \left(\sum_{\alpha=1}^2 \left(a_{\alpha}^{\dagger} a_{\alpha} + \frac{1}{2} \right) \right) \Phi &= \sum_{\ell_1, \ell_2} M_{\ell_1 \ell_2} \left\{ \left(\ell_1 - \frac{1}{2} \right) + \left(\ell_2 - \frac{1}{2} \right) \right\} |\ell_1, \ell_2\rangle \\ &= \sum_{\ell_1, \ell_2} \left(\ell_1 - \frac{1}{2} \right) \{ M_{\ell_1 \ell_2} |\ell_1, \ell_2\rangle + M_{\ell_2 \ell_1} |\ell_2, \ell_1\rangle \}. \end{aligned}$$

Therefore one obtains:

$$\begin{aligned} (2.2) \quad \Phi^{\dagger} \left(\sum_{\alpha=1}^2 \left(a_{\alpha}^{\dagger} a_{\alpha} + \frac{1}{2} \right) \right) \Phi &= \sum_{\ell_1, \ell_2} \left(\ell_1 - \frac{1}{2} \right) \{ M_{\ell_2 \ell_1}^{\dagger} M_{\ell_1 \ell_2} + M_{\ell_2 \ell_1} M_{\ell_1 \ell_2}^{\dagger} \} \\ &= \text{Tr} \{ M^{\dagger} E M + M E M^{\dagger} \}, \end{aligned}$$

where in the last line we have introduced the diagonal matrix

$$(2.3) \quad E_{\ell_1 \ell_2} = \left(\ell_1 - \frac{1}{2} \right) \frac{4\pi}{N} \delta_{\ell_1 \ell_2}.$$

Consider the action [8, 9]

$$\begin{aligned} S_{\text{LSZ}}(\Phi, \Phi^{\dagger}) &= \int_{\mathbb{R}_{\theta}^2} \left\{ \frac{1}{2} \Phi^{\dagger} \left(-\sigma D^2 - \tilde{\sigma} \tilde{D}^2 \right) \Phi + \frac{1}{2} \Phi \left(-\sigma D^2 - \tilde{\sigma} \tilde{D}^2 \right) \Phi^{\dagger} \right. \\ &\quad \left. + m_0^2 \Phi^{\dagger} \Phi + \frac{g_0^2}{2} \left(\Phi^{\dagger} \Phi \right)^2 \right\}. \end{aligned}$$

Evaluated at the symmetric point $\sigma = \tilde{\sigma} = \frac{1}{2}$ it becomes:

$$(2.4) \quad S_{\text{LSZ}}(\Phi, \Phi^{\dagger}) = N \text{Tr} \left\{ M^{\dagger} E M + M E M^{\dagger} + \hat{m}^2 M^{\dagger} M + \frac{\hat{g}^2}{2} \left(M^{\dagger} M \right)^2 \right\}$$

where we have introduced the dimensionless couplings

$$\hat{m}^2 = \left(\frac{2\pi\theta}{N} \right) m_0^2, \quad \hat{g}^2 = \left(\frac{2\pi\theta}{N} \right) g_0^2.$$

We have also used equations (2.1) and (2.2), and the matrix E defined in (2.3). In section 4 we will solve the matrix model with action (2.4).

Notice that, to regularize the integral, we truncate the matrix M to its top-left $N \times N$ block, which introduces a finite cutoff at short distance $\sqrt{\frac{2\pi\theta}{N}}$. The full theory is recovered in the large N limit. As expected from general features of noncommutative field theory (see e.g. [6]), the original noncommutativity of the phase space is eventually encoded in the noncommutativity of matrix multiplication. Consistently, the space-time integral becomes a trace.

3. SCALAR THEORY ON THE FUZZY SPHERE

We now discuss another noncommutative scalar field theory in two dimensions, this time on the fuzzy sphere [45], that leads to the LSZ matrix model (1.1).

Scalar theories on the fuzzy sphere have been extensively studied, mainly looking at them as regularized UV/IR mixing-free versions of commutative theories [46–48]. In the standard setting the kinetic sector prevents the integration of the angular degrees of freedom, and thus to reduce to an integral over eigenvalues. A perturbative approach in the kinetic term was proposed in [49, 50], equivalent to a high-temperature expansion, and the presence of phase transitions, together with a triple point, was suggested. This procedure was later generalized to $\mathbb{C}\mathbb{P}^N$ [51]. See [52] for a review, describing both theoretical predictions and numerical results, paying special attention to the phase transition. An extended scheme, allowing both a high-temperature (large-interaction) and small-interaction analysis was presented in [53], providing further understanding of the phase structure of the model. Other nonperturbative aspects are still under investigation: in [54, 55] the behaviour of correlation functions of the matrix model in the disordered phase is analyzed¹.

However, all these works lead to matrix models different from (1.1). Instead, we will start with a gauge theory in a noncommutative version of \mathbb{R}^3 which is “foliated” by fuzzy spheres [38]. The projection of the gauge theory onto each fuzzy sphere gives a noncommutative scalar theory which reduces to LSZ in the large radius limit. We stress that the model we consider was introduced in [32, 33] and it has the distinctive feature that it is not directly defined as a scalar theory on the fuzzy sphere, but it is derived in a roundabout way. The identification with the LSZ model at large radius is possible precisely because the kinetic operator of the scalar theory descends from a three-dimensional gauge theory, instead of being given by the adjoint action of the generators of $\mathfrak{su}(2)$, as the standard kinetic term for a scalar theory on the fuzzy sphere would be.

The rest of the section is dedicated to the proof of this equivalence. We first give a very brief review of the properties of the deformation of \mathbb{R}^3 known as \mathbb{R}_λ^3 . Then, the second subsection is dedicated to the construction of the gauge theory in \mathbb{R}_λ^3 , which is essentially a review of the setup of [32, 33] with minor modifications. Eventually, we reduce to a scalar theory on a fuzzy sphere of fixed radius and study the large radius limit.

3.1. From \mathbb{R}_λ^3 to the fuzzy sphere. We start with a noncommutative version of the three-dimensional Euclidean space, imposing the coordinates to satisfy the commutation relations of $\mathfrak{su}(2)$. Such space is known in the literature as \mathbb{R}_λ^3 . Then, as we will see, irreducible representations of $\mathfrak{su}(2)$ determine a foliation of \mathbb{R}_λ^3 in terms of fuzzy spheres. We refer to [37, 38] for detailed insights in \mathbb{R}_λ^3 .

Coordinates in \mathbb{R}_λ^3 are the generators $\{x^\mu\}_{\mu=1}^3$ of $\mathfrak{su}(2)$, up to a length scale factor λ , satisfying:

$$[x^\mu, x^\nu] = i\lambda\epsilon^{\mu\nu\rho}x^\rho.$$

Irreducible representation of $\mathfrak{su}(2)$ in terms of $N \times N$ matrices are labelled by non-negative half-integers n , with $N = (2n + 1)$. Whenever n is fixed the Casimir relation implies:

$$\vec{x}^2 := \sum_{\mu=1}^3 (x^\mu)^2 = \lambda^2 n(n + 1),$$

which corresponds to pick a sphere of fixed radius $r^2 = \lambda^2 n(n + 1)$, denoted by \mathbb{S}_n^2 . One could interpret this construction of the fuzzy sphere as analogous to the usual embedding of \mathbb{S}^2 into \mathbb{R}^3 , but replacing coordinates with noncommutative Hermitian operators.

¹For NC gauge theories on the fuzzy sphere, a procedure to reduce the partition function to a matrix model was put forward in [56].

It is well-known that the large N limit of the fuzzy sphere at fixed radius gives the commutative sphere. Conversely, the scaling limit $r \rightarrow \infty$ keeping the parameter $\theta = \frac{r^2}{n}$ fixed leads to the Moyal plane. We focus on this latter setting and write:

$$\lambda^2 = \frac{\theta}{n+1}, \quad r^2 = \theta n, \quad \theta \text{ fixed.}$$

As shown in [32, 36, 37], there exists a matrix basis on each fuzzy sphere, which allows to identify fields in \mathbb{R}_λ^3 with a tower of matrices of increasing size. Noncommutativity is then encoded in $N \times N$ matrix multiplication, with $N = 2n + 1$, at each level $n \in \frac{1}{2}\mathbb{N}$.

Integration over \mathbb{R}_λ^3 is defined as

$$\int_{\mathbb{R}_\lambda^3} F = 2\pi \sum_{n \in \frac{1}{2}\mathbb{N}} \lambda^3(n+1) \text{Tr} f^{(n)} = 2\pi\theta^{3/2} \sum_{n \in \frac{1}{2}\mathbb{N}} \frac{1}{\sqrt{n+1}} \text{Tr} f^{(n)},$$

where $f^{(n)}$ is the $N \times N$ matrix representation of the function F at level n . We have had to adapt the prescription of [37] to treat λ and r as functions of n . This integral has the expected behaviour at large N , giving the volume of a sphere of radius $\sim \sqrt{\theta n}$. Freezing the radial degree of freedom, the integral reduces to

$$\int_{\mathbb{S}_n^2} F = 2\pi\theta \text{Tr} f^{(n)}$$

on the n -th fuzzy sphere.

3.2. Gauge theory setup. At this point, we introduce the algebra of derivations on \mathbb{R}_λ^3 . We define the forms

$$\tau_\mu := \frac{x^\nu}{r\lambda} \delta_{\nu\mu},$$

which yield associated derivations defined as

$$(3.1) \quad \partial_\mu := \text{ad}_{\tau_\mu} = i[\tau_\mu, \cdot],$$

satisfying the ring relation

$$[\partial_\mu, \partial_\nu] = -\frac{1}{r} \epsilon_{\mu\nu\rho} \partial_\rho.$$

Our derivations only differ from [32, 33, 36] by a factor λ/r , which belongs to the centre of the algebra and does not change the relevant properties of ∂_μ .

It is also possible to introduce the 1-form Θ such that $\Theta(\partial_\mu) = -i\tau_\mu$. It satisfies

$$d\Theta(\partial_\mu, \partial_\nu) + [\Theta(\partial_\mu), \Theta(\partial_\nu)] = 0,$$

implying that Θ is a flat connection. The most general gauge-invariant action in this framework includes a term coupling the gauge field to Θ . However, in our analysis, it will follow from the definition (3.1) that such term would only provide a constant correction to the mass term, hence we will reabsorb it in the definition of bare mass. This is one of the small differences between the present setting and [32, 33].

To construct an Abelian gauge theory on \mathbb{R}_λ^3 , we need to consider the noncommutative analogue of an $\text{ad}(P)$ -bundle, for $P \rightarrow \mathbb{R}^3$ a principal $U(1)$ -bundle. This is given by the right module $\mathfrak{u}(1) \otimes \mathbb{R}_\lambda^3$ over \mathbb{R}_λ^3 . We introduce the anti-Hermitian gauge fields A_μ and the gauge covariant derivative

$$\nabla_\mu = \partial_\mu + A_\mu.$$

The curvature F^A has components:

$$F_{\mu\nu}^A = [\nabla_{\partial_\mu}, \nabla_{\partial_\nu}] - \nabla_{[\partial_\mu, \partial_\nu]} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu] + \frac{1}{r} \epsilon_{\mu\nu\rho} A_\rho.$$

We see that the curvature has the usual form plus an extra term inherited from the underlying noncommutativity. The group of gauge transformations is the group of unitary transformations of \mathbb{R}_λ^3 (see [32] and references therein), and the gauge field transforms in the usual way:

$$A_\mu \mapsto U^\dagger A_\mu U + U^\dagger \partial_\mu U,$$

for any unitary U . Furthermore, one can also consider the Θ -covariant derivative $\tilde{\nabla}$ associated to the 1-form Θ , which for any field f on \mathbb{R}_λ^3 satisfies:

$$(3.2) \quad \tilde{\nabla}_\mu f = \partial_\mu f - i\tau_\mu f = -if\tau_\mu,$$

where the second equality follows from the definition (3.1) of the derivation as a commutator with τ_μ . $\tilde{\nabla}$ is gauge invariant:

$$\tilde{\nabla}_\mu f \mapsto U^\dagger \circ \tilde{\nabla}_\mu (Uf) = -if\tau_\mu,$$

where we used (3.2). Besides, using again (3.1) we see that $i\tau_\mu$ transforms under gauge transformations as

$$i\tau_\mu \mapsto iU^\dagger \tau_\mu U - U^\dagger \partial_\mu U.$$

For any choice of gauge fields A_μ we can define the so-called covariant coordinates C_μ , defined as the components of the 1-form $\nabla - \tilde{\nabla}$,

$$C_\mu = \nabla_\mu - \tilde{\nabla}_\mu = A_\mu + i\tau_\mu.$$

The C_μ fields are clearly anti-Hermitian and behave under gauge transformations according to:

$$(\nabla_\mu - \tilde{\nabla}_\mu) f \mapsto U^\dagger A_\mu U f + iU^\dagger \tau_\mu U f,$$

that is, $C_\mu \mapsto U^\dagger C_\mu U$. This is because the $U^\dagger \partial_\mu U$ parts cancel between the two contributions with opposite sign. On the other hand, by the very definition of derivations, see (3.1), it stems that

$$U^\dagger i\tau_\mu U = U^\dagger \partial_\mu U + i\tau_\mu,$$

which implies

$$(3.3) \quad C_\mu \mapsto (U^\dagger A_\mu U + U^\dagger \partial_\mu U) + i\tau_\mu.$$

Therefore the gauge transformations leave unchanged the τ_μ -part and modify the gauge field A_μ in the ordinary way. In particular, we underline for later convenience that, since we are working in a matrix basis in which x^3 is diagonal, the field C_3 has, by its very definition, the eigenvalues of τ_3 multiplied by $\sqrt{-1}$ in its diagonal, plus the diagonal entries of A_3 . The off-diagonal entries of C_3 in the basis we are working are simply the off-diagonal entries of A_3 . Therefore, a gauge transformation acting on C_3 transforms the contribution by A_3 in the standard way without affecting the eigenvalues of τ_3 in the diagonal of C_3 .

At this point, two options disclose to set up an Abelian gauge theory: we ought to chose either A_μ or C_μ as the fundamental fields of the theory. They differ by a flat connection: we expect the two resulting theories to be related by a redefinition of the vacuum. The two different procedures have been carried out, respectively, in [36] and [32]. We follow the second approach, taking C_μ as variables, and we pursue the most general action such that:

- (i) it is gauge invariant and at most quartic in the fundamental variable;
- (ii) it does not involve tadpoles at classical order, which imposes not to include linear terms in the fundamental variable;
- (iii) it is positive definite.

The second condition is essential in order to have a stable vacuum. Taking such family of gauge-invariant actions and writing down the classical equations of motion, one finds out that the absolute minimum is the null configuration $C_\mu = 0$. Other local minima are present. However, due to the special form of the derivations in the present work, all the vacua in the centre of the noncommutative algebra are constant configurations. (Nevertheless, some other vacuum configurations may fall outside the centre of the algebra).

The action which satisfies the above mentioned three hypotheses is:

$$(3.4) \quad S[C] = \frac{2}{g_0^2} \int_{\mathbb{R}_\lambda^3} \left([C_\mu, C_\nu]^2 + \Omega (C_\mu C_\nu + C_\nu C_\mu)^2 - m_0^2 C^2 \right)$$

We rewrite it in terms of Hermitian fields as $C_\mu = i\Phi_\mu$ and remove the redundant degree of freedom fixing the gauge $\Phi_3 = \tau_3$. Notice that, according to (3.3) and subsequent discussion, this corresponds to the Coulomb gauge $A_3 = 0$ in terms of the standard gauge fields A_μ . This procedure could be done, indeed, in a BRST-invariant fashion, introducing a ghost/anti-ghost pair (c, \bar{c}) with grading $(-1, +1)$ together with a Stückelberg field b , with zero ghost number. We add to the action a gauge fixing term

$$(3.5) \quad \text{Tr} \{ b (\Phi_3 - \tau_3) + i\bar{c} [\Phi_3, c] \}$$

and integrate out the field b first, enforcing the gauge fixing condition $\Phi_3 = \tau_3$. Then the second term above reduces to $\bar{c} D_3 c$, hence the ghost and anti-ghost fields decouple. This is not an exhaustive discussion, and we refer to [32, 33] for more details on the definition of a BRST coboundary operator in the present theory. The proof that the gauge fixing term (3.5) is in fact BRST-exact with respect to the just mentioned BRST operator can be found in [32, Eq.s (3.6)-(3.9)]. The general formalism of BV/BRST quantization of noncommutative gauge theories is discussed in [57].

At this point, we rearrange the remaining scalar fields into complex ones Φ, Φ^\dagger , scaled by a factor $\sqrt{2}/g_0$, and restrict to the region of parameter space for which the potential is of the form $V(\Phi^\dagger \Phi)$, corresponding to the choice $\Omega = \frac{1}{3}$ in (3.4). We finally arrive at the action (see [33, Eq. (3.17)] and discussion around it)

$$(3.6) \quad S = \int_{\mathbb{R}_\lambda^3} \left(\Phi^\dagger \hat{K} \Phi + \Phi \hat{K} \Phi^\dagger + \frac{g_0^2}{2} \Phi^\dagger \Phi \Phi^\dagger \Phi \right),$$

with kinetic operator

$$\hat{K} := m_0^2 \text{Id} + \frac{8}{3} \tau_3 (\tau_3 - i\partial_3).$$

The product in (3.6) is nothing but the matrix multiplication when the fields are represented in the natural matrix basis at each level n . The sole difference between the action (3.6) and the one in [33] is the different scaling of the kinetic operator \hat{K} in the large radius limit, analyzed in the next subsection.

3.3. Large radius limit. At this point, we project the system onto a single fuzzy sphere. This means we fix a half-integer n and restrict to those states spanned by the n -th eigenstate of the quadratic Casimir operator, related to the radial coordinate. The remaining degree of freedom is the degeneracy at fixed n , labelled by $k = -n, \dots, n$. We recall that the radius is $\sqrt{\theta n}$, with θ to be kept fixed at large N , where $N = 2n + 1$. Fields Φ are projected onto $N \times N$ matrices ϕ , according to the foliation of \mathbb{R}_λ^3 . We then calculate the matrix elements appearing in the action (3.6):

$$\begin{aligned} \langle k | \hat{K} \phi | k' \rangle &= m_0^2 \langle k | \phi | k' \rangle + \frac{8}{3r^2 \lambda^2} \left(\langle k | (x^3)^2 \phi | k' \rangle + \langle k | x^3 [x^3, \phi] | k' \rangle \right) \\ &= \langle k | \phi | k' \rangle \left\{ m_0^2 + \frac{8k}{3\theta n} (2k - k') \right\}. \end{aligned}$$

We multiply this expression by a factor $2\pi\theta$ which will arise from the integral, and obtain:

$$2\pi\theta\langle k|\hat{K}\phi|k'\rangle = (2n+1)\langle k|\phi|k'\rangle \left\{ \hat{m}^2 + \frac{16\pi}{3} \frac{k(2k-k')}{n(2n+1)} \right\},$$

where we have defined the dimensionless parameter

$$\hat{m}^2 := \left(\frac{2\pi\theta}{2n+1} \right) m_0^2$$

by scaling the bare mass with the cutoff induced by finite N . Therefore,

$$2\pi\theta \operatorname{Tr} \left\{ \phi^\dagger \hat{K} \phi + \phi \hat{K} \phi^\dagger \right\} = (2n+1) \sum_{k,k'=-N}^N \left\{ \langle k'|\phi^\dagger|k\rangle \langle k|\phi|k'\rangle \left(\hat{m}^2 + \frac{16\pi}{3} \frac{k(2k-k')}{n(2n+1)} \right) + \langle k'|\phi|k\rangle \langle k|\phi^\dagger|k'\rangle \left(\hat{m}^2 + \frac{16\pi}{3} \frac{k(2k-k')}{n(2n+1)} \right) \right\}.$$

Switching the labels of the dummy variables k, k' in the last summand, we obtain

$$(2n+1) \sum_{k,k'=-n}^n \left\{ \hat{m}^2 \left(\langle k'|\phi^\dagger|k\rangle \langle k|\phi|k'\rangle + \langle k'|\phi|k\rangle \langle k|\phi^\dagger|k'\rangle \right) + \langle k'|\phi^\dagger|k\rangle \langle k|\phi|k'\rangle \frac{32\pi}{3} \frac{k^2 + k'^2 - kk'}{n(2n+1)} \right\}.$$

At this point, in order to compare with the LSZ model, we need to pass from the angular momentum to the harmonic oscillator description. This is done using the relation $k = n - \ell$ for $\ell = 0, 1, \dots, 2n = N - 1$. We thus define the rearranged matrix M as

$$M_{\ell_1 \ell_2} := \langle n - \ell_1 | \phi | n - \ell_2 \rangle.$$

In terms of the new indices we have

$$k^2 + k'^2 - kk' = n^2 - n(\ell_1 + \ell_2) + \ell_1^2 + \ell_2^2 - \ell_1 \ell_2.$$

Hence, denoting by S_n the restriction of the action (3.6) to a single fuzzy sphere, we get:

$$S_n = (2n+1) \sum_{\ell_1, \ell_2=0}^{2n} \left\{ \hat{m}^2 \left(M_{\ell_2 \ell_1}^\dagger M_{\ell_1 \ell_2} + M_{\ell_2 \ell_1} M_{\ell_1 \ell_2}^\dagger \right) + M_{\ell_2 \ell_1}^\dagger M_{\ell_1 \ell_2} \frac{32\pi}{3} \frac{n^2 - n(\ell_1 + \ell_2) + \ell_1^2 + \ell_2^2 - \ell_1 \ell_2}{n(2n+1)} \right\} + \left(\frac{2\pi\theta}{2n+1} \right) \frac{g_0^2}{2} \operatorname{Tr} \left\{ \left(M^\dagger M \right)^2 \right\}.$$

Eventually, rescaling g_0^2 into a dimensionless parameter

$$\hat{g}^2 := \left(\frac{2\pi\theta}{2n+1} \right) g_0^2$$

and splitting the second line into the sum of two terms, the action on the fuzzy sphere reads:

$$(3.7) \quad S_n = N \operatorname{Tr} \left\{ M^\dagger E' M + M E' M^\dagger + \left(\hat{m}^2 + \frac{8\pi}{3} \right) M^\dagger M + \frac{\hat{g}^2}{2} \left(M^\dagger M \right)^2 \right\},$$

where the matrix E' coming from the kinetic operator \hat{K} is

$$(3.8) \quad E'_{\ell_1 \ell_2} = \left(-\frac{32\pi}{3N} \left(\ell_1 + \frac{1}{4} \right) \right) \delta_{\ell_1 \ell_2} + \mathcal{O} \left(\frac{1}{N^2} \right).$$

Therefore, we find that, in the large N limit, this model coincides with the LSZ at the self-dual point, with magnetic field scaled by a factor $-\frac{8}{3}$. Note that the restriction to preferred points in parameter space, both for the LSZ and the model of [32], is crucial to obtain the result. Indeed, the large amount of symmetry at these preferred points produces cancellations which do not hold in general.

4. SOLUTION OF THE MATRIX MODEL

As we have seen in sections 2 and 3, different noncommutative field theories reduce to a matrix model of the form (1.1)

$$\mathcal{Z}_{\text{LSZ}}(E, \tilde{E}) = \int \mathcal{D}M \mathcal{D}M^\dagger \exp\left(-N \text{Tr} \left\{ M E M^\dagger + M^\dagger \tilde{E} M + \widehat{V}(M^\dagger M) \right\}\right)$$

in terms of $N \times N$ complex matrices, depending on the insertion of two external matrices. The potential $\widehat{V}(M^\dagger M)$ is a quadratic polynomial with dimensionless coefficients

$$(4.1) \quad \widehat{V}(M^\dagger M) = \widehat{m}^2 (M^\dagger M) + \frac{\widehat{g}^2}{2} (M^\dagger M)^2.$$

We also let the external fields E, \tilde{E} have arbitrary eigenvalues which, for convenience and consistently with the noncommutative field theory setting (recall Eq.(2.3)), we write as $\frac{4\pi}{N}\eta_\ell, \frac{4\pi}{N}\tilde{\eta}_\ell$ respectively, for $\ell = 1, \dots, N$. In particular, the original LSZ model, discussed in section 2, corresponds to external matrices with $\eta_\ell = \tilde{\eta}_\ell$ given by consecutive integers plus a constant shift, see Eq. (2.3). The unconventional presence of \tilde{E} explicitly breaks $U(N)$ symmetry, but the system is still tractable.

We now reduce this matrix model to an ordinary multiple integral in terms of the spectra of the external fields, and show that the results can be written in terms of the observables of $U(N)$ Chern–Simons theory in \mathbb{S}^3 , with $q = e^{-g_s}$ real.

4.1. General solution. As shown in [9], we can approach the solution using the singular value decomposition of a generic $N \times N$ complex matrix M :

$$M = U_1^\dagger \text{diag}(\lambda_1, \dots, \lambda_N) U_2,$$

with $U_{\alpha=1,2}$ unitary matrices and $\lambda_\ell \geq 0$.² The measure becomes

$$\mathcal{D}M \mathcal{D}M^\dagger = [dU_1] [dU_2] \prod_{\ell=1}^N dy_\ell \Delta_N [y]^2,$$

where $[dU_\alpha]$ is the invariant Haar measure over $U(N)$, and $y_\ell := \lambda_\ell^2$ and

$$(4.2) \quad \Delta_N [y] = \prod_{1 \leq \ell < \ell' \leq N} (y_\ell - y_{\ell'})$$

is the Vandermonde determinant.

As shown in [9], the use of this transformation implies that integrations over $U_{\alpha=1,2} \in U(N)$ decouple over the two types of external field terms. Denoting by $Y = \text{diag}(y_1, \dots, y_N)$, the angular degrees of freedom U_α can be integrated out using the Harish-Chandra–Itzykson–Zuber (HCIZ) formula [59, 60]:

$$(4.3) \quad \int_{U(N)} [dU_2] \exp\left\{-N \text{Tr} \left(E U_2^\dagger Y U_2 \right)\right\} = \mathcal{C}_N \frac{\det_{1 \leq \ell, \ell' \leq N} (e^{-4\pi\eta_\ell y_{\ell'}})}{\Delta_N [\eta] \Delta_N [y]},$$

²See [58] for comments on this parametrization with regards to the more usual one in terms of eigenvalues [21]. In any case, this transformation is much used and very useful when studying complex matrix models, such as the LSZ.

where we have used the explicit form of the eigenvalues of E and denoted

$$\mathcal{C}_N := (4\pi)^{-\frac{N(N-1)}{2}} \prod_{j=1}^{N-1} j! = (4\pi)^{-\frac{N(N-1)}{2}} G(N+1),$$

where $G(\cdot)$ is the Barnes G -function.

Analogous expression is obtained for integration over U_1 replacing E by \tilde{E} . For a potential $\widehat{V}(M^\dagger M)$ as in (4.1), one gets:

$$(4.4) \quad \begin{aligned} \mathcal{Z}_{\text{LSZ}}(E, \tilde{E}) &= \int_{[0, \infty)^N} \Delta_N[y]^2 \prod_{\ell=1}^N e^{-\left(\widehat{m}^2 y_\ell + \frac{\widehat{g}^2}{2} y_\ell^2\right)} dy_\ell \\ &\times \int [dU_2] \exp\left(-NEU_2^\dagger Y U_2\right) \int [dU_1] \exp\left(-N\tilde{E}U_1^\dagger Y U_1\right) \end{aligned}$$

Plugging HCIZ (4.3) into (4.4), we note that the partition function for the matrix model is:

$$\mathcal{Z}_{\text{LSZ}}(E, \tilde{E}) = \mathcal{C}_N^2 \int_{[0, \infty)^N} \Delta_N[y]^2 \frac{\det_{1 \leq \ell, \ell' \leq N} (e^{-4\pi\eta_\ell y_{\ell'}}) \det_{1 \leq \ell, \ell' \leq N} (e^{-4\pi\tilde{\eta}_\ell y_{\ell'}})}{\Delta_N[y]^2 \Delta_N[\eta] \Delta_N[\tilde{\eta}]} \prod_{\ell=1}^N e^{-\widehat{m}^2 y_\ell - \frac{\widehat{g}^2}{2} y_\ell^2} dy_\ell,$$

where we recall that η_ℓ (respectively $\tilde{\eta}_\ell$) stand for the eigenvalues of E (respectively \tilde{E}), up to a factor $\frac{4\pi}{N}$. This was already done in [9] but now, after applying the HCIZ formula, we will not expand the resulting determinants into sums of permutations of N objects, and cancel a Vandermonde squared instead. That is, after a suitable rescaling of the integration variables:

$$(4.5) \quad \mathcal{Z}_{\text{LSZ}}(E, \tilde{E}) = \frac{\mathcal{C}'_N}{\Delta_N[\eta] \Delta_N[\tilde{\eta}]} \int_{[0, \infty)^N} \det_{1 \leq \ell, \ell' \leq N} (e^{-\eta_\ell y_{\ell'}}) \det_{1 \leq \ell, \ell' \leq N} (e^{-\tilde{\eta}_\ell y_{\ell'}}) \prod_{\ell=1}^N e^{-m^2 y_\ell - \frac{g^2}{2} y_\ell^2} dy_\ell,$$

with coefficients redefined as:

$$(4.6) \quad m^2 = \frac{\widehat{m}^2}{4\pi}, \quad g^2 = \frac{\widehat{g}^2}{(4\pi)^2},$$

and

$$\mathcal{C}'_N = (4\pi)^{-N} \mathcal{C}_N^2 = \frac{G(N+1)^2}{(4\pi)^{N^2}} = \frac{2^{-N(N-1)} \pi^N}{\text{vol}(U(N))^2},$$

where $\text{vol}(\cdot)$ is the volume of the gauge group. Note that this normalization is essentially the square of the partition function of the Gaussian unitary ensemble (GUE) [21].

In the theory of non-intersecting Brownian motion, the determinants in (4.5) are very familiar. This whole theory of determinantal processes is known to be directly related to $U(N)$ Chern–Simons theory on \mathbb{S}^3 and with the Wess–Zumino–Witten (WZW) model [42], where such connection was shown to follow from specializations of the determinants

$$\det_{1 \leq \ell, \ell' \leq N} (e^{-\eta_\ell y_{\ell'}}), \quad \det_{1 \leq \ell, \ell' \leq N} (e^{-\tilde{\eta}_\ell y_{\ell'}})$$

in (4.5). However, it is more direct to show the relation through the corresponding matrix model formulation. Recall for this the definition of a Schur polynomial [61]

$$s_\mu(x_1, \dots, x_N) = \frac{\det_{1 \leq \ell, \ell' \leq N} (x_\ell^{\mu_{\ell'} + N - \ell'})}{\det_{1 \leq \ell, \ell' \leq N} (x_\ell^{N - \ell'})},$$

and hence we rewrite the scaled eigenvalues $\eta_\ell, \tilde{\eta}_\ell$ of the external matrices E, \tilde{E} , assuming they are integers, as:

$$(4.7) \quad \begin{aligned} \eta_\ell &= \mu_\ell + N - \ell, \\ \tilde{\eta}_\ell &= \nu_\ell + N - \ell, \end{aligned}$$

for $\ell = 1, \dots, N$. Without loss of generality, we can relabel the eigenvalues so that $\mu_1 \geq \mu_2 \geq \dots \geq \mu_N \geq 0$, and likewise for $\nu_1 \geq \dots \geq \nu_N$. Then, using

$$\prod_{1 \leq \ell < \ell' \leq N} (e^{y_\ell} - e^{y_{\ell'}})^2 = \prod_{1 \leq \ell < \ell' \leq N} \left(2 \sinh \left(\frac{y_\ell - y_{\ell'}}{2} \right) \right)^2 \prod_{m=1}^N e^{(N-1)y_m}$$

and reflecting the variables $y_\ell \mapsto -y_\ell$, we immediately have³:

$$(4.8) \quad \mathcal{Z}_{\text{LSZ}}(\mu, \nu) = \frac{(-1)^N \mathcal{C}'_N}{\Delta_N[\eta] \Delta_N[\tilde{\eta}]} \int_{(-\infty, 0]^N} \prod_{1 \leq \ell < \ell' \leq N} \left(2 \sinh \left(\frac{y_\ell - y_{\ell'}}{2} \right) \right)^2 \prod_{\ell=1}^N dy_\ell e^{\beta y_\ell - \frac{g^2}{2} y_\ell^2} \\ \times s_\mu(e^{y_1}, \dots, e^{y_N}) s_\nu(e^{y_1}, \dots, e^{y_N}),$$

with $\beta = m^2 - N + 1$. We henceforth write $\mathcal{Z}_{\text{LSZ}}(\mu, \nu)$ for $\mathcal{Z}_{\text{LSZ}}(E, \tilde{E})$, stressing the dependence on the partitions $\mu = (\mu_1, \dots, \mu_N)$ and $\nu = (\nu_1, \dots, \nu_N)$. Except for the integration domain, the expression (4.8) is close to the general version of the $U(N)$ Chern–Simons on \mathbb{S}^3 matrix model, with two different insertions of Schur polynomials, whose evaluation gives the topological invariant of a pair of unknots [14, 62] carrying the $U(N)$ representations μ and ν . It is worthwhile to mention that the same matrix model (4.8) is also related to the Hopf link invariant, but now with representations μ and ν^* , where $\nu^* := (\nu_1 - \nu_N, \nu_1 - \nu_{N-1}, \dots, 0)$. We give the details in Appendix A.3. Furthermore, if the external matrix E has positive integer eigenvalues while \tilde{E} has negative integers eigenvalues, or vice versa, the relation then is with the Hopf link invariant carrying representations μ and ν .

Instead, after a shift of variables in (4.8) and using the identity (A.1), we obtain the matrix model representation

$$\mathcal{Z}_{\text{LSZ}}(\mu, \nu) = \mathcal{A}_N(\mu, \nu) \int_{(-\infty, -\gamma]^N} \prod_{j=1}^N dx_j e^{-\frac{g^2}{2} \sum_{j=1}^N x_j^2} \prod_{1 \leq j < k \leq N} \left(2 \sinh \left(\frac{x_j - x_k}{2} \right) \right)^2 \\ \times s_\mu(e^{x_1}, \dots, e^{x_N}) s_\nu(e^{x_1}, \dots, e^{x_N}),$$

where $\gamma = \beta/g^2$ and

$$(4.9) \quad \mathcal{A}_N(\mu, \nu) := \frac{(-1)^N \mathcal{C}'_N}{\Delta_N[\eta] \Delta_N[\tilde{\eta}]} \exp \left(\frac{\gamma^2 N}{2} + \gamma (|\mu| + |\nu|) \right).$$

Notice also that the Vandermonde factors in the denominator of (4.9), which depend exclusively on the eigenvalues of the external matrices, can be written, using Weyl's denominator formula (see Appendix A), as

$$\begin{aligned} \Delta_N[\eta] &= \prod_{1 \leq \ell < \ell' \leq N} (\mu_\ell - \mu_{\ell'} - \ell + \ell') = G(N+1) \dim \mu, \\ \Delta_N[\tilde{\eta}] &= \prod_{1 \leq \ell < \ell' \leq N} (\nu_\ell - \nu_{\ell'} - \ell + \ell') = G(N+1) \dim \nu, \end{aligned}$$

³With our definition, η_ℓ and $\tilde{\eta}_\ell$ are non-negative integers, which is the physical choice motivated by the LSZ model. If we want them to be non-positive integers, we simply do not reflect y_ℓ and define $\eta_\ell, \tilde{\eta}_\ell$ with opposite sign. The discussion would be exactly the same, except for the integration domain being $[0, \infty)$.

These Barnes G -functions cancel in (4.9) against the ones coming from the double application of HCIZ formula. We finally obtain

$$(4.10) \quad \boxed{\mathcal{Z}_{\text{LSZ}}(\mu, \nu) = C_N(\gamma, |\mu|, |\nu|) \frac{\mathcal{Z}_{\text{CS}} \langle W_\mu W_\nu \mathbb{1}_{(-\infty, -\gamma]^N} \rangle_{\text{CS}}}{\dim \mu \dim \nu}},$$

where the relation between the partitions μ, ν and the external matrices of the generalized LSZ model is given in equation (4.7), and

$$C_N(\gamma, |\mu|, |\nu|) = \frac{(-1)^N e^{\frac{\gamma^2 N}{2} + \gamma(|\mu| + |\nu|)}}{(4\pi)^{N^2}}.$$

In formula (4.10), \mathcal{Z}_{CS} is the $U(N)$ Chern–Simons partition function on \mathbb{S}^3 , which is a quantum topological invariant also known as Witten–Reshetikhin–Turaev invariant [63–65], defined as the matrix model (1.3) and whose explicit evaluation we give below. Besides, W_μ is the trace of the holonomy of the gauge connection along an unknot inside \mathbb{S}^3 , taken in the $U(N)$ representation corresponding to the partition μ , and likewise for W_ν . In (4.10), the Chern–Simons coupling is $g_s = \frac{1}{g^2}$. In Chern–Simons theory, g_s is related to the Chern–Simons level k by $g_s = \frac{2\pi i}{N+k}$, while the real string coupling constant g_s is used when describing topological strings. That is the same type of description here, since g^2 is real. Finally, $\langle \cdots \rangle_{\text{CS}}$ in (4.10) means the average in the Chern–Simons matrix model (1.3), and $\mathbb{1}_{(-\infty, -\gamma]^N}$ is the N -dimensional indicator function

$$\mathbb{1}_{(-\infty, -\gamma]^N}(x_1, \dots, x_N) = \prod_{j=1}^N \mathbb{1}_{(-\infty, -\gamma]}(x_j) = \begin{cases} 1, & x_j \leq -\gamma \quad \forall j = 1, \dots, N, \\ 0, & \text{otherwise.} \end{cases}$$

Therefore, $\langle W_\mu W_\nu \mathbb{1}_{(-\infty, -\gamma]^N} \rangle_{\text{CS}}$ would correspond to the two-unknot invariant, but averaged only using variables $x_j \leq -\gamma$ (instead of $x_j \in \mathbb{R}$). Its relation with the actual invariant of a pair of unknots is further discussed in subsection 4.4 through the lenses of random matrix theory.

4.2. Quantum dimensions. An important particular case of the above general setting is when one of the partitions in (4.7) is void. That is, in terms of LSZ theory, one has an external matrix with the equispaced spectra and the other one generalized with a partition, see the definitions (2.3) and (4.7). This case corresponds, as we shall see, to quantum dimensions in the Chern–Simons interpretation [19]. Quantum dimension of a representation associated to the partition μ is given by the following hook-content formula [19, 61]

$$\dim_q \mu := \prod_{x \in \mu} \frac{[N + c(x)]_q}{[h(x)]_q},$$

where for each box $x \equiv (j, k)$ of the Young diagram determined by μ , the quantity $h(x) := \mu_j + \mu'_k - j - k + 1$ is the hook length, with the prime meaning conjugate diagram, and $c(x) := j - k$ is known as the content of the box x . The operation $[\cdot]_q$ denotes the symmetric q -number, that is

$$[n]_q = \frac{q^{n/2} - q^{-n/2}}{q^{1/2} - q^{-1/2}}.$$

In Chern–Simons theory on \mathbb{S}^3 , the unknot invariant is given by quantum dimensions [14, 19]. Since one of the two external matrices has harmonic oscillator spectrum the matrix model

above reduces to

$$\mathcal{Z}_{\text{LSZ}}(\mu, \emptyset) = \mathcal{A}_N(\mu, \emptyset) \int_{(-\infty, -\gamma]^N} \prod_{1 \leq j < k \leq N} \left(2 \sinh \left(\frac{x_j - x_k}{2} \right) \right)^2 \prod_{j=1}^N e^{-\frac{g^2}{2} x_j^2} dx_j \\ \times s_\mu(e^{x_1}, \dots, e^{x_N}),$$

whose evaluation leads to

$$(4.11) \quad \boxed{\mathcal{Z}_{\text{LSZ}}(\mu, \emptyset) = C_N(\gamma, |\mu|, 0) \frac{\mathcal{Z}_{\text{CS}} \langle W_\mu \mathbb{1}_{(-\infty, -\gamma]^N} \rangle_{\text{CS}}}{\dim \mu}}.$$

This should be compared with the unknot invariant computed in Chern–Simons theory, which differs from the present setting in the fact that the integral is taken over \mathbb{R}^N instead of $(-\infty, -\gamma]^N$. We will come back to this point in subsection 4.4. The exact evaluation of the unknot invariant gives [19]

$$\langle W_\mu \rangle = q^{-\frac{1}{2} C_2(\mu)} \dim_q \mu,$$

where $q = e^{-1/g^2}$. Besides, in the expression above the term

$$C_2(\mu) = (N+1)|\mu| + \sum_{\ell=1}^N (\mu_\ell^2 - 2\ell\mu_\ell)$$

is the $U(N)$ quadratic Casimir of the representation μ , labelled by the Young diagram associated to the partition μ , with μ_ℓ boxes in the ℓ -th row, with rows understood to be aligned on the left.

The appearance of quantum dimensions is interesting in that they appear as well in the study of noncommutative gauge theories through the analysis of WZW D-branes [39, 66]. However, as we have seen in section 2, only the simpler setting, where the two external matrices are equal and have harmonic oscillator spectra, is directly linked to a noncommutative scalar theory.

4.3. Chern–Simons and LSZ partition function. We have shown in section 2 that our study of noncommutative scalar field theory naturally leads to a LSZ matrix model with $E = \tilde{E}$ and spectra $\frac{4\pi}{N}(\ell - \frac{1}{2})$ for $\ell = 1, \dots, N$, see Eq. (2.3) (or the same but scaled and shifted, see (3.8)). Thus, we consider now the case in which both partitions are void: this corresponds to the two external matrices having harmonic oscillator spectra. In particular, from (4.7), we have that $\eta_\ell = N - \ell$ for $\ell = 1, \dots, N$. The fact that the two physical spectra have an overall energy shift only has an impact at the level of renormalization of the mass parameter. This follows immediately from a simple property of Schur polynomials, given in Appendix A.

Then one obtains the matrix model without Schur polynomial insertions, and the corresponding matrix integral is related to the one for the Chern–Simons partition function, given in (1.3) (recall that $g_s = 1/g^2$). The Chern–Simons matrix model has the exact solution [18]:

$$(4.12) \quad \mathcal{Z}_{\text{CS}} = \left(\frac{2\pi}{g^2} \right)^{N/2} N! e^{\frac{N(N+1)(N-1)}{6g^2}} \prod_{j=1}^{N-1} (1 - q^j)^{N-j},$$

with $q = e^{-1/g^2}$ as above. The product can also be written as a q -deformed Barnes function, which, in the limit $g \rightarrow \infty$ (which is $q \rightarrow 1$), reduces to the Barnes G -function. Then, the

LSZ matrix model partition function is

$$(4.13) \quad \mathcal{Z}_{\text{LSZ}}(\emptyset, \emptyset) = \frac{(-1)^N e^{\frac{\gamma^2 N}{2}}}{(4\pi)^{N^2}} \mathcal{Z}_{\text{CS}} \langle \mathbb{1}_{(-\infty, -\gamma]^N} \rangle_{\text{CS}}.$$

Notice that, apart from the simple prefactor, the ratio between the LSZ and the Chern–Simons partition function is the average $\langle \mathbb{1}_{(-\infty, -\gamma]^N} \rangle_{\text{CS}}$ of the N -dimensional indicator function. This aspect is further analyzed the next subsection.

4.4. Probabilistic interpretation. The main result (4.10) and its specializations (4.11) and (4.13) admit an interpretation in terms of probabilities of large deviations of the smallest or largest eigenvalue of Hermitian random matrices. Consider a generic weight function $\omega(x)$, and let \mathcal{Z}_ω be the associated Hermitian random matrix ensemble,

$$(4.14) \quad \mathcal{Z}_\omega = \int_{\mathbb{R}^N} \Delta_N[x]^2 \prod_{j=1}^N \omega(x_j) dx_j.$$

The probability that the largest eigenvalue of a random matrix in the ensemble (4.14) is smaller than a given threshold $s \in \mathbb{R}$ is

$$(4.15) \quad \begin{aligned} \text{Prob}_\omega(x_{\max} \leq s) &= \text{Prob}_\omega(x_1 \leq s, \dots, x_N \leq s) \\ &= \frac{1}{\mathcal{Z}_\omega} \int_{(-\infty, s]^N} \Delta_N[x]^2 \prod_{j=1}^N \omega(x_j) dx_j = \left\langle \mathbb{1}_{(-\infty, s]^N} \right\rangle_\omega, \end{aligned}$$

where in the last expression $\langle \cdot \rangle_\omega$ means the average in the ensemble (4.14) and $\mathbb{1}_{(-\infty, s]^N}$ is the N -dimensional indicator function. In the Coulomb gas picture, replacing the weight $\omega(x)$ by $\omega(x)\mathbb{1}_{(-\infty, s]}(x)$ introduces a hard wall placed at $x = s$ which leaves the charges on its left. See [67] for the large N limit of matrix models in presence of hard walls. For the GUE ensemble, the probabilities of large fluctuations at large N have been found in [24, 68].

We immediately see from equation (4.13) that

$$\frac{\mathcal{Z}_{\text{LSZ}}(\emptyset, \emptyset)}{\mathcal{Z}_{\text{CS}}} = \frac{(-1)^N e^{\frac{\gamma^2 N}{2}}}{(4\pi)^{N^2}} \text{Prob}_{\text{CS}}(x_{\max} \leq -\gamma),$$

hence the ratio between the partition function of the LSZ model and that of Chern–Simons theory is effectively computing the probability of large deviations of eigenvalues in the Chern–Simons ensemble (1.3) (or, strictly speaking, in the Stieltjes–Wigert ensemble [18], see below), up to a completely determined, parameter-dependent overall factor. More in general, formula (4.10) states that

$$\frac{\mathcal{Z}_{\text{LSZ}}(\mu, \nu)}{\mathcal{Z}_{\text{CS}}} = C_N(\gamma, |\mu|, |\nu|) \frac{\langle W_\mu W_\nu \rangle_{\text{CS}}}{\dim \mu \dim \nu} \text{Prob}_{\text{CS}; W_\mu W_\nu}(x_{\max} \leq -\gamma),$$

where by $\text{Prob}_{\text{CS}; W_\mu W_\nu}(\cdot)$ we mean the probability in the matrix ensemble computing the invariant of two unknots, that is, in the Chern–Simons ensemble (1.3) with the insertion of two Schur polynomials. This probability must be normalized by $\mathcal{Z}_{\text{CS}} \cdot \langle W_\mu W_\nu \rangle_{\text{CS}}$ and not only \mathcal{Z}_{CS} . Therefore, the LSZ matrix model with general assignment of the external matrices, divided by the Chern–Simons partition function, is proportional to the two-unknot invariant weighted by the probability of large deviations in the random matrix description of such topological invariant. The proportionality constant yields an elementary dependence on the size N and on the free parameters of the generalized LSZ theory.

We emphasize that the LSZ partition function does not compute the (typically small) fluctuations of the largest eigenvalue around the edge of the eigenvalue distribution. Equivalently, in the Coulomb gas picture, the LSZ partition function does not describe the fluctuations of

the rightmost charge around the equilibrium. What it gives is the probability of an atypically large fluctuation, with the greatest eigenvalue moving deep into the bulk of the eigenvalue distribution. This is a q -analogue of the large fluctuations in the GUE discussed in [24, 68] (see also [69, 70]). Besides, we underline that, differently from [24] where a large deviation from the equilibrium configuration is of order $\sim \sqrt{N}$, in the q -analogue the support of the eigenvalue density grows as Ng_s , thus large deviations from the equilibrium are of order $\sim N$.

Now, if we consider equation (4.5) without Schur insertions, $\mu = \emptyset = \nu$, then, changing variables $u_j = e^{y_j - (m^2 + 2N - 1)/g^2}$ [18], it is well-known that we have a standard-random matrix ensemble with a log-normal weight, named Stieltjes–Wigert (SW) ensemble. That is:

$$(4.16) \quad \mathcal{Z}_{\text{LSZ}}(\emptyset, \emptyset) = \mathcal{B}_N \int_{[s, \infty)^N} \prod_{1 \leq j < k \leq N} (u_j - u_k)^2 \prod_{j=1}^N e^{-\frac{1}{2g_s}(\ln u_j)^2} du_j,$$

for $g_s = 1/g^2$ and $s = e^{\frac{m^2 + 2N - 1}{g^2}}$, and with

$$\mathcal{B}_N := (4\pi)^{-N^2} \exp \left[\frac{N}{2g^2} (m^2 - 1) (m^2 + 2N - 1) \right].$$

Note that a SW ensemble is not centered around 0, as the weight is supported on $u \geq 0$. Therefore, up to a proportionality factor, the LSZ partition function $\mathcal{Z}_{\text{LSZ}}(\emptyset, \emptyset)$ normalized by the SW partition function gives the probability of atypically large fluctuations of the eigenvalues away from the left edge at $u = 0$, with the smallest eigenvalue deep into the bulk, $u_{\min} \geq s$, in a Stieltjes–Wigert ensemble:

$$\frac{\mathcal{Z}_{\text{LSZ}}(\emptyset, \emptyset)}{\mathcal{Z}_{\text{SW}}} = \mathcal{B}_N \text{Prob}_{\text{SW}}(u_{\min} \geq s).$$

To evaluate this quantity numerically at finite N , it is convenient to rewrite the integrals in the numerator and denominator of $\text{Prob}_{\text{SW}}(u_{\min} \geq s)$ as determinants [21], obtaining:

$$\begin{aligned} \text{Prob}_{\text{SW}}(u_{\min} \geq s) &= \frac{\det_{1 \leq j, k \leq N} \left[\int_s^\infty u^{j+k-2} e^{-\frac{1}{2g_s}(\ln u)^2} du \right]}{\det_{1 \leq j, k \leq N} \left[\int_0^\infty u^{j+k-2} e^{-\frac{1}{2g_s}(\ln u)^2} du \right]} \\ &= 2^{-N} \frac{\det_{1 \leq j, k \leq N} \left[e^{\frac{g_s}{2}(j+k-1)^2} \text{erfc} \left(\sqrt{\frac{g_s}{2}}(m^2 + 2N - j - k) \right) \right]}{\det_{1 \leq j, k \leq N} \left[e^{\frac{g_s}{2}(j+k-1)^2} \right]}. \end{aligned}$$

In the latter expression, $\text{erfc}(z) = 1 - \text{erf}(z)$ is the complementary error function. The denominator is known, and is readily extracted from Eq. (4.12). We obtain:

$$\begin{aligned} \text{Prob}_{\text{SW}}(u_{\min} \geq s) &= \frac{q^{\frac{N}{3}(2N^2-1)}}{2^N (1-q)^{\frac{N}{2}(N-1)} \prod_{j=1}^{N-1} (1-q^j)^{N-j}} \\ &\quad \times \det_{1 \leq j, k \leq N} \left[e^{\frac{g_s}{2}(j+k-1)^2} \text{erfc} \left(\sqrt{\frac{g_s}{2}}(m^2 + 2N - j - k) \right) \right], \end{aligned}$$

and we recall that $q = e^{-g_s}$ and the argument of erfc is related to s through $\ln s = g_s(m^2 + 2N - 1)$. We plot the logarithm of this probability at finite N in figure 1. It is also clear from the definition of s that the role of any fixed m^2 becomes less relevant as N is increased, unless m^2 itself is increased linearly with N . This aspect is shown in figure 2.

We stress once more that the present setting is very different from the study of (typical) small fluctuation of the largest or smallest eigenvalue around the edge, which are suppressed by inverse powers of N . See [24, 69] for thorough discussion on this point, and [71] for further insights in the theory of large deviations. In the more general case, $\mathcal{Z}_{\text{LSZ}}(\mu, \nu)$ is




FIGURE 1. $-\frac{1}{N^3} \ln \frac{\mathcal{Z}_{\text{LSZ}}(\emptyset, \emptyset)}{\mathcal{B}_N \mathcal{Z}_{\text{SW}}}$ as a function of the coupling g_s at $m^2 = 0$. The curves correspond to N from 2 to 7.




FIGURE 2. $-\frac{1}{N^3} \ln \frac{\mathcal{Z}_{\text{LSZ}}(\emptyset, \emptyset)}{\mathcal{B}_N \mathcal{Z}_{\text{SW}}}$ as a function of the coupling g_s at $m^2 = 2$ (left) and $m^2 = \frac{N}{2} + 1$ (right). The curves correspond to N from 2 to 7.

proportional to the probability of a large fluctuation of the eigenvalues away from the left edge in a SW ensemble deformed by the insertion of two Schur polynomials $s_\mu(u_1, \dots, u_N)$ and $s_\nu(u_1, \dots, u_N)$.

4.5. Large N limit. In the LSZ model, the parameter N regularizes the path integral of the NC field theory, as discussed in section 2. Therefore, it is natural to consider the large N limit of $\mathcal{Z}_{\text{LSZ}}(\emptyset, \emptyset)$, and more generally of $\mathcal{Z}_{\text{LSZ}}(\mu, \nu)$. In turn, the probabilistic interpretation of the Chern–Simons theory observables also calls for a large N analysis, from the perspective of large deviations theory [71].

The knowledge of the large N solution of the matrix model $\mathcal{Z}_{\text{LSZ}}(\mu, \nu)$ on either the complex matrix model side or on the side of the probability in Chern–Simons observables, would directly provide the solution on the other side. However, there are difficulties in solving the large N limit from both perspectives. As mentioned in subsection 4.1, and already observed in the original work [9], a difficulty in the study of the LSZ partition function, generalized to arbitrary external matrices E, \tilde{E} , comes from the presence of \tilde{E} , which prevents a reformulation of the model in terms of a single Hermitian matrix $M^\dagger M$. Therefore, it becomes hard to solve the large N limit of the matrix model explicitly, using for example the loop equations (that are zero-dimensional Schwinger–Dyson equations) [72] when $\tilde{E} \neq 0$. For $\tilde{E} = 0$, the large N solution has been found in [9], but this choice does not yield meaningful observables in Chern–Simons theory.

Taking the converse route, we could try to analyze the large N asymptotics of (4.16), adapting the argument of [24] to the present case. This would amount to solve a constrained extremization problem defined as follows. Write

$$t \equiv g_s N = N/g^2, \quad z \equiv \log s = g_s(m^2 + 2N - 1)$$

and take the large N 't Hooft limit of (4.16) keeping t and z fixed. Let $\rho_{\text{SW}}(u)$ be the constrained SW eigenvalue density at large N , supported on the interval $[z, L(z)]$. We have stressed the dependence of the upper bound $L(z)$ on z , and let the more standard dependence on the 't Hooft coupling t implicit. Then $\rho_{\text{SW}}(u)$ solves the saddle point equation

$$(4.17) \quad \text{P} \int_z^{L(z)} du' \frac{\rho_{\text{SW}}(u')}{u - u'} = \frac{1}{2t} \frac{\ln u}{u},$$

with the symbol $\text{P} \int$ meaning the Cauchy principal value integral. The upper boundary of the support $L(z)$ is fixed as a function of z and t by the normalization condition

$$\int_z^{L(z)} du \rho_{\text{SW}}(u) = 1.$$

This problem cannot be solved by the method of [24], because of the non-polynomial form of the right-hand side of (4.17). More precisely, the extremization problem described by the saddle point equation (4.17) does not satisfy the hypothesis of Tricomi's theorem. A complete solution at large N would entail an extension of the method applied in [24, 68] to q -ensembles.

There are, nevertheless, two simplifying limits in which the matrix model becomes tractable at large N :

- The $g^2 \rightarrow \infty$ limit. The interaction term in the LSZ action dominates, and the fields become non-dynamical. This corresponds to $g_s \rightarrow 0$, that is, $q \rightarrow 1$ from below. In this limit, we recover the Gaussian ensemble from the SW ensemble (up to an overall factor, readable from (4.12)), and the results of [24] directly apply to the present setting.
- The $g^2 \rightarrow 0$ limit. The quartic interaction should become tractable in standard perturbation theory in the LSZ field theory. From the Chern–Simons perspective, this corresponds to $g_s \rightarrow \infty$, equivalently $q \rightarrow 0$ from above, and simplifications take place.

In the first case, the eigenvalue density is [24]

$$\lim_{g_s \rightarrow 0^+} \rho_{\text{SW}}(u) = \frac{1}{2\pi} \sqrt{\frac{L(z) - u}{u - z}} [L(z) - z + 2u],$$

with $L(z) = \frac{2}{3}\sqrt{z^2 + 6} + \frac{1}{3}z$, and now $z \equiv \gamma/\sqrt{N}$. From [24, Eq. (58)] and a change of variables $x \mapsto \sqrt{2g_s}x$ we have

$$\begin{aligned} \ln \text{Prob}_{\text{CS}}(x_{\max} \leq -\gamma) &= \ln \text{Prob}_{\text{CS}}(x_{\min} \geq \gamma) \\ &\approx \frac{N^2}{2} \ln \frac{g_s}{2} - \frac{N^2}{54} \left[36z^2 - z^4 + z(z^2 + 15)\sqrt{z^2 + 6} + 27 \left[\ln 18 - 2 \ln \left(\sqrt{z^2 + 6} - z \right) \right] \right] \end{aligned}$$

in the $N \rightarrow \infty$ and $g_s \rightarrow 0$ limit. This formula together with (4.13) describes the large N limit of the LSZ model in the strong interaction regime $g^2 \rightarrow \infty$.

The converse limit $g_s \rightarrow \infty$, that is, $g^2 \rightarrow 0$, can be analyzed as well. Changing variables $x \mapsto \sqrt{2g_s}x$ in the CS ensemble and using

$$2 \sinh \left(\sqrt{\frac{g_s}{2}} (x_j - x_k) \right) \approx \exp \left(\sqrt{\frac{g_s}{2}} |x_j - x_k| \right)$$

when $g_s \rightarrow \infty$, we get

$$\mathcal{Z}_{\text{LSZ}}(\emptyset, \emptyset) = \frac{(-1)^N e^{\frac{\gamma^2 N}{2}}}{(4\pi)^{N^2}} \left(\frac{g_s}{2}\right)^{\frac{N^2}{2}} \int_{(-\infty, -\gamma]^N} \prod_{j=1}^N dx_j \exp \left[-\sum_{j=1}^N x_j^2 + \sqrt{\frac{g_s}{2}} \sum_{j \neq k} |x_j - x_k| \right].$$

The latter expression describes the partition function of a constrained one-dimensional Coulomb gas [73, 74], corresponding to placing a hard wall at $x = -\gamma$ in a jellium model [75]. Here we derive the large N asymptotics at leading order for this constrained model.

Assuming x grows as $x = \xi N^\alpha$ in the large N limit, for some $\alpha > 0$ and fixed ξ , the first term in the exponential grows as $N^{1+2\alpha}$, while the second term grows as $N^{2+\alpha}$. A non-trivial saddle point exists for $\alpha = 1$. Therefore, the large N limit in this large g_s regime is governed by the eigenvalue density $\rho_0(\xi)$ that solves the saddle point equation

$$\sqrt{\frac{g_s}{2}} \int_z^{L(z)} d\xi' \rho_0(\xi') \text{sign}(\xi - \xi') = \xi,$$

with, this time, $z = \gamma/N$, for the scaling of γ with N to be consistent with the growth of the eigenvalues. Splitting the integral in two pieces, with $\xi' < \xi$ and $\xi' > \xi$ respectively, and taking the derivative of the saddle point equation in the interior of the domain, $z < \xi < L(z)$, we find

$$\rho_0(\xi) = \frac{1}{\sqrt{2g_s}}.$$

The normalization condition then imposes

$$\int_z^{L(z)} d\xi \rho_0(\xi) = 1 \implies L(z) - z = \sqrt{2g_s}.$$

With this eigenvalue density we obtain

$$\ln(-1)^N \mathcal{Z}_{\text{LSZ}}(\emptyset, \emptyset) \approx -N^3 \left[z^2 + z\sqrt{2g_s} + \frac{g_s}{3} \right]$$

at leading order in N and in the large g_s regime.

The complete solution will then interpolate between this two limiting situations. The N^2 -behaviour will correspond to small 't Hooft coupling, $t \rightarrow 0$, while letting t grow linearly with N will give back the N^3 -behaviour.

4.6. Energy levels of the generalized LSZ model. To conclude this section, we briefly comment on the interpretation of the generalized result (4.10) in terms of the LSZ noncommutative scalar field theory [8, 9]. The absence of a preferred temporal direction in the Moyal plane prevents us from a meaningful Hilbert space picture of the partition function (4.13), and of its generalization (4.10) as well. However, we can still provide an interpretation in terms of energy density levels. The system in facts splits into two clearly separated (on the LSZ side) contributions: a Landau Hamiltonian density and a quartic interaction term. Throughout the solution, the former is encoded in the Vandermonde determinants, together with the Schur polynomials in the most general case, while the interaction appears as the Gaussian measure in (1.3).

In the pure LSZ case, with both external matrices with harmonic oscillator spectra, the Landau levels are $\frac{2\ell-1}{\theta}$, being θ the noncommutativity parameter (recall the details from section 2). Introducing equal partitions $\mu = \nu$ to generalize the external matrices would correspond to a distortion of the Landau spectrum. As an example, the insertion of an antisymmetric partition $\mu = (1, \dots, 1)$ shifts the whole spectra by one level in the negative direction; as shown in Appendix A, this corresponds to a renormalization of the mass. The symmetric partition $\mu = (N, 0, \dots, 0)$ determines the same spectrum, although obtained by taking the highest energy level and sending it to the bottom. Moreover, the triangular

partition $\mu = (N - 1, N - 2, \dots, 1, 0)$ sends all the Landau levels to the lowest one. More in general, any triangular partition with rows with decreasing number of boxes from $N - n_0$ to 1, $1 \leq n_0 \leq N$, and the remaining rows void, introduces a cutoff at the n_0 -th Landau level and projects all the higher energy states onto it. We underline that such cutoff, determined by the choice of partition μ , is intrinsically different from the naturally induced short-distance cutoff $\sqrt{\frac{2\pi\theta}{N}}$.

5. SOLUTION OF THE SUPERMATRIX MODEL

The goal of the present section is to extend the analysis of the LSZ matrix model developed in section 4 to the supermatrix model

$$\mathbf{Z}_{\text{sLSZ}}(E, \tilde{E}) = \int \mathcal{D}M \mathcal{D}M^\dagger \exp\left(- (N_1 + N_2) \text{STr} \left\{ MEM^\dagger + M^\dagger \tilde{E}M + \widehat{V}(M^\dagger M) \right\}\right)$$

introduced in Eq. (1.2). The integral is over complex supermatrices of size $(N_1 + N_2) \times (N_1 + N_2)$, and the external supermatrices E, \tilde{E} have $N_1 + N_2$ real eigenvalues each, which we write as $\frac{4\pi}{N_1 + N_2}(\eta_\ell, \xi_r)$ and $\frac{4\pi}{N_1 + N_2}(\tilde{\eta}_\ell, \tilde{\xi}_r)$ for E and \tilde{E} respectively. We use indices $\ell = 1, \dots, N_1$ and $r = 1, \dots, N_2$. The potential \widehat{V} has a quadratic and a quartic interaction, as in (4.1). We write the partition functions of ordinary matrix models with curly \mathcal{Z} and supermatrix models with bold \mathbf{Z} .

In the next subsection we present a few generalities about supermatrices, then we will extend the derivation of section 4 to the sLSZ supermatrix model, this time establishing a connection with ABJ(M) theory.

5.1. Supermatrix models. We now introduce supermatrices and supermatrix models [76, 77]. We consider the general case of $(N_1 + N_2) \times (N_1 + N_2)$ supermatrices, which can be defined in the block form

$$M = \begin{pmatrix} A & \psi \\ \chi & B \end{pmatrix},$$

where A and B are respectively $N_1 \times N_1$ and $N_2 \times N_2$ complex matrices with bosonic entries, and ψ and χ are respectively $N_2 \times N_1$ and $N_1 \times N_2$ matrices with fermionic entries. The supermatrix M is acted on by the unitary supergroup $U(N_1|N_2)_{\text{left}} \times U(N_1|N_2)_{\text{right}}$. The supertrace operation is

$$\text{STr} M = \text{Tr} A - \text{Tr} B,$$

and the integration measure $\mathcal{D}M \mathcal{D}M^\dagger$ is the product of Haar measures

$$\mathcal{D}M \mathcal{D}M^\dagger = \mathcal{D}A \mathcal{D}A^\dagger \mathcal{D}B \mathcal{D}B^\dagger \mathcal{D}\psi \mathcal{D}\chi.$$

We will need the supersymmetric version of the Harish-Chandra–Itzykson–Zuber formula (4.3), which reads [78, 79]

$$(5.1) \quad \int_{U(N_1|N_2)} [dU] \exp\left\{-(N_1 + N_2) \text{STr}(EUYU)\right\} \\ = \mathbf{C}_{N_1 N_2} \frac{\prod_{\ell=1}^{N_1} \prod_{r=1}^{N_2} (y_\ell - z_r)}{\Delta_{N_1}[y] \Delta_{N_2}[z]} \frac{\prod_{\ell=1}^{N_1} \prod_{r=1}^{N_2} (\eta_\ell - \xi_r)}{\Delta_{N_1}[\eta] \Delta_{N_2}[\xi]} \det_{1 \leq \ell, \ell' \leq N_1} \left(e^{-4\pi\eta_\ell y_{\ell'}} \right) \det_{1 \leq r, r' \leq N_2} \left(e^{4\pi\xi_r z_{r'}} \right),$$

where E and Y are supermatrices with eigenvalues $\frac{4\pi}{N}(\eta_\ell, \xi_r)$ and (y_ℓ, z_r) respectively, for $\ell = 1, \dots, N_1$ and $r = 1, \dots, N_2$, and the coefficient is

$$\mathbf{C}_{N_1 N_2} = \frac{(4\pi)^{N_1 N_2}}{(4\pi)^{\frac{N_1(N_1-1)}{2}} (-4\pi)^{\frac{N_2(N_2-1)}{2}}} G(N_1 + 1) G(N_2 + 1),$$

where $G(\cdot)$ is as usual the Barnes G -function. The factors Δ_N are Vandermonde determinants, introduced in (4.2). We note the appearance of the terms

$$\frac{\Delta_{N_1}[y]\Delta_{N_2}[z]}{\prod_{\ell=1}^{N_1}\prod_{r=1}^{N_2}(y_\ell - z_r)},$$

and likewise for (η, ξ) , which is the superdeterminant (called Berezinian) version of the Vandermonde. When $N_2 = 0$, formula (5.1) reduces to the well known HCIZ formula (4.3).

5.2. General solution. We now focus on the supermatrix model \mathbf{Z}_{sLSZ} defined in (1.2), and solve it as we have done in section 4 for the LSZ model.

We first rewrite the partition function as⁴

$$\begin{aligned} \mathbf{Z}_{\text{sLSZ}}(E, \tilde{E}) &= \int_{[0, \infty)^{N_1}} \int_{[0, \infty)^{N_2}} \frac{\Delta_{N_1}[y]^2 \Delta_{N_2}[z]^2}{\prod_{\ell=1}^{N_1} \prod_{r=1}^{N_2} (y_\ell - z_r)^2} \prod_{\ell=1}^{N_1} e^{-\hat{m}^2 y_\ell - \frac{g^2}{2} y_\ell^2} dy_\ell \prod_{r=1}^{N_2} e^{\hat{m}^2 z_r + \frac{g^2}{2} z_r^2} dz_r \\ &\quad \times \int_{U(N_1|N_2)} [dU_2] \exp(-NEU_2^\dagger Y U_2) \int_{U(N_1|N_2)} [dU_1] \exp(-N\tilde{E}U_1^\dagger Y U_1), \end{aligned}$$

and then apply the supersymmetric HCIZ formula (5.1) twice. As in section 4, we simplify the Jacobian with the denominator coming from (5.1), and obtain

$$\begin{aligned} \mathbf{Z}_{\text{sLSZ}}(E, \tilde{E}) &= \mathbf{C}_{N_1 N_2}^2 \frac{\prod_{\ell=1}^{N_1} \prod_{r=1}^{N_2} (\eta_\ell - \xi_r)(\tilde{\eta}_\ell - \tilde{\xi}_r)}{\Delta_{N_1}[\eta] \Delta_{N_1}[\xi] \Delta_{N_2}[\tilde{\eta}] \Delta_{N_2}[\tilde{\xi}]} \\ &\quad \times \int_{[0, \infty)^{N_1}} \prod_{\ell=1}^{N_1} e^{-\hat{m}^2 y_\ell - \frac{g^2}{2} y_\ell^2} dy_\ell \det_{1 \leq \ell, \ell' \leq N_1} (e^{-4\pi \eta_\ell y_{\ell'}}) \det_{1 \leq \ell, \ell' \leq N_1} (e^{-4\pi \tilde{\eta}_\ell y_{\ell'}}) \\ &\quad \times \int_{[0, \infty)^{N_2}} \prod_{r=1}^{N_2} e^{\hat{m}^2 z_r + \frac{g^2}{2} z_r^2} dz_r \det_{1 \leq r, r' \leq N_2} (e^{4\pi \xi_r z_{r'}}) \det_{1 \leq r, r' \leq N_2} (e^{4\pi \tilde{\xi}_r z_{r'}}). \end{aligned}$$

We now assume the eigenvalues of the external supermatrices E, \tilde{E} are $\frac{4\pi}{N_1+N_2}$ times integers, and we rewrite them in the form

$$(5.2) \quad \begin{aligned} \eta_\ell &= \mu_{1;\ell} + N_1 - \ell, & \xi_r &= \mu_{2;r} + N_2 - r \\ \tilde{\eta}_\ell &= \nu_{1;\ell} + N_1 - \ell, & \tilde{\xi}_r &= \nu_{2;r} + N_2 - r, \end{aligned}$$

which is the obvious extension of (4.7). In our conventions, the partitions with subindex 1 have rows labelled by $\ell = 1, \dots, N_1$ and partitions with subindex 2 have rows labelled by $r = 1, \dots, N_2$. We again recognize the Schur polynomials,

$$\begin{aligned} \mathbf{Z}_{\text{sLSZ}}(E, \tilde{E}) &= \mathbf{C}'_{N_1 N_2} \frac{\prod_{\ell=1}^{N_1} \prod_{r=1}^{N_2} (\eta_\ell - \xi_r)(\tilde{\eta}_\ell - \tilde{\xi}_r)}{\Delta_{N_1}[\eta] \Delta_{N_1}[\xi] \Delta_{N_2}[\tilde{\eta}] \Delta_{N_2}[\tilde{\xi}]} \\ &\quad \times \int_{[0, \infty)^{N_1}} \Delta_{N_1}[e^y]^2 s_{\mu_1}(e^{-y_1}, \dots, e^{-y_{N_1}}) s_{\nu_1}(e^{-y_1}, \dots, e^{-y_{N_1}}) \prod_{\ell=1}^{N_1} e^{-m^2 y_\ell - \frac{g^2}{2} y_\ell^2} dy_\ell \\ &\quad \times \int_{[0, \infty)^{N_2}} \Delta_{N_2}[e^z]^2 s_{\mu_2}(e^{z_1}, \dots, e^{z_{N_2}}) s_{\nu_2}(e^{z_1}, \dots, e^{z_{N_2}}) \prod_{r=1}^{N_2} e^{m^2 z_r + \frac{g^2}{2} z_r^2} dz_r \end{aligned}$$

where we defined the parameters m and g as in (4.6) to reabsorb the factor 4π , and

$$\mathbf{C}'_{N_1 N_2} = (4\pi)^{-(N_1+N_2)} \mathbf{C}_{N_1 N_2}^2 = \frac{G(N_1+1)^2 G(N_2+1)^2}{(4\pi)^{(N_1-N_2)^2}}.$$

⁴The Jacobian is the squared Vandermonde Berezinian, see [76, Sec. 4] and references therein.

We see that the present setting closely resembles what we obtained in section 4, now with two sets of integration variables and two pairs of Schur polynomials. In fact, the integrals over the two sets of variables are factorized, and we may give the result as a product of two copies of the result in section 4.

However, we follow a different path and assemble the two pairs of Schur polynomials into two supersymmetric Schur polynomials labelled by representations of the supergroup $U(N_1|N_2)$, see [80, 81] for definitions and properties. Irreducible $U(N_1|N_2)$ representations are classified in typical and atypical, and here we need the typical ones. The two typical representations that appear in our computations correspond to the Young diagrams

$$\begin{aligned}\mu &= (\kappa + \mu_1) \sqcup \lambda, \\ \nu &= (\tilde{\kappa} + \nu_1) \sqcup \tilde{\lambda},\end{aligned}$$

where κ and $\tilde{\kappa}$ are rectangular $N_1 \times N_2$ diagrams. This means that the Young diagram μ is composed by a rectangle κ , a Young diagram μ_1 on the right of it and another diagram λ below it, and analogously for the Young diagram ν . For these typical representations, the supersymmetric Schur polynomials decompose as [80, 81]

$$\begin{aligned}S_\mu(e^{y_1}, \dots, e^{y_{N_1}} | e^{z_1}, \dots, e^{z_{N_2}}) &= s_{\mu_1}(e^{y_1}, \dots, e^{y_{N_1}}) s_{\lambda'}(e^{z_1}, \dots, e^{z_{N_2}}) \prod_{\ell=1}^{N_1} \prod_{r=1}^{N_2} (e^{y_\ell} + e^{z_r}), \\ S_\nu(e^{y_1}, \dots, e^{y_{N_1}} | e^{z_1}, \dots, e^{z_{N_2}}) &= s_{\nu_1}(e^{y_1}, \dots, e^{y_{N_1}}) s_{\tilde{\lambda}'}(e^{z_1}, \dots, e^{z_{N_2}}) \prod_{\ell=1}^{N_1} \prod_{r=1}^{N_2} (e^{y_\ell} + e^{z_r}),\end{aligned}$$

where S_μ, S_ν are the supersymmetric Schur polynomials and $\lambda', \tilde{\lambda}'$ are the conjugate partitions to $\lambda, \tilde{\lambda}$. Therefore, identifying the generic $\lambda, \tilde{\lambda}$ to be μ'_2 and ν'_2 in our case, and following the same manipulations as in section 4, we rewrite \mathbf{Z}_{sLSZ} as

$$\begin{aligned}(5.3) \quad \mathbf{Z}_{\text{sLSZ}}(\mu, \nu) &= \mathbf{A}_{N_1, N_2}(\mu, \nu) \times \int_{(-\infty, -\gamma_1]^{N_1}} \int_{(\gamma_1, \infty]^{N_2}} \prod_{j=1}^{N_1} e^{-\frac{\gamma_1^2}{2} x_j^2} dx_j \prod_{r=1}^{N_2} e^{\frac{\gamma_2^2}{2} w_r^2} dw_r \\ &\times \frac{\prod_{1 \leq j < k \leq N_1} \left(2 \sinh\left(\frac{x_j - x_k}{2}\right)\right)^2 \prod_{1 \leq r < s \leq N_2} \left(2 \sinh\left(\frac{w_r - w_s}{2}\right)\right)^2}{\prod_{j=1}^{N_1} \prod_{r=1}^{N_2} \left(2 \cosh\left(\frac{x_j - w_r}{2}\right)\right)^2} \\ &\times S_\mu(e^{x_1}, \dots, e^{x_{N_1}} | e^{w_1}, \dots, e^{w_{N_2}}) S_\nu(e^{x_1}, \dots, e^{x_{N_1}} | e^{w_1}, \dots, e^{w_{N_2}}),\end{aligned}$$

with $\gamma_\alpha = m^2 - N_\alpha + 1$, $\alpha = 1, 2$, and the overall coefficient being

$$\begin{aligned}\mathbf{A}_{N_1, N_2}(\mu, \nu) &= (-1)^{N_1} \mathbf{C}'_{N_1 N_2} \frac{\prod_{\ell=1}^{N_1} \prod_{r=1}^{N_2} (\eta_\ell - \xi_r) (\tilde{\eta}_\ell - \tilde{\xi}_r)}{\Delta_{N_1}[\eta] \Delta_{N_1}[\xi] \Delta_{N_2}[\tilde{\eta}] \Delta_{N_2}[\tilde{\xi}]} \\ &\times \exp\left(\frac{\gamma_1^2 N_1 - \gamma_2^2 N_2}{2} + \gamma_1 (|\mu_1| + |\nu_1|) + \gamma_2 (|\mu_2| + |\nu_2|)\right) \\ &= (-1)^{N_1} \frac{\prod_{\ell=1}^{N_1} \prod_{r=1}^{N_2} (\eta_\ell - \xi_r)^2 (\tilde{\eta}_\ell - \tilde{\xi}_r)^2}{(4\pi)^{(N_1 - N_2)^2} \dim \mu \dim \nu} \\ &\times \exp\left(\frac{\gamma_1^2 N_1 - \gamma_2^2 N_2}{2} + \gamma_1 (|\mu_1| + |\nu_1|) + \gamma_2 (|\mu_2| + |\nu_2|)\right).\end{aligned}$$

For the second equality, we have written the contribution from the eigenvalues of the external supermatrices E, \tilde{E} in terms of the partitions μ, ν , and noted again that the Barnes G -functions coming from the dimensions of the representations cancel with those arisen from the supersymmetric HCIZ formula, precisely as in the ordinary matrix model of section 4.

To arrive at (5.3) we have first shifted both sets of variables and then reflected the first set. If we go back to equation (5.2) and define $\mu_{2;r}, \nu_{2;r}$ from $\xi_r, \tilde{\xi}_r$ with opposite sign, we get to an expression analogous to (5.3) but with both (x_1, \dots, x_{N_1}) and (w_1, \dots, w_{N_2}) integrated over the same domain. In each case, we have to first change variables and then insert the supersymmetric Schur factorization identity.

Comparing with [62], we rewrite the expression (5.3) in the form

$$(5.4) \quad \mathbf{Z}_{\text{sLSZ}}(\mu, \nu) = \mathbf{A}_{N_1, N_2}(\mu, \nu) \mathbf{Z}_{\text{ABJ}} \left\langle W_\mu W_\nu \mathbb{1}_{(-\infty, -\gamma_1]^{N_1}}(x) \mathbb{1}_{[\gamma_2, \infty)^{N_2}}(w) \right\rangle_{\text{ABJ}},$$

where $\langle \dots \rangle_{\text{ABJ}}$ is the average in the ABJ matrix model (1.4), and $\mathbb{1}_{(-\infty, -\gamma_1]^{N_1}}(x)$ (respectively $\mathbb{1}_{[\gamma_2, \infty)^{N_2}}(w)$) is the N_1 -dimensional (N_2 -dimensional) indicator function. Besides, W_μ is the trace of the holonomy of a superconnection along an equatorial circle inside \mathbb{S}^3 , in the representation μ of the supergroup $U(N_1|N_2)$, which describes a Wilson loop in ABJ(M) theory [82, 83]. Previous works on averages of supersymmetric Schur polynomials over the ABJ(M) ensemble include [84–86].

The probabilistic interpretation of the result (5.4) in terms of large deviations away from the equilibrium in the ABJ(M) matrix model with supersymmetric Schur insertions, follows directly from the discussion in subsection 4.4. In particular

$$\frac{\mathbf{Z}_{\text{sLSZ}}(\mu, \nu)}{\mathbf{Z}_{\text{ABJ}}} = \mathbf{A}_{N_1, N_2}(\mu, \nu) \langle W_\mu W_\nu \rangle_{\text{ABJ}} \text{Prob}_{\text{ABJ}; W_\mu W_\nu}(x_{\max} \leq -\gamma_1, w_{\min} \geq \gamma_2),$$

hence the supermatrix model (1.2) measures the probability of atypically large deviations from the equilibrium in the random matrix description of two supersymmetric Wilson loops carrying two $U(N_1|N_2)$ typical representations. The specialization to two void partitions implies

$$\frac{\mathbf{Z}_{\text{sLSZ}}(\emptyset, \emptyset)}{\mathbf{Z}_{\text{ABJ}}} = \mathbf{A}_{N_1, N_2}(\emptyset, \emptyset) \text{Prob}_{\text{ABJ}}(x_{\max} \leq -\gamma_1, w_{\min} \geq \gamma_2).$$

Remark. Throughout this section we defined and analyzed a supermatrix version of the LSZ model. It would be interesting to derive the supermatrix model (1.2) from an extension of the LSZ scalar theory to a noncommutative superspace.

Acknowledgements. The work of MT was partially supported by the Fundação para a Ciência e a Tecnologia (FCT) through its program Investigador FCT IF2014, under contract IF/01767/2014, during the first stages of this work. The work of LS was supported by the FCT through the doctoral scholarship SFRH/BD/129405/2017. The work is also supported by FCT Project PTDC/MAT-PUR/30234/2017. MT thanks Harold Steinacker and the Mathematical Physics group at University of Vienna, for discussions and warm hospitality during a short research visit at the end of 2016.

APPENDIX A. SCHUR POLYNOMIALS

Here we include explicit formulas involving Schur polynomials [61].

A.1. Spectral shift and rectangular Schur. A simple identity of Schur polynomials quickly shows what occurs if, in the case of a equispaced, harmonic oscillator spectra, we have a global overall shift in the spectrum (that is, a different zero point energy). If we have

a rectangular partition of length N , (l, l, \dots, l) which we denote by l^N then, assuming that λ is a partition of length equal or lower than N , it holds

$$s_{\mu+l^N}(e^{x_1}, \dots, e^{x_N}) = \prod_{i=1}^N e^{lx_i} s_{\mu}(e^{x_1}, \dots, e^{x_N}),$$

Therefore, an overall spectral shift by an integer l in one external matrix, corresponds to a renormalization of the mass parameter $\widehat{m}^2 \mapsto \widehat{m}^2 - l$.

Another useful identity is [61]

$$(A.1) \quad s_{\mu}(ce^{x_1}, \dots, ce^{x_N}) = c^{|\mu|} s_{\mu}(e^{x_1}, \dots, e^{x_N}).$$

A.2. Dimensions. The value of $s_{\lambda}(1, \dots, 1)$ gives the dimension of the irreducible representation of $U(N)$ with highest weight λ . Using Weyl's denominator formula

$$s_{\lambda}(1, \dots, 1) = \frac{\prod_{i < j} (\mu_i - \mu_j)}{\prod_{i < j} (i - j)},$$

where $\mu_i = \lambda_i + N - i$. Thus, it can also be written as

$$s_{\lambda}(1, \dots, 1) = \dim \lambda = \frac{1}{G(N+1)} \prod_{i < j} (\lambda_i - \lambda_j - i + j),$$

where $G(\cdot)$ is the Barnes G -function.

A.3. The Hopf link invariant. We consider the Chern–Simons matrix model with two Schur insertions, and show that it computes the Hopf link invariant [62]. First, to deal with the inversion of variables appearing in a Schur polynomial, we can use the identity [61]

$$s_{\nu}(x_1^{-1}, \dots, x_N^{-1}) = \prod_{j=1}^N x_j^{-\nu_j} s_{\nu^*}(x_1, \dots, x_N),$$

where the starred partition is defined as

$$\nu^* := (\nu_1 - \nu_N, \nu_1 - \nu_{N-1}, \dots, 0).$$

Then the following holds:

$$\begin{aligned} & \int_{[0, \infty)^N} \prod_{1 \leq j < k \leq N} \left(2 \sinh \left(\frac{z_j - z_k}{2} \right) \right)^2 s_{\mu}(e^{z_1}, \dots, e^{z_N}) s_{\nu}(e^{z_1}, \dots, e^{z_N}) \prod_{j=1}^N e^{-\frac{1}{2g_s} z_j^2} dz_j \\ &= \int_{[0, \infty)^N} \prod_{1 \leq j < k \leq N} \left(2 \sinh \left(\frac{z_j - z_k}{2} \right) \right)^2 s_{\mu}(e^{z_1}, \dots, e^{z_N}) s_{\nu^*}(e^{-z_1}, \dots, e^{-z_N}) \prod_{j=1}^N e^{-\frac{1}{2g_s} z_j^2 + \nu_1 z_j} dz_j \\ &= e^{\frac{N\nu_1^2 g_s}{2}} \int_{[-\nu_1 g_s, \infty)^N} \prod_{1 \leq j < k \leq N} \left(2 \sinh \left(\frac{w_j - w_k}{2} \right) \right)^2 \prod_{j=1}^N e^{-\frac{1}{2g_s} w_j^2} dw_j \\ & \quad \times s_{\mu}(e^{w_1 + \nu_1 g_s}, \dots, e^{w_N + \nu_1 g_s}) s_{\nu^*}(e^{-w_1 - \nu_1 g_s}, \dots, e^{-w_N - \nu_1 g_s}) \\ &= C_N''(|\mu|, |\nu^*|) \int_{[-\nu_1 g_s, \infty)^N} \prod_{1 \leq j < k \leq N} \left(2 \sinh \left(\frac{w_j - w_k}{2} \right) \right)^2 \prod_{j=1}^N e^{-\frac{1}{2g_s} w_j^2} dw_j \\ & \quad \times s_{\mu}(e^{w_1}, \dots, e^{w_N}) s_{\nu^*}(e^{-w_1}, \dots, e^{-w_N}), \end{aligned}$$

where $C_N''(|\mu|, |\nu^*|) := e^{\frac{N\nu_1^2 g_s}{2} + \nu_1 g_s (|\mu| - |\nu^*|)}$. Plugging this calculation in equation (4.8) provides an equivalent interpretation of $\mathcal{Z}_{\text{LSZ}}(\mu, \nu)$ in terms of the Hopf link invariant carrying representations μ and ν^* . We also note that, in this interpretation, setting $\nu = \mu$, which

corresponds to two external matrices with equal eigenvalues, would not lead to the diagonal part of the Hopf link invariant, due to the appearance of the starred partition.

For the sLSZ model instead there is no equivalent interpretation in terms of the Hopf link with one starred partition. This is because, if we invert one set of variables in (5.3), trading the partitions ν_1, ν_2 for ν_1^*, ν_2^* , the factorization of the supersymmetric Schur polynomial would bring one product with inverted variables, $\prod_{\ell, r} (e^{-x_\ell} + e^{-w_r})$, and we would not arrive at the ABJ matrix model.

Recall that Chern–Simons observables depend on framing [63]: the relation between such dependence and the modular matrices is given in [87]. Observables in the matrix model description are not in the canonical framing, and the (usual) Hopf link average is:

$$\langle W_{\mu\nu^*} \rangle_{\text{CS}} = (TST)_{\mu\nu^*}$$

in terms of the modular S, T matrices of the Wess–Zumino–Witten (WZW) model [88]. In the general case, the framed Hopf link comes as $T^{n_1}ST^{n_2}$; if $n_1 = n_2 = 0$, that is, the canonical framing in S^3 , it is exactly the modular S matrix, while if $n_1 + n_2 = 2$, the framing is the $U(1)$ -invariant Seifert framing [89]. This is the case of the matrix model description.

REFERENCES

- [1] H. S. Snyder, *Quantized space-time*, *Phys. Rev.* **71** (1947) 38.
- [2] N. Seiberg and E. Witten, *String theory and noncommutative geometry*, *JHEP* **09** (1999) 032 [[hep-th/9908142](#)].
- [3] F. Lizzi and R. J. Szabo, *Noncommutative geometry and string duality*, *PoS Corfu98* **073** (1999) [[hep-th/9904064](#)].
- [4] S. Minwalla, M. Van Raamsdonk and N. Seiberg, *Noncommutative perturbative dynamics*, *JHEP* **02** (2000) 020 [[hep-th/9912072](#)].
- [5] M. R. Douglas and N. A. Nekrasov, *Noncommutative field theory*, *Rev. Mod. Phys.* **73** (2001) 977 [[hep-th/0106048](#)].
- [6] R. J. Szabo, *Quantum field theory on noncommutative spaces*, *Phys. Rept.* **378** (2003) 207 [[hep-th/0109162](#)].
- [7] H. Steinacker, *Non-commutative geometry and matrix models*, *PoS QGQGS2011* (2011) 004 [[1109.5521](#)].
- [8] E. Langmann, R. J. Szabo and K. Zarembo, *Exact solution of noncommutative field theory in background magnetic fields*, *Phys. Lett.* **B569** (2003) 95 [[hep-th/0303082](#)].
- [9] E. Langmann, R. J. Szabo and K. Zarembo, *Exact solution of quantum field theory on noncommutative phase spaces*, *JHEP* **01** (2004) 017 [[hep-th/0308043](#)].
- [10] J. de Jong and R. Wulkenhaar, *Nonperturbative evaluation of the partition function for the real scalar quartic QFT on the Moyal plane at weak coupling*, *J. Math. Phys.* **60** (2019) 083504 [[1809.09453](#)].
- [11] H. Grosse, A. Hock and R. Wulkenhaar, *Solution of all quartic matrix models*, [[1906.04600](#)].
- [12] H. Grosse, A. Hock and R. Wulkenhaar, *Solution of the self-dual Φ^4 QFT-model on four-dimensional Moyal space*, *JHEP* **01** (2020) 081 [[1908.04543](#)].
- [13] M. Mariño, *Chern-Simons theory, matrix integrals, and perturbative three manifold invariants*, *Commun. Math. Phys.* **253** (2004) 25 [[hep-th/0207096](#)].
- [14] M. Mariño, *Chern-Simons theory, matrix models, and topological strings*, *Int. Ser. Monogr. Phys.* **131** (2005) 1.
- [15] C. Beasley and E. Witten, *Non-Abelian localization for Chern-Simons theory*, *J. Diff. Geom.* **70** (2005) 183 [[hep-th/0503126](#)].
- [16] M. Blau and G. Thompson, *Chern-Simons theory on S^1 -bundles: Abelianisation and q -deformed Yang-Mills theory*, *JHEP* **05** (2006) 003 [[hep-th/0601068](#)].
- [17] J. Källén, *Cohomological localization of Chern-Simons theory*, *JHEP* **08** (2011) 008 [[1104.5353](#)].
- [18] M. Tierz, *Soft matrix models and Chern-Simons partition functions*, *Mod. Phys. Lett.* **A19** (2004) 1365 [[hep-th/0212128](#)].
- [19] Y. Dolivet and M. Tierz, *Chern-Simons matrix models and Stieltjes-Wigert polynomials*, *J. Math. Phys.* **48** (2007) 023507 [[hep-th/0609167](#)].
- [20] M. Romo and M. Tierz, *Unitary Chern-Simons matrix model and the Villain lattice action*, *Phys. Rev.* **D86** (2012) 045027 [[1103.2421](#)].
- [21] M. L. Mehta, *Random Matrices*, Pure and Applied Mathematics. Elsevier Science, 2004.

- [22] K. A. Muttalib, Y. Chen, M. E. H. Ismail and V. N. Nicopoulos, *New family of unitary random matrices*, *Phys. Rev. Lett.* **71** (1993) 471.
- [23] S. M. Nishigaki, *Level spacings at the metal-insulator transition in the Anderson Hamiltonians and multifractal random matrix ensembles*, *Phys. Rev.* **E59** (1999) 2853 [[cond-mat/9809147](#)].
- [24] D. S. Dean and S. N. Majumdar, *Extreme value statistics of eigenvalues of Gaussian random matrices*, *Phys. Rev. E* **77** (2008) 041108 [[0801.1730](#)].
- [25] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, *$N=6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, *JHEP* **10** (2008) 091 [[0806.1218](#)].
- [26] O. Aharony, O. Bergman and D. L. Jafferis, *Fractional M2-branes*, *JHEP* **11** (2008) 043 [[0807.4924](#)].
- [27] A. Kapustin, B. Willett and I. Yaakov, *Exact Results for Wilson Loops in Superconformal Chern-Simons Theories with Matter*, *JHEP* **03** (2010) 089 [[0909.4559](#)].
- [28] M. Mariño and P. Putrov, *Exact Results in ABJM Theory from Topological Strings*, *JHEP* **06** (2010) 011 [[0912.3074](#)].
- [29] M. Aganagic, A. Klemm, M. Mariño and C. Vafa, *Matrix model as a mirror of Chern-Simons theory*, *JHEP* **02** (2004) 010 [[hep-th/0211098](#)].
- [30] N. Halmagyi and V. Yasnov, *The Spectral curve of the lens space matrix model*, *JHEP* **11** (2009) 104 [[hep-th/0311117](#)].
- [31] H. Awata, S. Hirano and M. Shigemori, *The Partition Function of ABJ Theory*, *PTEP* **2013** (2013) 053B04 [[1212.2966](#)].
- [32] A. Géré, T. Jurić and J.-C. Wallet, *Noncommutative gauge theories on \mathbb{R}_λ^3 : perturbatively finite models*, *JHEP* **12** (2015) 045 [[1507.08086](#)].
- [33] J.-C. Wallet, *Exact partition functions for gauge theories on \mathbb{R}_λ^3* , *Nucl. Phys.* **B912** (2016) 354 [[1603.05045](#)].
- [34] P. Vitale and J.-C. Wallet, *Noncommutative field theories on \mathbb{R}_λ^3 : Toward UV/IR mixing freedom*, *JHEP* **04** (2013) 115 [[1212.5131](#)].
- [35] P. Martinetti, P. Vitale and J.-C. Wallet, *Noncommutative gauge theories on \mathbb{R}_θ^2 as matrix models*, *JHEP* **09** (2013) 051 [[1303.7185](#)].
- [36] A. Géré, P. Vitale and J.-C. Wallet, *Quantum gauge theories on noncommutative three-dimensional space*, *Phys. Rev.* **D90** (2014) 045019 [[1312.6145](#)].
- [37] P. Vitale, *Noncommutative field theory on \mathbb{R}_λ^3* , *Fortsch. Phys.* **62** (2014) 825 [[1406.1372](#)].
- [38] A. B. Hammou, M. Lagraa and M. M. Sheikh-Jabbari, *Coherent state induced star product on $R^{**3}(\lambda)$ and the fuzzy sphere*, *Phys. Rev.* **D66** (2002) 025025 [[hep-th/0110291](#)].
- [39] A. Yu. Alekseev, A. Recknagel and V. Schomerus, *Brane dynamics in background fluxes and noncommutative geometry*, *JHEP* **05** (2000) 010 [[hep-th/0003187](#)].
- [40] D. J. Grabiner, *Brownian motion in a Weyl chamber, non-colliding particles, and random matrices*, *Ann. Inst. H. Poincaré Probab. Statist.* **35** (1999) 177 [[math/9708207](#)].
- [41] J. Baik and T. M. Suidan, *Random matrix central limit theorems for nonintersecting random walks*, *Ann. Probab.* **35** (2007) 1807 [[math/0605212](#)].
- [42] S. de Haro and M. Tierz, *Brownian motion, Chern-Simons theory, and 2-D Yang-Mills*, *Phys. Lett.* **B601** (2004) 201 [[hep-th/0406093](#)].
- [43] Y. Takahashi and M. Katori, *Noncolliding Brownian Motion with Drift and Time-Dependent Stieltjes-Wigert Determinantal Point Process*, *J. Math. Phys.* **53** (2012) 103305 [[1207.4351](#)].
- [44] E. Langmann and R. J. Szabo, *Duality in scalar field theory on noncommutative phase spaces*, *Phys. Lett.* **B533** (2002) 168 [[hep-th/0202039](#)].
- [45] J. Madore, *The Fuzzy sphere*, *Class. Quant. Grav.* **9** (1992) 69.
- [46] B. P. Dolan, D. O'Connor and P. Presnajder, *Matrix ϕ^{**4} models on the fuzzy sphere and their continuum limits*, *JHEP* **03** (2002) 013 [[hep-th/0109084](#)].
- [47] B. P. Dolan, D. O'Connor and P. Presnajder, *Matrix models on the fuzzy sphere*, in *NATO Conf. Proc. Stara Lesna 2002*, 2002, [[hep-th/0204219](#)].
- [48] X. Martin, *A Matrix phase for the ϕ^{**4} scalar field on the fuzzy sphere*, *JHEP* **04** (2004) 077 [[hep-th/0402230](#)].
- [49] D. O'Connor and C. Saemann, *Fuzzy Scalar Field Theory as a Multitrace Matrix Model*, *JHEP* **08** (2007) 066 [[0706.2493](#)].
- [50] D. O'Connor and C. Saemann, *A Multitrace matrix model from fuzzy scalar field theory*, in *Proceedings, 7th International Workshop on Supersymmetries and Quantum Symmetries (SQS'07): Dubna, Russia, July 30 - August 04, 2007*, pp. 148–155, 2008, [[0709.0387](#)].
- [51] C. Saemann, *The Multitrace Matrix Model of Scalar Field Theory on Fuzzy CP^n* , *SIGMA* **6** (2010) 050 [[1003.4683](#)].

- [52] M. Panero, *Quantum field theory in a non-commutative space: Theoretical predictions and numerical results on the fuzzy sphere*, *SIGMA* **2** (2006) 081 [[hep-th/0609205](#)].
- [53] A. P. Polychronakos, *Effective action and phase transitions of scalar field on the fuzzy sphere*, *Phys. Rev. D* **88** (2013) 065010 [[1306.6645](#)].
- [54] K. Hatakeyama and A. Tsuchiya, *Correlation functions and renormalization in a scalar field theory on the fuzzy sphere*, *PTEP* **2017** (2017) 063B01 [[1704.01698](#)].
- [55] K. Hatakeyama, A. Tsuchiya and K. Yamashiro, *Renormalization on the fuzzy sphere*, *PTEP* **2018** (2018) 063B05 [[1805.03975](#)].
- [56] H. Steinacker, *Quantized gauge theory on the fuzzy sphere as random matrix model*, *Nucl. Phys.* **B679** (2004) 66 [[hep-th/0307075](#)].
- [57] K. Bering and H. Grosse, *On Batalin-Vilkovisky Formalism of Non-Commutative Field Theories*, *Eur. Phys. J.* **C68** (2010) 313 [[0908.1229](#)].
- [58] T. R. Morris, *Checked surfaces and complex matrices*, *Nucl. Phys.* **B356** (1991) 703.
- [59] Harish-Chandra, *Differential operators on a semisimple Lie algebra*, *Amer. J. Math.* **79** (1957) 87.
- [60] C. Itzykson and J. B. Zuber, *The Planar Approximation. 2.*, *J. Math. Phys.* **21** (1980) 411.
- [61] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Oxford classic texts in the physical sciences. Clarendon Press, 1998.
- [62] T. Kimura, *Linking loops in ABJM and refined theory*, *JHEP* **07** (2015) 030 [[1503.01462](#)].
- [63] E. Witten, *Quantum Field Theory and the Jones Polynomial*, *Commun. Math. Phys.* **121** (1989) 351.
- [64] N. Yu. Reshetikhin and V. G. Turaev, *Ribbon graphs and their invariants derived from quantum groups*, *Commun. Math. Phys.* **127** (1990) 1.
- [65] N. Y. Reshetikhin and V. G. Turaev, *Invariants of three manifolds via link polynomials and quantum groups*, *Invent. Math.* **103** (1991) 547.
- [66] J. Pawelczyk and H. Steinacker, *A Quantum algebraic description of D branes on group manifolds*, *Nucl. Phys.* **B638** (2002) 433 [[hep-th/0203110](#)].
- [67] L. Chekhov, *Matrix models with hard walls: Geometry and solutions*, *J. Phys.* **A39** (2006) 8857 [[hep-th/0602013](#)].
- [68] D. S. Dean and S. N. Majumdar, *Large deviations of extreme eigenvalues of random matrices*, *Phys. Rev. Lett.* **97** (2006) 160201 [[cond-mat/0609651](#)].
- [69] S. N. Majumdar and G. Schehr, *Top eigenvalue of a random matrix: large deviations and third order phase transition*, *J. Stat. Mech.* **2014** (2014) P01012 [[1311.0580](#)].
- [70] A. Dhar, A. Kundu, S. N. Majumdar, S. Sabhapandit and G. Schehr, *Exact Extremal Statistics in the Classical 1D Coulomb Gas*, *Phys. Rev. Lett.* **119** (2017) 060601 [[1704.08973](#)].
- [71] P. Vivo, *Large deviations of the maximum of independent and identically distributed random variables*, *Eur. J. Phys.* **36** (2015) 055037 [[1507.05442](#)].
- [72] Y. Makeenko, *Loop equations in matrix models and in 2-D quantum gravity*, *Mod. Phys. Lett. A* **6** (1991) 1901.
- [73] F. D. Cunden, P. Facchi, M. Ligabò and P. Vivo, *Universality of the third-order phase transition in the constrained Coulomb gas*, *JSTAT* **5** (2017) 053303 [[1702.05071](#)].
- [74] F. D. Cunden, P. Facchi, M. Ligabò and P. Vivo, *Universality of the weak pushed-to-pulled transition in systems with repulsive interactions*, *J. Phys.* **A51** (2018) [[1711.09141](#)].
- [75] R. J. Baxter, *Statistical mechanics of a one-dimensional Coulomb system with a uniform charge background*, *Math. Proc. Cambridge Phil. Soc.* **59** (1963) 779–787.
- [76] S. A. Yost, *Supermatrix models*, *Int. J. Mod. Phys.* **A7** (1992) 6105 [[hep-th/9111033](#)].
- [77] L. Alvarez-Gaume and J. L. Mañes, *Supermatrix models*, *Mod. Phys. Lett.* **A6** (1991) 2039.
- [78] J. Alfaro, R. Medina and L. F. Urrutia, *The Itzykson-Zuber integral for $U(m/n)$* , *J. Math. Phys.* **36** (1995) 3085 [[hep-th/9412012](#)].
- [79] T. Guhr, *Gelfand-Tsetlin coordinates for the unitary supergroup*, *Commun. Math. Phys.* **176** (1996) 555.
- [80] A. Berele and A. Regev, *Hook Young-diagrams with applications to combinatorics and to representations of Lie-superalgebras*, *Adv. Math.* **64** (1987) 118.
- [81] E. M. Moens and J. van der Jeugt, *A determinantal formula for supersymmetric Schur polynomials*, *J. Algebraic Combin.* **17** (2003) 283.
- [82] N. Drukker and D. Trancanelli, *A Supermatrix model for $N=6$ super Chern-Simons-matter theory*, *JHEP* **02** (2010) 058 [[0912.3006](#)].
- [83] K.-M. Lee and S. Lee, *1/2-BPS Wilson Loops and Vortices in ABJM Model*, *JHEP* **09** (2010) 004 [[1006.5589](#)].
- [84] S. Matsumoto and S. Moriyama, *ABJ Fractional Brane from ABJM Wilson Loop*, *JHEP* **03** (2014) 079 [[1310.8051](#)].

- [85] S. Matsuno and S. Moriyama, *Giambelli Identity in Super Chern-Simons Matrix Model*, *J. Math. Phys.* **58** (2017) 032301 [1603.04124].
- [86] T. Furukawa and S. Moriyama, *Jacobi-Trudi Identity in Super Chern-Simons Matrix Model*, *SIGMA* **14** (2018) 049 [1711.04893].
- [87] L. C. Jeffrey, *Chern-Simons-Witten invariants of lens spaces and torus bundles, and the semiclassical approximation*, *Commun. Math. Phys.* **147** (1992) 563.
- [88] P. Di Francesco, P. Mathieu and D. Senechal, *Conformal Field Theory*, Graduate Texts in Contemporary Physics. Springer-Verlag, New York, 1997, 10.1007/978-1-4612-2256-9.
- [89] M. Blau and G. Thompson, *Chern-Simons Theory on Seifert 3-Manifolds*, *JHEP* **09** (2013) 033 [1306.3381].

(LS) GRUPO DE FÍSICA MATEMÁTICA, DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS, UNIVERSIDADE DE LISBOA, CAMPO GRANDE, EDIFÍCIO C6, 1749-016 LISBOA, PORTUGAL.

E-mail address: `lsantilli@fc.ul.pt`

(MT) DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS, ISCTE - INSTITUTO UNIVERSITÁRIO DE LISBOA, AVENIDA DAS FORÇAS ARMADAS, 1649-026 LISBOA, PORTUGAL.

E-mail address: `mtpaz@iscte-iul.pt`

(MT) GRUPO DE FÍSICA MATEMÁTICA, DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS, UNIVERSIDADE DE LISBOA, CAMPO GRANDE, EDIFÍCIO C6, 1749-016 LISBOA, PORTUGAL.

E-mail address: `tierz@fc.ul.pt`