

# The relationships of lifetime physical activity and diet with salivary cell telomere length in current ultra-endurance exercisers


Karen Birkenhead<sup>a,\*</sup>, Anna Kuballa<sup>a</sup>, Geoff P. Lovell<sup>a,b</sup>, Susan I. Barr<sup>c</sup> and Colin Solomon<sup>a</sup>

<sup>a</sup>*School of Sport and Behavioural Sciences, University of the Sunshine Coast, Maroochydore, DC, QLD, Australia*

<sup>b</sup>*Department of Sport, Hartpury University, Hartpury House, Gloucester, UK*

<sup>c</sup>*Food, Nutrition and Health, Faculty of Land and Food Systems, University of British Columbia, East Mall,*

View metadata, citation and similar papers at [core.ac.uk](https://core.ac.uk)

brought to you by  CORE

provided by Hartpury University Repository

Received 6 April 2020

Accepted 9 July 2021

Pre-press 30 July 2021

## Abstract.

**BACKGROUND:** Physical activity and a healthy diet may delay the aging process and ultra-endurance exercise is an extreme form of physical activity. Telomeres are protective DNA sequences located at the ends of eukaryotic chromosomes which shorten as we age.

**OBJECTIVE:** The aim of this study was to investigate the relationships of lifetime physical activity and diet with salivary cell telomere length in current ultra-endurance exercisers ( $n = 49$ ; % female = 37, age range 26–74 years).

**METHODS:** Physical activity and dietary intake were measured using the Lifetime Physical Activity and Diet Questionnaire (LPADQ) and salivary cell telomere length was measured using quantitative polymerase chain reaction.

**RESULTS:** In this group of current ultra-endurance exercisers there was no relationship between lifetime physical activity or diet (according to food category scores) and telomere length. In contrast to the expected age-related decrease in telomere length, there was no relationship between age and telomere length (95% confidence interval [CI]:  $-38.86, 14.54, p = 0.359$ ) in this group of current ultra-endurance exercisers.

**CONCLUSIONS:** The relationships of lifetime physical activity and diet with telomere length remain uncertain. It is possible that lifetime physical activity (including ultra-endurance exercise) and lifetime diet may independently, or in combination, contribute to a decrease in the rate of age-related telomere shortening in current ultra-endurance exercisers.

Keywords: Aging, extreme exercise, dietary intake, ultra-marathon, ultra-cycling

## 1. Introduction

Ultra-endurance exercise involves prolonged periods of physical activity covering a distance of more than the standard marathon (42.195 km) or with a duration greater than 6 hours [1]. The majority of ultra-runners are masters athletes (>35 years) with reported age of participants ranging from 18–81

years [2–4]. Currently, men make up the majority of participants in ultra-endurance events, although an increasing number of women participants over the past decade has been reported [5, 6]. Ultra-endurance exercisers tend to be well-educated and show a large range of training and running experiences [2, 4]. Worldwide participation in ultra-endurance exercise is increasing, yet there is minimal research on the individuals who engage in this form of physical activity. Specifically, there is minimal information on the physical activity and dietary habits of ultra-endurance exercisers or how this form of extreme exercise influences the aging process.

\*Corresponding author: Karen Birkenhead, School of Sport and Behavioural Sciences, University of the Sunshine Coast, Locked Bag 4, Maroochydore DC, QLD 4558, Tel.: +61 2 439091766; E-mail: [klbirkenhead@gmail.com](mailto:klbirkenhead@gmail.com); ORCID: 0000-0002-8311-7794

48 Aging is an obligatory component of human life  
49 and is thought to be determined by several fac-  
50 tors, including genetics, the environment and lifestyle  
51 choices [7, 8]. Telomeres are non-coding repeating  
52 segments of DNA (human sequence – TTAGGG)  
53 located at the ends of eukaryotic chromosomes [9].  
54 In most dividing cells telomeres shorten which can  
55 lead to apoptosis or replicative senescence and this  
56 progressive loss of cells can contribute to chronic  
57 inflammation, tissue aging and age-related diseases  
58 [10]. Telomere shortening is a natural part of the  
59 aging process, but shortening may be accelerated  
60 by lifestyle factors such as obesity [11], inactivity,  
61 smoking [12, 13], and psychological stress [14, 15].  
62 Physical activity and diet have been found to be asso-  
63 ciated with components of telomere biology with  
64 many studies showing positive correlations of reg-  
65 ular exercise and healthy eating with telomere length  
66 [16–21]. While other indices of biological aging have  
67 been reported, such as the epigenetic clock [22],  
68 telomere length has been used as a marker of bio-  
69 logical aging in a multitude of research topics and  
70 may provide information on the aging process at the  
71 molecular level [23, 24].

72 Physical activity is commonly assessed using ques-  
73 tionnaires which collect information on intensity,  
74 duration and/or type of exercise [25]. Although evi-  
75 dence suggests habitual physical activity may help  
76 preserve telomere length, it is unclear if this depends  
77 on the type, volume or intensity of exercise [26]. To  
78 our knowledge, only five studies have investigated  
79 telomere length in ultra-endurance exercisers with  
80 three showing longer telomeres and two showing  
81 no difference in telomere length in ultra-endurance  
82 exercisers [20, 27–30]. However, these studies have  
83 not assessed physical activity across the lifespan and  
84 there are no data that examine the influence of diet  
85 on telomere length in ultra-endurance exercisers.

86 Most studies investigating the relationship bet-  
87 ween diet and telomere length have used a food fre-  
88 quency questionnaire (FFQ) [31–37]. This tool has  
89 been used to investigate diet in relation to telomere  
90 length based on individual nutrients, foods and/or  
91 beverages [38–42] or to examine associations with  
92 dietary patterns [17, 43, 44]. However, food fre-  
93 quency questionnaires used in telomere research only  
94 extend as far back as one year and provide dietary  
95 information specific to that time period. Although  
96 there are two longitudinal studies that have collected  
97 dietary intake at baseline and measured telomere  
98 length 10 years later [40, 43], there are no stud-  
99 ies that include information on lifetime diet from

100 childhood to older age. Information on lifetime diet  
101 is important considering the dynamics of telomere  
102 length changes and attrition begin at birth and extend  
103 across the lifespan [45]. Studies show higher intakes  
104 of vegetables and/or fruit and adherence to some  
105 dietary patterns, such as the Mediterranean diet, are  
106 positively associated with telomere length [17, 36].  
107 Therefore, assessing the lifetime intake of healthy  
108 and less healthy foods could be a useful and logi-  
109 cal approach to investigating the impact of diet on  
110 telomere length.

111 Telomere length has most frequently been assessed  
112 in blood cells, particularly peripheral blood mononu-  
113 cleared cells (PMBC's) [46–49]. However, more  
114 recently, studies have used salivary cells to assess  
115 telomere length as they are an appropriate alternative  
116 to blood cells and provide a fast, simple to obtain,  
117 and non-invasive method for sample collection [50].  
118 Salivary cell telomere length is highly correlated with  
119 telomere length from whole blood and leukocytes  
120 [51, 52] and, therefore, it has been suggested telom-  
121 ere length of salivary cells can be used as a proxy for  
122 telomere length in other tissues [53]. Furthermore,  
123 genomic DNA from salivary cells has been found  
124 to be of similar quality to that obtained from blood  
125 cells [54]. The aim of the present study was to inves-  
126 tigate the relationships of lifetime physical activity  
127 and diet with salivary cell telomere length in current  
128 ultra-endurance exercisers.

## 129 2. Materials and methods

### 130 2.1. Study design

131 This study includes a subset of participants from a  
132 previous study that reported on the lifetime physical  
133 activity [4] and diet [55] of current ultra-endurance  
134 exercisers. Ultra-endurance exercisers from the pre-  
135 vious study were invited via email to take part in  
136 the present study. Current male and female ultra-  
137 endurance exercisers were recruited for this study  
138 from April 2016 to September 2016. A control group  
139 was not included as it was not possible to obtain a  
140 comparison group for which physical activity and  
141 diet were suitably controlled across the lifespan as  
142 the individual patterns could not have been matched.  
143 This study was conducted according to the National  
144 Statement and Human Research Ethics Guidelines  
145 [56] and approved by the University of the Sun-  
146 shine Coast Human Research Ethics Committee.  
147 Lifetime physical activity and diet were investigated

retrospectively using an online survey hosted on the SurveyMonkey® platform. Participants were provided with an information sheet, and provided informed consent by completing and submitting the online questionnaires and by returning a written, signed consent form with their saliva samples.

## 2.2. Participants

The details of participant recruitment have been described elsewhere [4]. Briefly, participants were recruited internationally through Facebook, the Ultra Listserv ([www.ultra@listserve.dartmouth.edu](http://www.ultra@listserve.dartmouth.edu)) and web pages associated with ultra-endurance organizations and events worldwide, including Australia, New Zealand, the United States of America and Canada. Inclusion criteria were being healthy,  $\geq 18$  years of age, able to complete an online questionnaire in English, and being a current ultra-endurance exerciser. To qualify as a current ultra-endurance exerciser, participants had to have completed at least one ultra-endurance event within the last five years and engage in, on average, at least five hours of running or cycling per week during the past year. The criterion of completion of at least one ultra-endurance event within the last five years allowed for the inclusion of individuals who identified as a current ultra-endurance exerciser (i.e. in training only) but may not participate in events on a regular basis. For the current study, sampling was restricted to Australian and North American participants who had completed all required components of the initial study ( $n = 86$ ). Fifty-five participants agreed to take part in the study and 50 (91%) saliva samples (33 men; 17 women) were received.

## 2.3. Questionnaires

Lifetime physical activity and diet were assessed using the Lifetime Physical Activity and Diet Questionnaire (LPADQ), details of which have been described elsewhere [55]. In brief, the questionnaire is a modified and combined version of the Lifetime Physical Activity Questionnaire (LPAQ) and the Lifetime Diet Questionnaire (LDQ) [57, 58]. Medical health history was collected with a Medical History Questionnaire (MHQ).

The LPAQ assesses physical activity over several life periods ranging from childhood to older age [58]. Participants identified all activities they engaged in on more than 10 occasions over each life period. They then estimated the total number of years, months and

hours they engaged in the activity. Similarly, the LDQ assesses dietary intake over five life periods (5–18 years, 19–30 years, 31–45 years, 46–60 years and 61–75 years). Using a four-point scale (i.e. rarely, 2–3 times a month; 2–3 times per week; daily), individuals were asked to recall the general frequency at which they consumed certain foods during each time period. The life periods in the LPAQ were modified from the original version to match the diet history life periods [58] and the 5–18 year life period was divided into 5–12 and 13–18 years.

Stressful life events were measured using the List of Threatening Events (LTE) questionnaire [59]. Participants identified if they experienced any of 12 stressful events during each of six life periods (0–12 years; 13–18 years; 19–30 years, etc.). If a stressful event occurred more than once, they were instructed to select as many times as needed. Participants indicated ‘yes’, ‘no’ or ‘not applicable’ to each experience and a lifetime score was obtained ranging from 0–72 (one point for each ‘yes’ with a higher score indicating higher lifetime stress). For the current study, participants received a link to the List of Threatening Events (LTE) questionnaire by e-mail.

## 2.4. Assessment of lifetime physical activity and diet

Assessment of lifetime physical activity has been described in detail elsewhere [4]. Briefly, physical activity was calculated as the total number of hours spent in each activity per life period. This was determined by multiplying the number of years, by the number of months and the number of hours per week spent in each activity. Each activity was multiplied by the intensity according to the metabolic equivalent (MET) as determined by the Compendium of Physical Activities [60]. All activities were summed to obtain a total number of MET-hours of activity for each life period. Finally, a total lifetime MET-hours was calculated by summing the total MET-hours per life period up to the participant’s current life period.

The assessment of lifetime diet has also been described elsewhere [55]. Briefly, there are 78 food items within the LDQ and these were grouped into two categories (healthy and less healthy). Fifty foods were included in the healthy food category: 17 vegetables, 13 fruits, 4 dairy or alternatives, 5 whole grains, 9 meat or alternatives, 1 alcohol [i.e. red wine] and 1 fat [i.e. olive oil]. Eleven foods were included in the less healthy food category: 6 sweets or sugar sweetened beverages, 1 takeaway food, 1 snack food,

246 1 processed meat and 2 saturated fats. As such, foods  
 247 were grouped into categories according to healthy  
 248 eating guidelines (healthy) and those considered  
 249 other/extra/processed foods (less healthy). Seventeen  
 250 food items were not included as there was not enough  
 251 evidence to support inclusion of these items in one  
 252 category over the other. A separate sub-category for  
 253 vegetables and fruit (30 items) was created. These  
 254 were investigated separately as several studies have  
 255 reported a benefit of vegetables and/or fruit on telomere  
 256 length [34, 36, 43, 44]. Each participant received  
 257 three scores per life period. A score per life period  
 258 was calculated by summing response frequencies (i.e.  
 259 from 1 rarely/never to 4 daily) for each food within  
 260 the category. Possible ranges were 50–200 (or 49–196  
 261 for childhood and adolescence) for the healthy food  
 262 category, 11–44 for the less healthy food category  
 263 and 30–120 for vegetables/fruit food category. Child-  
 264 hood and adolescence had one less item in the healthy  
 265 food category (i.e. excludes red wine) and, therefore,  
 266 a weighted score was calculated by multiplying the  
 267 score by 200 and dividing by 196. Finally, a weighted  
 268 lifetime score was calculated as the sum of the scores  
 269 for each life period multiplied by the number of years  
 270 spent in the respective life period, and then dividing  
 271 by the total number of years spent in all life periods.

## 272 2.5. Saliva samples

273 Purpose-made collection kits were mailed or hand  
 274 delivered to participants along with instructions for  
 275 providing saliva samples. Each kit included two  
 276 30 ml collection containers with 2 ml of a stabilizing  
 277 solution (RNAlater® Thermo Fisher Scientific). Partic-  
 278 ipants were instructed to provide 5 ml of saliva per  
 279 container and, upon receipt, samples were stored at  
 280 –20°C until DNA extraction. At the time of DNA  
 281 extraction, saliva samples were aliquoted into approx-  
 282 imately 750 µl volumes and one was used for  
 283 immediate DNA extraction. The remaining aliquots  
 284 were stored at –20°C for use in subsequent DNA  
 285 extractions if the initial isolation was unsuccessful  
 286 (i.e. poor DNA quality/poor yield). All assays were  
 287 performed by the same investigator, under identical  
 288 conditions.

## 289 2.6. Measurement of telomere length

290 Saliva samples were spun for 5 minutes (× 8000 g)  
 291 to obtain a cell pellet and to remove the supernatant.  
 292 Genomic DNA was extracted from salivary sam-  
 293 ples using the ISOLATE II kit (Bioline) following

294 the genomic DNA bench top protocol (www.  
 295 bioline.com/au/). During the pre-lysis step samples  
 296 were incubated for up to 18 hours depending on the  
 297 time required for complete lysis. Following isolation  
 298 DNA samples were stored at –20°C until further anal-  
 299 ysis. DNA was assessed for quality and quantity via  
 300 visualization on a 1% agarose gel and spectropho-  
 301 tometry (NanoDrop, Thermo Scientific).

302 Relative (rTL) and absolute telomere length (aTL)  
 303 were measured using Quantitative polymerase chain  
 304 reaction (qPCR). Relative TL was measured as pre-  
 305 viously described by Cawthon [61] and aTL was  
 306 measured using a protocol adapted from O’Callaghan  
 307 and Fenech [62]. These methods calculate telomere  
 308 length based on the ratio of telomere repeat copy  
 309 number to a single copy reference gene *36B4* which  
 310 encodes the acidic ribosomal phosphoprotein PO.  
 311 The reference gene *36B4* is not expected to change  
 312 expression (i.e. copy number) as a function of any  
 313 of the independent variables being measured. There-  
 314 fore, the reference gene provides a control level of  
 315 gene expression whereby a ratio <1 (or >1) indicates  
 316 the telomere copy number was higher (or lower) than  
 317 the reference gene. Standard curves for each gene  
 318 were created using a 10-fold serial dilution of known  
 319 quantities of a synthesized 84 mer oligonucleotide  
 320 telomere (TEL) standard containing 14 TTAGGG  
 321 repeats and a synthesized 75 mer single copy refer-  
 322 ence gene (SCG) standard, *36B4*.

323 To assist in the normalisation of DNA sample con-  
 324 centration, and reduce pipetting error, each partic-  
 325 ipant sample DNA was prepared to a consistent  
 326 concentration of 5 ng/µl. The SensiFAST SYBR No-  
 327 Rox kit (Bioline) was used for qPCR according to  
 328 the manufacturer’s instructions. A master mix was  
 329 prepared with a final concentration of 1 × SensiFAST  
 330 SYBR No-ROX mix and 400 nM each of forward  
 331 and reverse primers. The primers used were those  
 332 previously described by O’Callaghan and Fenech  
 333 [62] for both the telomere sequence and the *36B4*  
 334 reference gene. The telomere primers used were:  
 335 telomere forward primer (CGGTTTGGTTTGGGTTT  
 336 GGGTTTGGGTTTGGGTTTGGGTT), and telomere  
 337 reverse primer (GGCTTGCCTTACCCTTACC  
 338 CTTACCCTTACCCTTACCCT). Primers for the single  
 339 copy reference gene amplicon (*36B4*) consisted  
 340 of: *36B4* forward primer (CAGCAAGTGGGAAG-  
 341 GTGTAATCC) and *36B4* reverse primer (CCCA  
 342 TTCTATCATCAACGGGTACAA). In each sample  
 343 both *36B4* and *TEL* genes were analysed in tripli-  
 344 cate with two no template controls (NTC) included  
 345 in each qPCR run. The Rotor-Gene Q (Qiagen), was

used for qPCR analysis. Each run was carried out in 0.1 ml strip tubes with the 72 sample Rotor-Disc, using a final volume of 10  $\mu\text{l}$  (5.8  $\mu\text{l}$  of master mix and 4.2  $\mu\text{l}$  of participant DNA) to give a final concentration of 20 ng/ $\mu\text{l}$  of sample DNA. A 3-step cycle was used with cycling conditions of: 3 min at 95°C, followed by 40 cycles of 95°C for 5 sec, 60°C for 10 sec and 72°C 15 sec (with data acquisition), followed by a dissociation (melt) curve ramping from 72°C to 95°C. The same amplification and cycling parameters were carried out for both *TEL* and *36B4* genes. Triplicate samples all displayed a standard deviation of the Ct (cycle threshold) values of <1 Ct. Samples were inspected to ensure all participants' values were within the linear range, resulting in two participants being identified who showed unusually low amplifications. DNA extraction and qPCR were repeated for these two samples, resulting in one participant being removed from further analysis due to poor amplification. All qPCR assays were performed by the same investigator.

Relative TL was determined by the ratio of telomere repeat copy number (T) to the single copy gene copy number (36B4) as described by Cawthon [61] and presented as the telomere length to single copy gene ratio (T/S ratio). Absolute telomere length was measured by determining the number of TTAGGG hexamer repeats as described by O'Callaghan and Fenech [62] and is reported in bases. Briefly, a ratio of telomere repeat to single copy gene (36B4) was obtained by first dividing the 36B4 reaction value by 2 (as there are two copies of this gene present in every cell). The telomere length in kilo base (kb) per reaction was then calculated by dividing the TEL reaction value obtained in the qPCR by the 36B4 (single gene reference). This was converted to length per telomere (kb) by dividing by 92 (there are 46 chromosomes with a telomere at each end and, therefore, 92 telomeres in total in the human genome). Telomere length is reported in bases by multiplying by 1000.

### 2.7. Covariates

Data for eight pre-identified covariates were collected: age, sex, body mass index, smoking history, paternal age at birth, education, medical health history and stressful life events.

### 2.8. Statistical analysis

Descriptive statistics are displayed as mean and standard deviation. Total lifetime MET hours and

paternal age at birth were log transformed, and a reflect and square root transformation was applied to lifetime health eating category scores, to normalize the data. Pearson's correlation was used to assess the relationship between absolute and relative salivary cell telomere length and age. Multiple linear regression was used to investigate the independent contributions of age, total lifetime MET-hours, lifetime food category scores, lifetime stressful events scores, paternal age at birth and sex on telomere length. Data were screened to ensure all assumptions for the use of multiple linear regression were met. Statistical analyses were performed using SPSS version 24.0 with an alpha level of  $p < 0.05$ .

## 3. Results

### 3.1. Salivary cell DNA quality and quantity

Considering the unique nature of the study design and population, it was deemed important to include details on the quality and quantity of the DNA extracted from saliva samples. The quality of the DNA was very good as assessed by agarose gel electrophoresis and spectrophotometry which showed no evidence of DNA degradation. The average 260/280 ratio was 2.0 (pure DNA will have a 260/280 ratio of 1.7–2.0 [62]). The average quantity of DNA, as assessed by spectrophotometry, was  $42.7 \pm 23.7$  ng/ $\mu\text{l}$  (range 5.8–95 ng/ $\mu\text{l}$ ).

### 3.2. Participant characteristics

Descriptive statistics were calculated and displayed as mean and standard deviation. Participant characteristics are presented in Table 1. Sixty-five percent (32 of 49) were ultra-runners, 18% (9 of 49) were ultra-cyclists and 16% (8 of 49) were a combination (ultra-runner, ultra-cyclist and/or ultra-triathlete). Sixty-three percent of participants were men. Men were older and had a higher BMI than women ( $50.8 \pm 10.1$  years for men vs  $45.2 \pm 6.7$  years for women,  $p = .043$  and  $23.7 \pm 2.6$  kg/m<sup>2</sup> for men vs  $21.9 \pm 1.7$  kg/m<sup>2</sup> for women,  $p = 0.008$ ).

### 3.3. Predictors of telomere length

Results of the multiple linear regression are presented in Tables 2 and 3. None of the measured variables were statistically significant cross-sectional

Table 1  
Participant characteristics

Ultra-endurance exercisers ( $n = 49$ )	
Absolute telomere length (in bases)	1602 $\pm$ 572
Relative telomere length (T/S ratio <sup>*†</sup> )	1.41 $\pm$ 0.05
Age (years)	48.7 $\pm$ 9.2
Lifetime total MET hours	133072 $\pm$ 106911
Lifetime food category scores	
Healthy category	117 $\pm$ 5.3
Less healthy category	23.9 $\pm$ 2.1
Vegetables and fruit	63.9 $\pm$ 4.5
Lifetime stressful events scores	9 $\pm$ 3.5
Father's age at birth (years)	28.5 $\pm$ 7.4
Education [ $n$ , (%)]	
$\leq$ high school	4 (8)
Some college, but no degree	12 (24)
Associate or bachelor's degree	18 (37)
Postgraduate	15 (31)
Smoke history [ $n$ , (%)]	
Current smoker	1 (2)
Past smoker	14 (29)
Never smoked	34 (69)
Ethnicity [ $n$ , (%)]	
Caucasian	46 (94)
Other	3 (6)
Country [ $n$ , (%)]	
Australia	29 (59)
Canada	12 (25)
United States	8 (16)

Data presented as mean  $\pm$  standard deviation and  $n$  (%) for categorical variables. \*ratio of telomere repeat copy number (T) to the single copy gene copy number (36B4). †Log transformed.

437 predictors of absolute telomere length ( $p > 0.05$ ). Sex  
438 was a statistically significant cross-sectional predic-  
439 tor of relative telomere length ( $p < 0.05$ ) with men  
440 having longer telomeres than women (1.42 + 0.04  
441 for men vs 1.39 + 0.06 for women,  $p = 0.04$ ). There  
442 was no statistically significant difference in abso-  
443 lute telomere length between men and women  
444 (1643 + 517 for men vs 1532 + 666 for women,  
445  $p = 0.52$ ). Age was not statistically correlated with




Fig. 1. Scatter plot illustrating the raw uncontrolled association between age and absolute (a) and relative (b) salivary cell telomere length in ultra-endurance exercisers ( $n = 49$ ; age range 26–74). Relative telomere length was log transformed.

relative or absolute telomere length for all partici- 446  
pants (unadjusted raw data) (Fig. 1). 447

#### 4. Discussion 448

This is the first study to investigate relationships 449  
of lifetime physical activity and diet with telomere 450  
length in current ultra-endurance exercisers. Partici- 451  
pants for this study were drawn from a larger sample 452  
of ultra-endurance exercisers who provided detailed 453  
information on lifetime physical activity [4] and diet 454  
[55]. The ultra-endurance exercisers engaged in a 455  
variety of physical activities across the lifespan with 456

Table 2  
Multiple regression summary statistics for absolute telomere length ( $n = 49$ )

Variable	B	95% [CI]	B	Partial r	t	p
Age	-12.16	[-38.86, 14.54]	-0.232	-0.170	-0.932	0.359
Lifetime total MET hours <sup>†</sup>	-0.899	[-667.17, 665.37]	-0.001	-0.001	-0.003	0.998
Lifetime food category scores						
Healthy category <sup>‡</sup>	56.15	[-334.67, 446.97]	0.106	0.054	0.294	0.771
Less healthy category	-16.28	[-114.08, 81.51]	-0.073	-0.063	-0.341	0.736
Vegetables and fruit	-2.50	[-92.83, 87.83]	-0.021	-0.011	-0.057	0.955
Lifetime stressful events scores	45.68	[-15.31, 106.67]	0.315	0.274	1.532	0.136
Paternal age at birth <sup>†</sup>	-928.98	[-3443.43, 1585.47]	-0.148	-0.139	-0.756	0.456
Sex <sup>§</sup>	-301.63	[-801.99, 198.73]	-0.272	-0.223	-1.233	0.228

Significance set at  $p < 0.05$ . †Log transformed. ‡Reflect and square root transformed. §Coding for categorical variables is male = 0; female = 1.

Table 3  
Multiple regression summary statistics for relative telomere length<sup>†</sup> ( $n=49$ )

Variable	B	95% [CI]	<i>B</i>	Partial r	<i>T</i>	<i>p</i>
Age	-0.002	[-0.005, 0.000]	-0.414	-0.319	-1.813	0.08
Lifetime total MET hours <sup>†</sup>	0.020	[-0.042, 0.082]	0.129	0.124	0.675	0.505
Lifetime food category scores						
Healthy category <sup>‡</sup>	0.017	[-0.019, 0.054]	0.322	0.177	0.971	0.340
Less healthy category	-0.004	[-0.013, 0.005]	-0.170	-0.158	-0.861	0.396
Vegetables and fruit	0.002	[-0.006, 0.010]	0.168	0.091	0.495	0.625
Lifetime stressful events scores	0.005	[-0.001, 0.011]	0.345	0.321	1.825	0.078
Paternal age at birth <sup>†</sup>	-0.123	[-0.357, 0.110]	-0.194	-0.197	-1.080	0.289
Sex <sup>§</sup>	-0.066	[-0.112, -0.019]	-0.585	-0.473	-2.888	0.007*

\*Indicates significant difference at  $p < 0.05$ . <sup>†</sup>Log transformed. <sup>‡</sup>Reflect and square root transformed. <sup>§</sup>Coding for categorical variables is male = 0; female = 1.

total volumes (MET hours) that were, on average, higher than general populations [4]. In this group of ultra-endurance exercisers, whose exercise volumes varied considerably [4], there was no relationship between total lifetime physical activity and telomere length.

The lack of a relationship between food scores and telomere length was unexpected considering the evidence regarding the influence of diet, particularly vegetables and fruit, on telomere length [17, 36, 43, 44]. Similarly, the lack of associations with the less healthy category was not expected as foods that are highly processed, high in saturated fat and/or sugar, have been shown to be associated with shorter telomeres [33, 35, 42]. It has been suggested that certain foods (e.g. processed meat, saturated fat, high sugar foods), many of which are found in the less healthy category in the current study, are negatively associated with telomere length because they increase oxidative stress and inflammation [63]. Although speculative, it is possible that engagement in habitual physical activity, which may lower oxidative stress [64], could help minimize the negative influence of poor dietary choices on telomere attrition. However, the current study collected information on the frequency of eating and the total quantity of food or energy consumed is unknown and it is possible this may have affected the findings. Also, as this was the first study to use food categories developed from the LDQ to assess lifetime diet, it was not possible to compare the food category scores of the ultra-endurance exercisers in the current study to general populations.

A secondary finding from this study was the absence of a relationship between age and telomere length for both absolute and relative telomere length. This finding supports previous research with ultra-endurance runners where a negative correlation of

telomere length with age was not observed [20, 27] and provides some evidence that participating in regular physical activity across the lifetime, that includes ultra-endurance exercise, may help protect against telomere shortening. The lack of association between age and telomere length is of interest considering participants ranged in age from 26–74 years. As expected, the older participants ( $\geq 50$  years,  $n=23$ ) had accumulated significantly more lifetime hours of physical activity (that included ultra-endurance exercise) than the younger ultra-endurance exercisers ( $<50$  years,  $n=26$ ). Studies have reported a positive relationship between habitual physical activity and telomere length [19, 65], particularly amongst older, active individuals [66–68]. It is possible that the accumulating lifetime physical activity of this group, in particular the older participants, had a benefit on telomere length. However, to our knowledge, there are no studies with older ultra-endurance exercisers to help explain the lack of association between age and telomere length and more research with this population is needed.

Possible mechanisms that may explain the lack of associations of lifetime physical activity and diet with telomere length within this group of current ultra-endurance exercisers are the roles of these two variables in promoting general health and wellbeing. Both physical activity and diet may influence levels of oxidative stress and inflammation which are two key underlying mechanisms linked to telomere attrition [63, 69]. Research shows health behaviours tend to cluster by which active individuals tend to eat healthier diets [70] and, therefore, it may be the cumulative effect of health behaviours that benefit telomere length rather than individual components [32, 71, 72]. As such, it is possible the lifetime habits of this group, including physical activity and diet, contributed to preserving telomere length. This study

also included a measure of lifetime stress which is important to consider given the known association between psychological stress and telomere length [15, 73]. In the current study there was no correlation between lifetime stress and telomere length which may be due to the possible role of physical activity on minimising the impact of stress on telomere length. Regular physical activity may reduce the impact of psychological stress as observed in a study in which postmenopausal women with the healthiest behaviours (had the highest score related to sleeping, eating and exercise habits) did not experience the effect of major life stressors on accelerating telomere attrition [74]. Engaging in regular physical activity may explain the lack of relationship between lifetime psychological stress and telomere length in the group of ultra-endurance exercisers in this study. Furthermore, it is possible, as reported in previous studies with ultra-endurance exercisers, there is an upregulation of cellular components involved in telomere maintenance that helps protect against telomere attrition [28]. Although the lack of age-related decline in telomere length may suggest a protective effect of physical activity and diet on telomere length, factors such as small sample size, unmeasured covariates (i.e. genetic variances) and life period specific cofounders (e.g. health related issues, changes in lifestyle or socioeconomic status), need to be considered. The reason for a significant finding related to sex and relative, but not absolute, telomere length is unclear and may be due to a difference in the expression of the control gene between sexes. However, more research that includes both measures of telomere length, within the same population, is necessary.

To our knowledge this is the first study to report on the lifetime physical activity and diet of ultra-endurance exercisers in relation to telomere length. It is the first study to include a measure of diet when investigating telomere length in ultra-endurance exercisers. Detailed information on physical activity and diet were collected across several life periods which is important when investigating telomere length as changes typically take place over long periods of time (i.e. at least 1 year) and may be influenced by cumulative factors over a lifetime [75, 76]. This study is one of two studies that have measured salivary cell telomere length in ultra-endurance exercisers and provides new information on diet for this population. This is the first study to describe both relative and absolute telomere length within the same population of ultra-endurance exercisers, which allows for comparisons using both measures. Furthermore, the measurement

of absolute telomere length in this study provides the substantial opportunity for a more direct comparison to other studies that have also measured absolute telomere length.

#### 4.1. Limitations

This study has five known potential limitations. Firstly, this study may have lower than optimal statistical power due to the sample size and this may, in part, explain the lack of associations with telomere length. Secondly, this study collected retrospective physical activity and diet data and relied on participants' ability to recall information on habits from several decades ago. Thirdly, this study measured telomere length at one point in time and did not have repeated measures of telomere length at different time points across the lifespan. Therefore, there was no information on the rate of telomere attrition or telomere length at baseline. Fourthly, this study did not include a control group. However, it was not possible to obtain a comparison group for whom physical activity and diet were suitably controlled across the lifespan. Fifth, it was not possible to consider, and control for, all potential co-variables that could influence telomere length across the lifespan. It is possible there were unmeasured variables (i.e. genetic variances) that may have influenced the findings.

More research that assesses lifetime physical activity and diet in a larger sample of ultra-endurance exercisers would assist in explaining the role of these two variables across the lifespan on telomere length. Early life experiences can have a positive or negative effect on telomere length and future research investigating the diet of ultra-endurance exercisers and telomere length during different life stages is needed. This should include information on the type, intensity, duration and frequency of physical activity of ultra-endurance exercisers which would help explain how, and if, diet and physical activity interact in relationship to telomere length.

## 5. Conclusions

This study adds to the limited amount of research investigating telomere length in ultra-endurance exercisers and provides the first information on diet and telomeres for this population. The primary results showed no relationship of lifetime physical activity or diet with telomere length. The secondary



631 finding that showed an absence of telomere short-  
 632 ening with age is of interest and provides some  
 633 evidence for a potential protective role of physical  
 634 activity and/or diet across the lifetime on cellular  
 635 aging. It is possible that engaging in various forms  
 636 of physical activity throughout life, which includes  
 637 ultra-endurance exercise, may help delay telomere  
 638 shortening. It is also possible the lack of age-related  
 639 telomere shortening is independently due to lifetime  
 640 diet or due to an interactive effect between physi-  
 641 cal activity and diet. However, this process remains  
 642 unclear and requires further investigation. As par-  
 643 ticipation in ultra-endurance exercise continues to  
 644 increase worldwide and the number of older indi-  
 645 viduals participating in this form of physical activity  
 646 continues to grow, it is important to understand the  
 647 impact this form of extreme exercise has on cellular  
 648 aging and how diet may influence this process.

## 649 Acknowledgments

650 All authors assisted with study design. KB was  
 651 responsible for data collection, data analysis and writ-  
 652 ing of the manuscript drafts. All authors provided  
 653 input on data analysis and subsequent manuscript  
 654 drafts. GL provided statistical guidance. All authors  
 655 read and approved the final version of the manuscript.  
 656 KB reported no potential conflict of interest. AK  
 657 reported no potential conflict of interest. GL reported  
 658 no potential conflict of interest. SB reported no poten-  
 659 tial conflict of interest. CS reported no potential  
 660 conflict of interest. KB received a Fee Offset Schol-  
 661 arship and a Stipend Scholarship from the Australian  
 662 Government Research Training Program during this  
 663 project.

## 664 References

- 665 [1] Knechtle B. Ultramarathon runners: Nature or nurture? *Int J*  
 666 *Sports Physiol Perform.* 2012;7(4):310-2.  
 667 [2] Hoffman MD, Fogard K. Demographic characteristics  
 668 of 161-km ultramarathon runners. *Res Sports Med.*  
 669 2012;20(1):59-69.  
 670 [3] Hoffman MD, Krishnan E. Health and exercise-related  
 671 medical issues among 1,212 ultramarathon runners: Base-  
 672 line findings from the ultrarunners longitudinal tracking  
 673 (ULTRA) study. *PLoS One.* 2014;9(1):e83867.  
 674 [4] Birkenhead K, Lovell G, Barr SI, Solomon C. Changes  
 675 in physical activity across the lifetime of current ultra-  
 676 endurance exercisers. *J Phys Activity Res.* 2018;3(1):11-9.  
 677 [5] Cejka N, Rust CA, Lepers R, Onywera V, Rosemann T,  
 678 Knechtle B. Participation and performance trends in 100-  
 679 km ultra-marathons worldwide. *J Sports Sci.* 2014;32(4):  
 680 354-66.  
 681 [6] da Fonseca-Engelhardt K, Knechtle B, Rust CA, Knechtle  
 682 P, Lepers R, Rosemann T. Participation and performance  
 683 trends in ultra-endurance running races under extreme condi-  
 684 tions - 'Spartathlon' versus 'Badwater', *Extrem Physiol Med.*  
 685 2013;2(1):15.  
 686 [7] Kipling D. Telomeres, replicative senescence and human age-  
 687 ing. *Maturitas.* 2001;38(1):25-37.  
 688 [8] Tzanetakou IP, Katsilambros NL, Benetos A, Mikhailidis DP,  
 689 Perrea DN. "Is obesity linked to aging?" Adipose tissue and  
 690 the role of telomeres. *Ageing Res Rev.* 2012;11(2):220-9.  
 691 [9] Blackburn EH. Structure and function of telomeres. *Nature.*  
 692 1991;350(6319):569-73.  
 693 [10] Zhu Y, Liu X, Ding X, Wang F, Geng X. Telomere and  
 694 its role in the aging pathways: telomere shortening, cell  
 695 senescence and mitochondria dysfunction. *Biogerontology.*  
 696 2019;20(1):1-16.  
 697 [11] Dankel SJ, Loenneke JP, Loprinzi PD. The impact of over-  
 698 weight/obesity duration and physical activity on telomere  
 699 length: An application of the WATCH paradigm. *Obes Res*  
 700 *Clin Pract.* 2017;11(2):247-52.  
 701 [12] Latifovic L, Peacock SD, Massey TE, King WD. The influ-  
 702 ence of alcohol consumption, cigarette smoking, and physical  
 703 activity on leukocyte telomere length. *Cancer Epidemiol*  
 704 *Biomarkers Prev.* 2015;25(2):374-80.  
 705 [13] Sjögren P, Fisher R, Kallings L, Svenson U, Roos G,  
 706 Hellénius ML. Stand up for health—avoiding sedentary  
 707 behaviour might lengthen your telomeres: secondary out-  
 708 comes from a physical activity RCT in older people. *Br J*  
 709 *Sports Med* 2014;48(19):1407-9.  
 710 [14] Shalev I, Entringer S, Wadhwa PD, Wolkowitz OM,  
 711 Puterman E, Lin J, Epel ES. Stress and telomere biol-  
 712 ogy: A lifespan perspective. *Psychoneuroendocrinology.*  
 713 2013;38(9):1835-42.  
 714 [15] Epel ES. Psychological and metabolic stress: A recipe for  
 715 accelerated cellular aging? *Hormones.* 2009;8(1):7-22.  
 716 [16] Gong Y, Tian G, Xue H, Zhang X, Zhao Y, Cheng G. Higher  
 717 adherence to the 'vegetable-rich' dietary pattern is related to  
 718 longer telomere length in women. *Clin Nutr.* 2017.  
 719 [17] Crous-Bou M, Fung TT, Prescott J, Julin B, Du M, Sun Q,  
 720 Rexrode KM, Hu FB, De Vivo I. Mediterranean diet and  
 721 telomere length in Nurses' Health Study: Population based  
 722 cohort study. *BMJ.* 2014;349.  
 723 [18] Tucker LA. Physical activity and telomere length in U.S.  
 724 men and women: An NHANES investigation. *Prev Med.*  
 725 2017;100:145-51.  
 726 [19] Kim JH, Ko JH, Lee DC, Lim I, Bang H. Habitual  
 727 physical exercise has beneficial effects on telomere  
 728 length in postmenopausal women. *Menopause.* 2012;19(10):  
 729 1109-15.  
 730 [20] Denham J, Nelson CP, O'Brien BJ, Nankervis SA, Denniff M,  
 731 Harvey JT, Marques FZ, Codd V, Zukowska-Szczechowska  
 732 E, Samani NJ, Tomaszewski M, Charchar FJ. Longer leuko-  
 733 cyte telomeres are associated with ultra-endurance exercise  
 734 independent of cardiovascular risk factors. *PLoS One.*  
 735 2013;8(7).  
 736 [21] Cherkas LF, Hunkin JL, Kato BS, Richards JB, Gardner  
 737 JP, Surdulescu GL, Kimura M, Lu X, Spector TD, Aviv  
 738 A. The association between physical activity in leisure  
 739 time and leukocyte telomere length. *Arch Intern Med.*  
 740 2008;168(2):154-158.

- 741 [22] Frenk S, Housely J. Gene expression hallmarks of cellular  
742 ageing. *Biogerontology*. 2018;19:1-20.
- 743 [23] Aviv A. Telomeres and human aging: Facts and fibs, *Sci*  
744 *Aging Knowledge Environ*. 2004;(51):43.
- 745 [24] Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer  
746 G. The hallmarks of aging. *Cell*. 2013;153(6):1194-217.
- 747 [25] Pereira MA, Fitzer Gerald SJ, Gregg EW, Joswiak ML, Ryan  
748 WJ, Suminski RR, Utter AC, Zmuda JM. A collection of  
749 physical activity questionnaires for health-related research.  
750 *Med Sci Sports Exerc*. 1997;29(6 Suppl):S1-205.
- 751 [26] Ludlow AT, Ludlow LW, Roth SM. Do telomeres adapt to  
752 physiological stress? Exploring the effect of exercise on  
753 telomere length and telomere-related proteins, *Bio Med Res*  
754 *Int*. 2013;2013:1-15.
- 755 [27] Borghini A, Giardini G, Tonacci A, Mastorci F, Mercuri A,  
756 Sposta SM, Moretti S, Andreassi MG, Pratali L. Chronic  
757 and acute effects of endurance training on telomere length.  
758 *Mutagenesis*. 2015;30(5):711-6.
- 759 [28] Laye MJ, Solomon TPJ, Karstoft K, Pedersen KK, Nielsen  
760 SD, Pedersen BK. Increased shelterin mRNA expression in  
761 peripheral blood mononuclear cells and skeletal muscle fol-  
762 lowing an ultra-long-distance running event. *J Appl Physiol*.  
763 2012;112(5):773-81.
- 764 [29] J. Denham, Lack of association between PBMC telomere  
765 length and endurance exercise, *J Appl Biomed*. 2016;15(1):  
766 9-13.
- 767 [30] Denham J, O'Brien BJ, Prestes PR, Brown NJ, Charchar  
768 FJ. Increased expression of telomere-regulating genes in  
769 endurance athletes with long leukocyte telomeres. *J Appl*  
770 *Physiol*. 2016;120(2):148-58.
- 771 [31] García-Calzón S, Moleres A, Martínez-González MA,  
772 Martínez JA, Zalba G, Martí A. Dietary total antioxidant  
773 capacity is associated with leukocyte telomere length in a  
774 children and adolescent population. *Clin Nutr*. 2015;  
775 34(4):694-9.
- 776 [32] Shiels PG, McGlynn LM, MacIntyre A, Johnson PCD,  
777 da Batty GD, Burns H, Cavanagh J, Deans KA, Ford I,  
778 McConnachie A, McGinty A, McLean JS, Millar K, Sattar N,  
779 Tannahill C, Velupillai YN, Packard CJ. Accelerated telomere  
780 attrition is associated with relative household income,  
781 diet and inflammation in the pSoBid Cohort, *PLoS One*.  
782 2011;6(7).
- 783 [33] Song Y, You NY, Song Y, Kang MK, Hou L, Wallace R,  
784 Eaton CB, Tinker LF, Liu S. Intake of small-to-medium-chain  
785 saturated fatty acids is associated with peripheral leuko-  
786 cyte telomere length in postmenopausal women. *J Nutr*.  
787 2013;143(6):907-14.
- 788 [34] Tiainen AMK, Mannisto S, Blomstedt PA, Moltchanova E,  
789 Perala MM, Kaartinen NE, Kajantie E, Kananen L, Hovatta  
790 I, Eriksson JG. Leukocyte telomere length and its relation to  
791 food and nutrient intake in an elderly population. *Eur J Clin*  
792 *Nutr*. 2012;66(12):1290-4.
- 793 [35] Nettleton JA, Diez-Roux A, Jenny NS, Fitzpatrick AL, Jacobs  
794 Jr DR. Dietary patterns, food groups, and telomere length in  
795 the multi-ethnic study of atherosclerosis (MESA), *Am J Clin*  
796 *Nutr*. 2008;88(5):1405-12.
- 797 [36] Marcon F, Siniscalchi E, Crebelli R, Saieva C, Sera F,  
798 Fortini P, Simonelli V, Palli D. Diet-related telomere short-  
799 ening and chromosome stability. *Mutagenesis*. 2012;27(1):  
800 49-57.
- 801 [37] Puterman E, Lin J, Krauss J, Blackburn EH, Epel ES. Deter-  
802 minants of telomere attrition over 1 year in healthy older  
803 women: stress and health behaviors matter. *Mol Psychiatry*.  
804 2014;1-7.
- 805 [38] Cassidy A, De Vivo I, Liu Y, Han J, Prescott J, Hunter DJ,  
806 Rimm EB. Associations between diet, lifestyle factors, and  
807 telomere length in women. *Am J Clin Nutr*. 2010;91(5):  
808 1273-80.
- 809 [39] Tucker LA. Consumption of nuts and seeds and telomere  
810 length in 5,582 men and women of the national health  
811 and nutrition examination survey (NHANES), *J Nutr Health*  
812 *Aging*. 2017;21(3):233-40.
- 813 [40] Lee JY, Shin C, Baik I. Longitudinal associations between  
814 micronutrient consumption and leukocyte telomere length. *J*  
815 *Hum Nutr Diet*. 2016;30(2):236-243.
- 816 [41] Fretts AM, Howard BV, Siscovick DS, Best LG, Beresford  
817 SA, Mete M, Eilat-Adar S, Sotoodehnia N, Zhao J. Processed  
818 meat, but not unprocessed red meat, is inversely associated  
819 with leukocyte telomere length in the strong heart family  
820 study. *J Nutr*. 2016;146(10):2013-8.
- 821 [42] Leung CW, Laraia BA, Needham BL, Rehkopf DH, Adler  
822 NE, Lin J, Blackburn EH, Epel ES. Soda and cell aging:  
823 Associations between sugar-sweetened beverage consump-  
824 tion and leukocyte telomere length in healthy adults from  
825 the national health and nutrition examination surveys. *Am J*  
826 *Public Health*. 2014;104(12):2425-31.
- 827 [43] Lee JY, Jun NR, Yoon D, Shin C, Baik I. Association between  
828 dietary patterns in the remote past and telomere length. *Eur*  
829 *J Clin Nutr*. 2015;1-5.
- 830 [44] Gu Y, Honig LS, Schupf N, Lee JH, Luchsinger JA,  
831 Stern Y, Scarmeas N. Mediterranean diet and leukocyte  
832 telomere length in a multi-ethnic elderly population. *Age*.  
833 2015;37(2):24.
- 834 [45] Eisenberg DT. An evolutionary review of human telomere  
835 biology: The thrifty telomere hypothesis and notes  
836 on potential adaptive paternal effects. *Am J Hum Biol*.  
837 2011;23(2):149-67.
- 838 [46] Astuti Y, Wardhana A, Watkins J, Wulaningsih W, Network  
839 PR. Cigarette smoking and telomere length: A system-  
840 atic review of 84 studies and meta-analysis. *Environ Res*.  
841 2017;158:480-9.
- 842 [47] Mundstock E, Zatti H, Louzada FM, Oliveira SG, Guma  
843 FT, Parisi MM, Rueda AB, Machado DG, Stein RT,  
844 Jones MH, Sarria EE, Barbe-Tuana FM, Mattiello R. Effects  
845 of physical activity in telomere length: Systematic  
846 review and meta-analysis. *Ageing Res Rev*. 2015;22:  
847 72-80.
- 848 [48] Oliveira BS, Zunzunegui MV, Quinlan J, Fahmi H, Tu MT,  
849 Guerra RO. Systematic review of the association between  
850 chronic social stress and telomere length: A life course per-  
851 spective. *Ageing Res Rev*. 2016;26:37-52.
- 852 [49] Rafie N, Golpour Hamedani S, Barak F, Safavi SM, Miragha-  
853 jani M. Dietary patterns, food groups and telomere length:  
854 A systematic review of current studies, *Eur J Clin Nutr*.  
855 2017;71(2):151-8.
- 856 [50] Wren ME, Shirliff EA, Drury SS. Not all biofluids are  
857 created equal: chewing over salivary diagnostics and the  
858 epigenome. *Clin Ther*. 2015;37(3):529-39.
- 859 [51] Stout SA, Lin J, Hernandez N, Davis EP, Blackburn E, Carroll  
860 JE, Glynn LM. Validation of minimally-invasive sample col-  
861 lection methods for measurement of telomere length, *Front*  
862 *Aging Neurosci*. 2017;9:397.
- 863 [52] Mitchell C, Hobcraft J, McLanahan SS, Siegeld SR, Berg  
864 A, Brooks-Gunn J, Garfinkel I, Notterman D, Social

- 865 disadvantage, genetic sensitivity, and children's telomere  
866 length, *Proc Natl Acad Sci U S A*. 2014;111(16):5944-9.
- 867 [53] Friedrich U, Griese EU, Schwab M, Fritz P, Thon KP, Klotz  
868 U. Telomere length in different tissues of elderly patients.  
869 *Mech Ageing Dev*. 2000;119(3):89-99.
- 870 [54] Hansen TV, Simonsen MK, Nielsen FC, Hundrup YA.  
871 Collection of blood, saliva, and buccal cell samples in a  
872 pilot study on the Danish nurse cohort: Comparison of the  
873 response rate and quality of genomic DNA. *Cancer Epi-  
874 demiol Biomarkers Prev*. 2007;16(10):2072-6.
- 875 [55] Birkenhead KL, Barr SI, Lovell G, Solomon C. Changes in  
876 diet, and the relationship between diet and physical activ-  
877 ity within and across the lifetime of current ultra-endurance  
878 exercisers. *J Sports Med Phys Fitness*. 2019;59(5):798-807.
- 879 [56] Commonwealth of Australia. (2007). National statement on  
880 ethical conduct in human research 2007 (Updated May 2015).
- 881 [57] Hosking D, Danthiir V, Nettelbeck T, Wilson C. Assessing  
882 lifetime diet: Reproducibility of a self-administered, non-  
883 quantitative FFQ. *Public Health Nutr*. 2011;14(5):801-8.
- 884 [58] Chasan-Taber L, Erickson JB, McBride JW, Nasca PC,  
885 Chasan-Taber S, Freedson PS. Reproducibility of a self-  
886 administered lifetime physical activity questionnaire among  
887 female college alumnae. *Am J Epidemiol* 2002;155(3):282-9.
- 888 [59] Rosmalen JG, Bos EH, de Jonge P. Validation of the long-  
889 term difficulties inventory (LDI) and the list of threatening  
890 experiences (LTE) as measures of stress in epidemi-  
891 ological population-based cohort studies. *Psychol Med*.  
892 2012;42(12):2599-608.
- 893 [60] Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bas-  
894 sett Jr DR, Tudor-Locke C, Greer JL, Vezina J, Whitt-Glover  
895 MC, Leon AS. 2011 compendium of physical activities: A  
896 second update of codes and MET values. *Med Sci Sports  
897 Exerc*. 2011;43(8):1575-81.
- 898 [61] Cawthon RM. Telomere measurement by quantitative PCR,  
899 *Nucleic Acids Res*. 2002;30(10).
- 900 [62] O'Callaghan NJ, Fenech M. A quantitative PCR method  
901 for measuring absolute telomere length. *Biol Proced Online*.  
902 2011;13(3).
- 903 [63] Shivappa N, Wirth MD, Hurley TG, Hebert JR. Association  
904 between the dietary inflammatory index (DII) and telomere  
905 length and C-reactive protein from the national health  
906 and nutrition examination survey-1999-2002, *Mol Nutr Food  
907 Res*. 2017;61(4).
- 908 [64] Arsenis NC, You T, Ogawa EF, Tinsley GM, Zuo L. Physical  
909 activity and telomere length: Impact of aging and potential  
910 mechanisms of action. *Oncotarget*. 2017.
- 911 [65] Sassenroth D, Meyer A, Salewsky B, Kroh M, Norman K,  
Steinhagen-Thiessen E, Demuth I. Sports and exercise at  
different ages and leukocyte telomere length in later life-  
Data from the Berlin aging study II (BASE-II), *PLoS One*.  
2015;10(12):e0142131.
- [66] La Rocca TJ, Seals DR, Pierce GL. Leukocyte telomere  
length is preserved with aging in endurance exercise-trained  
adults and related to maximal aerobic capacity. *Mech Ageing  
Dev*. 2010;131(2):165-7.
- [67] Østhus IBØ, Sgura A, Berardinelli F, Alsnes IV, Brønstad  
E, Rehn T, Støbakk PK, Hatle H, Wisløff U, Nauman J.  
Telomere length and long-term endurance exercise: Does  
exercise training effect biological age? A pilot study. *PLoS  
One*. 2012;7(12):e52769.
- [68] Werner C, Fürster T, Widmann T, Pöss J, Roggia C, Hanhoun  
M, Scharhag J, Büchner N, Meyer T, Kindermann W, Haen-  
deler J, Böhm M, Laufs U. Physical exercise prevents cellular  
senescence in circulating leukocytes and in the vessel wall,  
*Circulation*. 2009;120(24):2438-47.
- [69] Ludlow AT, Roth SM. Physical activity and telomere biology:  
Exploring the link with aging-related disease prevention. *J  
Ageing Res*. 2011;2011.
- [70] Glanz K, Basil M, Maibach E, Goldberg J, Snyder D. Why  
Americans eat what they do: Taste, nutrition, cost, conve-  
nience, and weight control concerns as influences on food  
consumption. *J Am Diet Assoc*. 1998;98(10):1118-26.
- [71] Mirabello L, Huang WY, Wong JYY, Chatterjee N, Reding  
D, Crawford ED, De Vivo I, Hayes RB, Savage SA. The  
association between leukocyte telomere length and cigarette  
smoking, dietary and physical variables, and risk of prostate  
cancer, *Ageing Cell*. 2009;8(4):405-13.
- [72] Sun Q, Shi L, Prescott J, Chiuve SE, Hu FB, De Vivo I,  
Stampfer MJ, Franks PW, Manson JE, Rexrode KM. Healthy  
lifestyle and leukocyte telomere length in U.S. women, *PLoS  
One*. 2012;7(5):e38374.
- [73] Shalev I. Early life stress and telomere length: Investigating  
the connection and possible mechanisms: A critical survey of  
the evidence base, research methodology and basic biology.  
*Bioessays*. 2012;34(11):943-52.
- [74] Puterman E, Lin J, Blackburn E, O'Donovan A, Adler N,  
and Epel E. The power of exercise: Buffering the effect of  
chronic stress on telomere length, *PLoS One* 5(5):2010.
- [75] Puterman E, Epel E. An intricate dance: Life experi-  
ence, multisystem resiliency, and rate of telomere decline  
throughout the lifespan. *Soc Personal Psychol Compass*.  
2012;6(11):807-25.
- [76] Akkad A, Hastings R, Konje JC, Bell SC, Thurston H,  
Williams B. Telomere length in small-for-gestational-age  
babies. *Br J Obstet Gynaecol*. 2006;113(3):318-23.