



## LJMU Research Online

**Latour, M, Calamida, A, Husser, T-O, Kamann, S, Dreizler, S and Brinchmann, J**

**A stellar census in globular clusters with MUSE. A new perspective on the multiple main sequences of  $\omega$  Centauri**

<http://researchonline.ljmu.ac.uk/id/eprint/15481/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Latour, M, Calamida, A, Husser, T-O, Kamann, S, Dreizler, S and Brinchmann, J A stellar census in globular clusters with MUSE. A new perspective on the multiple main sequences of  $\omega$  Centauri. *Astronomy and Astrophysics*. ISSN 0004-6361 (Accepted)**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

LETTER TO THE EDITOR

# A stellar census in globular clusters with MUSE.

## A new perspective on the multiple main sequences of $\omega$ Centauri<sup>★,★★</sup>

M. Latour<sup>1</sup>, A. Calamida<sup>2</sup>, T.-O. Husser<sup>1</sup>, S. Kamann<sup>3</sup>, S. Dreizler<sup>1</sup>, and J. Brinchmann<sup>4,5</sup>

<sup>1</sup> Institute for Astrophysics, Georg-August-University, Friedrich-Hund-Platz 1, 37077 Göttingen, Germany, e-mail: marilyn.latour@uni-goettingen.de

<sup>2</sup> Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

<sup>3</sup> Astrophysics Research Institute, Liverpool John Moores University, 146 Brownlow Hill, Liverpool L3 5RF, United Kingdom

<sup>4</sup> Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, PT4150-762 Porto, Portugal

<sup>5</sup> Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands

Received xxxx, 2021; accepted xxxx, 2021

### ABSTRACT

**Context.**  $\omega$  Cen is a rare example of a globular cluster where the iron abundance of the stars spans more than one order of magnitude. Many spectroscopic investigations of its red-giant- and sub-giant- branches have revealed multiple peaks in the iron abundance distribution. The metallicity distribution of main-sequence (MS) stars is not well characterized yet, due to the faintness of the stars and lack of data. So far, almost all studies of MS stars are based on photometric measurements.

**Aims.** Our goal is to investigate the metallicity distribution of a statistically significant sample of MS stars in  $\omega$  Cen. In particular, we aim at revisiting the metallicity difference between the red and blue MS of the cluster.

**Methods.** We use MUSE spectra obtained for the central region of  $\omega$  Cen to derive metallicities for  $\approx 3000$  MS stars.

**Results.** We find that blue MS stars are on average  $\approx 0.1$  dex more metal-rich than their red counterparts. On the basis of this new estimate, we find that the two sequences can be fit on the Hubble Space Telescope color-magnitude diagram with two isochrones having the same global metallicity and age but a slightly higher helium abundance for the blue MS, i.e.  $\Delta Y \lesssim 0.1$ . Furthermore, we determine the average metallicity of the five main populations along  $\omega$  Cen MS and these estimates are consistent with expectations from previous photometric studies.

**Key words.** globular clusters: individual:  $\omega$  Centauri — Stars: abundances — Techniques: spectroscopic

## 1. Introduction

$\omega$  Cen is the most massive Galactic globular cluster (GGC), with a mass of  $3.5 \times 10^6 M_{\odot}$  (Baumgardt & Hilker 2018). The iron abundance of its stars spans more than one order of magnitude  $-2.2 \lesssim [Fe/H] \lesssim -0.6$  and light-element abundance variations are observed within groups of stars along this metallicity range (Norris & Da Costa 1995; Suntzeff & Kraft 1996; Calamida et al. 2009; Johnson & Pilachowski 2010; Marino et al. 2011; Husser et al. 2020). All these studies find a metallicity distribution for red-giant branch (RGB) stars with multiple main peaks around  $[Fe/H] \approx -1.7, -1.4, -1.0, -0.8$ . The large range of iron abundances observed among cluster RGB stars was also detected among sub-giant branch (SGB) and main sequence turn-off (MSTO) stars (Kayser et al. 2006; Hilker et al. 2004; Sollima et al. 2005; Stanford et al. 2006; Villanova et al. 2007; Pancino et al. 2011).

The chemical picture of  $\omega$  Cen's stars is so complex that there currently are different hypothesis on its formation. The two main hypothesis are that  $\omega$  Cen is the nucleus or the nuclear star cluster of a dwarf galaxy accreted by the Milky Way or the result of the merger of two or more clusters (Norris et al. 1997; Pancino et al. 2000; Bekki & Norris 2006; Massari et al. 2019; Ibata et al. 2019; Alfaro-Cuello et al. 2019; Calamida et al. 2020; Pfeffer et al. 2021). In particular, the merger of smaller clusters or a cluster and a nuclear stars cluster could have happened in a dwarf galaxy, where these encounters are more frequent than in the Galactic halo due to the lower velocity dispersion, and the system could have been later accreted by the Galaxy (Amaro-Seoane et al. 2013; Bekki & Tsujimoto 2016; Gavagnin et al. 2016; Pasquato & Chung 2016).

A bifurcation of the main sequence (MS) of  $\omega$  Cen into two main components, the so called blue MS (bMS) and red MS (rMS), was first revealed by Hubble Space Telescope (HST) photometry (Anderson 2002; Bedin et al. 2004). Although the discrete nature of the cluster's RGB was known from previous photometric investigations (Lee et al. 1999; Pancino et al. 2000), this was the first time that such a bimodal distribution of stars was observed along the MS of a GGC. Follow up spectroscopy of a handful of MS stars by Piotto et al. (2005, hereinafter PI05) indicated that the stars belonging to the bMS had a higher metallicity ( $+0.3 \pm 0.2$  dex) compared to stars found on the rMS. This reversal of the expected color progression with metallicity was

\* Based on observations collected at the European Organisation for Astronomical Research in the Southern Hemisphere, Chile (Program IDs 094.D-0142(B), 095.D-0629(A), 096.D-0175(A), 097.D-0295(A), 098.D-0148(A), 099.D-0019(A), 0100.D-0161(A), 0101.D-0268(A), 0102.D-0270(A), 0103.D-0204(A), and 0104.D-0257(B))

\*\* Table B.1 is only available at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (XXXX) or via <http://cdsarc.u-strasbg.fr/viz-bin/cat/J/A+A/XXX/Lzzz>

explained in terms of different helium abundances, i.e., the bMS stars would contain a much higher helium proportion than the stars on the rMS ( $\Delta Y \approx 0.12 - 0.15$ ; Norris 2004, PI05, King et al. 2012).


Numerous studies tried to characterize the properties of the bimodal MS, such as the radial distribution, age, and chemical composition (Sollima et al. 2007; King et al. 2012; Calamida et al. 2017, 2020; Tailo et al. 2016; Milone et al. 2017). All these studies were based on photometric investigations. In recent years, more subpopulations were identified along the MS from the initial two: a high-metallicity component called MSa, that can distinctly be followed through the sub-giant branch and onto the RGB (SGB-a, RGB-a), and Bellini et al. (2017b, hereinafter BE17) separated 15 subpopulations by using pseudo-color diagrams based on a combination of the  $F275W$ ,  $F336W$ , and  $F438W$  WFC3/HST filters. These subpopulations are divided into five main components in the CMDs of  $\omega$  Cen; the rMS, bMS, MSa, plus the newly identified MSd and MSe. The chemical properties of these populations were qualitatively assessed thanks to the use of magnitudes in multiple filters ranging from the near UV to the near-infrared (Bellini et al. 2017a). While the effect of *CNO* variations leaves a rather clear signature in the WFC3/HST  $F336W$  and  $F438W$  filters, the effects of helium, and especially iron are more subtle (see Sect. 5 of BE17). Although very insightful, the information that can be gained from photometry is limited and this is where spectroscopy proves essential to gain a clearer picture. While a handful of spectroscopic surveys of RGB, SGB and MSTO stars in  $\omega$  Cen have been carried out in the past, no spectroscopic studies of MS stars have been done since PI05. The faintness of the MS stars combined with the highly crowded field make such observations extremely challenging and time-consuming. Indeed, PI05 observed only 34 MS stars using the multi-object spectrograph FLAMES on the Very Large Array Telescope (VLT) for a total exposure time of 12h. The spectra of the bMS and rMS stars were then co-added, resulting in signal-to-noise ratios (S/N) of  $\approx 30$  at a spectral resolution  $R = 6400$  (3960-4560 Å). The authors analyzed the two spectra that were considered as representative of the bMS and rMS stars.

In this Letter, we present, after more than fifteen years, new spectroscopic observations of MS stars in  $\omega$  Cen. Our goal is to revisit the properties of the red and blue MS based on spectroscopic evidence and look at the metallicity distribution of the MS subpopulations identified by BE17. To achieve this, we use MUSE observations of  $\omega$  Cen obtained as part of the GTO time dedicated to GGCs (PIs: S. Dreizler, S. Kamann).

## 2. MUSE observations and spectral analysis

We use the MUSE GTO observations of  $\omega$  Cen obtained in wide field mode ( $1' \times 1'$ ) between April 2014 and March 2019. The spectra cover the 4750–9350 Å range with an average spectral resolution of  $\sim 2.5$  Å ( $R \sim 3000$ ), although this varies slightly across the wavelength range (Husser et al. 2016). A general description of the data reduction and spectral analysis is presented in Husser et al. (2016) and Kamann et al. (2018). Additional information on the MUSE data used for this work is included in App. A. In the following, we briefly provide some details specific to the analysis of the  $\omega$  Cen MS spectra.

Because each of the seven MUSE fields were observed multiple times, the individual spectra of each star are co-added after being corrected for their measured radial velocities (RVs). The combined spectra are then fitted using the spectrum fit-



**Fig. 1.**  $F606W, F438W - F814W$  CMD from Bellini et al. (2017a) photometric catalog. The selected samples of blue and red MS stars are marked with blue and red dots, respectively. Two isochores from the BASTI database with different helium content and same metallicity and age are overlotted.


ting framework SPEXSY<sup>1</sup> and the Göttingen spectral library of PHOENIX models (Husser et al. 2013). The best fit provides  $T_{\text{eff}}$  and  $[M/H]$ , while the  $\log g$  is fixed to the value provided by the isochores and the  $\alpha$ -enhancement is kept constant at  $[\alpha/Fe]=0.3$ . For  $\omega$  Cen, up to three sets of photometry are available per star: from Sarajedini et al. (2007), Bellini et al. (2017a), and from archival ACS observation for Field #7 (see Sect. 4 of Kamann et al. 2018). For each of these photometric sets, a best fitting isochrone from the PARSEC database (Marigo et al. 2017) is selected to simultaneously match the position of the MS, SGB and RGB. The  $\log g$  of a star is then determined by finding the nearest point on the isochrone and the average  $\log g$  is adopted when the star is present in more than one photometric catalog. We note that this approach differs from previous spectroscopic studies of brighter stars in GCs. In literature work,  $T_{\text{eff}}$  is usually obtained from empirical color-temperature relations (e.g. Alonso et al. 1999) and  $[Fe/H]$  is obtained from the direct fit of Fe lines in high-resolution spectra. We fit the whole spectrum of our stars where the strongest metallic features are the Mg and Ca triplets. For this reason, we keep the  $[M/H]$  nomenclature to refer to our metallicities, as it is closer to a global metallicity and is sensitive to star-to-star variations in Mg and Ca abundances.

We paid a particular attention to the uncertainties of the metallicity measurements and performed a calibration following a procedure similar to the one used for radial velocities in Kamann et al. (2016). For each star, we obtain a correction factor by which we multiply the formal uncertainty returned by the fitting procedure.

## 3. Selection of red and blue main sequence stars

In order to select our blue and red MS star samples we retrieved the WFC3/HST photometric catalog of Bellini et al. (2017a).

<sup>1</sup> <https://github.com/thusser/spexsy>



**Fig. 2.** Normalised metallicity distribution for the stars with  $S/N > 20$  belonging to the rMS (solid line, 920 stars) and the bMS (dashed line, 737 stars). The mean  $[M/H]$  value of each distribution is indicated in the legend. The horizontal bars on the top right corner show the median value ( $\pm 0.1$  dex) of the corrected  $[M/H]$  uncertainties.

**Table 1.** Properties of the rMS and bMS stars

MS	$S/N$	# of stars	Mean $[M/H]$	$\sigma$ $[M/H]$
rMS	$> 15$	1629	$-1.669 \pm 0.005$	$0.179 \pm 0.005$
rMS	$> 20$	920	$-1.672 \pm 0.007$	$0.172 \pm 0.006$
bMS	$> 15$	1358	$-1.573 \pm 0.006$	$0.202 \pm 0.005$
bMS	$> 20$	737	$-1.576 \pm 0.008$	$0.192 \pm 0.006$

We first used the  $F606W$ ,  $F438W - F814W$  CMD since it is sensitive to metallicity changes (or helium variations) and less to light-element abundance variations (Calamida et al. 2017). At this stage, we are interested in only selecting the two main groups along  $\omega$  Cen’s MS. We then included all stars in the magnitude range  $19.2 < F606W < 20.2$  along the blue and red MS, making sure that the two selected boxes were not overlapping in the  $F438W - F814W$  color. Stars on the MSa, lying on the reddest part of the MS in this CMD were left out of the selection (see Fig. 1). We selected stars in this magnitude range as a trade-off between obtaining a better separation of the bMS and rMS stellar populations and having reasonable  $S/N$  from the MUSE spectra. The two MS samples also have a very similar completeness level since they are spanning the same magnitude range. The selected bMS and rMS samples were also plotted on the  $F606W$ ,  $F336W - F438W$  and  $F606W$ ,  $F275W - F438W$  CMDs (these two colors are very sensitive to light-element abundance variations) to eliminate cross-contamination between the two groups. The two samples are shown in Fig. 1 as red dots (rMS) and blue dots (bMS) and include 17 646 and 16 967 stars, respectively.

## 4. Metallicity distributions of MS stars

### 4.1. The rMS and bMS

We matched the targets selected in Sect. 3 with our MUSE database to retrieve the combined spectra of the stars that are in the MUSE fields and fitted them as described in Sect. 2. We applied a few additional criteria to define our final sample of red and blue MS stars: we computed results for two different  $S/N$  cuts, 15 and 20, and we then removed stars with a membership


**Table 2.** Properties of MSs identified in Bellini et al. (2017b)

MS	$S/N$	# of stars	Mean $[M/H]$	$\sigma$ $[M/H]$
rMS	$> 20$	464	$-1.661 \pm 0.009$	$0.160 \pm 0.008$
rMS1	$> 15$	248	$-1.663 \pm 0.013$	$0.174 \pm 0.012$
rMS2	$> 15$	302	$-1.641 \pm 0.012$	$0.160 \pm 0.010$
rMS3	$> 15$	329	$-1.667 \pm 0.012$	$0.172 \pm 0.011$
bMS	$> 20$	341	$-1.557 \pm 0.011$	$0.186 \pm 0.009$
bMS1	$> 15$	313	$-1.663 \pm 0.013$	$0.187 \pm 0.011$
bMS2	$> 15$	169	$-1.517 \pm 0.016$	$0.170 \pm 0.014$
bMS3	$> 15$	231	$-1.413 \pm 0.012$	$0.148 \pm 0.011$
MSe	$> 20$	292	$-1.504 \pm 0.010$	$0.150 \pm 0.009$
MSd	$> 20$	78	$-1.228 \pm 0.017$	$0.122 \pm 0.014$
MSa	$> 20$	250	$-0.964 \pm 0.009$	$0.138 \pm 0.008$

probability below 50%. This is based on the combination of a given stars’ RV and  $[M/H]$  (see Kamann et al. 2016 for more details); in the case of  $\omega$  Cen, the high RV of the cluster ( $232 \text{ km s}^{-1}$ ) is the main discriminating factor. Finally, we eliminated stars for which the standard deviation of the  $\log g$  obtained from the different isochrones is larger than 0.1 dex; these stars have an inconsistent magnitude in one of the catalogs, giving them a different position in the CMDs. The last two criteria exclude only a small amount of stars. The number of stars left in our samples is shown in Table 1 and includes, in all cases, several hundreds.

The metallicity distributions of rMS and bMS stars with  $S/N > 20$  are illustrated in Fig. 2. The median  $S/N$  of these samples is 25. To estimate the average  $[M/H]$ , the dispersion, and their uncertainty, we used the Markov chain Monte Carlo (MCMC) ensemble sampler developed by Foreman-Mackey et al. (2013). We used a simple Gaussian model to reproduce the observed distributions and retrieve the most likely mean ( $\mu$ ) and standard deviation ( $\sigma$ ) values. The results are listed in Table 1 where the uncertainties indicate the 16<sup>th</sup> and 84<sup>th</sup> percentile of the posterior probability distributions (equivalent to  $1\sigma$ ). We find that the bMS stars are slightly more metal-rich than the rMS stars, the difference of  $+0.096 \pm 0.011$  dex has a significance of  $8\sigma$ . This is at the lower limit of the  $+0.3 \pm 0.2$  dex range obtained by PI05. Our result, based on a statistically significant sample of MS stars ( $>1000$  vs 34), clearly rules out a 0.3 dex difference between  $\omega$  Cen’s bMS and rMS. This value, and the consequence of bluer MS stars being highly helium-enhanced compared to redder ones, were assumed to interpret the CMDs and the origin of multiple populations in  $\omega$  Cen and many other GGCs in subsequent years (e.g., Milone et al. 2017). However, it is worth pointing out that the two sequences used in PI05 are likely not equivalent to ours, nor to BE17’s, because PI05 have a ratio of bMS to rMS stars  $\approx 1:3$  while our numbers, and those of BE17, are closer to a 1:1 ratio. We also find that the average metallicity of both MSs is not significantly affected by the  $S/N$  cut. Because the MCMC procedure takes into account the calibrated  $[M/H]$  uncertainties, the value of  $\sigma$  is representative of the intrinsic dispersion of metallicities within each population. The values obtained for our samples suggests the presence of a small metallicity spread and it is in fact plausible that the two populations are not strictly mono-metallic. After all, Villanova et al. (2014) found that it was not the case for the multiple SGBs of the cluster and Tailo et al. (2016) reproduced, via stellar population synthesis, the bMS and rMS by using a mixture of metal-poor and metal-intermediate stars having different helium content.





**Fig. 3.** Normalised metallicity distributions for the stars with  $S/N > 20$  belonging to the five MSs defined by BE17. The mean  $[M/H]$  value (dotted line) of the distributions and the number of stars included are indicated in each panel. The horizontal bars on top right corner show the median value of the corrected  $[M/H]$  uncertainties.

#### 4.2. The five main sequences of Bellini et al.

In addition to the identification of 15 subpopulations across the MS of  $\omega$  Cen, BE17 made qualitative predictions about the chemical composition (Fe, He, and N) of these populations based on their photometric properties. Since the population tags are available, we compared our metallicities with their predictions of iron abundances. We selected our spectroscopic samples for the five main components (MSa, bMS, rMS, MSd and MSe) following the same criteria as in the previous subsection. We list their properties in Table 2, and show their metallicity distributions in Fig. 3. In order of increasing average metallicity we find the following sequence : rMS - bMS - MSe - MSd - MSa. We obtain a difference of  $+0.104 \pm 0.014$  dex in metallicity between the rMS and bMS, in excellent agreement with the value obtained from our own selection. The metallicity progression between the five MSs is in line with the expectations of BE17. The authors attributed a similar Fe enrichment for the bMS and the MSe stars but we find a  $3\sigma$  evidence for the MSe stars to be more metal-rich than the bMS stars. These two populations are the ones with the closest mean  $[M/H]$ . The dispersion ( $\sigma$ ) obtained for each population suggests the presence of a small ( $<0.2$  dex) intrinsic spread in metallicity.

Finally, we investigate on whether these main populations are mono-metallic or not. Each of them is subdivided into two

to four subpopulations in BE17 depending on the distribution of their stars in the chromosome map (see App. C). We focus here on the divisions of the rMS and bMS while a discussion on the metallicities of all MSs is presented App. C. We used spectra with  $S/N > 15$  in order to increase the number of stars in each sample. Results are listed in Table 2 and the distributions are shown in Fig. C.1. We find a different behavior between the rMS and bMS subpopulations: while the three rMSs have very similar metallicity distributions and average  $[M/H]$ , the three bMSs clearly have different metallicities. The metallicity increases from bMS1 to bMS3 with these two subpopulations having a difference of  $0.25 \pm 0.018$  dex. In fact, the bMS1, with a mean  $[M/H] = -1.66$ , is as metal-poor as the rMS.


#### 4.3. Comparison with spectroscopic studies in literature

We compared our metallicity distribution for a selected sample of MS stars with the most recent high-resolution spectroscopy for  $\omega$  Cen RGB stars from Johnson et al. (2020, hereinafter JO20). We also added spectra for 855 RGBs presented in Johnson & Pilachowski (2010, hereinafter JO10), since the study of JO20 is biased towards metal-poor stars. We combined the two samples and obtained a list of 1250 RGBs with a  $[Fe/H]$  measurement. We compared MUSE metallicities,  $[M/H]$ , directly to JO20+JO10  $[Fe/H]$  values since the synthetic spectra used for the spectral fits are  $\alpha$ -enhanced ( $[\alpha/Fe] = 0.3$ ).

Fig. 4 shows the normalized MUSE metallicity distribution for 2899 MS stars in the magnitude range  $19.2 \leq F606W \leq 20.2$  compared to the normalized and combined JO20+JO10 metallicity distribution for 1250 RGB stars. The shapes of the distributions are in good agreement: they cover the same wide metallicity range from  $[Fe/H] \approx -2.5$  up to  $\approx -0.5$  with a fast rise at low metallicity and a tail towards higher values. This qualitative comparison shows that a main peak is present at  $[Fe/H] \approx -1.75$  and a secondary one at  $\approx -1.5$  for JO20+JO10. Two such peaks are also present in the MUSE sample, albeit at slightly more metal rich positions;  $[M/H] \approx -1.6$  and  $\approx -1.4$ . Both distributions also show a shoulder at metallicities lower than the main peak. The most metal-rich stars,  $[Fe/H] \geq -1.0$ , in the MUSE distribution are  $\approx 3\%$  of the sample, in good agreement with previous photometric and spectroscopic studies that identified the most metal-rich population of  $\omega$  Cen, the RGBa, corresponding to the MSa, to include  $\approx 3\%$  of the cluster stars (Pancino et al. 2003; Castellani et al. 2007; Bellini et al. 2017b). The systematic difference in metallicity of  $\approx 0.15$  dex between the two distributions is not unexpected. Systematic offsets with the iron abundances of JO10 have been reported in the past for RGB stars ( $-0.13$  dex by Marino et al. 2011 and  $+0.1$  dex by Mészáros et al. 2021). These differences were partly explained by the use of different  $T_{\text{eff}}$ . In our case, it must be kept in mind that we are looking at different stars (MS vs RGB) and are using a different technique to derive  $T_{\text{eff}}$  and  $[M/H]$ . In addition, knowing that the Mg abundances are varying by  $\approx 1$  dex in the cluster’s RGB stars (Mészáros et al. 2021), and assuming some variation in the MS stars as well, we can expect an additional broadening of our MS  $[M/H]$  distribution when compared to the RGB  $[Fe/H]$  one.

#### 4.4. Comparison with models

On the basis of the new spectroscopic metallicities derived for the bMS and rMS, a set of BASTI isochrones was used to reproduce  $\omega$  Cen MS. In particular, we adopted two  $\alpha$ -enhanced isochrones with the same age,  $t = 13$  Gyr, same global metallic-



**Fig. 4.** Comparison between the metallicities obtained from the MUSE spectra of MS stars and the  $[Fe/H]$  measured in RGB stars (Johnson & Pilachowski 2010; Johnson et al. 2020)

ity,  $Z=0.0006$  ( $[M/H] \approx -1.5$ ), and different helium abundances, canonical,  $Y = 0.246$  and  $Y \approx 0.32$  (Pietrinferni et al. 2006). A distance modulus of  $DM_0 = 13.72$  (Braga et al. 2016) and reddening  $E(B - V) = 0.11$  (Calamida et al. 2005) were assumed. This reddening value was converted into extinctions in WFC3/HST filters by using the Cardelli et al. (1989) reddening law and the available WFC3 filter throughputs<sup>2</sup>. The two isochrones are overplotted to the CMD in Fig. 1: the canonical helium isochrone (red line) fits very well the rMS of  $\omega$  Cen from the base of the RGB down to the lower MS, and the helium-enhanced one (blue) for the same metallicity and age nicely fits the bMS. We find that, within the uncertainties, the blue and red MS in  $\omega$  Cen can be reproduced by models with similar average metallicities and ages but with different helium contents, i.e.,  $\Delta Y \approx 0.08$ , which is significantly less than the  $\Delta Y = 0.12-0.15$  derived by PI05.

## 5. Conclusion

We use MUSE spectra of  $\approx 3000$  MS stars in  $\omega$  Cen to derive their global metallicity  $[M/H]$  and examine the metallicity distribution and average value of the different MSs that have been identified in the cluster. This is the first time since PI05 that metallicities of MS stars in  $\omega$  Cen are obtained from spectroscopic data. We find the bMS stars to be, on average, slightly more metal-rich than the rMS stars, with a difference of  $+0.10 \pm 0.015$  dex. This is the lower limit of the range obtained by PI05. However, when looking at the bMS and rMS subdivisions identified by BE17, we discover a more complex behaviour: the three rMS subpopulations share a similar  $[M/H]$  while the three bMS subpopulations have metallicities that differ by  $\approx 0.1$  dex each, with the bluest sequence, bMS1, having the same metallicity as the rMS. We can reproduce the position of rMS and bMS stars in the CMD with isochrones of the same age and metallicity ( $Z=0.0006$ ) and a helium difference  $\approx 0.08$ , a smaller he-

lium enhancement than claimed until now for  $\omega$  Cen bMS ( $\Delta Y \approx 0.12-0.15$ , Piotto et al. 2005; Norris 2004; King et al. 2012). We derived average metallicities for the all MSs identified in BE17 and our results for the five main populations confirm the qualitative expectation for iron abundances made by the authors.

*Acknowledgements.* We thank G. Bono and N. Neumayer for useful comments and S. Cassisi, M. Salaris and A. Pietrinferni for providing us appropriate isochrones to fit the presented CMD. We acknowledge funding from the Deutsche Forschungsgemeinschaft (grant DR 281/35-1 and KA 4537/2-1) and from the German Ministry for Education and Science (BMBF Verbundforschung) through grants 05A14MGA, 05A17MGA, 05A14BAC, and 05A17BAA. SK gratefully acknowledges funding from UKRI in the form of a Future Leaders Fellowship (grant no. MR/T022868/1). JB acknowledges support by Fundação para a Ciência e a Tecnologia (FCT) through research grants UIDB/04434/2020 and UIDP/04434/2020 and work contract 2020.03379.CEECIND. This research has made use of NASA's Astrophysics Data System Bibliographic Services.

## References

- Alfaro-Cuello, M., Kacharov, N., Neumayer, N., et al. 2019, *ApJ*, 886, 57  
 Alonso, A., Arribas, S., & Martínez-Roger, C. 1999, *A&AS*, 140, 261  
 Amaro-Seoane, P., Konstantinidis, S., Brem, P., & Catelan, M. 2013, *MNRAS*, 435, 809  
 Anderson, J. 2002, in *Astronomical Society of the Pacific Conference Series*, Vol. 265, *Omega Centauri, A Unique Window into Astrophysics*, ed. F. van Leeuwen, J. D. Hughes, & G. Piotto, 87  
 Baumgardt, H. & Hilker, M. 2018, *MNRAS*, 478, 1520  
 Bedin, L. R., Piotto, G., Anderson, J., et al. 2004, *ApJ*, 605, L125  
 Bekki, K. & Norris, J. E. 2006, *ApJ*, 637, L109  
 Bekki, K. & Tsujimoto, T. 2016, *ApJ*, 831, 70  
 Bellini, A., Anderson, J., Bedin, L. R., et al. 2017a, *ApJ*, 842, 6  
 Bellini, A., Milone, A. P., Anderson, J., et al. 2017b, *ApJ*, 844, 164  
 Braga, V. F., Stetson, P. B., Bono, G., et al. 2016, *AJ*, 152, 170  
 Calamida, A., Bono, G., Stetson, P. B., et al. 2009, *ApJ*, 706, 1277  
 Calamida, A., Stetson, P. B., Bono, G., et al. 2005, *ApJ*, 634, L69  
 Calamida, A., Strampelli, G., Rest, A., et al. 2017, *AJ*, 153, 175  
 Calamida, A., Zocchi, A., Bono, G., et al. 2020, *ApJ*, 891, 167  
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245  
 Castellani, V., Calamida, A., Bono, G., et al. 2007, *ApJ*, 663, 1021  
 Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306  
 Gavagnin, E., Mapelli, M., & Lake, G. 2016, *MNRAS*, 461, 1276  
 Hilker, M., Kayser, A., Richtler, T., & Willemsen, P. 2004, *A&A*, 422, L9  
 Husser, T.-O., Kamann, S., Dreizler, S., et al. 2016, *A&A*, 588, A148  
 Husser, T.-O., Latour, M., Brinchmann, J., et al. 2020, *A&A*, 635, A114  
 Husser, T.-O., Wende-von Berg, S., Dreizler, S., et al. 2013, *A&A*, 553, A6  
 Ibata, R. A., Bellazzini, M., Malhan, K., Martin, N., & Bianchini, P. 2019, *Nature Astronomy*, 3, 667  
 Johnson, C. I., Dupree, A. K., Mateo, M., et al. 2020, *AJ*, 159, 254  
 Johnson, C. I. & Pilachowski, C. A. 2010, *ApJ*, 722, 1373  
 Kamann, S., Husser, T. O., Brinchmann, J., et al. 2016, *A&A*, 588, A149  
 Kamann, S., Husser, T. O., Dreizler, S., et al. 2018, *MNRAS*, 473, 5591  
 Kamann, S., Wisotzki, L., & Roth, M. M. 2013, *A&A*, 549, A71  
 Kayser, A., Hilker, M., Richtler, T., & Willemsen, P. G. 2006, *A&A*, 458, 777  
 King, I. R., Bedin, L. R., Cassisi, S., et al. 2012, *AJ*, 144, 5  
 Lee, Y.-W., Joo, J.-M., Sohn, Y.-J., et al. 1999, *Nature*, 402, 55  
 Marigo, P., Girardi, L., Bressan, A., et al. 2017, *ApJ*, 835, 77  
 Marino, A. F., Milone, A. P., Piotto, G., et al. 2011, *ApJ*, 731, 64  
 Massari, D., Koppelman, H. H., & Helmi, A. 2019, *A&A*, 630, L4  
 Mészáros, S., Masseron, T., Fernández-Trincado, J. G., et al. 2021, *MNRAS*, 505, 1645  
 Milone, A. P., Marino, A. F., Bedin, L. R., et al. 2017, *MNRAS*, 469, 800  
 Norris, J. E. 2004, *ApJ*, 612, L25  
 Norris, J. E. & Da Costa, G. S. 1995, *ApJ*, 447, 680  
 Norris, J. E., Freeman, K. C., Mayor, M., & Seitzer, P. 1997, *ApJ*, 487, L187  
 Pancino, E., Ferraro, F. R., Bellazzini, M., Piotto, G., & Zoccali, M. 2000, *ApJ*, 534, L83  
 Pancino, E., Mucciarelli, A., Sbordone, L., et al. 2011, *A&A*, 527, A18  
 Pancino, E., Seleznev, A., Ferraro, F. R., Bellazzini, M., & Piotto, G. 2003, *MNRAS*, 345, 683  
 Pasquato, M. & Chung, C. 2016, *A&A*, 589, A95  
 Pfeffer, J., Lardo, C., Bastian, N., Saracino, S., & Kamann, S. 2021, *MNRAS*, 500, 2514  
 Pietrinferni, A., Cassisi, S., Salaris, M., & Castelli, F. 2006, *ApJ*, 642, 797  
 Piotto, G., Villanova, S., Bedin, L. R., et al. 2005, *ApJ*, 621, 777

<sup>2</sup> <https://www.stsci.edu/hst/instrumentation/wfc3/performance/throughputs>

- Sarajedini, A., Bedin, L. R., Chaboyer, B., et al. 2007, *AJ*, 133, 1658  
Sollima, A., Ferraro, F. R., & Bellazzini, M. 2007, *MNRAS*, 381, 1575  
Sollima, A., Pancino, E., Ferraro, F. R., et al. 2005, *ApJ*, 634, 332  
Stanford, L. M., Da Costa, G. S., Norris, J. E., & Cannon, R. D. 2006, *ApJ*, 647, 1075  
Suntzeff, N. B. & Kraft, R. P. 1996, *AJ*, 111, 1913  
Tailo, M., Di Criscienzo, M., D'Antona, F., Caloi, V., & Ventura, P. 2016, *MNRAS*, 457, 4525  
Villanova, S., Geisler, D., Gratton, R. G., & Cassisi, S. 2014, *ApJ*, 791, 107  
Villanova, S., Piotto, G., King, I. R., et al. 2007, *ApJ*, 663, 296  
Weilbacher, P. M., Palsa, R., Streicher, O., et al. 2020, *A&A*, 641, A28

## Appendix A: MUSE observations

We used observations from the seven most central fields as these regions are included in the spatial coverage of the Bellini et al. (2017a) photometric catalog. A mosaic of these fields is shown in Fig. 1 of Kamann et al. (2018) and on our project website<sup>3</sup>. The observations from 2018 and 2019 benefited from the use of the adaptive optic system installed on UT4 of the VLT. The data reduction is done with the standard MUSE pipeline (Weilbacher et al. 2020) and the spectral extraction is performed with the PAMPELMUSE software (Kamann et al. 2013). A summary of the observations used for this work is presented in Table A.1.

**Table A.1.** MUSE observations of  $\omega$  Cen


Field	RA	DEC	# Epochs		Total exp. time (s)
			non-AO	AO	
1	13:26:45.0	-47:29:09	8	5	1755
2	13:26:45.0	-47:28:24	7	5	1620
3	13:26:49.5	-47:29:09	7	6	1710
4	13:26:49.5	-47:28:24	7	6	1755
5	13:26:40.6	-47:28:31	7	6	3040
6	13:26:53.1	-47:29:01	7	6	3120
7	13:26:36.8	-47:27:54	6	6	3600

## Appendix B: Additional atmospheric properties of the MS stars

The adopted effective temperatures of the stars affect the resulting metallicities. It is stated as the main source of uncertainties on the metallicity difference between the rMS and bMS in PI05 as well as the source of the systematic offset between the  $[Fe/H]$  obtained by Marino et al. (2011) and previous investigations of RGB stars. As opposed to many spectroscopic investigations of stars in GCs, including that of PI05, our temperatures are not based on the photometric properties of the stars. We used the  $T_{\text{eff}}$  obtained from the isochrones as starting value for the spectral fitting, but the final temperature is obtained simultaneously with the  $[M/H]$  from the spectra themselves. The Balmer lines  $H_{\beta}$  and  $H_{\alpha}$  are the main temperature indicators while the Mg  $b$  and Ca triplet lines are the main metallicity indicators. The median errors on  $[M/H]$  and  $T_{\text{eff}}$ , once calibrated as explained in Sect. 2, are  $\pm 0.1$  dex and 70 K. The fact that the bMS stars are bluer than the rMS in the CMD is an indication that at a given magnitude they are also hotter than their rMS counterparts. We retrieve this general property from our fitting procedure as shown in the top panel of Fig. B.1. The median  $T_{\text{eff}}$  for the bMS and rMS stars are 5980 K and 5860 K respectively. We note here that our MS stars are significantly hotter than the stars analyzed by PI05 who adopted  $T_{\text{eff}}$  of 5400 K and 5200 K for the bMS and rMS respectively. This difference is mainly due to the fact that our stars are  $\approx 0.7$  magnitudes brighter than the PI05 sample, thus they are intrinsically hotter. Indeed, our isochrones have differences of  $\approx 500$  K between an  $F606W$  magnitude of 19.7 and 20.5. In the bottom panel of Fig. B.1 we plotted the rMS and bMS stars (in red and blue respectively) in the  $T_{\text{eff}}-[M/H]$  plane and the distribution of stars in this diagram does not suggest the presence of a correlation between the two parameters.

The atmospheric parameters ( $T_{\text{eff}}$ ,  $\log g$ ,  $[M/H]$ ) and coordinates of the stars included in our rMS, bMS and in the BE17

<sup>3</sup> <https://musegc.uni-goettingen.de/index.php/targets?name=musegc5139>



**Fig. B.1.** Top– Effective temperature distribution for the rMS and bMS  $S/N > 20$  samples. As expected, the bMS stars are on average hotter than the rMS stars. Bottom– Distribution of the rMS and bMS stars in the  $T_{\text{eff}}-[M/H]$  plane.

sequences are available in Table B.1 which is only available in electronic form at CDS and on our project homepage<sup>4</sup>

## Appendix C: Metallicities of the 15 MSs of Bellini et al. 2017

For the sake of completeness, we present in Table C.1 the properties for the 15 MSs identified in BE17. Unfortunately, some of these subpopulations have very little stars left in our spectroscopic samples. However, the subpopulations of the bMS and rMS are well populated enough to clearly show a different behaviour in Fig. C.1. As discussed in Sect. 4.2, while the rMS subpopulations share the same metallicity, this is not the case for the bMS. It is interesting to note that the bMS and rMS stars also have a different behavior in their respective chromosome map; the bMS stars show populations aligned from the lower left to the upper right, while the rMS stars are aligned from the lower right to the upper left (see Fig.C.2). It seems possible that variations in iron shape, at least part of, the chromosome map of the bMS stars and light elements variations are responsible for the different disposition of the rMS stars. This is further supported

<sup>4</sup> <https://musegc.uni-goettingen.de/>





**Table C.1.** Properties of the 15 MSs identified in Bellini et al. (2017b)

MS	$S/N$	# of stars	Mean $[M/H]$	$\sigma$ $[M/H]$
rMS1	> 15	248	$-1.663 \pm 0.013$	$0.174 \pm 0.012$
rMS2	> 15	302	$-1.641 \pm 0.012$	$0.160 \pm 0.010$
rMS3	> 15	329	$-1.667 \pm 0.012$	$0.172 \pm 0.011$
bMS1	> 15	313	$-1.663 \pm 0.013$	$0.187 \pm 0.011$
bMS2	> 15	169	$-1.517 \pm 0.016$	$0.170 \pm 0.014$
bMS3	> 15	231	$-1.413 \pm 0.012$	$0.148 \pm 0.011$
MSe1	> 15	246	$-1.508 \pm 0.013$	$0.161 \pm 0.011$
MSe2	> 15	261	$-1.510 \pm 0.013$	$0.163 \pm 0.011$
MSe3	> 15	27	$-1.464 \pm 0.023$	$0.073 \pm 0.034$
MSe4	> 15	46	$-1.409 \pm 0.028$	$0.148 \pm 0.027$
MSe3+4	> 15	73	$-1.429 \pm 0.019$	$0.126 \pm 0.019$
MSd1	> 15	51	$-1.289 \pm 0.021$	$0.115 \pm 0.020$
MSd2	> 15	53	$-1.178 \pm 0.020$	$0.114 \pm 0.018$
MSd3	> 15	45	$-1.161 \pm 0.024$	$0.130 \pm 0.022$
MSa1	> 15	315	$-0.970 \pm 0.009$	$0.140 \pm 0.007$
MSa2	> 15	31	$-0.862 \pm 0.027$	$0.136 \pm 0.025$

when we look at the result for the MSe1 and MSe2, two subpopulations with enough stars to provide a spectroscopic sample of decent size (see Fig. C.1, bottom panel). The MSe1 and MSe2 stars have similar metallicities and they are the only other subpopulations that align in the same way as the rMS in the chromosome maps. Based on their photometric properties, BE17 expected the stars of MSe1 to have a similar iron content as the rMS, which is not what we observe. We find the MSe1 stars to be more metal-rich than the rMS by  $\approx 0.15$  dex.

There is one last behavior worth mentioning concerning the populations of the bMS, MSd and MSa. According to BE17, they share similar photometric properties, which is illustrated in their Fig. 15 where the subpopulations are ordered in terms of  $\Delta\text{color}$  ( $X - m_{F438W}$  where  $X$  is a given filter) in the bluest filters ( $\lambda < 336$  nm) with respect to the rMS1. There is a clear progression from bluer to redder as follows: bMS1, bMS2, bMS3, MSd1, MSd2, MSd3, MSa1, and MSa2. From our average metallicities, we also see a progression of increasing metallicity in the same order, besides for MSd2 and MSd3 which have the same average  $[M/H]$  but also a small spectroscopic sample size. This is consistent with the fact that the bluest filter ( $F225W$ ) is especially sensitive to Fe variations.


**Fig. C.1.** Metallicity distributions of the subpopulations of the rMS and bMS as identified by BE17. For the MSe, we plotted the distributions for the two most populous groups: MSe1 and MSe2.



**Fig. C.2.** Chromosome maps of the five MSs. The subpopulations of each MS are plotted in different colors. The axis are the two rectified and parallelized pseudo-colors used for the construction of the chromosome maps in BE17.