Abstract ID: 95 for KRD 2020

## STEP BY STEP PREPARATION OF URINE SAMPLES FOR IN-SOLUTION, GEL-FREE PROTEOMIC STUDIES, SUITABLE FOR MALDI TOF MS AND LCMS QTOF MS

Aniza Pakeer<sup>1\*</sup>, M. Yusri Idorus<sup>2</sup>, Nosrihah Ismail<sup>3</sup>, Vanitha Mariappan<sup>4</sup>, Imad Mustafa al Deen<sup>1</sup>

<sup>1</sup>Dept of Basic Medical Sciences, Kulliyyah of Medicine, International Islamic University Malaysia, Campus Kuantan

<sup>2</sup>Institute for Medical Molecular Biotechnology, University Teknologi Mara Sg. Buloh

<sup>3</sup>Medical Biotechnology Laboratory, Faculty of Medicine, University Malaya Kuala Lumpur.

<sup>4</sup>Corresponding Author's email Address: <u>anizapo@iium.edu.my</u>

## **ABSTRACT**

Two-dimensional electrophoresis is an established method for isolating and separating proteins for proteomic analysis and identification of proteins. However, it requires a lot of optimization and it suffers from some ongoing concerns regarding quantitative reproducibility and limitations on its ability to study certain class of proteins. We propose a simpler method in preparing urine samples from acute melioidosis patients for in-solution proteomics using liquid chromatography coupled with mass spectrometry quadrupole time of flight (LCMS-QTOF MS) or matrix assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF MS). This method is relatively easier than 2-dimensional electrophoresis, affordable, reproducible, can be performed in less equipped laboratories for transport to specialized centers for proteomic studies and bioinformatic analysis.

Keywords: proteomics, in-solution, LCMS, urine, melioidosis, proteome