

Search > Results > Results > Results > Results > Results > Excellence of Financial Re... > The impact of film thickne... > Macrolactin A as a Novel I... > Societal emotional environ... > Anomaly Detection in ICS D... > Response surface analysis ... > Evaluation of microplastic... > Energy, exergy and economi... > Toxicity profile of Phaler... > Results > MRI Evaluation of Anterola... > MRI Evaluation of Anterola...

Free Full Text From Publisher

Full Text Links

Export

Add To Marked List

< 21 of 372 >

MRI Evaluation of Anterolateral Ligament of the Knee: A Cross-Sectional Study in Malaysia

By: [Kow, RY](#) (Kow, Ren Yi)¹; [Low, CL](#) (Low, Chooi Leng)²; [Siron-Baharom, KN](#) (Siron-Baharom, Khairul Nizam)¹; [Said, SNBS](#) (Said, Siti Nor Badriati Sheikh)³
View Web of Science ResearcherID and ORCID (provided by Clarivate)

CUREUS

Volume: 13 Issue: 6

Article Number: e15758

DOI: 10.7759/cureus.15758

Published: JUN 19 2021

Document Type: Article

Abstract

Introduction

After detailed anatomical delineation of the anterolateral ligament (ALL) of the knee, there is a surge in research on this anatomical structure. Owing to the anatomical variation and lack of experience in the identification of this structure, magnetic resonance (MR) evaluation of the ALL produces mixed results. It was aimed to evaluate the ALL using the routinely performed MR imaging of the knee and to determine any associated factors with ALL injuries.

Materials and methods

Thirty-six MR images of the knee from 31 patients from January 1, 2017, to June 30, 2017, are evaluated. MR sequences performed include T1-weighted, T2-weighted, proton density (PD), and PD fat saturation (FS). All MR images were double-read by two authors and approved by a consultant radiologist with more than 20 years of radiological experience. The ALL was divided into three portions: femoral, meniscal, and tibial, and the ALL was considered fully visualized when all three portions were seen on MR images.

Results

At least a portion of the ALL was visualized in 27 scans (75%), and it was fully visualized in 20 scans (55.6%). The femoral portion was the most commonly identified (75%), followed by the meniscal portion (69.4%) and the tibial portion (58.3%). ALLs were best visualized on coronal view in PD FS with the lateral inferior genicular artery as a guide to locate the bifurcation of the meniscal and tibial components.

Conclusion

The ALL can be visualized in routine 1.5-T MR imaging, either full delineation (55.6%) or partially visualized (75%). It is best characterized via a PD-weighted sequence with fat saturation on the coronal plane. The ALL injury was associated with an anterior cruciate ligament (ACL) injury.

Keywords

Author Keywords: anterolateral ligament; knee; anterior cruciate ligament; magnetic resonance imaging; injury

Keywords Plus: ANTERIOR CRUCIATE LIGAMENT; ANATOMY; ABNORMALITIES; INJURIES; TEARS; BAND

Author Information

Corresponding Address: Kow, Ren Yi (corresponding author)

▲ Int Islamic Univ Malaysia, Dept Orthopaed Traumatol & Rehabil, Kuantan, Malaysia

Affiliation

International Islamic University Malaysia

Addresses:

▲ ¹ Int Islamic Univ Malaysia, Dept Orthopaed Traumatol & Rehabil, Kuantan, Malaysia

Affiliation

International Islamic University Malaysia

▲ ² Int Islamic Univ Malaysia, Dept Radiol, Kuantan, Malaysia

Affiliation

International Islamic University Malaysia

Citation Network

In Web of Science Core Collection

0

Citations

🔔 Create citation alert

Cited References

29

View Related Records

You may also like...

Astur, DC; Santos, CV; Cohen, M; et al.

Characterization of Cruciate Ligament Impingement: The Influence of Femoral or Tibial Tunnel Positioning at Different Degrees of Knee Flexion

ARTHROSCOPY-THE JOURNAL OF ARTHROSCOPIC AND RELATED SURGERY

Angelini, FJ; Albuquerque, RFM; Hernandez, AJ; et al.

COMPARATIVE STUDY ON ANTERIOR CRUCIATE LIGAMENT RECONSTRUCTION: DETERMINATION OF ISOMETRIC POINTS WITH AND WITHOUT NAVIGATION CLINICS

Osti, M; Krawinkel, A; Benedetto, KP; et al.

Femoral and Tibial Graft Tunnel Parameters After Transtibial, Anteromedial Portal, and Outside-In Single-Bundle Anterior Cruciate Ligament Reconstruction AMERICAN JOURNAL OF SPORTS MEDICINE

Zantop, T; Petersen, W;

Arthroscopic filling of misplaced and wide bone tunnels after reconstruction of the anterior cruciate ligament with bone graft in patients with recurrent instability OPERATIVE ORTHOPADIE UND TRAUMATOLOGIE

Lee, YS; Lee, SW; Lee, BK; et al.

Analysis of tunnel widening after double-bundle ACL reconstruction KNEE SURGERY SPORTS TRAUMATOLOGY ARTHROSCOPY

See all

Use in Web of Science

Web of Science Usage Count

19



³ Hosp Tengku Ampuan Afzan, Dept Radiol, Kuantan, Malaysia

E-mail Addresses: renyi_kow@hotmail.com

Categories/Classification

Research Areas: General & Internal Medicine

+ See more data fields

0

Last 180 Days

[Learn more](#)

0

Since 2013

This record is from:
Web of Science Core Collection

Emerging Sources Citation Index (ESCI)

Journal information

CUREUS

eISSN: 2168-8184

Current Publisher: CUREUS INC, PO BOX 61002, PALO ALTO, CA 94306 USA

Research Areas: General & Internal Medicine

Web of Science Categories: Medicine, General & Internal

29 Cited References

Showing 29 of 29

[View as set of results](#)

(from Web of Science Core Collection)

1 [Pathogenesis of the segond fracture: Anatomic and MR imaging evidence of an iliotibial tract or anterior oblique band avulsion](#)

[Campos, JC](#); [Chung, CB](#); (...); [Resnick, D](#)

May 2001 | Radiology

[Full Text at Publisher](#) ...

147

Citations

26

References

[Related records](#)