

RESEARCH

Open Access



Coenzyme Q10 suppresses apoptosis of mouse pancreatic β -cell line MIN6

Keisuke Sumi^{1†}, Tsuyoshi Okura^{1*†}, Youhei Fujioka¹, Masahiko Kato², Takeshi Imamura³, Shin-ichi Taniguchi⁴ and Kazuhiro Yamamoto²

Abstract

Background: In mitochondrial diabetes, apoptosis of β -cells caused by mitochondrial stress plays an important role in impaired insulin secretion. Several studies have reported that coenzyme Q10 (CoQ10) has therapeutic effects on mitochondrial diabetes, but no reports have examined the fundamental effectiveness or mechanism of CoQ10 in mitochondrial diabetes. We previously reported in a Japanese article that CoQ10 has protective effects on pancreatic β -cells against mitochondrial stress using mouse pancreatic β -cell line MIN6 and staurosporine (STS). Here, we report that CoQ10 protects MIN6 cells against apoptosis caused by STS and describe the more detailed apoptotic cascade.

Methods: Apoptosis of MIN6 cells was induced by 0.5 μ M STS treatment for specific periods with or without 30 μ M CoQ10. The apoptosis cascade in MIN6 cells was then investigated using WST-8 assays, annexin-V staining, western blotting, and DNA degradation analysis.

Results: Sixteen hours of 0.5 μ M STS treatment led to 47% cell viability, but pretreatment with 30 μ M CoQ10 resulted in significantly higher viability of 76% ($P < 0.01$). CoQ10 also prevented translocation of phosphatidylserine from the inner leaflet to the outer leaflet of the cell membrane. CoQ10 prevented cytochrome c release from mitochondria and activation of caspase-3.

Conclusion: We concluded that CoQ10 protects pancreatic β -cells through anti-apoptotic effects against STS treatment.

Keywords: MIN6, Coenzyme Q10, Staurosporine, Apoptosis, Mitochondrial diabetes

Background

Hyperglycemia occurs because of impaired insulin action in diabetes as a result of insulin resistance and impaired insulin secretion. In insulin resistance, although insulin is secreted, its effect decreases. Conversely, in impaired insulin secretion, insulin secretion itself decreases. Mitochondria are important for insulin secretion of pancreatic β -cells. Impaired insulin secretion occurs in mitochondrial diabetes, a hereditary disorder of mitochondrial functions [1]. Mitochondrial diabetes accounts for 1% of diabetes in Japan. Although insulin secretion depends on the function

and quantity of pancreatic β -cells, these cells have a low proliferation ability, and their quantitative and qualitative decreases are strongly affected by apoptosis [2]. Apoptosis of β -cells is caused by oxidative stress, lipotoxicity, and cytokines. Furthermore, it has been reported that pancreas β -cells decrease as a result of apoptosis in BHE/cdb rats, a model of mitochondrial diabetes [3]. These results suggest that mitochondrial dysfunction and apoptosis have an important role in impaired insulin secretion. Other than insulin injection, treatments have not been established for mitochondrial diabetes, although several studies have shown that coenzyme Q10 (CoQ10) has therapeutic effects on mitochondrial diabetes [4]. However, to the best of our knowledge, no reports have examined the fundamental effectiveness or mechanism of CoQ10 in mitochondrial diabetes. Reactive oxygen species produced by mitochondria cause mitochondrial disorders [5]. A report

*Correspondence: ohkura@grape.med.tottori-u.ac.jp

[†]Keisuke Sumi and Tsuyoshi Okura contributed equally to this work

¹Division of Endocrinology and Metabolism, Molecular Medicine and Therapeutics, Faculty of Medicine, Tottori University, 36-1 Nishi-chou, Yonago, Tottori 683-8504, Japan

Full list of author information is available at the end of the article



has demonstrated the effectiveness of CoQ10 to treat a central symptom of mitochondrial disease [6]. The antioxidant and ATP supplying effects of CoQ10 may protect central nerves [7]. Therefore, the mechanism of CoQ10 treatment in mitochondrial diabetes might be protective effects from apoptosis of pancreatic β -cells. We previously reported in a Japanese article that CoQ10 has protective effects on pancreatic β -cells against mitochondrial stress using mouse pancreatic β -cell line MIN6 and staurosporine (STS), a strong inducer of apoptosis and mitochondrial stress by producing reactive oxygen species [8, 9]. Here, we report that CoQ10 protects MIN6 cells against apoptosis caused by STS and describe the apoptotic cascade.

Methods

Materials

Mouse insulinoma cell line MIN6 was kind gift from Dr. Makoto Shigeto (Division of Diabetes and Endocrinology Department of Medicine Kawasaki Medical School), Kohei Kaku (Division of Diabetes and Endocrinology Department of Internal Medicine Kawasaki Medical School), and Jun-ichi Miyazaki (Division of Stem Cell Regulation Research, Osaka University Graduate School of Medicine). Fetal bovine serum (FBS) was obtained from Thermo Fisher Scientific, Inc. (Kanagawa, Japan). Phosphate buffered saline (PBS), Dulbecco's modified Eagle's medium (DMEM), penicillin, and streptomycin were obtained from Sigma-Aldrich, Inc. (St. Louis, MO). CoQ10 (oxidized form) was obtained from Yokohama Oils & Fats Industry (Kanagawa, Japan). Z-VAD-FMK (carbobenzoxy-valyl-alanyl-aspartyl-[O-methyl]-fluoromethylketone) was obtained from Peptide Institute, Inc. (Osaka, Japan). STS and polyacrylamide gels were obtained from Wako Pure Chemical Industries (Osaka, Japan). A protease inhibitor cocktail was obtained from BioVision, Inc. (Milpitas, CA). Tris/Glycine/SDS electrophoresis buffer was obtained from Bio-Rad Laboratories, Inc. (Hercules, CA). Western Blotting Detection Reagent was obtained from GE Healthcare (Tokyo, Japan). Polyvinylidene fluoride (PVDF) microporous membranes were obtained from Merck Millipore (Billerica, MA). Cell lysis buffer, an anti- β -actin antibody, anti-rabbit IgG, and annexin V assay kit were obtained from MBLCo., Ltd. (Aichi, Japan). An anti-caspase-3 antibody was obtained from Cell Signaling Technology (Tokyo, Japan). A cytochrome c releasing apoptosis assay kit was obtained from Funakoshi Co., Ltd. (Tokyo, Japan). An apoptotic DNA ladder kit was obtained from Roche Diagnostics (Tokyo, Japan). Cell Counting Kit-8 was obtained from Dojindo Laboratories (Kumamoto, Japan).

Cell culture

MIN6 cells were cultured in 60 ml flasks, 96-well plates, or on glass coverslips in DMEM (high-glucose; 4.5 g/l) with

15% FBS, 75 μ g/ml penicillin, and 50 μ g/ml streptomycin at 37 °C in a humidified atmosphere with 5% CO₂. At 80% confluence, the cells were divided into four groups: CoQ10, Z-VAD, STS, and control. The CoQ10 group was treated with 30 μ M CoQ10 for 4 h before induction of apoptosis. Z-VAD-FMK is a pan-caspase inhibitor and anti-apoptotic agent. The Z-VAD group was treated with 30 μ M Z-VAD-FMK for 1 h before induction of apoptosis. STS and control groups were not pre-treated. Except for the control group, each group was treated with 0.5 μ M STS for the specified periods to induce apoptosis.

WST-8 assay

Cell viability was compared among the four groups after apoptotic stimulation using Cell Counting Kit-8 that determines cell viability through reduction of a water-soluble tetrazolium salt, WST-8. WST-8 is reduced by dehydrogenases in cells to produce a yellow formazan dye. The amount of formazan dye generated by the activities of dehydrogenases in cells is directly proportional to the number of living cells. Except for the control group, MIN6 cells were treated in 96-well plates (100 μ l medium per well) with 0.5 μ M STS for 16 h. Then, 10 μ l Cell Counting Kit-8 solution was added to each well, followed by incubation for 2 h. The number of living cells was determined by absorbance at 450 nm using fluorescence microplate reader. Several experiments were performed, and average values were calculated.

Annexin-V staining

At the early stage of apoptosis, cells lose their phospholipid membrane asymmetry and expose phosphatidylserine (PS) at the cell surface. This process can be monitored using annexin-V, a Ca²⁺-dependent, phospholipid-binding protein with high affinity for PS, which is useful for identifying apoptotic cells with exposed PS. For annexin-V staining, MIN6 cells were cultured on glass coverslips. After 6 h of treatment with STS, cells were washed with PBS and then stained with green fluorescent protein-labeled annexin-V for 5 min while protected from light, according to the manufacturer's protocol. The cells were then washed with PBS and treated with methanol for 10 min. The methanol was removed, and propidium iodide (PI) was added, a red fluorescent nuclear counterstain. Samples were observed under a fluorescence microscope. Methanol-treated samples (all cells) were stained with PI (red), whereas early apoptotic cells were stained with annexin-V (green) and PI. We counted respectively 140, 113, 208 and 235 cells in Control, STS, CoQ10 and Z-VAD group. Thus, we were able to calculate the ratio of early apoptotic cells.

SDS-polyacrylamide gel electrophoresis (PAGE) and western blot analysis



To detect activation of caspase-3 and cytochrome c release from mitochondria after treatment with STS, cells were washed and then lysed in cell lysis buffer containing a protease inhibitor cocktail. The cell lysates (10 µg protein per sample) were then separated by 12.5% SDS-PAGE and electrotransferred onto a PVDF transfer membrane. Membranes were blocked with Tris-buffered saline containing with 0.05% Tween 20 and 5% nonfat dried milk and then incubated with primary antibodies for 1 h at room temperature. Primary antibodies were against caspase-3 (1:1000 dilution), cytochrome c (1:200 dilution), and β-actin (1:1000 dilution). The membrane was then incubated with secondary antibodies for appropriate times (30–60 min). Specific proteins were visualized by the enhanced chemiluminescence method using western blotting detection reagent.

DNA degradation analysis

After 14 h of treatment with STS, isolation of DNA from cells of each group was performed according to a standard procedure using a commercially available apoptotic DNA ladder kit. DNA fragmentation analysis was performed using a 1.5% agarose gel.

Statistical analysis

Data are expressed as the mean ± standard deviation. Comparisons of individual treatments in WST-8 assays were conducted using the Games-Howell post hoc test following one-way analysis of variance. Comparisons of individual treatments in annexin-V analysis were



conducted using the Chi square test. $P < 0.05$ was considered as statistically significant.


Results

Comparing of cell viability

We first examined cell viability after STS treatment using WST-8 assays (Fig. 1). After 16 h of STS treatment, the STS group had 47% cell viability, but the CoQ10 group had significantly higher viability of 76% ($P < 0.01$ vs. STS group). The Z-VAD group had 60% cell viability ($P < 0.01$ vs. STS group). These results indicate that CoQ10 has a protective effect on MIN-6 cells against STS treatment, and CoQ10 has the same or a stronger protective effect than Z-VAD-FMK.

DNA fragmentation

Next, we examined DNA fragmentation to determine whether cell death was caused by apoptosis (Fig. 2). Electrophoresis of genomic DNA from MIN6 cells treated with STS for 14 h revealed the characteristic laddering pattern, indicating that MIN6 cell death was caused by apoptosis. However, DNA fragmentation was inhibited by CoQ10 and Z-VAD, suggesting that CoQ10 had a protective effect against apoptosis.




Annexin-V analysis

Next, we investigated the detailed apoptosis cascade. During apoptosis, characteristic biochemical changes are observed. Translocation of PS from the inner leaflet to the outer leaflet of the cell membrane is one of the characteristic biochemical changes in the early stage of apoptosis. We investigated exposure of PS using annexin-V staining together with PI staining. After 6 h of STS treatment, 15% of cells in the STS group were positive for annexin-V. Conversely, only 1% of cells in the CoQ10 group and 3% of cells in the Z-VAD group were positive for annexin-V (Fig. 3).

Release of cytochrome c

Cytochrome c release from mitochondria is a central event in apoptotic signaling. Cytochrome c usually exists between mitochondrial inner and outer membranes, but it is released to the cytoplasm by apoptotic stimulation. Then, cytochrome c together with Apaf-1 activate caspase-9, leading to the apoptotic caspase cascade. We investigated the release of cytochrome c from mitochondria by western blotting using a commercially available kit. After 14 h of STS treatment, cytochrome c release was increased in the STS group. However, cytochrome c release in the other groups was lower than that in the STS group (Fig. 4). These results indicated that cytochrome c was released into the cytoplasm from mitochondria by STS treatment.




Caspase-3

We also investigated activation of caspase-3 by western blotting. After 12 h of STS treatment, cleaved caspase-3, which is the activated form of caspase-3, was strongly detected in the STS group. Conversely, activation of caspase-3 was inhibited in CoQ10 and Z-VAD groups (Fig. 5).

Discussion

In the pathophysiology of diabetes including mitochondrial diabetes, apoptosis of β -cells plays important roles in impaired insulin secretion. Although the detailed fundamental mechanism of CoQ10 in mitochondrial diabetes is unknown, therapeutic effects of CoQ10 on mitochondrial



diabetes have been reported in several studies [4]. We hypothesized that CoQ10 has protective effects against apoptosis in pancreatic β -cells. In this study, we induced apoptosis of MIN6 cells using STS, a mitochondrial stress agent [9]. In WST-8 assays, 0.5 μ M STS treatment for 16 h induced the death of about half of the cells. Although cell death might be caused by both apoptosis and necrosis, we showed translocation of PS from the inner leaflet to the outer leaflet of the MIN6 cell membrane, release of cytochrome c from mitochondria, activation of caspase-3, and DNA fragmentation of MIN6 cells. Therefore, MIN6 cell death induced by STS treatment was caused by apoptosis in this study. Based on these results, β -cells were protected by CoQ10 from apoptosis caused by STS treatment. In addition, this protective effect was equivalent to that of Z-VAD-FMK, an anti-apoptotic agent, suggesting that CoQ10 may be an effective anti-apoptotic agent. The fundamental effectiveness or mechanism of CoQ10 in mitochondrial diabetes have not been reported. However, the clinical effectiveness of CoQ10 against a central symptom of mitochondrial disease has been reported [6]. The mechanism is an antioxidant action and stimulates ATP supply from mitochondria [7]. In our study, CoQ10 had an anti-apoptotic action in β -cells. We believe that the protective mechanism of CoQ10 is a direct effect on mitochondria through ATP supply, but elucidation of the detailed mechanism is needed in the future. In mitochondrial diabetes, a hereditary disorder of mitochondrial functions causes impaired insulin secretion [1]. Conversely, in type-2 diabetes, mitochondrial dysfunction caused by various kinds of stress, such as oxidative stress, plays important roles in the pathological progression [10]. Hodgson et al. [11] reported that CoQ10 treatment improves blood pressure and glycaemic control in type-2 diabetes patients. This finding suggests that CoQ10 may have protective effects on pancreatic β -cells against mitochondrial stress in the clinic. In Japan,

CoQ10 as a ubiquinone is applicable to insurance for heart failure, but not diabetes. However, CoQ10 is already generally used as a supplement. Therefore, if evidence that CoQ10 has effective effects on mitochondrial or type-2 diabetes accumulates, CoQ10 may be adopted clinically.

There are limitations in our study. First, STS treatment is non-physiological stimulation. Further studies are needed to clarify whether CoQ10 has protective effects against apoptosis induced by physiological stimulation. Second, we only investigated MIN6 cells. Experiments using an animal model are necessary to examine the clinical effectiveness of CoQ10. However, we revealed a part of the underlying mechanism of CoQ10 in protection of pancreatic β -cells from apoptosis caused by mitochondrial stress. Elucidation of the more detailed mechanism may facilitate clinical application of CoQ10.

Conclusion

CoQ10 protects pancreatic β -cells through anti-apoptotic effects against STS treatment.

Abbreviations

MIN6: mouse insulinoma 6; CoQ10: coenzyme Q10; STS: staurosporine; ATP: adenosine triphosphate; FBS: fetal bovine serum; PBS: phosphate buffered saline; DMEM: Dulbecco's modified Eagle's medium; Z-VAD-FMK: carbobenzoxy-valyl-alanyl-aspartyl-(O-methyl)-fluoromethylketone; PVDF: polyvinylidene fluoride; DNA: deoxyribonucleic acid; WST-8: water soluble tetrazolium salts 8; PS: phosphatidylserine; PI: propidium iodide; SDS: sodium dodecyl sulfate; PAGE: polyacrylamide gel electrophoresis.

Authors' contributions

KS and TO conceived of the presented idea. KS, TO and YF performed the experiments. MK, TI, ST and KY supervised the findings of this work. All authors discussed the results and contributed to the final manuscript. All authors read and approved the final manuscript.

Author details

¹ Division of Endocrinology and Metabolism, Molecular Medicine and Therapeutics, Faculty of Medicine, Tottori University, 36-1 Nishi-chou, Yonago, Tottori 683-8504, Japan. ² Division of Cardiovascular Medicine, Endocrinology and Metabolism, Department of Molecular Medicine and Therapeutics, Faculty of Medicine, Tottori University, Yonago, Tottori 683-8504, Japan. ³ Division of Molecular Pharmacology, Faculty of Medicine, Tottori University, Yonago, Tottori 683-8504, Japan. ⁴ Department of Regional Medicine, Faculty of Medicine, Tottori University, Yonago, Tottori 683-8504, Japan.

Acknowledgements

We thank Prof. Masao Izawa for his excellent technical assistance (Department of Obstetrics and Gynecology, Tottori University Faculty of Medicine, Yonago, Japan). We thank M. Arico from Edanz Group (<http://www.edanzediting.com/ac>) for editing a draft of this manuscript.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Consent for publication

Not applicable.

Ethics approval and consent to participate

Not applicable.

Funding

No specific funding has been provided for this study.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 12 April 2018 Accepted: 5 June 2018

Published online: 14 June 2018

References

- Ballinger SW, Shoffner JM, Hedaya EV, Trounce I, Polak MA, Koontz DA, et al. Maternally transmitted diabetes and deafness associated with a 10.4 kb mitochondrial DNA deletion. *Nat Genet.* 1992;1:11–5.
- Butler AE, Janson J, Bonner-Weir S, Ritzel R, Rizza RA, Butler PC, et al. Beta-cell deficit and increased beta-cell apoptosis in humans with type 2 diabetes. *Diabetes.* 2003;52:102–10.
- Saleh MC, Fatehi-Hassanabad Z, Wang R, Nino-Fong R, Wadowska DW, Wright GM, et al. Mutated ATP synthase induces oxidative stress and impaired insulin secretion in beta-cells of female BHE/cdb rats. *Diabetes Metab Res Rev.* 2008;24:392–403.
- Suzuki S, Hinokio Y, Ohtomo M, Hirai M, Hirai A, Chiba M, et al. The effects of coenzyme Q10 treatment on maternally inherited diabetes mellitus and deafness, and mitochondrial DNA 3243(A to G) mutation. *Diabetologia.* 1998;41:584–8.
- Richter C. Oxidative damage to mitochondrial DNA and its relationship to ageing. *Int J Biochem Cell Biol.* 1995;27:647–53.
- Ogasahara S, Yorifuji S, Nishikawa Y, Takahashi M, Wada K, Hazama T, et al. Improvement of abnormal pyruvate metabolism and cardiac conduction defect with coenzyme Q10 in Kearns-Sayre syndrome. *Neurology.* 1985;35:372–7.
- Frei B, Kim MC, Ames BN. Ubiquinol-10 is an effective lipid-soluble antioxidant at physiological concentrations. *Proc Natl Acad Sci USA.* 1990;87:4879–83.
- Sumi K. Coenzyme Q10 suppresses the apoptosis of mouse pancreatic beta-cell line MIN6. *J Yonago Med Assoc.* 2013;64:33–9.
- Seleznev K, Zhao C, Zhang XH, Song K, Ma ZA. Calcium-independent phospholipase A2 localizes in and protects mitochondria during apoptotic induction by staurosporine. *J Biol Chem.* 2006;281:22275–88.
- Ma ZA, Zhao Z, Turk J. Mitochondrial dysfunction and β -cell failure in type 2 diabetes mellitus. *Exp Diabetes Res.* 2012. <https://doi.org/10.1155/2012/703538>.
- Hodgson JM, Watts GF, Playford DA, Burke V, Croft KD. Coenzyme Q10 improves blood pressure and glycaemic control: a controlled trial in subjects with type 2 diabetes. *Eur J Clin Nutr.* 2002;56:1137–42.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

