Bibliography on HIV/AIDS in Ethiopia and Ethiopians in the Diaspora: The 2015 Update

Paul J. Converse¹, Damen Haile Mariam², Helmut Kloos³, Wubegzier Mekonnen², Mesfin Samuel Mulatu⁴, Mirgissa Kaba²

Introduction

This thirteenth annual update of the HIV/AIDS literature on Ethiopia includes references to studies covering all major public health aspects of this infection in Ethiopia and Ethiopians in the diaspora. All references are again listed under eight main headings, as follows: 1) basic biomedical research 2) epidemiological, behavioral, socio-economic and cultural research 3) impacts research 4) treatment, care and clinical research 5) prevention research 6) health services and policy research 7) health informatics and evaluation research and 8) research on Ethiopians in the diaspora. The text preceding each list of references briefly summarizes patterns and trends and highlights and major findings of studies presenting new approaches, concepts and tools.

We hope that this annual update will, like previous issues, help to identify and encourage research in neglected, but relevant and promising areas in HIV/AIDS epidemiology, prevention, control, care and support. Because of the increasing complexity of the pertinent issues and relationships at hand and the fact that we do not cross reference any entries in our updates, we encourage readers interested in any one area of research to review also other sections in this update.

We used the same methods as in previous updates to identify and classify references. As such, literature searches was made using keywords "Ethiopia AND HIV AND 2015" and "Ethiopia AND AIDS AND 2015" in PubMed, CINHAL, EconLit, EMBASE, Global Health, POPLINE, PsycINFO, Social Services Abstracts, Sociological Abstracts, and other major databases that archive relevant published articles, dissertation, and reports from multiple sources. We made additional online searches on major national and regional HIV/AIDS resource centers and international organizations, mostly http://www.etharc.org and http://unaids.org. The number of references in this update (468, 110 fewer than in the 2014 update, probably due to fewer conferences and abstracts) includes 86 references from 4 major conferences, including the 26th Annual Conference of the Ethiopian Public Health Association (51 references), and a large number of published articles (329). But there were only 47 theses (62 fewer than in 2014), including 37 Master theses and 3 PhD dissertation (no degrees were identified for 7 theses); all of them were from Addis Ababa University. It was not possible to access theses from Jimma, Gondar, Hawassa and Mekele universities as in 2013. The remaining references were unpublished reports by different agencies and organizations.

We used the following PubMed search terms: Ethiopia AND HIV AND 2015[dp] or Ethiopia AND AIDS AND 2015[dp]. The findings indicate a linear increase in published articles from 2005 through 2015. The increase in conference presentations in recent years (see the trend in the 2014 Update for International AIDS Conference abstracts) appears to continue to be followed by an increase in presentation materials which are becoming full-length manuscripts. It is important to note that the trend in publications may be a product of both increased productivity and the expansion of the number and origins of journals indexed in PubMed. For example, the Ethiopian Journal of Health Sciences was indexed in PubMed only recently, permitting work done in Ethiopia to be counted in Figure 1. Other African journals that publish HIV/AIDS related articles from Ethiopia, (e.g. the Ethiopian Journal of Health Sciences and East African journals) have also been indexed in PubMed recently, although it is not clear when they were first included in the database.

¹Center for Tuberculosis Research, Johns Hopkins University, School of Medicine, Baltimore, MD 21231, E-mail-pconver1@jhmi.edu, USA;

²School of Public Health, College of Health Sciences, Addis Ababa University, E-mail- D.H. damen_gmail.com, W.M wubegzierm@gmail.com, M.K mirgissk@yahoo.com, P.O. Box 32812, Addis Ababa, Ethiopia;

³Department of Epidemiology and Biostatistics, University of California, Medical Center, San Francisco, San Francisco, E-mail- helmutkloss@gmail.com, CA 94143, USA;

⁴4169 Idlewood Parc Courtm, E-mail- mesfin.mulatu@yahoo.com, Tucker, Georgia 30084, USA




Figure 1: Publications Cited in PubMed Concerning Ethiopia and HIV or AIDS




Figure 2: Publications Cited in PubMed Concerning Ethiopia and HIV, AIDS, Tuberculosis, Malaria, and Helminthes

Similarly, Figure 2 presents trends in publications of articles on other common infections, including tuberculosis (TB) and malaria. Unlike previous updates, which included publication trends on schistosomiasis (until 2012), this update includes publication trends in helminth infections because of the growing relevance of parasitic infections in health outcomes of people living with HIV/AIDS in Ethiopia (see Section 2). For Figure 2, the PubMed search terms were: Ethiopia AND HIV AND 2015[dp]; Ethiopia AND AIDS AND 2015[dp]; Ethiopia malaria AND 2015[dp]; Ethiopia AND AND tuberculosis AND 2015[dp]; Ethiopia AND helminth AND 2015[dp]. In 2014, there was a large increase in the number of publications concerning HIV/ AIDS, and/or tuberculosis and Ethiopia. These numbers remained stable in 2015. (The authors note that this figure in the 2014 update was truncated at 150, thus not reflecting the true number). The number of papers concerning malaria continues at a steady level with 83 publications in 2015 similar to the 87 in 2014. Within the last four years, publication on helminthes appears to be increasing though it is down from the very high number in 2014.

Section 1: Basic Biomedical Research

This section covers laboratory-based biomedical research, including studies on HIV structure, replication, and host immune responses; co-infection with other agents; development and testing of laboratory procedures; and other related laboratory studies.

Only 11 articles were classified in the Biomedical Research category in 2015. Of these, seven were concerning tuberculosis, including one originally presented as a conference abstract. Thus, there were six independent studies related to TB. Of the remainder, Mudie et al. (7) presented an abstract documenting the protective effects for rat liver of an extract of Nigella sativa (black cumin, tiqur azmud) after exposure to HAART drugs. Mulu et al. (8) noted the absence of protease inhibitor resistance mutations in the prevalent HIV-1C strain or clade in 45 untreated patients. Shimelis and Tadesse (9), working in Hawassa, evaluated the performance of a kit to detect both syphilis and HIV infection and found positive and negative predictive values approaching or achieving a 100% performance. Tessema et al. (11) reported the declining trend of sulfadoxine-pyrimethamine (SP)-associated resistance mutations in isolates of *Plasmodium falciparum* in Pawe, Ethiopia, following discontinuation of SP use, and noted the possibility that cross-resistance to trimethoprim/sulfamethoxazole, used for treating HIVassociated opportunistic infections, might counteract the decline.

Returning to the tuberculosis studies, Ayana et al. (1) found that acid-fast microscopy for identifying TB bacilli performed at 8 laboratories \geq 500 km from Addis Ababa often reported false negative results that may be due to lack or reagents or problems in the work environment;

HIV seropositivity can also increase the rate of apparent negative results. Geleta et al. (2,3), working in Jigjiga >600 km from Addis, showed that diagnostic sensitivity can be increased through the use of the PCR-based Xpert MTB/RIF assay, which also can detect resistance to rifampin, but that this assay may not be as sensitive as culture in liquid or agar media, although culture is also prone to contamination. Getachew et al. (4) documented that auramine 0-based fluorescence microscopy for detection of Mtb in HIV-positive individuals is far more sensitive than traditional Ziehl-Neelsen acid-fast microscopy. Korma et al. (5) noted the increased frequency of extrapulmonary TB that has occurred since the onset of the HIV epidemic. Spoligotyping analysis suggested, however, that there are clusters of epidemiologically related strains possibly indicative of recent transmission. Although rifampin-resistance was detected, there were no multi-drug resistant strains found in the 200 cases studied. Manalebh et al. (6) used a crosssectional design to evaluate the acid-fast sputum smear microscopy performance of public-private mix (PPM) clinics and other health care facilities in West Amhara Zone. The quality of this microscopy underlies the effectiveness of directly observed treatment programs for TB control. They observed, like Ayana et al. (1) that facilities were often cramped and the quality of staining reagents was often below standard as was the ability to detect low numbers of acid-fast bacilli. They concluded that regular external quality assessments are the best way to enhance laboratory performance of the PPM facilities, in particular. Testing plasma samples of 195 HIV+/TB+, 170 HIV-/TB+ patients and 31 controls for levels of the inflammatory markers, neopterin and C-reactive protein (CRP) that were correlated to CD4 lymphocyte count before and after anti-TB treatment, Skogmar et al. (10) found that both biomarkers were significantly elevated and that neopterin levels, in particular, had negative correlation with CD4 levels, suggesting that inflammation may account, in part, for the reduced levels but that neither marker was predictive of counts below 100 per ml.

- 1. Ayana DA, Kidanemariam ZT, Tesfaye HM, Milashu FW. External quality assessment for acid fast bacilli smear microscopy in eastern part of Ethiopia. BMC Res Notes. 2015 Oct 05;8:537. Epub 2015/10/07.
- Geleta DA, Megerssa YC, Gudeta AN, Akalu GT, Debele MT, Tulu KD. Xpert MTB/RIF assay for diagnosis of pulmonary tuberculosis in sputum specimens in remote health care facility. BMC Microbiology. 2015 2015 2016-04-16;15.
- Geleta DA, Megerssa YC, Negussie A. Evaluation of Xpert MTB/RIF Assay for diagnosis of pulmonary tuberculosis in sputum specimens in remote health care facility at Kharamara Hospital, Jijiga, eastern Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.

- Getachew K, Abebe T, Kebede A, Mihret A, Melkamu G. Performance of LED Fluorescence Microscopy for the diagnosis of pulmonary tuberculosis in HIV positive individuals in Addis Ababa, Ethiopia. Tuberculosis Research and Treatment. 2015;2015:794064. Epub 2015/12/22.
- Korma W, Mihret A, Hussien J, Anthony R, Lakew M, Aseffa A. Clinical, molecular and drug sensitivity pattern of mycobacterial isolates from extrapulmonary tuberculosis cases in Addis Ababa, Ethiopia. BMC Infect Dis. 2015 Oct 26;15:456. Epub 2015/10/28.
- Manalebh A, Demissie M, Mekonnen D, Abera B. The quality of sputum smear microscopy in publicprivate mix directly observed treatment laboratories in West Amhara Region, Ethiopia. PLoS One. 2015 Apr 2015 2015-04-11;10(4).
- Mudie K, Seifu D, Challa F, Abebe A, Debella A, et al. Hepatoprotective activity of aqueous seed extract of *Nigella sativa* against highly active antiretroviral therapy induced hepatotoxicity in rats. 51st Medical Conference, Ethiopian Medical Association; February 26-28, 2015; Bahir Dar, 2015.
- Mulu A, Maier M, Liebert UG. Lack of integrase inhibitors associated resistance mutations among HIV-1C isolates. Journal of Translational Medicine. 2015 December 01;13 (1) (no pagination)(377).
- Shimelis T, Tadesse E. The diagnostic performance evaluation of the SD BIOLINE HIV/syphilis Duo rapid test in southern Ethiopia: a cross-sectional study. BMJ Open. 2015 Apr 23;5(4):e007371. Epub 2015/04/25.
- Skogmar S, Schön T, Balcha TT, Sturegård E, Jansson M, Björkman P. Plasma levels of neopterin and C-reactive protein (CRP) in tuberculosis (TB) with and without HIV coinfection in relation to CD4 cell count. PLoS One. 2015 Dec 2015 2015-12-05;10(12).
- Tessema SK, Kassa M, Kebede A, Mohammed H, Leta GT, Woyessa A, et al. Declining trend of *Plasmodium falciparum* dihydrofolate reductase (dhfr) and dihydropteroate synthase (dhps) mutant alleles after the withdrawal of sulfadoxinepyrimethamine in north western Ethiopia. PLoS One. 2015 Oct 2015 2015-10-10;10(10).

Section 2: Epidemiological, Behavioral, Socio-Economic and Cultural Research

This section includes studies on the epidemiology of HIV and other opportunistic infections, AIDS and related diseases, and risk and protective behaviors. It also covers research on the biological, psychosocial, socio-economic, cultural, structural, and other contextual determinants of HIV transmission and prevention.

This section contains 138 references: 71 (51.4%) published articles, 39 (28.3%) conference abstracts, 25 (18.1%) masters theses, 2 (1.4%) doctoral dissertations, and 1 (0.7%) report. The broad categories of research

interest within this section are very similar to those of the previous update. In this update, the primary areas of research focus are: 1) HIV prevalence and factors associated with HIV infection; 2) prevalence and impacts of opportunistic infections and other comorbidities; 3) sexual and other risk behaviors for HIV infection; 4) reproductive health knowledge, attitudes, and perception; 5) occupational risk exposure and protection; 6) violence and harmful traditional practices; and 7) other existing and emerging research areas.

HIV seroprevalence and its determinants were examined in several studies using different approaches, in different settings, and among a variety of population groups (40, 54, 62, 69, 74, 77, 83, 85, 87, 93, 104, 111). Of special note are two studies that linked HIV seroprevalence with broader socio-economic drivers of risk for HIV infection. Lakew and colleagues (85) used data from the 2011 Ethiopian demographic and Health Survey (DHS) to examine the link between HIV seroprevalence and multiple social factors, including wealth, age, occupation, knowledge of HIV, use of alcohol or khat, migration, religion, location of residence including administrative region, education, mass media exposure, gender, number of lifetime sexual partners, and marital status. The authors found high levels of HIV prevalence (10-21%) in some selected geographic hotspots in central, eastern and western geographic clusters of Ethiopia. In addition, HIV prevalence was associated with higher levels of wealth, education, and lifetime sexual partners. Furthermore, increased odds of HIV infection were observed among younger adults, merchants, urban residents and females than among older, unemployed, rural residents and males, respectively. Another analysis by Hargreaves and colleagues (74) used multiple years of DHS data from eastern and southern African countries and revealed different patterns of relationship between level of education and HIV prevalence (74). This study confirmed Lakew and colleagues' finding that in Ethiopia and Malawi, HIV prevalence was higher among more educated women, although the reverse relationship was found in Lesotho, Kenya and Zimbabwe. The authors also found that in Ethiopia HIV prevalence fell significantly from 2003-2005 to 2008-2012 only among adult women with no and secondary education. Another study in this area was by Pegurri and colleagues (117), who set out to estimate the burden of HIV in older children and the prevalence of HIV in orphans and vulnerable children (OVC) households. The researchers used data derived from the Ethiopian DHS of 2005 and 2011 and antenatal HIV sentinel surveillance, and the Spectrum/ Estimation and Projection Package (EPP), a software used globally to produce HIV estimates. The analysis estimated that there were 160,000 HIV-positive children under 15 years old and 800,000 orphans due to AIDS in 2013. An HIV prevalence of 11.9% was also found among almost 10,000 OVC under 18 years old. The authors concluded that there is a large population of children living with HIV in Ethiopia, the majority of

Ethiop. J. Health Dev. 2016;30(3)

them were vertically infected and never identified nor linked into treatment. Overall, these highlighted studies underscore the need for continued research on the patterns and trends of HIV prevalence in the country with a focus on the most vulnerable groups of the population in order to design better HIV prevention, treatment, and care programs.

Consistent with previous updates, a large number of studies, in this section, are devoted to research on opportunistic infections and other comorbidities among people living with HIV infections. The incidence or prevalence of, risk factors for and impact of viral, bacterial, or parasitic infections among people living with HIV continued to be of significant interest to researchers. As in previous updates, the most frequently studied coinfections were HIV and tuberculosis (11, 14, 21, 41, 52, 98, 107). Ayalew and colleagues (14) found a high incidence rate of tuberculosis infection (4.9% per year) among HIV-positive children. In another study with 499 HIV-positive persons in southern Ethiopia, Fekadu and co-researchers (52) found that 18.2% of them had tuberculosis. Other researchers studied persons with known or suspected tuberculosis and determined prevalence of co-infection with HIV. For example, Belay and colleagues (21) conducted one of the earliest studies on the co-occurrence of HIV and tuberculosis among pastoral communities. In this study, among Afar pastoralists with bacteriologically confirmed pulmonary tuberculosis, 40.4% were co-infected with HIV. In addition to co-occurrence of HIV and tuberculosis, other studies focused on risk factors and the impact of coinfection on immune response, treatment, and health outcomes (11, 41, 52, 98, 107). In three studies. researchers examined prevalence and risk factors for coinfection of HIV with hepatitis B or C virus (4, 29, 136). Researchers were also interested in other co-infections including HIV and other sexually transmitted infections such as syphilis (29, 46, 59, 94, 115), HIV and intestinal parasites (11, 56, 84), and HIV and other infectious conditions (43, 84, 109, 116, 138). There has also been an interest in non-communicable co-morbidities among people living with HIV infection, including diabetes (100), anemia (44, 45), and cognitive impairment (105). Amare and colleagues (11) published a review paper that summarizes the burden imposed by HIV/AIDS, tuberculosis, chronic intestinal parasites, and multiple micronutrient deficiency in Ethiopia. This paper provides a brief review of evidence about the bi-directional and interactive nature of these conditions and their role in health outcomes of people living with HIV infection.

As with previous updates, and with recognition of the strong and syndemic relationships between HIV and tuberculosis, several studies focusing on the prevalence, risk factors, and impacts of tuberculosis are included in this update (9, 10, 22, 24, 30, 36, 47, 90, 98, 120, 126, 127, 130). Three of these studies focused on the growing problem of multi-drug resistant tuberculosis (90, 98,

130). In one of the studies, the overall prevalence multidrug resistant tuberculosis was 5.7% among new and 13.9% among previously treated smear-positive tuberculosis patients (90). Given that multi-drug tuberculosis is increasing globally and it has been associated with poor health outcomes among people living with HIV infection, continued research in this area to document prevalence, identify risk factors, develop effective prevention and control measures will be warranted. In addition to tuberculosis, studies on other health problems associated with HIV, such as infections with intestinal parasites among tuberculosis patients (1, 2), hepatitis C infection in Africa (111), and pediatric leishmaniasis (49) are included in this update.

A large number of studies in this section dealt with the patterns of sexual, drug use, and other risk behaviors for HIV acquisition and spread, and the demographic and psychosocial factors associated with these risk behaviors. Sexual risk behavior studies covered such topics as sexual initiation, pre-marital sexual activity, sex with multiple partners, and unprotected sex among youth, primarily those in high schools and colleges across Ethiopia (13, 17, 18, 23, 25, 27, 31, 35, 38, 58, 60, 68, 72, 81, 85, 93, 96, 102, 103, 108, 119, 123, 125, 128, 129, 134, 135). Some studies focused on sexual risk behaviors, associated factors, and health outcomes (e.g., HIV infection, unintended pregnancy) among commercial sex workers (3, 58, 90, 106, 122, 134, 135). There were also studies of migrant workers (18, 129). Of particular note is the study by Tiruneh and colleagues (129), which documented high levels of risky sexual behaviors among seasonal migrant laborers in northwestern Ethiopia. In this study, 74% of seasonal laborers reported sexual intercourse with commercial sex workers, 69% reported multiple sexual partners, and 49% admitted to unprotected sex within the preceding six months. This is a significant finding given that migrant labor is likely to become a prominent commodity with the growing economy of the country. Substance use as a potential behavioral risk for HIV infection is still understudied, with only two studies reporting khat chewing among youth in this update (70, 132). The prevalence and potential impact of alcohol and other illicit substance use is still poorly understood. There is a need for further study that examines a wide range of behavioral risk factors for HIV, including the interplay between substance use and sexual risk among vulnerable populations, such as commercial sex workers, migrant workers, and young people.

Although research on knowledge, attitudes, and perceptions around HIV/AIDS has declined, some relevant studies have still been reported. These studies covered the assessment of the link between comprehensive knowledge of HIV/AIDS and risk behaviors among college students (113), knowledge of pregnant women about mother-to-child transmission of HIV (28), HIV/AIDS knowledge and its relations with stigma among women (50), knowledge and misconception of young women toward sexually transmitted infection and condom use (63), knowledge of healthcare workers about tuberculosis control (64), and perception of risk for HIV infection and its connection with utilization of voluntary counseling and testing services among college students (117). Other relevant topics that have been studied include knowledge and awareness about broader reproductive health issues among early adolescents (7), sexual discourse, socialization, and communication, including parentadolescent communication about sex (55, 60, 101, 121), construction of masculinity and health behaviors (18), and assessment of religious attitudes towards sexuality among youth (101).

In this update, there appears to be a resurgence of interest in examining the sexual, physical, and other forms of violence and harmful traditional practices and their connections to poor health outcomes including increased risk for HIV infection and reproductive health problems. A large number of studies presented findings on the prevalence, associated factors, and/or health-related consequences of sexual abuse and gender-based violence against children, adolescent girls, maids, commercial sex workers, refugee women, and young women in high schools or colleges (6, 8, 15, 37, 53, 57, 82, 86, 89, 91, 92, 99, 114, 123). Lifetime sexual violence has been reported to be high. For example, in a study on 369 high school female students, Mekuria and colleagues (92) found a lifetime rate of rape among 11% of the students. In another study, Alemayehu and co-researchers (8) reported very high levels of sexual and physical violence among commercial sex workers including rape (44%), unwanted touch (62%), pressure to have sex without a condom (56%), and physical harm (46%). Researchers have also documented the immediate and long-term consequences of violence including post-stress traumatic disorders (37) and intergenerational psychosocial effects on survivors' children (15). Two traditional practices that are indirectly tied to HIV infection are early marriage and female genital cutting. These practices were the subject of investigation in four studies (5, 12, 32, 39); all focused on the magnitude and reproductive health consequences of these practices.

Six studies reported research on healthcare related practices that might contribute to HIV acquisition or transmission among healthcare staff or patients (20, 34, 75, 78, 88, 97). Bekele and colleagues (20) conducted a survey with a randomly selected sample of 362 healthcare workers to learn about their attitude, reporting behaviours, and management practices in relation to needle stick and sharp injuries. The authors found that 37.1% of their respondents had experienced injuries at work but 58.7 % of such injuries were not reported to appropriate management. The authors concluded that occupational needle stick and/or sharps injuries are common among healthcare workers in the study area and

recommended provision of on-the-job training on the risk of occupational exposure to pathogens and the need for timely reporting and management of injuries. In a mixed methods study of the practices of health facilities in Dessie, Mesfin and co-researchers (97) reported widespread unsafe practices in handling injection and sharp objects that predisposed the patient, the health worker, and the community to risk of accidental needle stick injuries. Burssa and colleagues (34) provided evidence that indicated a decline in transfusion transmitted blood borne infections among blood donors, although the authors acknowledged that such infections continue to occur at an overall prevalence rate of 6.9% among blood donors - prevalence of HIV, HBV, HCV, and syphilis transmission being 0.9%, 4.3%, 0.7%, and 1.0% respectively. Overall, these studies underscore the need for further research into the prevalence and determinants of unsafe practices and programs designed to prevent occupational risk to HIV and other blood bone infections in the country.

Other studies explored issues covered in previous updates including HIV/AIDS related stigma (19, 50, 131), fertility desires of people living with HIV (51, 71), exposure to explicit sexual content and its associations with sexual behavior (48, 73) and sexual minority rights (133). Two studies covered relatively new areas - one on the potential effects of tourism on tribal communities (124) and the other on the relationships between physical exercise and HIV/AIDS. Further research on these and other broader social determinants of HIV/AIDS are warranted.

- 1. Abate E, Belayneh M, Idh J, Diro E, Elias D, Britton S, et al. Asymptomatic helminth infection in active tuberculosis is associated with increased regulatory and Th-2 responses and a lower sputum smear positivity. PLoS Negl Trop Dis. 2015 Aug;9(8):e0003994. Epub 2015/08/08.
- Abate E, Belayneh M, Idh J, Diro E, Elias D, Britton 2. S, et al. Asymptomatic helminth infection in active tuberculosis is associated with increased regulatory and Th-2 responses and a lower sputum smear positivity. PLoS Neglected Tropical Diseases. 2015 Aug 2015 2015-10-09;9(8).
- Abebe F, Abdo A, Gudeta D, Hagos H, Caswell G. 3. Empowering each other: young people who sell sex in Ethiopia. a case study from the Link Up Project. Hove, United Kingdom, International HIV / AIDS Alliance, Link Up, [2015]. 2015.
- 4. Abera B, Zenebe Y, Mulu W, Kibret M, Kashu G. Seroprevalence of hepatitis B and C viruses and risk factors in HIV-infected children at Felege Hiwot Referral Hospital, Ethiopia. 26th Annudal Conference of the Ethiopian Public Health Association; February 5-6, 2015; Bahir Dar, 2015.
- Abera L, Dibaba Y, Alemu T. Magnitude of female 5. genital mutilation and associated birth complications among reproductive age women and first birth in

Basketo Special Woreda, SNNPR, Ethiopa. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.

- Admas M, Tebeje B, Megersa H. Predictors and consequences of sexual violence among female students of College of Public Health and Medical Sciences, Jimma University, southwest Ethiopia, 2014. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 7. Admassu S. Awareness of sexual and reproductive health mong very young (10-14 years) in shool adolescents in Addis Ababa. [MPH Thesis]: Addis Ababa University; 2015.
- Alemayehu M, Yohannes G, Damte A, Fantahun A, Gebrekirstos K, Tsegay R, et al. Prevalence and predictors of sexual violence among commercial sex workers in northern Ethiopia. Reprod Health. 2015 May 23;12:47. Epub 2015/05/24.
- Ali S, Amlak AH, Heinrich N, Hoelscher M, Loescher T, Rachow A. Tuberculosis among Ethiopian inmates. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Ali S, Haileamlak A, Wieser A, Pritsch M, Heinrich N, Loscher T, et al. Prevalence of Pulmonary Tuberculosis among Prison Inmates in Ethiopia, a Cross-Sectional Study. PLoS One. 2015;10(12):e0144040. Epub 2015/12/08.
- Amare B, Moges B, Mulu A, Yifru S, Kassu A. Quadruple burden of HIV/AIDS, tuberculosis, chronic intestinal parasitoses, and multiple micronutrient deficiency in Ethiopia: A summary of available findings. BioMed Research International. 2015 (no pagination) (598605).
- Ashenafi W. Assessment of health and sociodemographic problems of early marriage among ever married women in Kers District, East Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Asres A, Abebe W. Risky sexual practices among private college students: a cross-sectional study at Central University College, Addis Ababa, Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 14. Ayalaw SG, Alene KA, Adane AA. Incidence and predictors of tuberculosis among HIV positive children at University of Gondar Referral Hospital, northwest Ethiopia: a Retrospective Follow-Up Study. International scholarly research notices. 2015;2015:307810. Epub 2015/01/01.
- 15. Ayentu E. Psychosocial consequences of child abuse on abuse survivor female children: the case of three children rehabilitation centers in Addis Ababa. [MA Thesis]: Addis Ababa University; 2015.
- 16. Barreiro P, Tisiano G, Fano H, Yohannes T, Gosa A, Fruttero E, et al. Prevalence of HIV, HBV and HCV positive serology in blood donors in a rural general

hospital in Ethiopia. Tropical Medicine and International Health. 2015 September;20:416.

- Barsisa A. Construction of masculinity, health risk behaviors and help seeking attitudes among undergraduate students in selected universities in Ethiopia. [PhD Thesis]: Addis Ababa University; 2015.
- Befirdu J, Seme A. Unprotected sex and contraceptive use: perceptions among Ethiopian female migrants to Arab States. 26th Annual Confeence of the Ethiopian Public Health Association February 26-28, 2016; Bahir Dar, 2015.
- 19. Bekalu MA, Eggermont S. Socioeconomic and socioecological determinants of AIDS stigma and the mediating role of AIDS knowledge and media use. Journal of Communication in Healthcare. 2015;8(4):316-324.
- 20. Bekele T, Gebremariam A, Kaso M, Ahmed K. Attitude, reporting behavour and management practice of occupational needle stick and sharps injuries among hospital healthcare workers in Bale Zone, southeast Ethiopia: a cross-sectional study. Journal of Occupational Medicine and Toxicology. 2015 03 Dec;10 (1) (no pagination)(42).
- 21. Belay M, Bjune G, Abebe F. Prevalence of tuberculosis, HIV, and TB-HIV co-infection among pulmonary tuberculosis suspects in a predominantly pastoralist area, northeast Ethiopia. Global Health Action. 2015;8:27949.
- 22. Belay M, Legesse M, Mihret A, Bekele Y, Ottenhoff THM, Franken LM, et al. Pro- and antiinflammatory cytokines against Rv2031 are elevated during latent tuberculosis: a Study in cohorts of tuberculosis patients, household contacts and communityc controls in an endemic setting. PLoS One. 2Apr 2015 2015-04-26;10(4).
- 23. Benti T, E. AN, Anteneh K. Prevalence of premarital sexual practice and associated factors among undergraduate health science students of Madawalabu University, Bale Goba, south east Ethiopia: an institution based cross sectional study. Pan African Medical Journal. 2015;20(209).
- 24. Berg S, Schelling E, Hailu E, Firdessa R, Gumi B, Erenso G, et al. Investigation of the high rates of extrapulmonary tuberculosis in Ethiopia reveals no single driving factor and minimal evidence for zoonotic transmission of *Mycobacterium bovis* infection. BMC Infectious Diseases. 2015 March 03;15 (1) (no pagination)(112).
- Berhan Y, Berhan A. A meta-analysis of risky Sexual behaviour among male youth in developing countries. AIDS Research and Treatment. 2015:580961. Epub 2015/02/25.
- 26. Berhe W, Adama MY, Ferede AS, Ashebir Y, Haile F, Afework M, et al. Infant mortality and causes of infant deaths in rural Ethiopia: a population-based cohort of 3684 births. BMC Public Health. 2015 2016-04-16;15.

- 27. Beyene A, Muhiye A, Getachew Y, Hiruye A, Mamo D, H/Mariam D. Assessment of the magnitude of teenage pregnancy and its associated factors among teenage females visiting Gassosa General Hospital for health care services. 51st Medical Conference, Ethiopian Medical Association; February 5-6, 2015; Addis Ababa 2015.
- Birhane T, Assefa Tessema G, Addis Alene K, Dadi AF. Knowledge of pregnant women on mother-tochild transmission of HIV in Meket District, Northeast Ethiopia. J Pregnancy. 2015:960830. Epub 2015/03/06
- 29. Birhaneselassie M. Sero-prevalence of hepatitis, HIV and syphils among blood donors at Yirgalem Hsopital, southern Ethiopia: trends over a period of 5 years. 51st Medical Conference, Ethiopian Medical Association; February 26-28, 2015; Bahir Dar, 2015.
- 30. Birlie A, Tesfaw G, Dejene T, Woldemichael K. Time to death and associated factors among tuberculosis patients in Dangila Woreda, northwest Ethiopia. PLoS ONE. 2015 01 Dec;10 (12) (no pagination)(e0144244).
- Bogale A, Seme A. Premarital sexual practices and their predictors among in-school youth of Shendi Town, West Gojam Zone, northwestern Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 32. Bogale D, Markos D, Kaso M. Prevalence of early marriage and its consequences among reproductive age women in Bale Zone, Ethiopia: a cross-sectional study. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Burnett D, Aronson J, Asgary R. Oral health status, knowledge, attitudes and behaviours among marginalized children in Addis Ababa, Ethiopia. Journal of child health care : for professionals working with children in the hospital and community. 2016 Jun;20(2):252-261. Epub 2015/02/26.
- 34. Burssa DB, Demewoz HD, Shiferaw MS. A paradigm shift in the prevalence of transfusion transmitted infections among blood donors in Bahir Dar Blood Bank, northwest Ethiopia. Vox Sanguinis. 2015 June;109:230.
- 35. Dadi AF, Teklu FG. Risky sexual behavior and associated dactors among grade 9-12 students in Humera Secondary School, western Zone of Tigray, northwest Ethiopia, 2014. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Dangisso MH, Datiko DG, Lindtjorn B. Spatiotemporal analysis of smear-positive tuberculosis in the Sidama Zone, southern Ethiopia. PLoS ONE. 2015;10(6).
- 37. Daniel S. Prevalence of post-traumatic stress disorders among sexually abused children at Safe House. [MA Thesis]: Addis Ababa University; 2015.

- 38. Debebe W, Solomon S. Sexual risk behaviors and its associated factors among undergraduate students in Madda Walabu University, southeast Ethiopia: a facility based cross sectional study. Epidemiology (open access). 2015;5(4).
- 39. Dejenu G, Workineh S, Degu G. Early marriage: perceptions, practice and determinants among female children from 2009 to 2013 in Sinane District, northwest Ethiopia. 26th Annual Conference of thee Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar 2015.
- 40. Demarco T, Rountree W, Hora B, Chen Y, Keinonen S, Racz L, et al. HIV-1 subtype C is significantly more infectious than other subtypes. Journal of the International AIDS Society. 2015;18:112.
- Deribe K, Yami A, Deribew A, Mesfin N, Colebunders R, Van Geertruyden JP, et al. Predictors of mortality among tuberculosis/HIV-coinfected persons in southwest Ethiopia: a case-control study. J Int Assoc Provid AIDS Care. 2015 May-Jun;14(3):269-273. Epub 2013/08/24.
- 42. Dewo Z. Qualitative study on adolescent trafficking for primary activity and domestic labour in Addis Ababa. . 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 43. Diro E, van Griensven J, Mohammed R, Colebunders R, Asefa M, Hailu A, et al. Atypical manifestations of visceral leishmaniasis in patients with HIV in north Ethiopia: a gap in guidelines for the management of opportunistic infections in resource poor settings. The Lancet Infectious Diseases. 2015 01 Jan;15(1):122-129.
- Enawgaw B, Alem M, Melku M, Addis Z, Terefe B, Yitayaw G. Prevalence and associated risk factors of anemia among HIV-infected children attending Gondar University Hospital, northwest Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 5-6, 2015; Bahir Dar, 2015.
- 45. Enawgaw B, Alem M, Melku M, Addis Z, Terefe B, Yitayew G. Prevalence and associated risk factors of anemia among HIV infected children attending Gondar University Hospital, Northwest Ethiopia: a cross sectional study. BMC Hematology. 2015;15:12. Epub 2015/09/29.
- 46. Endris M, Deressa T, Belyhun Y, Moges F. Seroprevalence of syphilis and human immunodeficiency virus infections among pregnant women who attend the University of Gondar Teaching Hospital, nrthwest Ethiopia: across sectional study. BMC Infectious Diseases. 2015 March 03;15 (1) (no pagination)(111).
- 47. Ephrem T, Mengiste B, Mesfin F, Godana W. Determinants of active pulmonary tuberculosis in Ambo Hospital, West Ethiopia. African journal of primary health care & amp; family medicine. 2015;7(1):doi: 10 4102/phcfm v4107i4101 4608. Epub 2015/10/16.

- 48. Ermias D. Pornographic consumption and its association with early sexual abuse among high school students of Gulele Sub-City, Addis Ababa, Ethiopia.: Addis Ababa University; 2015.
- Ermias D, Lynen L, Berhane G, Abraham A, Wubishet L, Zewdu B, et al. Clinical aspects of paediatric visceral leishmaniasis in north-west Ethiopia. Tropical Medicine and International Health. 2015;20(1):8-16.
- Eshetu G, Dula E. HIV/AIDS knowledge and stigma among women of reproductive age in Ethiopia. African Journal of AIDS Research. 2015;14(3):191-199.
- 51. Fassika A, Worku A, Yayehirad A. Fertility desire and associated factors among clients on highly active antiretroviral treatment at Finoteselam Hospital northwest Ethiopia: a cross sectional study. Reproductive Health. 2015;12(69).
- 52. Fekadu S, Teshome W, Alemu G. Prevalence and determinants of Tuberculosis among HIV infected patients in south Ethiopia. J Infect Dev Ctries. 2015 Aug 29;9(8):898-904. Epub 2015/09/01.
- 53. Gedilu E. Assessment of prevalence of sexual abuse and predisposing condition among children treated at Tikur Anbessa Specialized Hospital, Department of Pediatrics and Child Health. [MSc Thesis]: Addis Ababa University; 2015.
- 54. Gelaye A, Gidey B, Abera Feleke A. HIV prevalence among exposed infants in Amhara Region, Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 55. Gertachew Y. Persuation techniques in the formation of sexual intimacy among first year students of the School of Commerce at Addis Ababa University. [MA Thesis]: Addis Ababa University; 2015.
- 56. Getachew M. Prevalence and determinants of sexual violence among female huse maids in selected junior secondary night schools: cross sectional study, Addis Ababa. [MSc Thesis]: Addis Ababa University; 2015.
- Getaneh L. Sex business in Addis Ababa. [MA Thesis]: Addis Ababa University; 2005.
- 58. Getu Y. Prevalence of urinary tract infection associated bacterial pathogen among HIV-positive patients attending ALERT Center: a cross sectional hospital based study. [MPH Thesis]: Addis Ababa University; 2015.
- 59. Gezahegn T. Sex and sexual health talk among Debre Birhan University students, North Shewa, Ethiopia: a qualitative study. 26th Annual Conference of the Ethiopian Public Health Association February 26-28, 2015; Bahir Dar, 2015.
- 60. Girgis S, Longhurst R, Cheng L. Infant oral mutilation: fiction, myths and facts. International Journal of Oral and Maxillofacial Surgery. 2015 October;44:e47.
- 61. Girma M. Study of plasma HIV viral RNA level among HIV-infected pregnant mothers who are on

antiretroviral prophylaxis and its impact on HIV transmission to their infants. [MSc Thesis]: Addis Ababa University; 2015.

- 62. Girmatsion F. Knowledge and misconception of young women toward sexual transmitted infection and condom use in northern Ethiopia: cross sectional study. Journal of Public Health and Epidemiology. 2015;7(4):138-144.
- 63. Gizaw GD, Aderaw Z, Kibret KT. Assessment of knowledge and practice of health workers towards tuberculosis infection control and associated factors in public health facilities on Addos Ababa: crosssectional study. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 64. Godefay H, Byass P, Graham WJ, Kinsman J, Mulugeta A. Risk factors for maternal mortality in rural Tigray, northern Ethiopia: a case-control study. PLoS One. 2015;10(12):e0144975. Epub 2015/12/19.
- 65. Gurmu E, Etana D. HIV/AIDS knowledge and stigma among women of reproductive age in Ethiopia. African Journal of AIDS Research. 2015 03 Jul;14(3):191-199.
- 66. Gutu B. Prevalence of HIV preventive sexual behavior and associated factors using behavioral model among provate college students in Ambo Town, Oromia Region, Ethiopia. 51st Medical Conference, Ethiopian Medical Association; February 5-6, 2015; Addis Ababa, 2015.
- 67. Habesha T, Aderaw Z, Lakew S. Assessment of exposure to sexually explicit materials and factors associated with exposure among preparatory school youths in Hawassa City, Southern Ethiopia: a cross-sectional institution based survey. Reprod Health. 2015 Sep 14;12:86. Epub 2015/09/16.
- Habte E, Yami A, Alemseged F, Abdissa Y, Deribe K, Memiah P, et al. Predictors of HIV serodiscordance among couples in southwestern Ethiopia. J Int Assoc Provid AIDS Care. 2015 May-Jun;14(3):234-240. Epub 2013/05/24.
- 69. Haile D, Lakew Y. Khat chewing practice and associated factors among adults in Ethiopia: further analysis using the 2011 Demographic and Health Survey. PLoS One. 2015 Jun 2015 2015-06-20;10(6).
- 70. Haile F. Assessment of fertility desire among on ART people living with HIV in Arba Minch and Zuria Woreda facilities attending care and treatment in ART unit. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 71. Hailu H. Assessment of religious attitude towards sexuality among youths in Addis Ababa: Addis Ababa University; 2015.
- 72. Hailu M. Assessment of unsafe sexual practice and determinant factors among high school students in Arba Minch Town, Gamo Gofa Zone, SNNPR, Ethiopia. 26th Annual Conference of the Ethiopian *Ethiop. J. Health Dev.* 2016;30(3)

Public Health Association; February 26-28, 2015; Bahir Dar, 2015.

- 73. Hargreaves JR, Davey C, Fearon E, Hensen B, Krishnaratne S. Trends in socioeconomic inequalities in HIV prevalence among young people in seven countries in eastern and southern Africa. PLoS ONE. 2015 20 Mar;10 (3) (no pagination)(e0121775).
- 74. Holana DN. Assessment of prevalence and determinants of needs stick injuries among health professionals in Addis Ababa. [MPH Thesis]: Addis Ababa University; 2015.
- 75. Hunchak C, Teklu S, Meshkat N, Meaney C, Puchalski Ritchie L. Patterns and predictors of early mortality among emergency department patients in Addis Ababa, Ethiopia. BMC Res Notes. 2015 Oct 24;8:605. Epub 2015/10/27.
- 76. Huruy K, Maier M, Mulu A, Liebert UG. Limited increase in primary HIV-1C drug resistance mutations in treatment naive individuals in Ethiopia. Journal of Medical Virology. 2015 Jun;87(6):978-984. Epub 2015/02/05.Kebede G, Tefera Y, Kisi T. Occupational exposure to HIV/AIDS risky conditions and post exposure prophylaxis utilization among health care workers in Gondar City, northwest Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 77. Kebede T. Challenges of being a child household head in child headed households affected by HIV/AIDS in Addis Ababa: the case of six child houshold heads in Gulele Sub City.: Addis Ababa University; 2015.
- Kebrid D. The relationship between physical exercise and HIV/AIDS, a systematic review and meta analysis. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 79. Kenyon CR, Tsoumanis A, Schwartz IS. HIV Prevalence Correlates with high-risk sexual behavior in Ethiopia's Regions. PLoS One. 2015;10(10):e0140835. Epub 2015/10/27.
- Kibru N. Coverage analysis of human trafficking: a focus on Addis Admas, Addis Zemen and reporter newspapers. [MA Thesis]: Addis Ababa University; 2015.
- Kidane HG. Seroprevalence of HIV among major gynecologic patients at Ghandi Memorial Hospital. [MSc Thesis]: Addis Ababa University; 2015.
- 82. Kiros H, Nibret E, Munshea A, Kerisew B, Adal M. Prevalence of intestinal protozoan infections among individuals living with HIV/AIDS at Felegehiwot referral hospital, Bahir Dar, Ethiopia. International Journal of Infectious Diseases. 2015 June 01;35:80-86.
- 83. Lakew Y, Benedict S, Haile D. Social determinants of HIV infection, hotspot areas and subpopulation groups in Ethiopia: evidence from the National

Demographic and Health Survey in 2011. BMJ Open. 2015;5(11):e008669.

- 84. Leta T, Feleke A, Dersseb L. Assessment of violence associated factors among rural high school female students in Hadiua Zone, Southern Nation and Nationalities Peoples' Region, Ethiopia, 2013. 26th Annual Conference of the Ethiopian Public Health Conference; February 26-28, 2015; Bahir Dar, 2015.
- 85. Marye S. Assessing point prevalence estimate of causes for IV related emergency room visits in all Addis Ababa public hospitals, [MSc Thesis]: Addis Ababa University; 2015.
- 86. Medeksa L. Needle-stick injuries and contributing factors among healthcare workers in public health facilities in Jigjiga Zone, eastern Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 87. Megersa T. Gender based violence in the rural setting of Arsi: causes and consequence analysis in Hetosa District. [MA Thesis]: Addis Ababa University; 2015.
- Mekonnen F, Tessema B, Moges F, Gelaw A, Eshetie S, Kumera G. Multidrug resistant tuberculosis: prevalence and risk factors in districts of Metema and West Armachiho, Northwest Ethiopia. BMC Infect Dis. 2015 Oct 26;15:461. Epub 2015/10/28.
- 89. Mekuria A, Abera M, Nigussie A. Childhood sexual abuse experiences and its associated factors among female high school students in Arba Minch Town, Gamo Gofa Zone, southern Ethiopia. 26th Annul Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 90. Mekuria A, Nigussie A, Abera M. Childhood sexual abuse experiences and its associated factors among adolescent female high school students in Arbaminch town, Gammo Goffa Zone, southern Ethiopia: a mixed method study. BMC Int Health Hum Rights. 2015 Aug 18;15:21. Epub 2015/08/19.
- 91. Melhado L. Parental HIV status linked to sexual debut In Sub-Saharan Africa. International Perspectives on Sexual and Reproductive Health. 2015 Jun 2015 2015-08-15;41(2):110.
- 92. Melku M, Kebede A, Addis Z. Magnitude of HIV and syphilis seroprevalence among pregnant women in Gondar, northwest Ethiopia: a cross-sectional study. HIV/AIDS (Auckland, NZ). 2015;7:175-182. Epub 2015/06/18.
- Mengsteab M. Exploratory study on trafficking-led prostitution: personal narratives of street prostitution in Addis Ababa. [MA Thesis]: Addis Ababa University; 2015.
- 94. Menna T, Ali A, Worku A. The trend of HIV/AIDS related mortality among primary and secondary school teachers in Addis Ababa, Ethiopia: using a verbal autopsy method. Ethiopian Medical Journal. 2015 01 Apr;53(2):49-56.
- 95. Mesfin Haile K, Teferi G, Efrem E, Tsige Gebre M. Assessment of health care providers injection *Ethiop. J. Health Dev.* 2016;30(3)

practices in Dessie District, north-eastern Ethiopia: facility based study. Journal of Pharmaceutical and Scientific Innovation. 2015;4(2):100-107.

- 96. Moges Y, H/Mariam D. Assessment of the association between HIV/AIDS and multi-drug resistance tuberculosis: a systenatic review with meta-analysis. 26th Annual Conference of the Ethiopan Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 97. Mohamed F. Sexual and gender based violance against refugee women in the refugee settlements. [MA Thesis]: Addis Ababa University; 2015.
- 98. Mohammed AE, Shenkute TY, Gebisa WC. Diabetes mellitus and risk factors in human immunodeficiency virus-infected individuals at Jimma University Specialized Hospital, southwest Ethiopia. Diabetes, Metabolic Syndrome and Obesity : Targets and Therapy. 2015;8:197-206. Epub 2015/05/01.
- 99. Mullu G, Abebe N. Assessment of parent adolescent communication on sexual and reproductive health issues associated with factors in Alemata High School, northern Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association February 26-28, 2015 Bahir Dar, 2015.
- 100.Mullu G, Abebe N, Berhane E. Prevalence or premarital sexual practice and associated factors among Alamata High School and Preparatory School adolescents, northern Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 101.Mullu G, Tsegaye G, Abebe N, Bogale W, Tadesse T, Amare D, et al. Age at first sexual initiation and associated risk factors among Debre Markos University students, northwest Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2005; Bahir Dar, 2015.
- 102.Mulu W, Yimer M, Zenebe Y, Abera B. Common causes of vaginal infections and antibiotic susceptibility of aerobic bacterial isolates in women of reproductive age attending at Felegehiwot Referral Hospital, Ethiopia: a cross sectional study. BMC Womens Health. 2015 May 13;15:42. Epub 2015/05/15.
- 103.Mulugeta B, Bjune G, Abebe F. Prevalence of tuberculosis, HIV, and TB-HIV co-infection among pulmonary tuberculosis suspects in a predominantly pastoralist area, northeast Ethiopia. Global Health Action. 2015;8(27949).
- 104.Mulugeta S, Tesfaye M, Hanlon C. Cognitive impairment and factors associated with it among people living with HIV in Jimma, southwest Ethioipa: a cross-sectional study. 26th Anual Conference of the Ethiopian Public Health Association; February 26-29, 2015 Bahir Dar, 2015.
- 105.Ngugi EN, Roth E, Mastin T, Nderitu MG, Yasmin S. Female sex workers in Africa: epidemiology overview, data gaps, ways forward. SAHARA J :

Journal of Social Aspects of HIV/AIDS Research Alliance. 2012;9(3):148-153. Epub 2012/12/15.

- 106.Nigussie M. Incidence and factors predicting active TB occurrence among patients enrolled in ART: retrospective cohort study, Addis Ababa. [MPH Thesis]: Addis Ababa University; 2015.
- 107.Omori R, Chemaitelly H, Abu-Raddad LJ. Dynamics of non-cohabiting sex partnering in sub-Saharan Africa: a modelling study with implications for HIV transmission. Sex Transm Infect. 2015 Sep;91(6):451-457. Epub 2015/03/10.
- 108.Padovese V, Racalbuto V, Barnabas GA, Morrone A. Operational research on the correlation between skin diseases and HIV infection in Tigray Region, Ethiopia. International Journal of Dermatology. 2015 01 Oct;54(10):1169-1174.
- 109.Pegurri E, Konings E, Crandall B, Haile-Selassie H, Matinhure N, Naamara W, et al. The missed HIVpositive children of Ethiopia. PLoS ONE. 2015 16 Apr;10 (4) (no pagination) (e0124041).
- 110.Riou J, Ait Ahmed M, Blake A, Vozlinsky S, Brichler S, Eholie S, et al. Hepatitis C virus seroprevalence in adults in Africa: a systematic review and meta-analysis. Journal of Viral Hepatitis. 2016 Apr;23(4):244-255. Epub 2015/10/20.
- 111.Ritmeijer K. Challenges of visceral leishmaniasis (VL) and HIV co-infection. Tropical Medicine and International Health. 2015 September;20:7-8.
- 112.Sahile Z, Mekuria M, Yared A. Comprehensive HIV/AIDS knowledge and sexual behavior among universitysStudents in Ambo, central Ethiopia: implication to improve intervention. Journal of Sexually Transmitted Diseases. 2015;2015:890202. Epub 2015/09/01.
- 113.Shimekaw B. Sexual violence in Bahir Dar: a crossectional institution based survey among private college female students. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 114.Shimelis T, Lemma K, Ambachew H, Tadesse E. Syphilis among people with HIV infection in southern Ethiopia: sero-prevalence and risk factors. BMC Infect Dis. 2015 Apr 17;15:189. Epub 2015/04/18.
- 115.Sibani D. Prevalence of oral *Candida albicans* carriage or infection in HIV sero positive adult patientss in the Armed Forces General and Teaching Hospital, Addis Ababa, Ethiopia. [MSc Thesis]: Addis Ababa Univerity; 2015.
- 116.Sisay S, Erku W, Medhin G, Woldeyohannes D. Perception of high school students on risk for acquiring HIV and utilisation of voluntary counselling and testing (VCT) service for HIV in Debre-Berhan town, Ethiopia: a quantitative crosssectional study. Sexually Transmitted Infections. 2015 September;91:A95.
- 117.Tadelle M, Fantahun N, Abebe L. Exclusive breastfeeding and maternal employment, a comparitive study among emlpoyed and non *Ethiop. J. Health Dev.* 2016;30(3)

employed mothers. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.

- 118. Tadesse G, Yakob B. Risky sexual behaviors among female youth in Tiss Abay, a semi-urban area of the Amhara Region, Ethiopia. PLoS One. 2015;10(3):e0119050. Epub 2015/03/05.
- 119. Tadesse H, Gotu B. Statitsical analysis of tuberculosis distribution and identifying risky areas: the case of North Shoa Zone, Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 120.Tadesse R. Sexual socialization of adolescents in Addis Ababa: the case of adolescents in Ayer Tena Secondary Preparatory School. [MA Thesis]: Addis Ababa University; 2015.
- 121.Tamene MM, Tessema GA, Beyera GK. Condom utilization among female sex workers in Gondar Town, northwest Ethiopia 2014. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 122.Tarkegn D. Assessment of the prevalence and associated factors of sexual violence among high school female students in Dilla Town. [MPH Thesis]: Addis Ababa University; 2015.
- 123. Tefera T. Tourism in Hamar, southwestern Ethiopia: stakeholders, interactions, and implication. [PhD Thesis]: Addis Ababa University; 2015.
- 124. Teferra TB, Erena AN, Kebede A. Prevalence of premarital sexual practice and associated factors among undergraduate health science students of Madawalabu University, Bale Goba, south east Ethiopia: institution based cross sectional study. Pan Afr Med J. 2015;20:209. Epub 2015/06/27.
- 125.Tenna E, Bezatu M, Frehiwot M, Wanzahun G. Determinants of active pulmonary tuberculosis in Ambo Hospital, west Ethiopia. African Journal of Primary Health Care and Family Medicine. 2015;7(1).
- 126.Tesema C, Tadesse T, Gebrehiwot M, Tsegaw A, Weldegebreal F. Environmental and host-related determinants of tuberculosis in Metema district, north-west Ethiopia. Drug, Healthcare and Patient Safety. 2015;7:87-95. Epub 2015/06/13.
- 127.Tilahun M, Worku A, Bogale A. Sexual initiation and factors associated with it among Addis Ababa University undergraduate students, Addis Ababa, Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 128. Tiruneh K, Wasie B, Gonzalez H. Sexual behavior and vulnerability to HIV infection among seasonal migrant laborers in Metema District, northwest Ethiopia: a cross-sectional study. BMC Public Health. 2015 Feb 11;15:122. Epub 2015/04/18.
- 129.Titiyos A, Jerene D. The yield of sceening symptomatic contacts of multidrug-resistant tuberculosis cases at a tertiary hospital in Addis Ababa, Ethiopia. 26th Annual Conference of the

Ethiopian Public Health Association; February 5-6, 2015; Bahir Dar, 2015.

- 130.Tsai AC. Socioeconomic gradients in internalized stigma among 4,314 persons with HIV in sub-Saharan Africa. AIDS and Behavior. 2015 01 Feb;19(2):270-282.
- 131.Tsegaye G. Prevalence rate of substance use, its association with selected socio-demographic factors and its impacts on education: the case of Halaba secondary school students. Indian Journal of Health and Wellbeing. 2015 Aug 2015 2015-10-03;6(8):752-756.
- 132.Tsegaye S. The human rights approach to sexual minority rights: the life experiences of gay, lesbian and bisexual Ethiopians living in Addis Ababa. [MA Thesis]: Addis Ababa University; 2015.
- 133.W/Giorgis Y. Exploring the general health situation and risk behaviors of street based female sex workers in Addis Ababa, Ethiopia.: Addis Ababa University; 2015.
- 134.Weldegebreal R, Melaku YA, Alemayehu M, Gebrehiwot TG. Unintended pregnancy among female sex workers in Mekelle city, northern Ethiopia: a cross-sectional study. BMC Public Health. 2015 Jan 31;15:40. Epub 2015/02/01.
- 135.Yimer FG, Asfaw HA, Adhena BM. Hepatitis B virus co-infection and its predisposing factors among HIV-positives at Karamara Hospital: cross-sectional study between ART naive and ART initiated. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 136.Yimer T, Gobena T, Egata G. Magnitude of domestic violence and associated factors among pregnant womenin Hulet Ejju Enessir District, northwest Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 137.Zenebe Y, Tibebu M, Tulu B, Mekonnen D, Mekonnen Z. Methicillin resistant *Staphylococcus aureus* and its associated factors among HIV positive paediatric patients in Amhara National Regional State, Ethiopia: a cross ectional study design. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.

Section 3: Impact Research

This section covers studies on the demographic, social, psychological, and economic impacts of HIV/AIDS on individuals, families, communities, institutions or the nation.

The HIV impacts examined in this Update include mental disorders associated with HIV/AIDS and tuberculosis (1,5,8,11,15), psychosocial adjustments of sexually abused persons (2,4), stress among caregivers of orphans (7), a comparative study of resilience and coping behavior of children affected by war and HIV/AIDS (13), the effect of HIV disclosure on ANC attendants (12), a

Ethiop. J. Health Dev. 2016;30(3)

longitudinal study of the impacts of maternal mortality on infant and child survival (9) and different levels of society (11), the potential contribution of gardening to increasing food security of HIV affected families (16) and a historical study of socio-behavioral factors and mourning during the early era of the AIDS epidemic (17).

Whereas most of these issues have been covered in previous updates, this particular update is unique in presenting four types of impacts. First, two studies examined the role of psychological disorders on healthseeking behavior and tuberculosis treatment outcome (1,15). Identifying these barriers need to be considered in optimizing long-term outcomes of TB as well as HIV care and treatment. The increase in the number of HIVinfected persons on ART and the persistently high TB rates make such studies particularly pertinent. Second, a longitudinal, multi-country study of the comprehensive impacts of maternal mortality at all levels of society and in areas not only in health but also economic opportunities and poverty (9) highlighted the need to understand these relationships in planning postmillennium strategies and programs. Third, Moucheraud et al. (10) examined the short-term impact of maternal mortality on infant survival, showing that the risk of infants dying within a month of their mother's death was 46 times higher than among infants whose mothers lived. In view of the currently declining AIDS mortality, there is an increasing need to determine the impact of ART on the wellbeing of mothers, infants and children. Fourth, the first gendered study of health related quality of life among PLHIV receiving ART, females had significantly lower quality of life scores than males in regard to perceived stigma, psychological support from family members and clinical parameters (14). These results indicate the need for measuring and monitoring clinical care of PLHIV receiving ART by using quality of life assessment tools. Also noteworthy is the study that reported low level of social support received and high stress experienced by caregivers of HIV/AIDS-affected orphans (7). In view of the continuing increase in the number of orphans and vulnerable children on ART in Ethiopia and the heavy reliance on kinship care, what needs to be considered is the author's recommendation that policy makers and social service providers should explicitly acknowledge the role of kinship care and strengthen their support that can reduce stress levels in care givers.

- Ambaw F, Mayston R, Hanlon C, Alem A. Depression among patients with tuberculosis: determinants, course and impact on pathways to care and treatment outcomes in a primary care setting in southern Ethiopia--a study protocol. BMJ Open. 2015 Jul 08;5(7):e007653. Epub 2015/07/15.
- 2. Belete K. Interpersonal reationship of rape victims who reside in Association for Sanctuary and

Development (AWSAD) [MA Thesis]: Addis Ababa University; 2015.

- 3. Dagnew Z, Jara D, Ketema K. Level of impacts of currently applied HIV and AIDS intervention practices 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 4. Dejene M. Female child sexual abuse and its association with traumatic symptoms and psychosocial adjustments in selected Ethiopian secondary schools. [PhD Thesis]: Addis Ababa University; 2015.
- Duko B, Gebeyehu A, Ayano G. Prevalence and correlates of depression and anxiety among patients with tuberculosis at Wolaita Sodo University Hospital and Sodo Health Center, Wolaita Sodo, south Ethiopia, cross sectional study. BMC Psychiatry. 2015 Sep 14;15:214. Epub 2015/09/16.
- Hunchak C, Teklu S, Meshkat N, Meaney C, Lisa Puchalski R. Patterns and predictors of early mortality among emergency department patients in Addis Ababa, Ethiopia. BMC Research Notes. 2015 2015 2016-04-16;8.
- Kassahun M. Social support and stress of kin caregivers of orphaned children in Addis Ababa, Ethiopia: implications for practice and policy. International Journal of Social Welfare. 2015 Oct;24(4):388-398.
- Mahir M, Bezatu M, Yadeta D, Wanzahun G. Prevalence of depression and associated factors among HIV patients seeking treatments in ART clinics at Harar town, eastern Ethiopia. Journal of AIDS and Clinical Research. 2015;6(6).
- Miller S, Belizan JM. The true cost of maternal death: individual tragedy impacts family, community and nations. Reprod Health. 2015 Jun 17;12:56. Epub 2015/06/18.
- Moucheraud C, Worku A, Molla M, Finlay JE, Leaning J, Yamin A. Consequences of maternal mortality on infant and child survival: a 25-year longitudinal analysis in Butajira Ethiopia (1987-2011). Reprod Health. 2015 May 06;12 Suppl 1:S4. Epub 2015/05/23.
- 11. Salilih SZ, Mosse NA. Common mental disorders and associated factors among HIV-infected individuals at ART Clinic of Debre Markos, Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 12. Tamiru B. Predictors and effects of HIV serostatus disclosure for pregnant women attending ANC. [MSc Thesis]: Addis Ababa University; 2015.
- 13. Tefera B. Resilience and coping compared between war and HIV/AIDS affected children and implications: experiences of children in eastern African countries. Indian Journal of Community Psychology. 2015 Sep;11(2):201-219.
- Tesfay A, Gebremariam A, Gerbaba M, Abrha H. Gender differences in health related quality of life *Ethiop. J. Health Dev.* 2016;30(3)

among people living with HIV on highly active antiretroviral therapy in Mekelle Town, northern Ethiopia. Biomed Res Int. 2015;2015:516369. Epub 2015/01/30.

- Tola HH, Shojaeizadeh D, Garmaroudi G, Tol A, Yekaninejad MS, Ejeta LT, et al. Psychological distress and its effect on tuberculosis treatment outcomes in Ethiopia. Glob Health Action. 2015;8:29019. Epub 2015/11/27.
- 16. Tsomondo D. A garden grown. PDN ; Photo District News. 2015 Feb 2015 2016-08-13;35(2):88-93.
- 17. Woubshet D. The Calendar of Loss: Race, Sexuality, and Mourning in the Early Era of AIDS. Baltimore, MD: Johns Hopkins University Press; US; 2015.

Section 4: Health Services and Health Policy Research This section includes reports on research and programmatic activities that aimed at expanding and improving the healthcare system including such issues as expansion of services for people living with HIV/AIDS, health resource economics and management, healthcare staff training, and national as well as international policies, laws, and guidelines for the provision of services and the protection of people living with HIV/AIDS, women, children, and other vulnerable groups.

This section is comprised of 43 journal articles, three published abstracts, two editorials, two conference presentations, two masters theses, and one PhD dissertation, one book, and one declaration document that deal with a range of issues related to health services that include: local commitment and innovation, adherence to guidelines, knowledge of treatment standards, cost assessment for HIV/AIDS, tuberculosis, STI, and reproductive health issues (1-56).

A number of articles in this category (1, 2, 4, 5, 12, 17, 20, 22, 32, 34, 38, 45, 48, 50, 54, 56) have dealt with strategies and health services related factors (including the referral system) for the promotion of the utilization of HIV/TB/STI and SHRH services. According to Abebe and Asnake (1), integrated multi-media interventions are important for the successful promotion of SRH related services. Abrahim et al. (2) highlight the importance of the efficiency of the referral system, while Alemayehu & Godana (4) emphasize the knowledge and practice of clinicians regarding syndromic management of STIs in the health seeking behaviors of clients. Ameyan and et al. (5) report the need for addressing numerous barriers to attract FSWs to HIV counseling and testing centers; while Deme and et al. (12) highlight the importance of the ANC program in rural areas in Ethiopia in detecting cases of hypertension, anemia, as well as HIV and STI. The findings of Dutta et al. (17) also emphasize the importance of symptomatic tuberculosis screening practice. According to Holcombe et al. (20) and Rehnstrom Loi et al. (38) attitudes and social as well as gender based reservations of health workers are

important in the provision of abortion services. The findings of Imiru (22) and Muntean et al. (34) indicate the inadequacy of information dissemination for creating awareness on youth reproductive health services. Miller et al. (32) reported the low quality of care provided to sick children by HEWs. Shiferaw et al. (45) advocate a single visit approach as a sustainable cervical prevention strategy; while Soressa (48) highlights the importance of peer education programs for engaging young people and other stakeholders in SRHR services. Thomas et al. (50) report on an organizational network analysis for improving referrals and integrating family planning and HIV, while Wissow et al. (54) describe a model for integrating mental health into hospital-based HIV treatment services in Ethiopia. According to Yakob and Ncama (56), issues such as financial fairness, perceived transportation convenience, employment status, and distance from the health facility are important factors to consider for improving quality of HIV treatment services.

Local and international level partnerships are among the main themes of papers in this category (6-8, 10, 11, 13, 14, 40, 43, 47). In terms of local commitment and innovation, an editorial for the Bulletin of the WHO (6) highlights the need for support and understanding by partner organizations and donors for local innovations for sustainable development; while according to a "Lancet World Report" (7) Ethiopia's political commitment along with an army of thousands of health extension workers is helping it to successfully tackle tuberculosis. Another editorial for the Bulletin of the WHO (8) highlights the needs for local government support to WHO's effort in introducing guidelines on the role, education and integration of community-based practitioners. Datiko and et al. (10) report the needs to develop context embedded strategies to support and motivate health extension workers within community based TB control approaches, while De Rosis (11) views the organization of the fight against HIV/AIDS in Ethiopia as the complex result of the interplay between government strategies and the efforts of affected people themselves to face their condition by seeking care and support. Similarly, Derbew et al. (13), in an abstract published in Annals of Global Health, describe the collaboration between the medical and nursing professions and its relevance for advancing health workforce capacity to meet the efforts of achieving the President's Emergency Plan for AIDS Relief (PEPFAR) goals in Ethiopia of meeting the health care needs of persons with HIV within the health care systems of developing countries. Dida and et al. (14) highlight the needs for discussion between health workers and students as being important factors for the utilization of reproductive health services among university students in southeast Ethiopia. According to Kok et al. (24), health extension workers' relationships with the community and health sector can be constrained as a result of inadequate support systems, lack of trust, communication and dialogue and differing expectations. The results of a pilot project reported by Robinson (40,

41) show the importance of integration of religious leaders in health education for increasing the number of ANC visits including the level of understanding about HIV transmission. Lönnroth et al. (26)'s action framework also highlight the needs for multi-sectorial approaches for addressing the social determinants of TB. Lunsford et al. (27) present cases for supproting close-tocommunity HIV/AIDS related services providers through a community health system approach. The Maputo Plan of Action (2016-2030) (43) also calls for African governments, civil society, the private sector, and all multi-sectoral development partners to join forces and redouble efforts for implementation of SRHR. Smith et al. (47) highlight the need to support national and international actors for sustained and strategic family planning advocacy.

There are some articles in this category (9, 18, 25, 33, 39) that dealt with costing, financing as well as inequalities with regard to HIV/AIDS and SHRH services. Cooke (9) highlighted the potential of low cost IVF for contribution toward reducing the impact of infertility, particularly in low resource economies. Grosso et al. (18) reported underfunding and fewer resources on HIV programs serving MSM, especially in countries that criminate these groups; while according to Resch et al. (39), the prospect for HIV/AIDS funding in lower income countries, such as Mozambique and Ethiopia, would remain heavily dependent on donor funds. According to Laurence et al. (25), global cost related data for MDR-TB treatment are very limited as opposed to those for DS-TB treatment. Muluneh et al. (33) reported the existence of wealthbased inequalities in the provision of FP services in Ethiopia despite efforts to provide contraceptives for free at all public health facilities.

Issues related to unmet and client needs and utilization of family planning as well as TB related services were also subjects for some articles (23, 30, 31, 51, 55, 57). According to Balew et al. (23), there is a big variation in family planning use both at the individual and group levels; while Feyissa and Melka (30) reported high unmet need for modern FP in HIV-positive married women in western Ethiopia. Low case notification rates for childhood TB were reported by Mesay et al. (31) and Tulloch et al. (51), found that poor rural populations and women are at high risk of unmet health needs in terms of TB diagnoses. Woldie et al. (55) examined the nature of client needs and community centered home based care services provided for PLHIV and OVC in Ethiopia.

Two articles (3, 19) also indicated the decreasing trend in the prevalence of HIV/AIDS. Takele Mena (3) demonstrated a statistically significant decline in the total and HIV/AIDS related mortality between November 2005 and October 2013 among teachers at public primary and secondary schools in Addis Ababa. Haile Amlak's editorial (19) commended Ethiopia's progress in achieving MDG goals, especially in indicators related to child health and HIV/AIDS.

There were three articles (37, 44, 46) that paid attention to quality of laboratories for HIV/AIDS, TB, and STI. Putoto et al. (37) presented a harmonized model for African health laboratories with a list of procedures to challenge the major health problems – HIV/AIDS, malaria, tuberculosis (TB) – at each level of pyramidal health system. Shiferaw et al. (44) highlighted the need for strengthening external quality assessment programs and technical support for tuberculosis diagnostic laboratory systems. Sinishaw et al. (46) emphasized on the need for strengthening supply chain management for uninterrupted TB diagnostic service.

Adoption of approaches and guidelines for treatment were also subjects of a number of articles (15, 16, 50) within this category. Doherty and et al. (15) stressed that progress towards ending the AIDS epidemic by 2030 depends on the adoption and implementation of global guidelines with evidenced based approaches to treating people with HIV. According to Dutta et al. (16), the projected number of people receiving ART across three scenarios suggests that countries are unlikely to meet the 90-90-90-treatment target (81% of people living with HIV on ART by 2020) unless they adopt a test-and-offer approach and increase ART coverage. Summoro et al. (49) emphasized the need for improving prescribing patterns in developing countries including the reduction of injection use to prevent transmission of infections like HIV and other blood-borne pathogens.

There are also a few articles (21, 28, 29) related to motivation and incentives of health workers. According to Hotchkiss et al. (21) both financial and non-financial factors are important determinants of health worker motivation in the Ethiopian context. Maes (28) showed how positive change within HIV/AIDS treatment programs requires ethnographic analysis of how CHWs exercise capacities, also highlighting (29) the international admiration given to the salaried HEW system of Ethiopia.

Importance of family planning services to development by Osotimehin (36) is highlighted in the paper while effectiveness of PMTCT intervention by Abadula (42) was reported. A book by Anke van der Kwaak & Madeleen Wgelin-Schuringa (eds) (52) presented strategies, approaches, and tools to mainstream concerns of gender-equality in the formulation of health policy and practice.

1. Abebe S, Asnake M. Using integrated multimedia intervention as a key strategy for promoting sexual reproductive health (SRH) issues: experience of integrated family health program (IFHP) in Ethiopia. 26th Annual Conference of the Ethiopan Public Health Association; February 26-28, 2015; Bahir Dar2015.

- 2. Abrahim O, Linnander E, Mohammed H, Fetene N, Bradley E. A patient-centered understanding of the referral system in Ethiopian primary health care units. PLoS ONE. 2015;10(10).
- Adilo TM. The effect of HIV/AIDS on the education sector and the response to mitigate the epidemic in Addis Ababa, Ethiopia. [MPH Thesis]: Addis Ababa University; 2015.
- Alemayehu A, Godana W. Knowledge and practice of clinicians regarding syndromic management of sexually transmitted infections in public health facilities of Gamo Gofa Zone, south Ethiopia. Journal of Sexually Transmitted Diseases. 2015;2015:310409. Epub 2015/11/26.
- 5. Ameyan W, Jeffery C, Negash K, Biruk E, Taegtmeyer M. Attracting female sex workers to HIV testing and counselling in Ethiopia: a qualitative study with sex workers in Addis Ababa. African Journal of AIDS Research. 2015 03 Apr;14(2):137-144.
- Balcha T, Getahun H, Admasu K. Local innovations and country ownership for sustainable development. Bulletin of the World Health Organization. 2015 Nov 2015 2015-11-11;93(11):742.
- Burki T. Ethiopia could be a model country for tuberculosis control. The Lancet. 2015 2015 Dec 05 2016-01-18;386(10010):2241-2242.
- Campbell J, Admasu K, Soucat A, Tlou S. Maximizing the impact of community-based practitioners in the quest for universal health coverage. Bulletin of the World Health Organization. 2015 Sep 01;93(9):590-590A. Epub 2015/10/20.
- Cooke ID. Development and status of low-cost IVF. International Journal of Gynecology and Obstetrics. 2015 October;131:E45.
- Datiko DG, Yassin MA, Tulloch O, Asnake G, Tesema T, Jamal H, et al. Exploring providers' perspectives of a community based TB approach in Southern Ethiopia: implication for community based approaches. BMC Health Services Research. 2015 Nov 09;15:501. Epub 2015/11/11.
- De Rosis C. The Organization of the Fight against HIV/AIDS in Ethiopia: rallying around Afflictions. Northeast African Studies. 2015 2015 2015-06-08;15(1):67-111.
- 12. Deme C, Edao B, Jaya G, Reyes F, Alegria I, Barreiro P, et al. Antenatal care program including screening of syphilis, hepatitis B virus infection and HIV in pregnant women attending in a rural hospital southern Ethiopia. Tropical Medicine and International Health. 2015 September;20:281.
- Derbew M, Lulseged S, Haile Mariam D. Collaborations between MEPI and NEPI at Addis Ababa University. Annals of Global Health. 2015 January-February;81 (1):145.
- 14. Dida N, Darega B, Takele A. Reproductive health services utilization and its associated factors among

Madawalabu University students, southeast Ethiopia: cross-sectional study. BMC Res Notes. 2015 Jan 17;8:8. Epub 2015/01/18.

- 15. Doherty M, Beusenberg M, Asamoah-Odei E, Lule F, Pendse R, Ghidinelli M, et al. Rapid uptake and adoption of the WHO 2013 consolidated ARV guideline recommendations: paving the way to achieving the 90/90/90 global target. Journal of the International AIDS Society. 2015;18:26-27.
- 16. Dutta A, Barker C, Kallarakal A. The HIV Treatment gap: estimates of the financial resources needed versus available for scale-up of antiretroviral therapy in 97 countries from 2015 to 2020. PLoS Medicine. 2015 Nov;12(11):e1001907; discussion e1001907. Epub 2015/11/26.
- 17. Gebregeorgis GB, Alemneh M, Negese D, Kassie Y, Assefa M, Ayalew W, et al. Poor symptomatic tuberculosis screening practice in quarter of health centers in Amhara Region, Ethiopia. 26th Annual Conference of the Ethiopan Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Grosso A, Ryan O, Tram KH, Baral S. Financing the response to HIV among gay men and other men who have sex with men: case studies from eight diverse countries. Glob Public Health. 2015;10(10):1172-1184. Epub 2015/07/04.
- Haileamlak A. Ethiopia successfully attaining the millennium development goals. Ethiopian Journal of Health Sciences. 2015 Apr;25(2):109-110. Epub 2015/07/01.
- 20. Holcombe SJ, Berhe A, Cherie A. Personal beliefs and professional responsibilities: Ethiopian midwives' attitudes toward providing abortion services after legal reform. Studies in Family Planning. 2015 Feb 2015 2016-12-01;46(1):73-95.
- 21. Hotchkiss DR, Banteyerga H, Tharaney M. Job satisfaction and motivation among public sector health workers: evidence from Ethiopia. Hum Resour Health. 2015 Oct 29;13:83. Epub 2015/10/30.
- 22. Imiru S. Effectiveness of youth reproductive health services in government based youth centers to promote youth reproductive health. [MA Thesis]: Addis Ababa University; 2015.
- 23. Jembere Gizachew B, Cho Y, Kim CT, Ko W. Structural Determinants in Family Planning Service Utilization in Ethiopia: EDHS 2011 Analysis. BioMed Research International. 2015 2015 2015-12-28.
- 24. Kok MC, Kea AZ, Datiko DG, Broerse JE, Dieleman M, Taegtmeyer M, et al. A qualitative assessment of health extension workers' relationships with the community and health sector in Ethiopia: opportunities for enhancing maternal health performance. Hum Resour Health. 2015 Sep 30;13:80. Epub 2015/10/02.
- 25. Laurence YV, Griffiths UK, Vassall A. Costs to health services and the patient of treating tuberculosis: a systematic literature review. *Ethiop. J. Health Dev.* 2016;30(3)

Pharmaco Economics. 2015 Sep;33(9):939-955. Epub 2015/05/06.

- Lönnroth K, Migliori GB, Abubakar I, D'Ambrosio L, de Vries G, Diel R, et al. Towards tuberculosis elimination: an action framework for low-incidence countries. The European Respiratory Journal. 2015 Apr;45(4):928-952. Epub 2015/03/21.
- Lunsford SS, Fatta K, Stover KE, Shrestha R. Supporting close-to-community providers through a community health system approach: case examples from Ethiopia and Tanzania. Hum Resour Health. 2015 Mar 28;13:12. Epub 2015/04/18.
- Maes K. "Volunteers are not paid because they are priceless": community health worker capacities and values in an AIDS treatment intervention in urban Ethiopia. Medical Anthropology Quarterly. 2015 Mar 2015 2015-03-18;29(1):97.
- 29. Maes K, Closser S, Vorel E, Tesfaye Y. A women's development army: narratives of community health worker investment and empowerment in rural Ethiopia. Studies in Comparative International Development. 2015 Dec 2015 2015-12-03;50(4):455-478.
- Melka AS, Feyissa TR. Demand for modern family planning among married women living with HIV in western Ethiopia. Contraception. 2015 October;92 (4):386-387.
- Mesay HD, Daniel Gemechu D, Lindtjorn B. Low case notification rates of childhood tuberculosis in southern Ethiopia. BMC Pediatrics. 2015 2015 2016-04-16;15.
- 32. Miller NP, Amouzou A, Hazel E, Degefie T, Legesse H, Tafesse M, et al. Assessing the quality of sick child care provided by community health workers. PLoS One. 2015 Nov 2015 2015-11-14;10(11).
- 33. Muluneh Y, Zakus D, Yehualashet T, Muluked D, Mesganaw F. Paving the way for universal family planning coverage in Ethiopia: an analysis of wealth related inequality. International Journal for Equity in Health. 2015;14(77).
- 34. Muntean N, Kereta W, Mitchell KR. Addressing the sexual and reproductive health needs of young people in Ethiopia: an analysis of the current situation. African Journal of Reproductive Health. 2015 Sep 2015 2015-11-21;19(3):87-99.
- Oshone A. The African Union's responsibility to protect and respond to conflict related sexual violence. [MA Thesis]: Addis Ababa University; 2015.
- Osotimehin B. Family planning as a critical component of sustainable global development. Global Health Action. 2015 2015 2016-01-18;8.
- Putoto G, Cortese A, Pecorari I, Musi R, Nunziata E. Harmonization of clinical laboratories in Africa: a multidisciplinary approach to identify innovative and sustainable technical solutions. Diagnosis. 2015 01 Jun;2(2):129-135.

- 38. Rehnstrom Loi U, Gemzell-Danielsson K, Faxelid E, Klingberg-Allvin M. Health care providers' perceptions of and attitudes towards induced abortions in sub-Saharan Africa and Southeast Asia: a systematic literature review of qualitative and quantitative data. BMC Public Health. 2015 Feb 12;15:139. Epub 2015/04/18.
- 39. Resch S, Ryckman T, Hecht R. Funding AIDS programmes in the era of shared responsibility: An analysis of domestic spending in 12 low-income and middle-income countries. The Lancet Global Health. 2015 01 Jan;3(1):e52-e61.
- 40. Robinson E. Religious leaders as health educators: a pilot project in northern Ethiopia. Annals of Global Health. 2015 January-February;81 (1):168.
- 41. Robinson E. Incorporating religious leaders into the HIV care continuum in northern Ethiopia: evaluation of a pilot project and development of a scale up plan with a focus on sustainability. Annals of Global Health. 2015 January-February;81 (1):62-63.
- 42. Seid MA. Assessment of effectiveness of prevention of mother to child transmisiosn of HIV-positive women in Arsi University, Assela Health and Referral Hospital. [MSc Thesis]: Addis Ababa University; 2015.
- 43. Session AUCoMoHS. Universal access to comprehensive sexual and reproductive health services in Africa. Maputo Plan of Action for the Operationalisation of the Continental Policy Framework for Sexual and Reproductive Health and Rights, 2007-2010. Addis Ababa: African Union, 2006.
- 44. Shiferaw MB, Hailu HA, Fola AA, Derebe MM, Kebede AT, Kebede AA, et al. Tuberculosis laboratory diagnosis quality assurance among public health facilities in West Amhara Region, Ethiopia. PLoS One. 2015 Sep 2015 2015-10-10;10(9).
- 45. Shiferaw N, Salvador-Davila G, Kassahun K, Tilahun Y, Weldegebreal T, Ahmed I, et al. Client satisfaction with the single-visit approach service for cervical cancer prevention in Ethiopia. International Journal of Gynecology and Obstetrics. 2015 October;131:E557-E558.
- 46. Sinishaw MA, Gebregergs GB, Shiferaw MB. Distribution and availability of essential tuberculosis diagnostic items in Amhara Region, Ethiopia. PLoS One. 2015;10(12):e0141032. Epub 2015/12/08.
- 47. Smith E, Musila R, Murunga V, Godbole R. An assessment of family planning decision makers' and advocates' needs and strategies in three East African countries. International Perspectives on Sexual and Reproductive Health. 2015 Sep 2015 2015-10-30;41(3):136-144.
- 48. Soressa M. Smart investment: young peer educators serving as a bridge to link demand and supply of sexual and reproductive health services in Southern Region, Ethiopia. International Journal of Gynecology and Obstetrics. 2015 October;131:E77.

- 49. Summoro TS, Gidebo KD, Kanche ZZ, Woticha EW. Evaluation of trends of drug-prescribing patterns based on WHO prescribing indicators at outpatient departments of four hospitals in southern Ethiopia. Drug Design, Development and Therapy. 2015;9:4551-4557. Epub 2015/08/27.
- Thomas JC, Reynolds HW, Alterescu X, Bevc C, Tsegaye A. Improving referrals and integrating family planning and HIV services through organizational network strengthening. Health Policy Plan. 2016 Apr;31(3):302-308. Epub 2015/07/03.
- 51. Tulloch O, Theobald S, Morishita F, Datiko DG, Asnake G, Tesema T, et al. Patient and community experiences of tuberculosis diagnosis and care within a community-based intervention in Ethiopia: a qualitative study. BMC Public Health. 2015 Feb 25;15:187. Epub 2015/04/18.
- 52. van der Kwaak A, Wegelin-Schuringa M. Gender and Health: Policy and Practice. a Global Sourcebook. Amsterdam, Netherlands, KIT Publishers, 2006.
- 53. Weldegebreal F, Seyoum B, Oljira L. Factors affecting acceptance of provider-initiated HIV testing and counseling services among outpatient clients in selected health facilities in Harar town, eastern Ethiopia. HIV/AIDS - Research and Palliative Care. 2015 15 May;7:157-165.
- 54. Wissow LS, Tegegn T, Asheber K, McNabb M, Weldegebreal T, Jerene D, et al. Collaboratively reframing mental health for integration of HIV care in Ethiopia. Health Policy and Planning. 2015 Jul;30(6):791-803.
- 55. Woldie M, Sudhakar M, Feyissa GT, Aaantje C, Quinlan T. Community home based care: an overview of client needs, actors and services provided in Ethiopia. 26th Annual Conference of the Ethiopian Public health Association; February 26-28, 2015; Bahir Dar, 2015.
- 56. Yakob B, Ncama BP. Perceived quality of HIV treatment and care services in Wolaita Zone of southern Ethiopia: a cross-sectional study. BMJ Open. 2015;5(12).

Section 5: Prevention Research

This section includes reports on research and programmatic activities that aimed at provision of prevention services targeted at HIV/AIDS and related opportunistic infections. Information and behavioral change communication, provision of voluntary testing and counseling and prevention of mother-to-child transmission, community mobilization, and other riskreduction efforts against HIV/AIDS are studies included in this section.

This section contains 46 references, considerably fewer than in the 2014 Update. Most references fell again into the testing and counseling category (1,3,5,9,10,12,13,15,17,21,22,27,28,29,33,34, 43-45), followed by studies of information, education and communication (IEC) (7,14,20,30,32, 35,39,40, the use of contraceptives (8,19,25,37,40,41), infant and child feeding (23,24,26,45), family planning (4,16,47), and 6 studies on various other subjects including the effectiveness of preventive therapy in TB (6) and integration of preeclampsia and anemia for ANC clients (11). In a study of provider-initiated HIV counseling and testing (PICT), 27% of respondents said that one should be tested anytime and only 8% stated that testing should be done when one is sick, indicating that counseling and testing has been able to increase knowledge about HIV and its prevention and treatment in recent years (1). The continued interest by researchers in HIV testing and counseling is encouraging since testing and counseling service utilization is still relatively low in Ethiopia and inadequately understood.

Social stigma was identified as a major barrier to PICT by pregnant women but more rural than urban women in Assosa Town accepted PICT (2). Marelign and Shikur (30) reported socio-demographic barriers to husbands participating in PMTCT commonly found by earlier investigators. A study of urban ANC clients identified discrimination by their husbands as a barrier to the use of Option B+ PMTCT (10). Use of voluntary counseling and testing services (VCT) in a northwestern highland community was similarly impeded by stigmatization and discrimination but use rates increased 4.2 fold if husbands came with their wives to clinics (12). HIVpositive pregnant women in hospitals in Tigray Region, on the other hand, reported that HIV status disclosure, in addition to counseling on medication, was a positive predictor of adherence to Option B+ PMTCT (15). In a community-based study of HIV-positive mothers in Addis Ababa, peer support from other PLHIV, faith and increasing optimism were reported the main facilitators of PMTCT uptake (17). One of the first studies of the timing and predictors of loss to follow-up in the Option+ PMTCT program revealed that younger women, those attending hospitals rather than health centers, patients starting ART on the day of their diagnosis, and missing CD4 information at ART initiation were lost to follow-up (33). Workagegn et al., using the health belief model, showed that the low rate of PMTCT services in government health centers in Addis Ababa was due to the perception that the service was not beneficial (45). Several studies reported on the use of TB tests (5) and testing services (6, cervical cancer screening (9,21).

The remaining studies focused on IEC, family planning, and infant and child feeding, among others. IEC studies dealt with HIV communication disclosure of HIV positive status (22), parent/adolescent sexual and reproductive health communication (14), and the use of newspapers and peer education in high schools in influencing sexual behavior (30,32) and of community conversation in facilitating the utilization of testing services (35). The potential use of traditional songs as media in HIV prevention was explored by Bekalu and Eggermont (7), who identified 23 AIDS songs serving that purpose. Of the two studies of family planning (4,15), the one by Adamchak et al. (4) developed a system to monitor family planning and HIV service integration using existing data from health facilities in Ethiopia, Rwanda, Tanzania and Uganda. The system is expected to allow for monitoring the integration of family planning and HIV services but requires further improvements in daily and active client counts and recording. Three studies examined attitudes and other factors associated with infant and child feeding practices among HIV-positive mothers (23,26,47) and one study among non-infected mothers (24). Whereas the infant and young child feeding practices reported by Worku from Gondar Zone met the WHO recommendations (46), complementary feeding practices in southern Ethiopian communities were considered inadequate (23). The two MA theses focusing on breastfeeding (24,26) were not available online. Norheim et al. (36) analyzed the decline in mortality world-wide and found a 6.8% decline in under-5 mortality and a decline in TB and malaria mortality by half between 1990 and 2010, three years before the MDG4. They discuss these changes within the context of broadly based socioeconomic changes and government health programs and estimated that the remaining risks of premature deaths, particularly injuries and non-communicable diseases, will be halved again between 2010 and 2030.

- Abdurahman S, Seyoum B, Oljira L, Weldegebreal F. Factors affecting acceptance of provider-initiated HIV testing and counseling services among outpatient clients in selected health facilities in Harar Town, Eastern Ethiopia. HIV/AIDS (Auckland, NZ). 2015;7:157-165. Epub 2015/06/02.
- 2. Abebe AL. Interpretative phenomenological analysis of street children's resilience: striving to stay healthy in an unhealthy environment. [MA Thesis]: Addis Ababa University; 2015.
- Abtew S, Awoke W, Asrat A. Acceptability of provider-initiated HIV testing as an intervention for prevention of mother to child transmission of HIV and associated factors among pregnant women attending at Public Health Facilities in Assosa town, northwest Ethiopia. BMC Res Notes. 2015 Nov 09:8:661. Epub 2015/11/11.
- Adamchak SE, Okello FO, Kabore I. Developing a system to monitor family planning and HIV service integration: results from a pilot test of indicators. The Journal of Family Planning and Reproductive Health Care. 2016 Jan;42(1):24-29. Epub 2015/10/02.
- Asmar S, Drancourt M. Rapid culture-based diagnosis of pulmonary tuberculosis in developed and developing countries. Frontiers in Microbiology. 2015;6:1184. Epub 2015/11/19.
- 6. Assebe LF, Reda HL, Wubeneh AD, Lerebo WT, Lambert SM. The effect of isoniazid preventive therapy on incidence of tuberculosis among HIV-

infected clients under pre-ART care, Jimma, Ethiopia: a retrospective cohort study. BMC Public Health. 2015;15:346.

- 7. Bekalu MA, Eggermont S. Aligning HIV/AIDS communication with the oral tradition of Africans: a theory-based content analysis of songs' potential in prevention efforts. Health Communication. 2015 May;30(5):441-450.
- 8. Bekele T, Walabu M, Gebremariam A, Tura P. Contraceptive discontinuation and associated factors among women on marriage in Agarfa District, Bale Zone, southeast Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Belete N, Tsige Y, Mellie H. Willingness and acceptability of cervical cancer screening among women living with HIV/AIDS in Addis Ababa, Ethiopia: a cross sectional study. Gynecologic Oncology Research and Practice. 2015;2:6. Epub 2015/01/01.
- BG/Egziabher A. Assessment of knowledge and attitude about PMTCT Option B+ and associated factors among ANC clients in Dessie Twon, South Wollo Amhara Regional State. [MA thesis]: Addis Ababa University; 2015.
- 11. Birhanu Z, Kebede Y, Chapleau G, Dicksin K. Integrating strategies for the prevetion of preclampsia and anemia into community-based programs in Ethiopia: A formative assessment reslt. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahis Dar, 2015.
- 12. Bruke G, Bizuayehu HM, Muluken T. Voluntary HIV counseling and testing service utilization among pregnant mothers in north west Ethiopia in 2014. Journal of AIDS and Clinical Research. 2015;6(3).
- 13. Deksissa ZM, Tesfamichael FA, Ferede HA. Prevalence and factors associated with VIA positive result among clients screened at Family Guidance Association of Ethiopia, south west area office, Jimma model clinic, Jimma, Ethiopia 2013: a crosssectional study. BMC Res Notes. 2015 Oct 29;8:618. Epub 2015/10/31.
- 14. Dessie Y, Berhane Y, Worku A. Parent-adolescent sexual and reproductive health communication is very limited and associated with adolescent poor behavioral beliefs and subjective norms: evidence from a community based cross-sectional study in eastern Ethiopia. PLoS One. 2015;10(7):e0129941. Epub 2015/07/15.
- Ebuy H, Yebyo H, Alemayehu M. Level of adherence and predictors of adherence to the Option B+ PMTCT programme in Tigray, northern Ethiopia. International Journal of Infectious Diseases. 2015 April 01;33:e123-e129.
- 16. Feyssa MD, Tsehay YB, Tadesse AW. Unmet need for family planning among women in HIV/AIDS care at antiretroviral treatment clinic in south Ethiopia: a challenge to prevention of mother to *Ethiop. J. Health Dev.* 2016;30(3)

child transmission. Journal of AIDS and Clinical Research. 2015;6(6).

- Fleek KA. Perspectives of HIV+ women on the prevention of mother to child transmission of HIV in Addis Ababa, Ethiopia. Dissertation Abstracts International: Section B: The Sciences and Engineering. 2015;76(4-B(E)):No pagination specified.
- 18. Gebrehiwot TG, Ashebir Y, Belayneh M, G/Mariam S, Abebe Z, Birhan E, et al. The experience of Mekelle University STI Confidential Clinic on provision of promotive, preventive and curative intervention for commercial sex workers. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Gelagay AA, Koye DN, Yeshita HY. Demand for long acting contraceptive methods among married HIV positive women attending care at public health facilities at Bahir Dar City, northwest Ethiopia. Reprod Health. 2015 Aug 27;12:76. Epub 2015/08/28.
- 20. Genet M, Sebsibie G, Gultie T. Disclosure of HIV seropositive status to sexual partners and its associated factors among patients attending antiretroviral treatment clinic follow up at Mekelle Hospital, Ethiopia: a cross sectional study. BMC Research Notes. 2015;8:109.
- 21. Getinet M, Gelaw B, Sisay A, Mahmoud EA, Assefa A. Prevalence and predictors of Pap smear cervical epithelial cell abnormality among HIV-positive and negative women attending gynecological examination in cervical cancer screening center at Debre Markos referral hospital, East Gojjam, northwest Ethiopia. BMC Clin Pathol. 2015;15:16. Epub 2015/09/25
- Haftamu E, Henock Y, Mussie A. Level of adherence and predictors of adherence to the Option B+ PMTCT programme in Tigray, northern Ethiopia. International Journal of Infectious Diseases. 2015;33:123-129.
- Haile D, Belachew T, Berhanu G, Setegn T, Biadgilign S. Complementary feeding practices and associated factors among HIV positive mothers in southern Ethiopia. Journal of Health, Population, and Nutrition. 2015 May 01;34:5. Epub 2016/01/31.
- 24. Hailu H. Assessment of knowledge, attitude and practice of breast feeding among mothers attending public health institutions in Addis Ababa. [MPH Thesis]: Addis Ababa University; 2015.
- 25. Hounton S, Barros AJD, Amouzou A, Shiferaw S, Maïga A, Akinyemi A, et al. Patterns and trends of contraceptive use among sexually active adolescents in Burkina Faso, Ethiopia, and Nigeria: evidence from cross-sectional studies. Global Health Action. 2015 2015 2016-01-18;8.
- 26. Ketema Z. Associated factors of exclusinve bearst feeding among HIV-positive women in selcted health facilities of Adama City Administrative

Region, Ethiopia. [MPH Thesis]: Addis Ababa University; 2015.

- 27. Klaus K, Baldwin J, Izurieta R, Naik E, Seme A, Corvin J, et al. Reducing PMTCT attrition: perspectives of HIV+ women on the prevention of mother-to-child HIV services in Addis Ababa, Ethiopia. Ethiopian Medical Journal. 2015 01 Apr;53(2):91-104.
- Legesse B. Quality and implementation barriers of prevention of mother to child transmission of HIV program in Ethiopia: a cross-sectional study. Value in Health. 2015 May;18 (3):A275-A276.
- 29. Marelign T, Shikur M. Male partners' involvement in the prevention of mother-to-child transmission of HIV and associated factors in Arba Minch town and Arba Minch Zuria Woreda, southern Ethiopia. BioMed Research International. 2015;763876(10).
- Melese E, Abebe S. The effectiveness of newspaper towards influencing behavior change of taxi community in Addis Ababa: the case of "Sechento" Newspaper. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 31. Menna T, Ali A, Worku A. Factors associated with HIV counseling and testing and correlations with sexual behavior of teachers in primary and secondary schools in Addis Ababa, Ethiopia. HIV/AIDS - Research and Palliative Care. 2015 30 Jun;7:197-208.
- 32. Menna T, Ali A, Worku A. Effects of peer education intervention on HIV/AIDS related sexual behaviors of secondary school students in Addis Ababa, Ethiopia: a quasi-experimental study. Reprod Health. 2015 Sep 07;12:84. Epub 2015/09/09.
- Mitiku I, Arefayne M, Mesfin Y, Gizaw M. Factors associated with loss to follow-up among women in Option B+ PMTCT programme in northeast Ethiopia: a retrospective cohort study. Journal of the International AIDS Society. 2016 2016;19(1):20662.
- 34. Molla G, Huruy A, Mussie A, Wondowosen T. Factors associated with HIV counseling and testing among males and females in Ethiopia: evidence from Ethiopian Demographic and Health Survey data. Journal of AIDS and Clinical Research. 2015;6(3).
- 35. Nigatu YT, Abera S, Mekonnen MG, Melesse WN. The role of the community conversation approach in facilitating HIV/AIDS competence and utilisation of testing services in Africa: the case of Ethiopia. African Journal of AIDS Research. 2015 02 Oct;14(4):295-301.
- 36. Norheim OF, Jha P, Admasu K, Godal T, Hum RJ, Kruk ME, et al. Avoiding 40% of the premature deaths in each country, 2010-30: review of national mortality trends to help quantify the UN sustainable development goal for health. Lancet (London, England). 2015 Jan 17;385(9964):239-252. Epub 2014/09/23.
- Pulerwitz J, Hughes L, Mehta M, Kidanu A, Verani F, Tewolde S. Changing gender norms and reducing *Ethiop. J. Health Dev.* 2016;30(3)

intimate partner violence: results from a quasiexperimental intervention study with young men in Ethiopia. American Journal of Public Health. 2015 Jan 2015 2015-02-14;105(1):132-137.

- 38. Salih NA, Metaferia H, Reda AA, Biadgilign S. Premarital sexual activity among unmarried adolescents in northern Ethiopia: a cross-sectional study. Sexual and Reproductive Healthcare. 2015 01 Mar;6(1):9-13.
- Sambe MM. Students talk about their HIV/AIDS education courses: a case of Addis Ababa, Ethiopia. Journal of International Education Research. 2015 2015 2015-03-03;11(1):35-n/a.
- 40. Takele M, Ahmed A, Alemayehu W. Effects of peer education intervention on HIV/AIDS related sexual behaviors of secondary school students in Addis Ababa, Ethiopia: a quasi-experimental study. Reproductive Health. 2015;12(84).
- Tamene MM, Tessema GA, Beyera GK. Condom utilization and sexual behavior of female sex workers in northwest Ethiopia: a cross-sectional study. Pan Afr Med J. 2015;21:50. Epub 2015/09/26.
- 42. Terefe G, Antehun A. Consistent condom use and preference of HIV risk preventive interventions among undergraduate university students in southern Ethiopia: a cross sectional study. Journal of AIDS and Clinical Research. 2015;6(4).
- Tilahun M, Mohamed S. Male partners' involvement in the prevention of mother-to-child transmission of HIV and associated factors in Arba Minch Town and Arba Minch Zuria Woreda, southern Ethiopia. BioMed Research International. 2015;2015 (no pagination)(763876).
- 44. Titiyos A, Jerene D, Enquselasie F. The yield of screening symptomatic contacts of multidrug resistant tuberculosis cases at a tertiary hospital in Addis Ababa, Ethiopia: Descriptive cross-sectional study. 26th Annual Conference of the Ethiopian Public Health Association; February 26-18, 2015; Bahir Dar, 2015.
- 45. Workagegn F, Kiros G, Abebe L. Predictors of HIVtest utilization in PMTCT among antenatal care attendees in government health centers: institutionbased cross-sectional study using health belief model in Addis Ababa, Ethiopia, 2013. HIV/AIDS -Research and Palliative Care. 2015 13 Jul;7:215-222.
- 46. Worku A, Ejigu T, Berhanu D. Assessment of infant and young child feeding pracices in HIV-positive mothers and its associated factors in South Gondar Zone, northwest Ethiopia. 26th Annual Conference of the Ethiopain Public Health Association; February 5-6, 2015; Bahir Dar, 2015.
- 47. Yigzaw M, Zakus D, Tadesse Y, Desalegn M, Fantahun M. Paving the way for universal family planning coverage in Ethiopia: an analysis of wealth related inequality. International Journal for Equity in Health. 2015 14 Sep;14 (1): .

Section 6: Treatment, care, and Clinical Research

This section includes studies on the characteristics and clinical course of HIV infection and opportunistic infections, treatment to AIDS and opportunistic infections, effects and outcomes associated with treatment, clinical and non-clinical care and supportive services provided to people living with HIV/AIDS.

The Treatment, Care and Clinical Research section of this update pulled all relevant literature. The major ones were on HIV co-morbidity with long established TB and relatively new studies in the series of update such as Leishmaniasis, malaria, anemia and diabetes. Besides, studies on ART adherence, treatment of opportunistic infections, care and support, domains related to fertility, nutrition and intestinal parasites in connection to HIV and ART, gender differentials to quality of life and survival, pediatric ART, outcomes of ART. These works were publications in peer review journals, conference abstracts and student thesis work. Of the 162 materials accessed, 124 were published documents, 31 were conference abstracts and the remaining 7 were master's thesis all of which were from Addis Ababa University. Some of the major entries under this section are highlighted below.

In this section of the update, HIV co-morbidity with TB, malaria, Leishmaniasis, anemia and diabetes are predominate – 23% of all collection under this section. The studies focused on case detection; prevalence of specific co-infection; magnitudes; risk factors, predictors and determinants. The majority of the collection were on TB-HIV co-infection (2,8,11,17,20,28-9,31,39,59,60,67,68,86,90,102,106,111,112,120,1232,14 4,151), while there are studies on malaria (13,78); Leishmaniasis, (51,52,64), anemia (26,61,67,69,114,142) and diabetes (154).

Next set of collection constituting 15% of all collection under this section focuses on the outcomes of ART and HAART. The studies emphasized mainly on the outcome of ART use including survival and mortality, immunologic failure, predicators and determinants, implications of late start of treatment. The studies were references under 27.30.32.35.37.43.92. 101,102,109,119,122,125,127,136,137,145,148,149, 151,156,160-62. Nearly similar proportion of studies (14%) included in this update has shaded light on drug outcomes focusing more particularly on dose, utility, efficacy, adverse reaction, determinants and implications of use of isoniazid preventive therapy combined with ART or ART alone (1,38,42,53,54,57,58,66,74,77, 82,94,105,117,124,126,127,143 and 159).

Similar proportion of studies (14%) was found to be on adherence to treatment. These studies focus on predictors and determinants of adherence role of disclosure in adherence and couple of studies one focusing on use of traditional medicine in adherence and text message as reminder. Specific details on adherence related studies are found under reference numbers 10,12,34,36,41,46, 49,50,65,79,87,93,95,99,103,107,132,134,139,146, 147,152 and 158.

Unlike previous updates, this particular update has relatively more studies that focus on pediatric HIV. The studies focused mainly on immune response, impairments, recovery, immunologic failure, survival and mortality in connection to ART. These studies are found in references: 21,33,47,55,73,89,107,138,140.

This update has identified handful of studies that have focused on effects of nutrient supplements among patients on ART, determinants of dietary diversity, prevalence and determinants of anemia among HIV infected patients on ART, impact of HAART on nutritional status (4,14,19,45,56,88 and 141)

Studies on ART associated opportunistic infecitons were found to focus on incidence, prevalence, determinants and risks, CD4 counts, management, resistence to specific infecitons as referenced in 5,9,16,83,100,115,116,118,128,131 and 157. There were some studies that were found in reference to sexual behaviour, fertility desire, choice of contraceptive methods and pregnancy outcome among women living with the virus and/or are on ART as spelt out in references 3,15,24,44,133,135.

A collection of studies on deworming, prevalence of intestinal parasites and consequent compromised immunity were included in this update and shaded light on intestinal parasites and ART. These studies are referenced under 6,81,84,85,110,113 and 121. Care and support is an important component of this section where lived experiences in caring for orphans and vulnerable children, support pathways, role of FBOs, determinants, barriers and income generating activities were referenced in 40,48,62,63,70,80,97 and 104.

The current update has new research outcomes that generated useful evidence on the domains of cervical cancer among those who are living with the virus. The studies documented on determinants of precancerous cervical and VIA postive leisions as well as prevneiton of cervical cacner using single vist approach among HIV positive women. The references are 71,72 and 91. Similar few studies on gender differentials in quality of life, survival are presented here 18,22 and 75.

Finally, there are different studies on ART with specific interest on eligibility, admission to service, serice delivery and treatement strategy, quality of service and life, beleifs. These studies are found in references: 25,96,98,108,150,153 and 155.

1. Abate E, Elias D, Getachew A, Alemu S, Diro E, Britton S, et al. Effects of albendazole on the clinical

outcome and immunological responses in helminth co-infected tuberculosis patients: a double blind randomised clinical trial. International Journal for Parasitology. 2015 01 Feb;45(2-3):133-140.

- Abay SM, Deribe K, Reda AA, Biadgilign S, Datiko D, Assefa T, et al. The effect of early initiation of antiretroviral therapy in TB/HIV-coinfected patients: a systematic review and meta-analysis. J Int Assoc Provid AIDS Care. 2015 Nov-Dec;14(6):560-570. Epub 2015/08/21.
- 3. Abbawa F, Awoke W, Alemu Y. Fertility desire and associated factors among clients on highly active antiretroviral treatment at Finoteselam Hospital northwest Ethiopia: a cross sectional study. Reproductive Health. 2015 11 Aug;12 (1) (no pagination)(69).
- Abdissa A, Olsen MF, Yilma D, Tesfaye M, Girma T, Christiansen M, et al. Lipid-based nutrient supplements do not affect efavirenz but lower plasma nevirapine concentrations in Ethiopian adult HIV patients. HIV medicine. 2015 Aug;16(7):403-411. Epub 2015/05/15.
- Abebe N, Mulu G. Prevalence of opportunistic infections and associated factors among HIVpositive patients taking antiretroviral therapy in Debre Markos Referral Hospital, northwest Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association February 26-28, 2015; Bahir Dar, 2015.
- 6. Abossie A, Petros B. Deworming and the immune status of HIV positive pre-antiretroviral therapy individuals in Arba Minch, Chencha and Gidole hospitals, Southern Ethiopia. BMC Res Notes. 2015 Sep 28;8:483. Epub 2015/09/30.
- 7. Addis GA, Negese DK, Hedija Yenus Y. Demand for long acting contraceptive methods among married HIV positive women attending care at public health facilities at Bahir Dar City, northwest Ethiopia. Reproductive Health. 2015;12(76).
- Addisu A, Dagim A, Tadele E, Adissu A, Mussie A, Filmon K. CD4 cell count trends after commencement of antiretroviral therapy among HIV-infected patients in Tigray, northern Ethiopia: a retrospective cross-sectional study. PLoS ONE. 2015;10(3).
- Adejumo OA, Malee KM, Ryscavage P, Hunter SJ, Taiwo BO. Contemporary issues on the epidemiology and antiretroviral adherence of HIVinfected adolescents in sub-Saharan Africa: a narrative review. J Int AIDS Soc. 2015;18:20049. Epub 2015/09/20.
- Adelman MW, Tsegaye M, Kempker RR, Alebachew T, Haile K, Tesfaye A, et al. Intensified tuberculosis case finding among HIV-infected persons using a WHO symptom screen and Xpert MTB/RIF. International Journal of Tuberculosis and Lung Disease. 2015 01 Oct;19(10):1197-1203.
- 11. African Union. Roadmap on shared responsibility and global solidarity for AIDS, TB and malaria *Ethiop. J. Health Dev.* 2016;30(3)

response in Africa. Addis Ababa, Ethiopia, African Union, 2011.

- 12. Alelgen T, Abebaw A, Berhane K, Tekeste Z. Evaluation of adherence to antiretroviral therapy among HIV/AIDS patients in Amhara Regional State, North Shewa Zone, Ethiopia. 26th Annual conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Alemayehu G, Melaku Z, Abreha T, Alemayehu B, Girma S, Tadesse Y, et al. Burden of malaria among adult patients attending general medical outpatient department and HIV care and treatment clinics in Oromia, Ethiopia: a comparative cross-sectional study. Malaria Journal. 2015 15 Dec;14 (1) (no pagination)(1029).
- Aleme A, Mohammed Y. Traditional consumption, therapeutic value and its derived dairy products of dromedary camel (*Camelus dromedaries*) milk in Somali Regional State, eastern Ethiopia: a review. Global Journal of Animal Scientific Research. 2015;3(1):240-246.
- Alemu FM, Yalew AW, Fantahun M, Ashu EE. Antiretroviral Therapy and Pregnancy Outcomes in Developing Countries: A Systematic Review. International Journal of MCH and AIDS. 2015;3(1):31-43. Epub 2015/01/01.
- 16. Alemu J, Tsegaye A. Highly active antiretroviral therapy improved hematologic parameters of immunodeficiency virus-infected adult individuals at Black Lion Specialized Hospital, Addis Ababa, Ethiopia. American Journal of Clinical Pathology. 2015 October;144:A217.
- 17. Amante TD, Ahemed TA. Risk factors for unsuccessful tuberculosis treatment outcome (failure, default and death) in public health institutions, eastern Ethiopia. Pan Afr Med J. 2015;20:247.
- Amanuel T, Abebe G, Mulusew G, Hailay A. Gender differences in health related quality of life among people living with HIV on highly active antiretroviral therapy in Mekelle town, northern Ethiopia. BioMed Research International. 2015;516369(25).
- 19. Amare Tariku W, Melkie Edris Y, Teresa Kisi B, Melaku Kindie Y. Factors associated with dietary diversity among HIV positive adults (>=18 years) attending ART clinic at Mettema hospital, northwest Ethiopia: cross-sectional study. Journal of AIDS and Clinical Research. 2015;6(8).
- Amogne W, Aderaye G, Habtewold A, Yimer G, Makonnen E, Worku A, et al. Efficacy and Safety of Antiretroviral Therapy Initiated One Week after Tuberculosis Therapy in Patients with CD4 Counts < 200 Cells/muL: TB-HAART Study, a Randomized Clinical Trial. PLoS One. 2015;10(5):e0122587. Epub 2015/05/13.
- 21. Asefa F, Edessa D, Sheikhahmed J. Predictors of early mortality in HIV-infected children initiated with ART at Hiwot Fana Specialized University

Hospital. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.

- 22. Asefa H, Dube L, Ewnetu H. Gender as a risk factor for survival among HIV patients on antiretroviral therapy: a systematic review protocol. JBI database of systematic reviews and implementation reports. 2015 Sep 16;13(8):14-20. Epub 2015/10/13.
- 23. Asfaw A, Ali D, Eticha T, Alemayehu A, Alemayehu M, Kindeya F. CD4 cell count trends after commencement of antiretroviral therapy among HIV-infected patients in Tigray, northern Ethiopia: a retrospective cross-sectional study. PLoS ONE. 2015 27 Mar;10 (3) (no pagination)(e0122583).
- 24. Asfaw HM. Effect of antiretroviral therapy (ART) on fertility and contraceptive preference of HIV-positive women in Addis Ababa, Ethiopia. [MPH Thesis]: Addis Ababa University; 2015.
- 25. Asrie F, Gelaw B, Alemu M, Moges F, Awoke T. Determination of eligibility to antiretroviral therapy in resource limited settings using total lymphocyte counts, hemoglobin and body mass index among HIV-positive patients. 26th Annual Conference of the Ethiopian Pubic Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Assefa M, Abegaz WE, Shewamare A, Medhin G, Belay M. Prevalence and correlates of anemia among HIV infected patients on highly active antiretroviral therapy at Zewditu Memorial Hospital, Ethiopia. BMC Hematology. 2015;15:6. Epub 2015/06/06.
- Assefa Y, Lynen L, Kloos H, Hill P, Rasschaert F, Hailemariam D, et al. Long-term outcomes and their determinants in patients on antiretroviral treatment in Ethiopia, 2005/6-2011/12: a retrospective cohort study. Journal of Acquired Immune Deficiency Syndromes. 2015 01 Dec;70(4):414-419.
- 28. Ayele HT, Mourik MS, Debray TP, Bonten MJ. Isoniazid prophylactic therapy for the prevention of tuberculosis in HIV infected adults: a systematic review and meta-analysis of randomized trials. PLoS One. 2015;10(11):e0142290. Epub 2015/11/10.
- Ayele HT, van Mourik MS, Bonten MJ. Effect of isoniazid preventive therapy on tuberculosis or death in persons with HIV: a retrospective cohort study. BMC Infect Dis. 2015 Aug 13;15:334. Epub 2015/08/14.
- 30. Ayele W, Mulugeta A, Desta A, Rabito FA. Treatment outcomes and their determinants in HIV patients on anti-retroviral treatment program in selected health facilities of Kembata and Hadiya Zones, Southern Nations, Nationalities and Peoples Region, Ethiopia. BMC Public Health. 2015;15:826.
- Balcha TT, Skogmar S, Sturegard E, Bjorkman P, Winqvist N. Outcome of tuberculosis treatment in HIV-positive adults diagnosed through active versus passive case-finding. Glob Health Action. 2015;8:27048. Epub 2015/03/31.

- 32. Bayou B, Sisay A, Kumie A. Assessment of the magnitude and associated factors of immunological failure among adult and adolescent HIV-infected patients in St. Luke and Tulubolo Hospital, Oromia Region, Ethiopia. Pan Afr Med J. 2015;21:291. Epub 2015/11/21.
- Bekele Y, Amu S, Bobosha K, Lantto R, Nilsson A, Endale B, et al. Impaired phenotype and function of T follicular helper cells in HIV-1-infected children receiving ART. Medicine. 2015 Jul;94(27):e1125. Epub 2015/07/15.
- 34. Belayihun B, Negus R. Antiretroviral treatment adherence rate and associated factors among people living with HIV in Dubti Hospital, Afar Regional State, East Ethiopia. International Scholarly Research Notices. 2015;2015:187360. Epub 2015/01/01.
- 35. Belayneh M, Giday K, Lemma H. Treatment outcome of human immunodeficiency virus and tuberculosis co-infected patients in public hospitals of Eastern and Southern Zone of Tigray Region, Ethiopia. The Brazilian Journal of Infectious Diseases. 2015 Jan-Feb;19(1):47-51. Epub 2014/12/04.
- 36. Berbeto TM, Berihun D. Predictors of loss to followup in patients living with HIV/AIDS after initiation of antiretroviral therapy in southwest Ethiopia: a cohort study. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 37. Beyene MB, Beyene HB. Predictors of late diagnosis of HIV among HIV positive adults coming for initial CD4 T-cell count to public health facilities, northern Ethiopia. Tropical Medicine and International Health. 2015 September;20:171.
- 38. Bezabhe WM, Bereznicki LR, Chalmers L, Gee P, Kassie DM, Bimirew MA, et al. Adverse drug reactions and clinical outcomes in patients initiated on antiretroviral therapy: a prospective cohort study from Ethiopia. Drug Safety. 2015 02 Jul;38(7):629-639.
- Biadglegne F, Rodloff AC, Sack U. Review of the prevalence and drug resistance of tuberculosis in prisons: a hidden epidemic. Epidemiol Infect. 2015 Apr;143(5):887-900. Epub 2014/11/08.
- 40. Biru M, Lundqvist P, Molla M, Jerene D, Hallstrom I. Surviving overwhelming challenges: family caregivers' lived experience of caring for a child diagnosed with HIV and enrolled in antiretroviral treatment in Ethiopia. Issues in Comprehensive Pediatric Nursing. 2015;38(4):282-299. Epub 2015/09/17.
- Bucciardini R, Fragola V, Abegaz T, Lucattini S, Halifom A, Tadesse E, et al. Retention in care of adult HIV patients initiating antiretroviral therapy in Tigray, Ethiopia: a prospective observational cohort study. PLoS ONE. 2015 04 Sep;10 (9) (no pagination)(e0136117).

- 42. Chen I, Poirot E, Newman M, Kandula D, Shah R, Hwang J, et al. An assessment of the supply, programmatic use, and regulatory issues of single low-dose primaquine as a *Plasmodium falciparum* gametocytocide for sub-Saharan Africa. Malaria Journal. 2015 15 May;14 (1) (no pagination)(714).
- 43. Damtew B, Mengistie B, Alemayehu T. Survival and determinants of mortality in adult HIV/Aids patients initiating antiretroviral therapy in Somali Region, Eastern Ethiopia. Pan Afr Med J. 2015;22:138. Epub 2016/02/19.
- Demissie K, Asfaw S, Abebe L, Kiros G. Sexual behaviors and associated factors among antiretroviral treatment attendees in Ethiopia. HIV/AIDS (Auckland, NZ). 2015;7:183-190. Epub 2015/06/18.
- 45. Dereje G, Getnet M, Gemechu K, Tewodros E, Fituma F, Temesgen. Food insecurity and its associated factors among people living with HIV/AIDS receiving anti-retroviral therapy at Butajira Hospital, southern Ethiopia. Journal of Nutrition and Food Sciences. 2015;5(2).
- Deribe K. Adherence to antiretroviral therapy among children aged 2-14 years in Tikur Anbessa Specialized Hospital, Addis Ababa, Ethiopia. [MPH Thesis]: Addis Ababa University; 2015.
- 47. Desta A, Fesseha G. Factors affecting survival of HIV positive children taking antiretroviral therapy at Adama Referral Hospital and Medical College, Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Desta MK, Linsk N. Pathways and motivations for raising relatives' children orphaned by HIV/AIDS in Ethiopia. Journal of HIVAIDS & Social Services. 2 Oct/Dec 2015 2016-08-16;14(4):392-404.
- 49. Digaffe T. Drop out of newly diagnosed HIV infected adults from routine pre-ART care in a public hospital, Harar Town, eastern Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association February 26-28, 2015; Bahir Dar, 2015.
- 50. Diribe K. Adherence to antiretroviral therapy among children aged 2-12 in Tikur Anbessa Specialized Hospital, Addis Ababa, Ethiopia: Addis Ababa University; 2015.
- 51. Diro E, Lynen L, Assefa M, Takele Y, Mengesha B, Adem E, et al. Impact of the use of a rapid diagnostic test for visceral leishmaniasis on clinical practice in Ethiopia: a retrospective study. PLoS Neglected Tropical Diseases. 2015 12 May;9 (5) (no pagination)(e0003738).
- 52. Diro E, Lynen L, Gebregziabiher B, Assefa A, Lakew W, Belew Z, et al. Clinical aspects of paediatric visceral leishmaniasis in north-west Ethiopia. Tropical Medicine and International Health. 2015 01 Jan;20(1):8-16.
- 53. Diro E, Lynen L, Ritmeijer K, Alves F, Mohamed R, Abongomera C, et al. Use of pentamidine as secondary prophylaxis to prevent visceral *Ethiop. J. Health Dev.* 2016;30(3)

leishmaniasis relapse in HIV infected patients. 51st Medical Conference, Ethiopian Medical Association;; February 5-6, 2015; Addis Ababa2015.

- 54. Diro E, Ritmeijer K, Boelaert M, Alves F, Mohammed R, Abongomera C, et al. Use of pentamidine as secondary prophylaxis to prevent visceral leishmaniasis relapse in HIV-infected patients. Tropical Medicine and International Health. 2015 September;20:217.
- 55. Ebissa G, Deyessa N, Biadgilign S. Predictors of early mortality in a cohort of HIV-infected children receiving high active antiretroviral treatment in public hospitals in Ethiopia. AIDS Care -Psychological and Socio-Medical Aspects of AIDS/HIV. 2015 03 Jun;27(6):723-730.
- 56. Ebissa G, Deyessa N, Biadgilign S. Impact of highly active antiretroviral therapy on nutritional and immunologic status in HIV-infected children in the low-income country of Ethiopia. Nutrition. 2016 Jun;32(6):667-673. Epub 2016/02/16.
- 57. Edessa D, Likisa J. Comparison of effects of isoniazid preventive therapy plus ART alone on the risks of all-cause mortality among HIV-infected individuals in Addis Ababa, Ethiopia: a cohort study. 51st Medical Conference, Ethiopian Medical Association; February 5-6, 2015; Addis Ababa, 2015.
- Edessa D, Likisa J. A Description of Mortality Associated with IPT plus ART Compared to ART Alone among HIV-Infected Individuals in Addis Ababa, Ethiopia: A Cohort Study. PLoS One. 2015;10(9):e0137492. Epub 2015/09/09.
- Ejeta E, Birhanu T, Wolde T. Tuberculosis outcomes among TB/HIV co-infected cases treated under directly observed treatment of short course in western Ethiopa. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015, Bahir Dar, 2015.
- 60. Ejeta E, Chala M, Arega G, Ayalsew K, Tesfaye L, Birhanu T, et al. Outcome of tuberculosis patients under directly observed short course treatment in western Ethiopia. Journal of Infection in Developing Countries. 2015 01 Jul;9(7):752-759.
- 61. Enawgaw B, Tesfaye Z. Prevalence of anemia before and after initiation of highly active antiretroviral therapy among HIV-positive patients in northwest Ethiopia: a retrospective study. . 26th Annual Conference of the Ethiopian Public Health Association; February 5-6, 2015; Bahir Dar, 2015.
- 62. Endeshaw M, Alemu S, Andrews N, Dessie A, Frey S, Rawlins S, et al. Involving religious leaders in HIV care and treatment at a university-affiliated hospital in Ethiopia: application of formative inquiry. Glob Public Health. 2015 Aug 10:1-16. Epub 2015/08/11.
- 63. Erango MA, Ayka ZA. Psychosocial support and parents' social life determine the self-esteem of orphan children. Risk Management and Healthcare Policy. 2015 13 Oct;8:169-173.

- 64. Ermias D, Griensven Jv, Rezika M, Colebunders R, Mesfin A, Asrat H, et al. Atypical manifestations of visceral leishmaniasis in patients with HIV in north Ethiopia: a gap in guidelines for the management of opportunistic infections in resource poor settings. Lancet Infectious Diseases. 2015;15(1):122-129.
- 65. Fikadu F. Assessment of HIV serostatus discloser and its influence on aherence to ARV treatment among HIV-infected adolescents in Addis Ababa, Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 66. Finkelstein JL, Gala P, Rochford R, Glesby MJ, Mehta S. HIV/AIDS and lipodystrophy: implications for clinical management in resource-limited settings. J Int AIDS Soc. 2015;18:19033. Epub 2015/01/20.
- 67. Fiseha D, Demissie M. Assessment of Directly Observed Therapy (DOT) following tuberculosis regimen change in Addis Ababa, Ethiopia: a qualitative study. BMC Infect Dis. 2015 Sep 30;15:405. Epub 2015/10/02.
- 68. Fiseha T, Gebru T, Gutema H, Debela Y. Tuberculosis treatment outcome among HIV Coinfected patients at Mizan-Aman General Hospital, southwest Ethiopia: a retrospective study. Journal of Bioengineering & Biomedical Sciences. 2015. 2015-11-16;5(1):1.
- 69. Gedefaw L, Yemane T, Sahlemariam Z, Yilmar D. Anemia and risk factors in HAART native and HAART experienced positive participation in southwest Ethiopia: a comparitive study. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 70. Gelaw YA, Senbete GH, Adane AA, Alene KA. Determinants of late presentation to HIV/AIDS care in Southern Tigray Zone, northern Ethiopia: an institution based case-control study. AIDS Research and Therapy. 2015;12(40).
- 71. Gessesse Z, Tadesse Z, Alemayehu M, Hiruye A, Getachew Y, Derbew M, et al. Determinant factors of visual inspection with acetic acid (via) positive lesions among HIV positive women in Mekelle Hospital, northern Ethiopia: a case control study. Ethiopian Medical Journal. 2015 01 Jul:17-24.
- 72. Gessesse Z, Tadesse Z, Alemayehu M, Hiruye A, Getachew Y, Mamo D. Determinants of pre cancerous cervical lesions among HIV positive women in Mekele Hospital: unmatched case control study. 51st Medical Conference, Ethiopian Medical Association; February 5-6, 2015; Addis Ababa, 2015.
- 73. Girma A. Assessment of immuno-virological response to ART in children <15 years of age in Adama Referral Hospital in Adama Town, Oromia Regioonal State, central Ethiopia. [MSc Thesis]: Addis Ababa University; 2015.
- 74. Girma E, Likisa J. Comparison of effects of isoniazid preventive therapy plus ART and ART alone on the rise of all-cause mortality among HIV-

infected individuals in Addis Ababa, Ethiopia: a cohort study. 26th Annual Conference of the Ethiopian Public Health Association; February 26-18, 2015; Bahir Dar, 2015.

- 75. Gizaw A, Tilahun T, Tesfay A. Gender disparities in late presentation and survival among HIV patients on antiretroviral therapy in public health facilities in Arba Minch Town, southern Ethiopia: eight years retrospective cohoty study. 26th Annual Conference of the Ethiopian Public Health Conference; February 26-28, 2015; Bahir Dar, 2015.
- 76. Gize A, Mathewos B, Moges B. Establishment of normal reference intervals for Cd3+, Cd4+ Cd8+ and Cd4+ to Cd8+ ratio of T-lymphocytes in HIVnegative adults from University of Gondar Hosital, nothwest Ethiopia. 26th Annual Conference of the EEthiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 77. Gleason RL, Caulk AW, Seifu D, Parker I, Vidakovic B, Getenet H, et al. Current efavirenz (EFV) or ritonavir-boosted lopinavir (LPV/r) use correlates with elevate markers of atherosclerosis in HIV-infected subjects in Addis Ababa, Ethiopia. PLoS ONE. 2015 27 Apr;10 (4) (no pagination)(e0117125).
- 78. Guda A, Zenebe M, Tesfay A, Alemayehu B, Girma S, Yehualashet T, et al. Burden of malaria among adult patients attending general medical outpatient department and HIV care and treatment clinics in Oromia, Ethiopia: a comparative cross-sectional study. Malaria Journal. 2015;14(501).
- 79. Gultie T, Genet M, Sebsibie G. Disclosure of HIVpositive status to sexual partner and associated factors among ART users in Mekelle Hospital. HIV/AIDS (Auckland, NZ). 2015;7:209-214. Epub 2015/07/18.
- Habtamu T, Yazie C, Minuyelet J. Establishment of model home garden for home consumption and income generation of poor and HIV victim women at Bahir Dar, Ethiopia. Journal of Agricultural Science and Technology A. 2015;5(3):183-189.
- 81. Habteselassie A, Ali A, Birhaneselassie M, Tadesse G. Hematological outcomes of children before and after highly active antiretroviral therapy at Zewditu Memorial Hospital. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 82. Habtewold A, Makonnen E, Amogne W, Yimer G, Aderaye G, Bertilsson L, et al. Is there a need to increase the dose of efavirenz during concomitant rifampicin-based antituberculosis therapy in sub-Saharan Africa? The HIV-TB pharmagene study. Pharmacogenomics. 2015;16(10):1047-1064. Epub 2015/04/02.
- 83. Habtewold T, Yarlagadda R, Wolde-Mariam M. Assessment of therapeutic management of opportunistic infections among adults taking art in Jimma University specialized hospital, Jimma, Ethiopia. Pharmanest. 2015;6(2):2721-2726.

- Habtom K, Endalkachew N, Abaineh M, Bizuayehu K, Melaku A. Prevalence of intestinal protozoan infections among individuals living with HIV/AIDS at Felegehiwot Referral Hospital, Bahir Dar, Ethiopia. International Journal of Infectious Diseases. 2015;35:80-86.
- 85. Hailu AW, S GS, Merid Y, Gebru AA, Ayene YY, Asefa MK. The case control studies of HIV and Intestinal parasitic infections rate in active pulmonary tuberculosis patients in Woldia General Hospital and Health Center in North Wollo, Amhara Region, Ethiopia. International journal of pharma sciences. 2015 May-Jun;5(3):1092-1099. Epub 2016/03/22.
- 86. Hassen MS. Risk factor and resistance pattern of MDR TB among suspected cases of Oromia Region. European Respiratory Journal Conference: European Respiratory Society Annual Congress. 2015;46(no pagination).
- Hiko D, Wondafrash B, Abdulahi M. Predictors of non-adherence with concurrent treatment among HIV/TB co-infected patients: case control study. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 88. Hussen S, Belachew T, Hussien N. Nutritional status and its effect on treatment outcome among patients taking HAART in Ethiopia: cohort study. . 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Isaacson G, Melaku A. Results of pediatric tympanoplasty on short-term surgical missions. The Laryngoscope. 2016 Jun;126(6):1464-1469. Epub 2015/08/01.
- 90. Kassa D, de Jager W, Gebremichael G, Alemayehu Y, Ran L, Fransen J, et al. The effect of HIV coinfection, HAART and TB treatment on cytokine/chemokine responses to Mycobacterium tuberculosis (Mtb) antigens in active TB patients and latently Mtb infected individuals. Tuberculosis (Edinburgh, Scotland). 2016 Jan;96:131-140. Epub 2015/12/04.
- 91. Kassahun K, Shiferaw N, Tesfamichael T, Bayssa Y, Lulu K, Weldegebreal T, et al. Cervical cancer prevention using the single-visit approach in lowresource settings: experience from pathfinder international's Addis Tesfa project in Ethiopia. International Journal of Gynecology and Obstetrics. 2015 October;131:E268.
- 92. Kassie DM, Tigabu T, Kassaye M, Feleke G, Kokeb M, Mesfin N. Experience of antiretroviral treatment failure screening at Gondar University Hospital: innovative approach for resource limited settings. 51st Medical Conference, Ethiopian Medical Association; February 5-6, 2015; Addis Ababa2015.
- 93. Kebede M, Zeleke A, Asemahagn M, Fritz F. Willingness to receive text message medication reminders among patients on antiretroviral treatment in North West Ethiopia: A cross-sectional study.

Ethiop. J. Health Dev. 2016;30(3)

BMC Medical Informatics and Decision Making. 2015 Aug 13;15:65. Epub 2015/08/14.

- 94. Kedir MS, Gemeda DH, Suleman S. Treatment outcomes of nevirapine- versus efavirenz-based highly active antiretroviral therapy regimens among antiretroviral-naive adult patients in Ethiopia: a cohort study. Therapeutic Innovation & Regulatory Science. 2015 May 2015 2015-05-18:49(3):443-449.
- 95. Ketema AK, Weret ZS. Assessment of adherence to highly active antiretroviral therapy and associated factors among people living with HIV at Debre Birhan referral hospital and health center, northeast Ethiopia: a cross-sectional study. HIV/AIDS -Research and Palliative Care. 2015 11 Mar;7:75-81.
- 96. Kondo M, Woldegiorgis A, Tilahun H, Harris C. Characterization of inpatient admission within a large HIV treatment program in Ethiopia. Annals of Global Health. 2015 January-February;81 (1):128-129.
- 97. Kulkarni S, Hoffman S, Gadisa T, Melaku Z, Fantehun M, Yigzaw M, et al. Identifying perceived barriers along the HIV care continuum: findings from providers, peer educators, and observations of provider-patient interactions in Ethiopia. Journal of the International Association of Providers of AIDS Care. 2015 2015 Jul 14:[10] p.
- Kumela K, Amenu D, Chelkeba L. Comparison of anti-retroviral therapy treatment strategies in prevention of mother-to-child transmission in a teaching hospital in Ethiopia. Pharmacy Practice. 2015;13(2):1-10.
- 99. Legesse B, Shewamene Z. Traditional medicine use and its impact on antiretroviral therapy adherence among HIV patients in Ethiopia. Value in Health. 2015 May;18 (3):A241.
- 100.Lemma MT, Zenebe Y, Tulu B, Mekonnen D, Mekonnen Z. Methicillin resistant *Staphylococcus aureus* among HIV infected pediatric patients in northwest Ethiopia: carriage rates and antibiotic coresistance profiles. PLoS One. 2015;10(9):e0137254. Epub 2015/10/01.
- 101.Lenjisa JL, Wega SS, Lema TB, Ayana GA. Outcomes of highly active antiretroviral therapy and its predictors: a cohort study focusing on tuberculosis co-infection in south west Ethiopia. BMC Res Notes. 2015 Sep 15;8:446. Epub 2015/09/17.
- 102.Lessells RJ, Swaminathan S, Godfrey-Faussett P. HIV treatment cascade in tuberculosis patients. Current Opinion in HIV and AIDS. 2015 Nov;10(6):439-446. Epub 2015/09/10.
- 103.Letta S, Demissie A, Oljira L, Dessie Y. Factors associated with adherence to antiretroviral therapy (ART) among adult people living with HIV and attending their clinical care, eastern Ethiopia.[Erratum appears in BMC Int Health Hum Rights. 2016;16:8 . BMC Int Health Hum Rights. 2015;15:33.

- 104.Lifson AR, Workneh S, Hailemichael A, Demisse W, Slater L, Shenie T. Implementation of a peer HIV community support worker program in rural Ethiopia to promote retention in care. J Int Assoc Provid AIDS Care. 2015 Oct 30. Epub 2015/11/01.
- 105.Lifson AR, Workneh S, Hailemichael A, Demissie W, Slater L, Shenie T. Perceived social support among HIV patients newly enrolled in care in rural Ethiopia. AIDS Care. 2015 2015;27(11):1382-1386.
- 106.Maru M, Mariam SH, Airgecho T, Gadissa E, Aseffa A. Prevalence of tuberculosis, drug susceptibility testing, and genotyping of mycobacterial isolates from pulmonary tuberculosis patients in Dessie, Ethiopia. Tuberculosis Research and Treatment. 2015;2015:215015. Epub 2015/07/17.
- 107.Mekonnen D, Derbie A, Desalegn E. TB/HIV coinfections and associated factors among patients on directly observed treatment short course in northeastern Ethiopia: a 4 years retrospective study. BMC Res Notes. 2015 Nov 11;8:666. Epub 2015/11/13.
- 108. Mekuria LA, Prins JM, Yalew AW, Sprangers MA, Nieuwkerk PT. Retention in HIV care and predictors of attrition from care among HIV-infected adults receiving combination anti-retroviral therapy in Addis Ababa. PLoS One. 2015;10(6):e0130649. Epub 2015/06/27.
- 109.Mekuria LA, Sprangers MA, Prins JM, Yalew AW, Nieuwkerk PT. Health-related quality of life of HIVinfected adults receiving combination antiretroviral therapy in Addis Ababa. AIDS Care. 2015 2015;27(8):934-945.
- 110.Melaku Z, Lamb MR, Wang C, Lulseged S, Gadisa T, Ahmed S, et al. Characteristics and outcomes of adult Ethiopian patients enrolled in HIV care and treatment: a multi-clinic observational study. BMC Public Health. 2015 May 03;15:462. Epub 2015/05/03.
- 111.Mengist HM, Taye B, Tsegaye A. Intestinal parasitosis in relation to CD4+ T cells levels and anemia among HAART initiated and HAART naive pediatric HIV patients in a model ART center in Addis Ababa, Ethiopia. PLoS ONE. 2015;10(2).
- 112.Mengistu Z, Getaw G. Treatment outcome of multidrug/extensive drug resistant tuberculosis among patients treated at St. Peter Hospital in Addis Ababa. . 26 Annual Conferene of the Ethiopian Public Health Association; Februay 26-28, 2015; Bahis Dar, 2015.
- 113.Meressa D, Hurtado RM, Andrews JR, Diro E, Abato K, Daniel T, et al. Achieving high treatment success for multidrug-resistant TB in Africa: initiation and scale-up of MDR TB care in Ethiopia an observational cohort study. Thorax. 2015 01 Dec;70(12):1181-1188.
- 114.Mihireti H, Taye B, Tsegaye A. Intestinal parasitosis in relation to CD4=T cell levels and anemia among HAART-initiated and non-HAART iniitiated pediatric HIV patients in Zewditu Memorial *Ethiop. J. Health Dev.* 2016;30(3)

Hospital, Addis Ababa, Ethiopia. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.

- 115.Mihiretie H, Taye B, Tsegaye A. Magnitude of anemia and associated factors among pediatric HIV/AIDS patients attending Zewditu Memorial Hospital ART Clinic, Addis Ababa, Ethiopia. Anemia. 2015;2015:479329. Epub 2015/04/17.
- 116.Misleer D. Effect of highly active antiretroviral therapy on incidence and opportunistic infections amomng HIV positive adults in public health facilities in Arba Minch: a retrospective cohort study. 26th Annudal Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 117.Mitiku H, Weldegebreal F, Teklemariam Z. Magnitude of opportunistic infections and associated factors in HIV-infected adults on antiretroviral therapy in eastern Ethiopia. HIV/AIDS (Auckland, NZ). 2015;7:137-144. Epub 2015/05/23.
- 118.Mohamed Z, Kote M, Gizaw A. The effect of isoniazid preventive prophylaxis on immunological response rate among highly active antiretroviral therapy patients in Addis Ababa, Ethiopia: a retrospective cohort study. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 119.Molla G. Prevalence of symptomatic and asymptomatic bacteriuria and their antimicrobial sustceptibility patterns among adult HIV/AIDSinfected patients on HAART and HAART-naive attending Tikur Anbessa and Zeditu hospitals, Addis Ababa, Ethiopia.: Addis Ababa University; 2015.
- 120.Muhula SO, Peter M, Sibhatu B, Meshack N, Lennie K. Effects of highly active antiretroviral therapy on the survival of HIV-infected adult patients in urban slums of Kenya. Pan Afr Med J. 2015;20:63. Epub 2015/06/20.
- 121.Mulisa G, Workneh T, Hordofa N, Suaudi M, Abebe G, Jarso G. Multidrug-resistant *Mycobacterium tuberculosis* and associated risk factors in Oromia Region of Ethiopia. International Journal of Infectious Diseases. 2015;39:57-61.
- 122.Mulu A, Anagaw B, Gelaw A, Ota F, Kassu A, Yifru S. Effect of deworming on Th2 immune response during HIV-helminths co-infection. J Transl Med. 2015 Jul 18;13:236. Epub 2015/07/19.
- 123.Mulu A, Maier M, Liebert UG. Low incidence of HIV-1C acquired drug resistance 10 years after rollout of antiretroviral therapy in Ethiopia: a prospective cohort study. PLoS ONE. 2015 29 Oct;10 (10) (no pagination)(0141318).
- 124.Mulu W, Mekonnen D, Yimer M, Admassu A, Abera B. Risk factors for multidrug resistant tuberculosis patients in Amhara National Regional State. African Health Sciences. 2015 Jun;15(2):368-377. Epub 2015/07/01.
- 125.Neogi U, Haggblom A, Singh K, Rogers LC, Rao SD, Amogne W, et al. Factors influencing the

efficacy of rilpivirine in HIV-1 subtype C in lowand middle-income countries. The Journal of Antimicrobial Chemotherapy. 2016 Feb;71(2):367-371. Epub 2015/11/01.

- 126.Olsen MF, Kaestel P, Tesfaye M, Abdissa A, Yilma D, Girma T, et al. Physical activity and capacity at initiation of antiretroviral treatment in HIV patients in Ethiopia. Epidemiology and Infection. 2015 Apr 2015 2015-04-29;143(5):1048-1058.
- 127.Osafo C, Raji YR, Burke D, Tayo BO, Tiffin N, Moxey-Mims MM, et al. Human heredity and health (H3) in Africa kidney disease research network: a focus on methods in sub-Saharan Africa. Clinical Journal of the American Society of Nephrology. 2015 07 Dec;10(12):2279-2287.
- 128.Oumer F, Parekh M, Schneider J, Bekele A, Bayisa T, Sherman CB, et al. Utility of fiberoptic bronchoscopy at Tikur Anbessa specialized hospital, Addis Ababa, Ethiopia. American Journal of Respiratory and Critical Care Medicine Conference: American Thoracic Society International Conference, ATS. 2015;191(no pagination).
- 129.Reepalu A, Balcha TT, Yitbarek T, Jarso G, Sturegard E, Bjorkman P. Screening for cryptococcal antigenemia using the lateral flow assay in antiretroviral therapy-naive HIV-positive adults at an Ethiopian hospital clinic. BMC Research Notes. 2015;8:702.
- 130.Seboxa T, Amogne W, Abebe W, Tsegaye T, Azazh A, Hailu W, et al. High mortality from blood stream infection in Addis Ababa, Ethiopia, is due to antimicrobial resistance. PLoS ONE. 2015 01 Dec;10 (12) (no pagination)(e0144944).
- 131.Senbete GH, Adane AA, Gelaw YA. Determinants of late presentation to HIV/AIDS care among people living with HIV receiving care inhealth institutions of southern Tigray Zone, northern Ethiopia: An institution based case control study. 26th Annual Conference of the Ethiopian Public Health Association; Bahir Dar, 2015.
- 132.Setegn T, Takele A, Gizaw T, Nigatu D, Haile D. Predictors of mortality among adult antiretroviral therapy users in southeastern Ethiopia: retrospective cohort study. AIDS Research and Treatment. 2015;2015:148769. Epub 2015/03/31.
- 133.Sharew G, Azage M. Predictors of HIV/AIDS Related Ocular Manifestations among HIV/AIDS Patients in Felege Hiwot Referral Hospital, northwest Ethiopia. Journal of Ophthalmology. 2015;2015:965627. Epub 2015/05/23.
- 134.Shaweno T, Shaweno D. When are patients lost to follow-up in pre-antiretroviral therapy care? a retrospective assessment of patients in an Ethiopian rural hospital. Infectious Diseases of Poverty. 2015;4:27. Epub 2015/06/03.
- 135.Shewamene Z, Legesse B, Tsega B, Bhagavathula AS, Endale A. Consistent condom use in HIV/AIDS patients receiving antiretroviral therapy in northwestern Ethiopia: implication to reduce *Ethiop. J. Health Dev.* 2016;30(3)

transmission and multiple infections. HIV/AIDS (Auckland, NZ). 2015;7:119-124. Epub 2015/05/01.

- 136.Shiferaw L, Asrat D, Lemessa O, Yadeta D. Factors associated with adherence to antiretroviral therapy (ART) among adult people living with HIV and attending their clinical care, eastern Ethiopia. BMC International Health and Human Rights. 2015;15(33).
- 137.Sileshe M. Utilization of modern contraceptive methods and associated factors among reproductive age women on highly active antoretroviral therapy in Gambella Town, Ethiopia. [MA Thesis]: Addis Ababa University; 2015.
- 138.Soboka M, Tesfaye M, Feyissa GT, Hanlon C. Khat use in people living with HIV: a facility-based crosssectional survey from south west Ethiopia. BMC Psychiatry. 2015 April 03;15 (1) (no pagination)(69).
- 139. Taddesse D, Jamieson D, Cochrane L. Strengthening public health supply chains in Ethiopia: PEPFARsupported expansion of access and availability. Development in Practice. 2015 2015 Oct 3:25(7):1043-1056.
- 140. Tadele A. Assessment of health outcomes and predictors of survival of patients om highly active antiretroviral therapy at Debre Markos Hospital, North West Ethiopia: a retrospective cohort study. 26th Annual Conference of te Ethiopian Public Health Association; February 26-28, 20215; Bahir Dar, 2015.
- 141. Tadesse BT, Foster BA, Berhan Y. Cross sectional characterization of factors associated with pediatric HIV status disclosure in southern Ethiopia. PLoS ONE. 2015;10(7):e0132691.
- 142. Tariku A, Tegabu D, Meshesha M. Knowledge discovery for antiretroviral therapy adherence prediction: the case of Bahis Dar Felege Hiwot Referral Hospital. 26th Annual Conference of the Ehiopian Public Health Association; February 5-6, 2015; Bahir Dar, 2015.
- 143.Teasdale CA, Fayorsey R, Melaku Z, Chege D, Casalini C, Sebastian T, et al. Immune recovery at 5 years on ART in HIV+ children from four African countries. Topics in Antiviral Medicine. 2015 July;23:420.
- 144. Tekle Mariam K, Baraki N, Kedir H. Nutritional statue and its association with mortality in adult patients enrolled on ART at Fiche Hospital in north Shoa, Oromia Region, Ethiopia: a five year retrospective cohort study. 26th Annual Conference; February 26-28, 2015; Bahir Dar, 2015.
- 145. Teklemariam Z, Mitiku H, Mesfin F. Prevalence of anemia and nutritional status among HIV-positive children receiving antiretroviral therapy in Harar, eastern Ethiopa. HIV/AIDS (Auckland, NZ). 2015;7:191-196. Epub 2015/06/20.
- 146. Tekola-Ayele F, Adeyemo A, Aseffa A, Hailu E, Finan C, Davey G, et al. Clinical and pharmacogenomic implications of genetic variation in a southern Ethiopian population. The

pharmacogenomics journal. 2015 Feb;15(1):101-108. Epub 2014/07/30.

- 147. Tesfahuneygn G, Medhin G, Legesse M. Adherence to anti-tuberculosis treatment and treatment outcomes among tuberculosis patients in Alamata District, northeast Ethiopia. BMC Research Notes. 2015;8:503.
- 148. Teshome W, Asefa A. Correction: predictors of immunological failure of antiretroviral therapy among HIV infected patients in Ethiopia: a matched case-control study. PLoS One. 2015 May 2015 2015-05-16;10(5).
- 149. Teshome W, Belayneh M, Moges M, Endriyas M, Mekonnen E, Ayele S, et al. Who takes the medicine? adherence to antiretroviral therapy in southern Ethiopia. Patient Prefer Adherence. 2015;9:1531-1537. Epub 2015/11/26.
- 150. Teshome W, Belayneh M, Moges M, Mekonnen E, Endrias M, Ayele S, et al. Do loss to follow-up and death rates from ART care vary across primary health care facilities and hospitals in south Ethiopia? A retrospective follow-up study. HIV/AIDS (Auckland, NZ). 2015;7:167-174. Epub 2015/06/13.
- 151.Teshome W, Tefera A. Detection of immunological treatment failure among HIV infected patients in Ethiopia: a retrospective cohort study. BMC Immunology. 2015 September 16;16 (1) (no pagination)(55).
- 152. Teshome Yimer Y, Yalew AW. Magnitude and predictors of anti-retroviral treatment (ART) failure in private health facilities in Addis Ababa, Ethiopia. PLoS One. 2015;10(5):e0126026. Epub 2015/05/07.
- 153.Tessema SB, Adane MM. Assessment of antiretroviral treatment (ART) care service provision in Tigray Region health centers, North Ethiopia. BMC Health Services Research. 2015 Sep 10;15:368. Epub 2015/09/12.
- 154.Titiyos A, Jerene D, Enquselasie F. The yield of screening symptomatic contacts of multidrugresistant tuberculosis cases at a tertiary hospital in Addis Ababa, Ethiopia. BMC Research Notes. 2015 2015 2016-04-16;8.
- 155. Tola HH, Shojaeizadeh D, Garmaroudi G, Tol A, Yekaninejad MS, Ejeta LT, et al. Psychological distress and its effect on tuberculosis treatment outcomes in Ethiopia. Global Health Action. 2015;8(29019).
- 156.Tsega B, Srikanth BA, Shewamene Z. Determinants of non-adherence to antiretroviral therapy in adult hospitalized patients, Northwest Ethiopia. Patient Prefer Adherence. 2015;9:373-380. Epub 2015/03/19.
- 157.Tymejczyk O, Hoffman S, Kulkarni SG, Gadisa T, Lahuerta M, Remien RH, et al. HIV care and treatment beliefs among patients initiating antiretroviral treatment (ART) in Oromia, Ethiopia. AIDS and Behavior. 2015 2015 Sep 7:[11] p.
- 158. Tzur F, Chowers M, Agmon-Levin N, Mekori YA, Hershko AY. Increased Prevalence of diabetes *Ethiop. J. Health Dev.* 2016;30(3)

mellitus in a non-obese adult population: HIVinfected Ethiopians. The Israel Medical Association Journal (IMAJ). 2015 Oct;17(10):620-623. Epub 2015/12/17.

- 159. Villadsen SF, Negussie D, GebreMariam A, Tilahun A, Friis H, Rasch V. Antenatal care strengthening for improved quality of care in Jimma, Ethiopia: an effectiveness study. BMC Public Health. 2015;15:360.
- 160. Walleglign MA. Survival status among people living with HIV/AIDS on highly active antiretroviral therapy at Federal Police Referral Hospital, Ethiopia: retrospective cohort study. [MPH Thesis]: Addis Ababa University; 2015.
- 161.Wang T, Hoag SW, Eng ML, Polli J, Pandit NS. Quality of antiretroviral and opportunistic infection medications dispensed from developing countries and Internet pharmacies. Journal of Clinical Pharmacy and Therapeutics. 2015 01 Feb;40(1):68-75.
- 162. Worku M, Gashe F. Non adherence to antidepressant drugs and its effect on relapse and recurrence of depression. Pharmanest. 2015;6(2):2760-2764.
- 163. Yadesa TM, Gudina EK, Angamo MT. Antimicrobial use-related problems and predictors among hospitalized medical in-patients in southwest Ethiopia: prospective observational study. PLoS One. 2015;10(12):e0138385. Epub 2015/12/10.
- 164. Yimer YT, Yalew AW. Magnitude and predictors of anti-retroviral treatment (ART) failure in private health facilities in Addis Ababa, Ethiopia. PLoS ONE. 2015;10(5).
- 165. Yirdaw KD, Hattingh S. Prevalence and predictors of immunological failure among HIV patients on HAART in southern Ethiopia. PLoS One. 2015;10(5):e0125826. Epub 2015/05/12.
- 166. Yitibarek LT. Incidence of treatment failure and predictors among HIV/AIDS adult patients on ART at Debre Tabor Hospital, northwest Ethiopia: a retrospective follow-up study 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.

Section 7: Health Informatics Monitoring, and Evaluation Research

This section deals with the use of modeling and prediction, quantitative assessment, analytic and communication methods and research concerning monitoring and evaluation of HIV/AIDS programs. It covers the systematic application of information, computer science, and technology for HIV/AIDS prevention, care, research and evaluation (1-27). More than three quarter of the studies in this section focus on development of assessment, the analytic and communication tools, mostly models, epidemiological, diagnostic and scaling methods, and the remaining studies dealt with monitoring and evaluating HIV/AIDS, TB and malaria.

Ababeye (1) revealed that pathways to change (PtC) appear effective in ensuring that behavior, rather than information, becomes the focus of behavioral interventions. The study indicated that PtC works by changing the way health promoters think and work; enables to understand how peer educators see their job and more importantly how they perform their job. Bekalu and Eggermont (6) used the integrative model of behavioral prediction to assess the impact of exposure to HIV/AIDS-related media content on HIV testing intention. On the other hand, a study by Nigatu et al. (18) documented providing sexual and reproductive health (SRH) information to university students and peer educators through short message service (SMS). Girma et al. (12) indicated signs and predictors of improvements for stigma against people with HIV/AIDS in rural Ethiopia.

Efforts were also made to model different aspects of the HIV/AIDS pandemic. Abraha and Nigatu (2) modelled trends of health and health related indicators in Ethiopia (1995-2008) while Shemamane (23) fitted a statistical model for CD4-cell measurements of adult patients after the start of highly active ART in Yekatit 12 Hospital. Parametric modeling of survival data based on HIV-infected adult patients under HAART in Zewditu Referral Hospital was done by Legesse (16). Mathematical analysis of a model for AVL-HIV co-endemicity was also done by Hussaini et al.(14). Granich at al.(15) analyzed trends in AIDS deaths, new Infections and ART coverage in the top 30 countries with the highest AIDS mortality burden; 1990-2013.

Moreover, monitoring and evaluation were also done by few researchers. Intensified tuberculosis case finding among HIV-infected persons using a WHO symptom screen and Xpert((R)) MTB/RIF was done by Adelman et al. (3). Assefa et al. (4) conducted a cross sectional study to evaluate routine contact investigation in Addis Ababa, Ethiopia which is a missed opportunity to prevent tuberculosis in Children. Another study by Barber-Madden et al. (6) demonstrated final performance evaluation of Ethiopia food by prescription [Vienna, Virginia]. Global epidemiology of HIV among female sex workers with a special focus on its influence of structural determinants was done by Shannon et al. (22). Short and Goldberg (24) did an estimate of the prevalence of children living with HIV-infected adults for 23 Countries in sub-Saharan Africa.

When it comes to improving HIV/AIDS care Beyene and Beyene (8) assessed predictors of late HIV diagnosis among adult people living with HIV/AIDS who undertake an initial CD4 T cell evaluation, Northern Ethiopia whereas Bizuayehu et al.(8) did an assessment of duration of staying free from acquiring rehappening opportunistic infections among pre-ART people living with HIV/AIDS between 2008 and 2013. HIV treatment scale-up and HIV-related stigma in Sub-Saharan Africa was done by Chan et al. (9) using a longitudinal crosscountry analysis. A community based cohort study in Southern Ethiopia documented follow-up of chronic coughers improves tuberculosis case finding Woldesemayat et al.(25).

A couple of studies were focused on the health care financing of HIV/AIDS programs. A quantitative study was conducted by Chin et al. (10) on PEPFAR funding and reduction in HIV infection rates in 12 focus Sub-Saharan African countries. Forsythe et al. (11) explored the past, present and future of HIV/ AIDS and resource allocation. The tracking of the Global Fund HIV/AIDS resources used for sexual and reproductive health service integration in Ethiopia was also done by Mookherji et al. (17). Moreover, a study by Johansson et al. (15) dealt with health gains and financial protection from pneumococcal vaccination and pneumonia treatment in Ethiopia using an extended cost-effectiveness analysis.

In additon, certain studies were focused on assessing the quality of diagnostic procedures. For instance, a study by Shanks et al. (19) assessed the impact of visceral leishmaniasis for accounting false positivity in HIV tests, while another study by Shanks et al. (20) conducted a novel alternative for confirmation testing in resource limited settings by employing dilution testing using rapid diagnostic tests in a HIV diagnostic algorithm. Another study by Shanks et al. (21) did an evaluation of HIV testing algorithms in Ethiopia by assessing the role of the tie-breaker algorithm and weakly reacting test lines in contributing to a high rate of false positive HIV diagnoses. Another study recommeded an effort should be exerted towards host-directed therapies for tuberculosis Zumla et al.(26).

- 1. Ababeye B. Changing how youth peer educators promote behavior change: the case of the Pathways to Change Game. 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- Abraha MW, Nigatu TH. Modeling trends of health and health related indicators in Ethiopia (1995-2008): a time-series study. Health Research Policy and Systems. 2009 Dec 13;7:29. Epub 2009/12/17.
- Adelman MW, Tsegaye M, Kempker RR, Alebachew T, Haile K, Tesfaye A, et al. Intensified tuberculosis case finding among HIV-infected persons using a WHO symptom screen and Xpert((R)) MTB/RIF. The International Journal of Tuberculosis and Llung Disease. 2015 Oct;19(10):1197-1203. Epub 2015/10/16.
- Assefa D, Klinkenberg E, Yosef G. Cross Sectional Study Evaluating Routine Contact Investigation in Addis Ababa, Ethiopia: A Missed Opportunity to Prevent Tuberculosis in Children. PLoS One. 2015;10(6):e0129135. Epub 2015/06/18.
- 5. Barber-Madden R, Beyero M, Hansch S. Final performance evaluation of Ethiopia Food by

Prescription. [Vienna, Virginia], International Business and Technical Consultants, 2015 Mar 25., 2015.

- 6. Bekalu MA, Eggermont S. Exposure to HIV/AIDSrelated media content and HIV testing intention: applying the integrative model of behavioral prediction. Mass Communication & Society. 2015 Mar;18(2):144-164.
- Beyene MB, Beyene HB. Predictors of late HIV diagnosis among adult people living with HIV/AIDS who undertake an initial CD4 T cell evaluation, northern Ethiopia: a case-control study. PLoS ONE. 2015 08 Oct;10 (10) (no pagination)(e0140004).
- 8. Bizuayehu HM, Abyu DM, Aweke AM. Assessment of duration of staying free from acquiring rehappening opportunistic infections among pre-ART people living with HIV/AIDS between 2008 and 2013. Biomed Res Int. 2015;2015:146306. Epub 2015/02/17.
- Chan BT, Tsai AC, Siedner MJ. HIV Treatment scale-up and HIV-relatedstigma in Sub-Saharan Africa: a longitudinal cross-country analysis. Am J Public Health. 2015 Aug;105(8):1581-1587. Epub 2015/06/13.
- Chin RJ, Sangmanee D, Piergallini L. PEPFAR funding and reduction in HIV infection rates in 12 focus Sub-Saharan African countries: a quantitative analysis. International Journal of MCH and AIDS. 2015;3(2):150-158. Epub 2015/01/01.
- Forsythe S, Stover J, Bollinger L. The past, present and future of HIV, AIDS and resource allocation. BMC Public Health. 2009 Nov 18;9 Suppl 1:S4. Epub 2009/12/16.
- 12. Girma E, Abebe L, Morankar SN. Stigma against people with HIV/AIDS in rural Ethiopia, 2005-2011: signs and predictors of improvements. 26th Annual Conference of the Ethiopian Public Health Association February 26-28, 2015; Bahir Dar, 2015.
- 13. Granich R, Gupta S, Hersh B, Williams B, Montaner J, Young B, et al. Trends in AIDS deaths, new infections and ART coverage in the top 30 countries with the highest AIDS mortality burden; 1990-2013. PLoS One. 2015;10(7):e0131353. Epub 2015/07/07.
- Hussaini N, Lubuma JM, Barley K, Gumel AB. Mathematical analysis of a model for AVL-HIV coendemicity. Mathematical Biosciences. 2016 Jan;271:80-95. Epub 2015/11/26.
- 15. Johansson KA, Memirie ST, Pecenka C, Jamison DT, Verguet S. Health gains and financial protection from pneumococcal vaccination and pneumonia treatment in Ethiopia: results from an extended cost-effectiveness analysis. PLoS One. 2015 Dec 2015 2015-12-12;10(12).
- Legesse H. Parametric modeling of survival data based on HIV-infected adult patients under HAARt: a case at Zewditu referral hospital.: Addis Ababa University; 2015.
- 17. Mookherji S, Ski S, Huntington D. Tracking Global Fund HIV/AIDS resources used for sexual and *Ethiop. J. Health Dev.* 2016;30(3)

reproductive health service integration: case study from Ethiopia. Global Health. 2015 May 27;11:21. Epub 2015/05/28.

- 18. Nigatu T, McNabb M, Collins C, Koreta W. Providing sexual and reproductive health (SRH) information to university students and peer educators through short message service (SMS). 26th Annual Conference of the Ethiopian Public Health Association; February 26-28, 2015; Bahir Dar, 2015.
- 19. Shanks L, Ritmeijer K, Piriou E, Ruby Siddiqui M, Kliescikova J, Pearce N, et al. Accounting for false positive HIV tests: is visceral leishmaniasis responsible? PLoS ONE. 2015 10 Jul;10 (7) (no pagination)(e0132422).
- 20. Shanks L, Siddiqui MR, Abebe A, Piriou E, Pearce N, Ariti C, et al. Dilution testing using rapid diagnostic tests in a HIV diagnostic algorithm: a novel alternative for confirmation testing in resource limited settings. Virology Journal. 2015;14.
- 21. Shanks L, Siddiqui MR, Kliescikova J, Pearce N, Ariti C, Muluneh L, et al. Evaluation of HIV testing algorithms in Ethiopia: the role of the tie-breaker algorithm and weakly reacting test lines in contributing to a high rate of false positive HIV diagnoses. BMC Infectious Diseases. 2015 February 03;15 (1) (no pagination)(769).
- Shannon K, Strathdee SA, Goldenberg SM, Duff P, Mwangi P, Rusakova M, et al. Global epidemiology of HIV among female sex workers: influence of structural determinants. Lancet (London, England). 2015 Jan 03;385(9962):55-71. Epub 2014/07/26.
- 23. Shemamane E. Statistical modeling of CD4-cell measurements of adult patients after the start of highly active ART: the case of Yekatit 12 Hospital. [MSc Thesis]: Addis Ababa University; 2015.
- 24. Short SE, Goldberg RE. Children living with HIVinfected adults: estimates for 23 countries in sub-Saharan Africa. PLoS One. 2015;10(11):e0142580. Epub 2015/11/18.
- 25. Woldesemayat EM, Datiko DG, Lindtjørn B. Follow-up of chronic coughers improves tuberculosis case finding: results from a communitybased cohort study in southern Ethiopia. PLoS One. 2015 Feb 2015 2015-02-28;10(2).
- 26. Zumla A, Maeurer M, Chakaya J, Hoelscher M, Ntoumi F, Rustomjee R, et al. Towards host-directed therapies for tuberculosis. Nature Reviews Drug Discovery. 2015 Aug;14(8):511-512. Epub 2015/07/18.

Section 8: Diaspora Research

This section includes studies on HIV/AIDS among Ethiopians in the Diaspora and of Ethiopian health professionals in the Diaspora contributing to HIV/AIDS interventions in Ethiopia.

Two of the seven studies in this section present evidence of HIV transmission among Africa-born persons in Canada (1,2) and in a county in Washington State in the USA (5). Although HIV transmission among these populations has been recognized in these and earlier studies, absence of information in local health records and in the national census on the national origin of African patients does not permit analysis of HIV prevalence by nationalities. A study of attitudes, perceptions and practices surrounding HIV testing among East African immigrant women living in Washington DC revealed many cultural and personal barriers to voluntary testing. Most women were tested inadvertently or as part of applying for visas or employment (3). The persistence of cultural and demographic barriers to HIV testing in the diaspora was also reported from Ethiopian immigrants in Sweden (6). Grossman et al. (4) found that the number of HIV-infected Ethiopia-born persons living in Israel increased 1.3-fold between 2005 and 2013. Their phylogenetic studies of HIV indicated cross-ethnic spread of the virus through various independent introductions by different immigrant groups. Martinez Ortiz et al. (7) carried out one of the few studies of HIV among Ethiopian children in the diaspora. They found that 4.8% of 251 children, mostly below five years old, who arrived in Spain between 2006 and 2010, were HIVinfected.

- Brophy J, Lee T, Sauve L, Bitnun A, Singer J, Kakkar F, et al. Geographic origin trends among HIV+ mothers and children in Canada and impact on vertical HIV transmission rates. Journal of the International AIDS Society. 2015;18:117.
- Brophy JC, Sauve LJ, Bitnun A, Lee T, Kakkar F, Vaudry W, et al. Geographic origins of HIV+ mothers giving birth in Canada (1990-2013). Canadian Journal of Infectious Diseases and Medical Microbiology. 2015 March-April;26:81B.
- 3. De Jesus M, Carrete C, Maine C, Nalls P. Attitudes, perceptions and behaviours towards HIV testing among African-American and East African immigrant women in Washington, DC: implications for targeted HIV testing promotion and communication strategies. Sexually Transmitted Infections. 2015 Dec 2015 2016-04-19;91(8):569.
- 4. Grossman Z, Avidor B, Mor Z, Chowers M, Levy I, Shaha E, et al. A population-structured HIV epidemic in Israel: roles of risk and ethnicity. PLoS ONE. 2015 24 Aug;10 (8) (no pagination)(e0135061).
- 5. Kerani R, Herbeck J, Buskin S, Dombrowski J, Bennett A, Barash E, et al. Evidence of local HIV transmission in the African community of King County, Washington. Journal of the International AIDS Society. 2015;18:118-120.
- 6. Lindkvist P, Johansson E, Hylander I. Fogging the issue of HIV barriers for HIV testing in a migrated population from Ethiopia and Eritrea. BMC Public Health. 2015 Feb 05;15:82. Epub 2015/02/06.
- 7. Martinez Ortiz A, Dominguez Pinilla N, Wudineh M, Gonzalez-Granado LI. International adoption

from Ethiopia in a 5-year period. [Spanish]. Anales de Pediatria. 2015 01 May;82(5):302-307.

Section 9: Previous bibliographies

This section lists the previous year's update and potentially other bibliographies that were published during 2015. Another 12 bibliographies were published between 2003 and 2014 in this journal.

Haile Mariam D, Kloos H, Converse PJ, Mekonnen W, Mulatu MS, Kaba M, Beyene A. Bibliography on HIV/AIDS in Ethiopia and Ethiopians in the Diaspora: The 2014 Update. Ethiop J Health Dev 2015; 29(1): 43-78.

Section 10: Selected Websites Featuring HIV/AIDS in Ethiopia

- Federal HIV/AIDS Prevention and Control Office of Ethiopia: <u>http://hapco.gov.et</u>
- 2. The Ethiopian Journal of Health Development): http://www.ejhd.org
- 3. Ethiopian AIDS Resources Center: <u>http://www.etharc.org</u>
- 4. Family Health International: <u>http://www.fhi360.org</u> /countries/ethiopia
- 5. Christian Relief and Development Association: www.crdaethiopia.org
- Johns Hopkins University Center for Clinical Global Health Education: <u>http://main.ccghe.net/CCG/</u> <u>country/ethiopia</u>
- 7. People to People Organization: <u>http://www.peoplepeople.org</u>
- 8. Save the Children: <u>http://www.savethechildren.</u> org/site/c.8rKLIXMGIpI4E/b.6234245/k. <u>A159/HIVAids_Programs.htm?msource=weilpres05</u> <u>11#Ethiopia</u>

- 9. United Nations Children's Fund (UNICEF): http://www.unicef.org/ethiopia/hiv aids 464.html
- 10. United Nations Development Program (UNDP):<u>http://www.undp.org/content/undp/en/home</u>/ourwork/hiv-aids/Projects-initiatives/hiv-epidemicethiopia-case-study-transformational-change/
- 11. United Nations Joint Program on AIDS (UNAIDS):<u>http://www.unaids.org/en/Regionscountries/Countries/Ethiopia</u>
- 12. United States Agency for International Development: <u>http://www.usaid.gov/ethiopia/global-health</u>
- 13. United States Centers for Disease Control and Prevention (CDC): <u>http://www.cdc.gov/globalaids/</u> <u>Global-HIV-AIDS-at-CDC/countries/Ethiopia/</u>
- 14. AIDS Portal: <u>http://www.aidsportal.org/web/</u> guest/ethiopia
- 15. University of California, San Francisco HIV In Site: http://hivinsite.ucsf.edu/global?page=cr09-et-00
- 16. The International Technical Training and Education Center on HIV (I-TECH) of the University of Washington: <u>http://www.go2itech.org/itech?page=co-03-00</u>
- 17. The International Center for AIDS Care and Treatment Programs (ICAP) at Columbia University's Mailman School of Public Health: http://icap.columbia.edu/where-we-work/ethiopia
- 18. World Health Organization: <u>http://www.who.int/</u> <u>countries/eth/en/</u>
- 19. Management Sciences for Health's Ethiopia Network for HIV/AIDS Treatment, Care and Support (ENHAT-CS) Project: <u>http://www.msh.org/our-work/projects/ethiopianetwork-for-hivaids-treatment-care-support</u>
- 20. The Twinning Center: <u>http://www.</u> <u>twinningagainstaids.org/ethiopia.html</u>