

Heliotherapy: A South African perspective

Y Moosa, D J Esterhuyse

The research objective was to gather and collate data that will enable dermatologists to quantify exposure to solar radiation so that they can give accurate advice to patients using heliotherapy, thereby minimising harm from sun exposure. Other patients can also be advised regarding ultraviolet index (UVI) and sun safety.

The concept of minimal erythema dose per hour (MED/h) may be useful in future research into solar radiation and its effects on skin cancer.

S Afr Med J 2010; 100: 728-733.

In South Africa, dermatologists are often faced with a patient who requires phototherapy but, for financial or practical reasons (distance to a specialist centre, disruption of schooling, etc.), is unable to attend phototherapy clinics. Heliotherapy (the medicinal use of natural sunlight) has been practised for centuries. The main indication for heliotherapy is psoriasis, but it has also been used for treating atopic dermatitis and vitiligo; theoretically, it could be adapted to treat the pruritus from renal failure and HIV.¹⁻³

Current approaches

Several approaches to using sunlight to treat skin eruptions have been described. In the Dead Sea basin, physicians have advocated 10 - 20 minutes' sun exposure twice daily, followed by daily increments of 10 minutes until a maximum of 3 - 6 hours per day is reached. The recommended period of treatment ranges from 3 to 4 weeks.⁴ This method is largely generic, with daily adjustments based on individual responses to ultraviolet radiation (UVR) and the season.^{5,6} Its disadvantage is that variations in solar ultraviolet type B (UVB) due to changes in the thickness of the ozone layer, daily variations in cloud cover, rain, aerosol content of the air, precipitable water and season are not accounted for.^{7,8} Exposure to ultraviolet type A (UVA) is also not considered for the same reasons.

An alternative approach is to determine the minimal erythema dose (MED) of the patient in the phototherapy unit. Heliotherapy is then started at 70% of the MED, with increases of 10 - 15% of MED for skin types I to III and 15 - 20% of MED for skin types IV and V with each treatment. Patients undergo 3 - 5 treatments per week; the duration of treatment is at the doctor's discretion.^{8,9}

Avraham Kushelevsky *et al.*¹⁰ devised the concept of MED per hour (MED/h) to overcome these issues when using heliotherapy in the Dead Sea basin. The concept differs from MED (energy per unit area); the latter varies according to skin type, and is the minimum single dose of UVR required to produce erythema after 24 hours at an exposed site.¹¹ In contrast, MED/h is defined as the dose causing minimal redness of the average Fitzpatrick skin type 2 after 1 hour of irradiation.

In South Africa, although dermatologists advise suitable patients to expose themselves to sunlight, there is no scientific consensus regarding time and duration of exposure, which can be misleading and vague for patients. We aimed not to replace the controlled environment of the phototherapy unit but to provide less-privileged patients with more specific, evidence-based advice on the therapeutic use of sunlight, using the concept of MED/h.

Research setting

UVB data were collected at different times of the day throughout the year from meteorological stations in three major South African cities: east coast of South Africa in Durban, latitude 29.9 S, longitude 31.0 E, sea level; west coast of South Africa in Cape Town (latitude 34.0 S, longitude 18.6 E, sea level); and inland at the headquarters of the South African Weather Service (SAWS) in Pretoria (latitude 25.8 S, longitude 28.3 E, altitude 1 541 m).

The instrumentation to measure UVB radiation at all three sites was identical, consisting of a Model 501A UV-Biometer (Solar Light Co., Glenside, PA, USA). All UVR meters are positioned to monitor UVR intensity on a horizontal surface. The Biometer measures the biological effectiveness of UVB radiation in units of MED/h. Data were collected from readings taken over 11 years in Cape Town and Pretoria, and 6 years in Durban. The average hourly UVB values for times from 07h00 - 18h00 were calculated for each week of the year.

Results and applications

The average MED/h between 07h00 and 18h00 throughout the 52 weeks of the year for the three cities is shown in Tables I - III and Figs 1 - 3. In the figures, series 1 is the average of the values between 07h00 and 08h00, series 2 represents the values between 08h00 and 09h00, etc. All values in Tables I - III are expressed in MED/h. This information can be used to give precise instructions to the patient regarding therapeutic exposure to sunlight, depending on skin type. For skin type I, a starting dose of 0.02 J/cm² is recommended, while for skin types II - III, skin type IV and for skin type VI the starting doses are 0.03 J/cm², 0.05 J/cm² and 0.06 J/cm², respectively. The following equation is then used to calculate the exact exposure time required at different times of the day and year using the data from

Table I. Common conversions (adapted from www.solarlight.com)

MED (minimal erythema dose) conversions

1 MED/h = 0.05833 W/m²

1 MED = 210 J/m²

1 MED/h = 2.33 UV index

Radiance conversions

1 mW/cm² = 10 W/m²

Department of Dermatology, Steve Biko Academic Hospital and University of Pretoria
Yaaseen Moosa, MB BCH, FC Derm (SA), DTM&H

South African Weather Service, Pretoria
D J Esterhuyse

Corresponding author: Y Moosa (ymoosa75@gmail.com)

Table II. Definition of weeks in terms of calendar days

Week No.	Start	End
1	1 Jan	7 Jan
2	8 Jan	14 Jan
3	15 Jan	21 Jan
4	22 Jan	28 Jan
5	29 Jan	4 Feb
6	5 Feb	11 Feb
7	12 Feb	18 Feb
8	19 Feb	25 Feb
9	26 Feb	5 Mar
10	6 Mar	12 Mar
11	13 Mar	19 Mar
12	20 Mar	26 Mar
13	27 Mar	2 Apr
14	3 Apr	9 Apr
15	10 Apr	16 Apr
16	17 Apr	23 Apr
17	24 Apr	30 Apr
18	1 May	7 May
19	8 May	14 May
20	15 May	21 May
21	22 May	28 May
22	29 May	4 Jun
23	5 Jun	11 Jun
24	12 Jun	18 Jun
25	19 Jun	25 Jun
26	26 Jun	2 Jul
27	3 Jul	9 Jul
28	10 Jul	16 Jul
29	17 Jul	23 Jul
30	24 Jul	30 Jul
31	31 Jul	6 Aug
32	7 Aug	13 Aug
33	14 Aug	20 Aug
34	21 Aug	27 Aug
35	28 Aug	3 Sep
36	4 Sep	10 Sep
37	11 Sep	17 Sep
38	18 Sep	24 Sep
39	25 Sep	1 Oct
40	2 Oct	8 Oct
41	9 Oct	15 Oct
42	16 Oct	22 Oct
43	23 Oct	29 Oct
44	30 Oct	5 Nov
45	6 Nov	12 Nov
46	13 Nov	19 Nov
47	20 Nov	26 Nov
48	27 Nov	3 Dec
49	4 Dec	10 Dec
50	11 Dec	17 Dec
51	18 Dec	24 Dec
52	25 Dec	31 Dec

Tables I - III, bearing in mind that 1 MED/h is the equivalent of an irradiance of 0.05833 W/m², or 0.00583 mW/cm² (Table I):

$$\text{Irradiation time (seconds)} = \text{starting dose (J/cm}^2\text{)} \times 1000 / \text{irradiance (mW/cm}^2\text{)}^{12}$$

For example, consider a patient with psoriasis and Fitzpatrick skin type III skin seen in the 3rd week of January in Pretoria, for whom

daily exposure at 10h00 is convenient. The starting dose for this skin type is 0.03 J/cm². The MED/h for 10h00 - 11h00 on the 3rd week of January is 3.32 (UVB data Table III), which is equivalent to 0.019 mW/cm². Therefore, the exposure time at 10h00 is calculated as follows:

$$\begin{aligned} \text{Irradiation time (seconds)} &= 0.03 \times 1000 / 0.019 = 1578 \text{ seconds} \\ &= 26.3 \text{ minutes.} \end{aligned}$$

This patient can be advised to expose himself for 26.3 minutes a day at 10h00 in week 3 of January. The exposure times for subsequent weeks can be calculated in a similar manner, and the patient be given a written regimen. The doctor prescribing the heliotherapy can therefore quantify and adjust exposure to sunlight.

The UV data presented are also valuable in calculating the UV index (UVI), a concept with which the general public is familiar in preventing overexposure to solar radiation during outdoor activities. One MED/h equates to a UVI of 2.33, so in the example given above, the psoriatic patient will be exposed to a UVI of $3.32 \times 2.33 = 7.7$.

Discussion

The logistical limitation of carrying out phototherapy in South Africa, where much of the population live in isolated rural areas and lack the means for regular travel, led to this study. We aimed to assist dermatologists in employing a more scientific approach in the use of sunlight – a free, widely available, natural therapeutic resource. Heliotherapy (or climatotherapy) has been used extensively to treat thousands of patients in the Dead Sea basin, Switzerland and Sweden for many decades.¹²⁻¹⁴ Adapting this experience for use in South Africa, where many patients have little therapeutic alternative, is entirely possible and can be done in a controlled and quantifiable manner, taking into account seasonal and even hourly variations in UVR.

As with conventional phototherapy, heliotherapy requires precautions. A thorough history and physical examination should precede any consideration for heliotherapy, and should only be considered if the patient is unable to attend specialist phototherapy centres. Any potentially photosensitising drugs should be noted, as should a history of skin cancer, previous exposure to ionising radiation and skin type. A test for serum antinuclear antibodies may be indicated if there is a suggestion of an underlying photosensitive connective tissue disorder. The patient should be advised that the development of erythema is an integral part of the treatment. Strict compliance with treatment protocol should be emphasised, and solar exposure should be limited to the early morning and late afternoon hours.⁵ Avoidance of unnecessary sun exposure should be stressed, and patients should be advised to report any adverse events immediately, wear sunglasses, apply sunscreen to face and hands, and cover the genitalia during exposure to natural sunlight.^{10,13}

Absolute contraindications to heliotherapy include xeroderma pigmentosum and other disorders predisposing to sun-induced cancers. Caution is appropriate if there is a past history of non-melanoma skin cancer, immunosuppression, pemphigus, pemphigoid, cataracts, aphakia or photosensitivity.^{2,10} A family history of melanoma is also reason for caution.

Acute side-effects of heliotherapy include sunburn, pruritus and a polymorphous light eruption-like response, which is usually transient and affects mainly patients with skin types II and III.¹⁴

Little is known about the chronic side-effects of long-term sun exposure and in particular the cumulative dose of solar UVR predisposing to the development of cutaneous malignancies.¹⁵ An important aspect of this work is that exposure to sunlight can be quantified, paving the way for further research into this area. Another useful application of this work is the ability to calculate UVI,

Table III. UVB data for Cape Town, Durban and Pretoria in MED/h

	Durban																											
	Week No.																											
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26		
7 am	0.56	0.47	0.48	0.42	0.41	0.37	0.32	0.31	0.27	0.27	0.23	0.2	0.19	0.17	0.14	0.12	0.1	0.09	0.08	0.06	0.05	0.05	0.04	0.03	0.03	0.03	0.03	
8 am	1.28	1.12	1.12	1.07	1.09	0.96	0.92	0.88	0.81	0.83	0.73	0.66	0.59	0.58	0.51	0.45	0.39	0.35	0.32	0.28	0.24	0.22	0.19	0.18	0.17	0.16	0.16	
9 am	2.23	2	2.1	2.04	2	1.87	1.81	1.74	1.61	1.67	1.5	1.32	1.24	1.21	1.09	1	0.86	0.78	0.76	0.64	0.57	0.53	0.47	0.44	0.43	0.41	0.41	
10 am	3.21	2.75	3.1	3.02	3.01	2.81	2.68	2.56	2.54	2.52	2.28	2.08	1.98	1.89	1.67	1.55	1.41	1.27	1.2	1.01	0.92	0.89	0.78	0.76	0.73	0.71	0.71	
11 am	3.85	3.27	3.62	3.66	3.74	3.48	3.45	3.13	3.12	3.17	2.88	2.64	2.42	2.25	2.07	1.95	1.71	1.58	1.52	1.3	1.16	1.11	0.98	0.95	0.93	0.93	0.93	
12 pm	3.98	3.34	3.82	3.7	3.92	3.63	3.67	3.23	3.34	3.32	2.97	2.71	2.55	2.23	2.14	2.03	1.72	1.64	1.51	1.3	1.2	1.13	0.98	0.97	0.97	0.96	0.96	
1 pm	3.52	2.94	3.36	3.34	3.53	3.28	3.33	2.89	2.98	2.89	2.6	2.3	2.15	1.98	1.78	1.68	1.42	1.32	1.23	1.07	0.96	0.89	0.79	0.77	0.77	0.78	0.78	
2 pm	2.6	2.15	2.47	2.53	2.6	2.41	2.48	2.06	2.18	2.04	1.81	1.59	1.44	1.3	1.16	1.07	0.9	0.83	0.76	0.64	0.58	0.54	0.48	0.47	0.46	0.47	0.47	
3 pm	1.51	1.27	1.46	1.51	1.51	1.47	1.43	1.2	1.25	1.15	1	0.85	0.75	0.67	0.55	0.52	0.43	0.37	0.33	0.29	0.24	0.22	0.19	0.19	0.18	0.2	0.2	
4 pm	0.65	0.57	0.67	0.66	0.66	0.68	0.65	0.48	0.53	0.45	0.39	0.29	0.25	0.22	0.16	0.15	0.11	0.1	0.08	0.06	0.06	0.05	0.04	0.04	0.04	0.05	0.05	
5 pm	0.21	0.19	0.23	0.21	0.2	0.21	0.2	0.12	0.14	0.11	0.09	0.06	0.04	0.03	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0	0	0	0	0	0	
6 pm	0.04	0.04	0.04	0.03	0.03	0.03	0.03	0.02	0.02	0.01	0	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Week No.																											
	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	52	
7 am	0.03	0.03	0.03	0.04	0.05	0.06	0.07	0.08	0.11	0.13	0.17	0.2	0.24	0.27	0.31	0.36	0.4	0.51	0.47	0.52	0.58	0.6	0.58	0.6	0.54	0.58	0.58	
8 am	0.16	0.17	0.17	0.18	0.2	0.25	0.28	0.31	0.38	0.43	0.51	0.58	0.64	0.67	0.74	0.82	0.96	1.12	1.07	1.13	1.24	1.22	1.24	1.3	1.15	1.29	1.29	
9 am	0.44	0.44	0.44	0.47	0.51	0.61	0.64	0.68	0.8	0.89	0.99	1.08	1.24	1.27	1.32	1.48	1.63	1.86	1.79	1.83	2.07	1.98	2.11	2.25	2.03	2.24	2.24	
10 am	0.76	0.74	0.73	0.8	0.88	1	1.07	1.12	1.25	1.37	1.49	1.56	1.81	1.79	1.84	2.1	2.25	2.47	2.52	2.57	2.89	2.76	2.9	3.12	2.8	3.21	3.21	
11 am	0.97	0.97	0.96	1.04	1.13	1.3	1.35	1.42	1.55	1.64	1.76	1.88	2.11	2.15	2.1	2.44	2.69	2.79	2.83	2.94	3.26	3.14	3.43	3.6	3.3	3.69	3.69	
12 pm	0.99	1.02	1	1.11	1.18	1.35	1.39	1.45	1.58	1.66	1.72	1.87	2.05	2.1	1.98	2.45	2.56	2.75	2.68	2.8	3.1	3.13	3.35	3.55	3.22	3.78	3.78	
1 pm	0.81	0.84	0.83	0.9	0.98	1.13	1.14	1.18	1.32	1.32	1.39	1.48	1.67	1.68	1.64	1.99	2	2.25	2.28	2.36	2.49	2.57	2.85	2.97	2.67	3.25	3.25	
2 pm	0.49	0.52	0.52	0.56	0.61	0.7	0.72	0.76	0.84	0.84	0.88	0.96	1.03	1.11	1.08	1.28	1.33	1.49	1.58	1.67	1.72	1.72	2.05	2.19	1.87	2.39	2.39	
3 pm	0.21	0.22	0.23	0.25	0.27	0.31	0.32	0.35	0.39	0.39	0.41	0.47	0.49	0.56	0.53	0.64	0.69	0.77	0.84	0.93	0.95	0.97	1.1	1.24	1	1.39	1.39	
4 pm	0.05	0.05	0.06	0.07	0.07	0.09	0.92	0.1	0.11	0.12	0.12	0.14	0.16	0.19	0.18	0.24	0.25	0.29	0.32	0.38	0.38	0.41	0.46	0.55	0.44	0.62	0.62	
5 pm	0	0	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.04	0.04	0.05	0.05	0.07	0.08	0.1	0.11	0.13	0.14	0.18	0.15	0.19	0.19	
6 pm	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.03	0.03	0.03	0.03	

Table III. Continued

Cape Town		Week No.																										
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	
7 am	0.3	0.28	0.24	0.2	0.17	0.16	0.12	0.09	0.07	0.05	0.05	0.03	0.02	0.01	0.01	0.01	0.01	0.01	0	0	0	0	0	0	0	0	0	0
8 am	0.92	0.89	0.83	0.74	0.68	0.65	0.54	0.47	0.39	0.34	0.31	0.24	0.2	0.16	0.13	0.1	0.1	0.1	0.08	0.07	0.06	0.04	0.04	0.03	0.02	0.02	0.02	0.02
9 am	1.92	1.89	1.86	1.67	1.6	1.55	1.37	1.23	1.1	0.98	0.93	0.75	0.66	0.56	0.48	0.39	0.38	0.32	0.28	0.27	0.27	0.21	0.19	0.15	0.15	0.14	0.13	0.13
10 am	3.14	3	3.02	2.76	2.64	2.71	2.47	2.27	2.06	1.86	1.79	1.49	1.33	1.18	1.04	0.84	0.85	0.75	0.65	0.62	1	0.78	0.74	0.63	0.39	0.37	0.36	0.36
11 am	4.11	3.99	4.02	3.78	3.71	3.78	3.52	3.28	2.91	2.74	2.61	2.27	2.02	1.82	1.6	1.32	1.34	1.16	1.02	1	1.22	0.94	0.9	0.77	0.82	0.77	0.76	0.76
12 pm	4.6	4.5	4.65	4.4	4.44	4.4	4.11	3.88	3.5	3.38	3.11	2.72	2.49	2.22	1.94	1.65	1.57	1.39	1.22	1.22	0.94	0.9	0.77	0.82	0.77	0.77	0.76	0.76
1 pm	4.52	4.52	4.68	4.47	4.49	4.44	4.21	3.97	3.54	3.44	3.13	2.78	2.48	2.22	1.95	1.62	1.5	1.37	1.21	1.15	0.88	0.88	0.73	0.76	0.73	0.73	0.76	0.76
2 pm	3.89	3.89	4.09	3.98	4	3.93	3.69	3.45	3.14	2.93	2.69	2.37	2.09	1.87	1.57	1.3	1.16	1.05	0.91	0.88	0.66	0.66	0.65	0.55	0.55	0.57	0.56	0.56
3 pm	2.91	2.95	3.04	3	2.99	2.92	2.72	2.53	2.26	2.03	1.88	1.56	1.37	1.23	1.03	0.8	0.67	0.61	0.51	0.48	0.36	0.36	0.35	0.3	0.29	0.3	0.31	0.31
4 pm	1.76	1.78	1.84	1.8	1.85	1.75	1.59	1.46	1.26	1.1	0.98	0.79	0.66	0.57	0.45	0.33	0.27	0.23	0.18	0.16	0.11	0.11	0.1	0.09	0.09	0.09	0.1	0.1
5 pm	0.8	0.83	0.84	0.82	0.83	0.77	0.67	0.59	0.09	0.41	0.33	0.24	0.19	0.15	0.11	0.07	0.05	0.04	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
6 pm	0.23	0.25	0.24	0.24	0.23	0.2	0.16	0.13	0.1	0.07	0.05	0.03	0.01	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	52	
7 am	0.03	0.03	0.03	0.04	0.05	0.06	0.07	0.08	0.11	0.13	0.17	0.2	0.24	0.27	0.31	0.36	0.4	0.51	0.47	0.52	0.58	0.58	0.6	0.58	0.6	0.54	0.58	
8 am	0.16	0.17	0.17	0.18	0.2	0.25	0.28	0.31	0.38	0.43	0.51	0.58	0.64	0.67	0.74	0.82	0.96	1.12	1.07	1.13	1.24	1.24	1.22	1.24	1.3	1.15	1.29	
9 am	0.44	0.44	0.44	0.47	0.51	0.61	0.64	0.68	0.8	0.89	0.99	1.08	1.24	1.27	1.32	1.48	1.63	1.86	1.79	1.83	2.07	1.98	1.98	2.11	2.25	2.03	2.24	
10 am	0.76	0.74	0.73	0.8	0.88	1	1.07	1.12	1.25	1.37	1.49	1.56	1.81	1.79	1.84	2.1	2.25	2.47	2.52	2.57	2.89	2.76	2.9	3.12	2.8	3.21	3.21	
11 am	0.97	0.97	0.96	1.04	1.13	1.3	1.35	1.42	1.55	1.64	1.76	1.88	2.11	2.15	2.1	2.44	2.69	2.79	2.83	2.94	3.26	3.14	3.43	3.6	3.3	3.69	3.69	
12 pm	0.99	1.02	1	1.11	1.18	1.35	1.39	1.45	1.58	1.66	1.72	1.87	2.05	2.1	1.98	2.45	2.56	2.75	2.68	2.8	3.1	3.13	3.35	3.55	3.22	3.78	3.78	
1 pm	0.81	0.84	0.83	0.9	0.98	1.13	1.14	1.18	1.32	1.32	1.39	1.48	1.67	1.68	1.64	1.99	2	2.25	2.28	2.36	2.49	2.57	2.85	2.97	2.67	3.25	3.25	
2 pm	0.49	0.52	0.52	0.56	0.61	0.7	0.72	0.76	0.84	0.84	0.88	0.96	1.03	1.11	1.08	1.28	1.33	1.49	1.58	1.67	1.72	1.72	2.05	2.19	1.87	2.39	2.39	
3 pm	0.21	0.22	0.23	0.25	0.27	0.31	0.32	0.35	0.39	0.39	0.41	0.47	0.49	0.56	0.53	0.64	0.69	0.77	0.84	0.93	0.95	0.97	1.1	1.24	1	1.39	1.39	
4 pm	0.05	0.05	0.06	0.07	0.07	0.09	0.92	0.1	0.11	0.12	0.12	0.14	0.16	0.19	0.18	0.24	0.25	0.29	0.32	0.38	0.38	0.41	0.46	0.55	0.44	0.62	0.62	
5 pm	0	0	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.04	0.04	0.05	0.05	0.07	0.08	0.1	0.11	0.13	0.14	0.18	0.15	0.19	0.19	
6 pm	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.03	0.03	0.03	0.03	

Table III. Continued

Pretoria

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
	Week No.																									
7 am	0.61	0.53	0.47	0.44	0.45	0.39	0.29	0.31	0.29	0.26	0.24	0.21	0.19	0.18	0.17	0.15	0.13	0.11	0.1	0.08	0.07	0.06	0.05	0.05	0.04	0.04
8 am	1.53	1.32	1.27	1.16	1.27	1.09	0.86	0.98	0.95	0.89	0.83	0.76	0.71	0.68	0.64	0.58	0.55	0.46	0.44	0.38	0.33	0.3	0.27	0.26	0.23	0.22
9 am	2.69	2.36	2.34	2.14	2.39	2.13	1.71	1.89	1.99	1.84	1.68	1.62	1.54	1.48	1.41	1.33	1.25	1.05	1.05	0.93	0.83	0.77	0.73	0.69	0.63	0.63
10 am	3.6	3.36	3.32	3.07	3.54	3.08	2.76	2.81	2.97	2.82	2.59	2.48	2.42	2.3	2.24	2.11	1.94	1.71	1.69	1.53	1.38	1.33	1.25	1.2	1.12	1.13
11 am	4.28	4	3.84	3.71	4.34	3.72	3.37	3.47	3.67	3.44	3.23	3.05	2.86	2.88	2.74	2.61	2.41	2.12	2.1	1.9	1.75	1.67	1.66	1.57	1.47	1.49
12 pm	4.29	4.16	4.04	3.8	4.51	3.81	3.5	3.7	3.85	3.62	3.33	3.09	2.94	2.88	2.78	2.63	2.48	2.13	2.1	1.86	1.73	1.7	1.66	1.56	1.48	1.51
1 pm	3.65	3.58	3.67	3.27	3.94	3.48	3.29	3.24	3.33	3.09	2.91	2.75	2.47	2.42	2.24	2.25	2.04	1.79	1.7	1.48	1.38	1.36	1.35	1.27	1.2	1.23
2 pm	2.69	2.81	2.69	2.55	2.81	2.71	2.37	2.45	2.43	2.25	2.11	1.81	1.81	1.68	1.51	1.47	1.29	1.16	1.04	0.9	0.84	0.83	0.82	0.78	0.74	0.76
3 pm	1.57	1.78	1.71	1.5	1.66	1.61	1.36	1.48	1.39	1.31	1.22	1	0.98	0.86	0.74	0.7	0.6	0.51	0.45	0.39	0.35	0.35	0.34	0.33	0.31	0.33
4 pm	0.79	0.8	0.76	0.7	0.79	0.73	0.59	0.62	0.59	0.51	0.46	0.35	0.33	0.26	0.21	0.2	0.16	0.13	0.1	0.09	0.08	0.07	0.07	0.07	0.07	0.07
5 pm	0.22	0.23	0.22	0.23	0.23	0.21	0.16	0.16	0.14	0.11	0.09	0.06	0.05	0.03	0.02	0.02	0.01	0.01	0	0	0	0	0	0	0	0
6 pm	0.03	0.03	0.02	0.02	0.03	0.02	0.01	0.01	0.01	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Week No.																									
7 am	0.04	0.04	0.04	0.05	0.05	0.07	0.08	0.09	0.13	0.16	0.18	0.22	0.3	0.33	0.38	0.44	0.54	0.56	0.63	0.6	0.7	0.63	0.66	0.66	0.61	0.6
8 am	0.23	0.23	0.25	0.26	0.27	0.34	0.36	0.39	0.52	0.59	0.59	0.7	0.87	0.93	0.97	1.08	1.34	1.37	1.43	1.39	1.57	1.4	1.48	1.52	1.4	1.38
9 am	0.63	0.64	0.67	0.69	0.72	0.88	0.91	0.94	1.16	1.29	1.25	1.4	1.72	1.74	1.76	1.91	2.28	2.36	2.41	2.28	2.59	2.34	2.45	2.59	2.47	2.45
10 am	1.13	1.16	1.21	1.22	1.28	1.53	1.54	1.68	1.85	2.01	1.98	2.13	2.52	2.55	2.57	2.73	3.11	3.23	3.27	3.01	3.44	3.2	3.41	3.54	3.42	3.55
11 am	1.5	1.53	1.6	1.61	1.71	2	2.01	2.17	2.34	2.49	2.4	2.59	3.07	2.97	3.02	3.07	3.51	3.72	3.7	3.39	3.8	3.71	3.98	3.98	4	4.03
12 pm	1.59	1.59	1.67	1.67	1.8	2.03	2.06	2.19	2.37	2.56	2.43	2.64	2.97	2.95	2.91	3.02	3.39	3.52	3.6	3.25	3.62	3.84	4.1	4.1	4.1	4.33
1 pm	1.28	1.35	1.38	1.4	1.52	1.7	1.71	1.84	2.02	2.16	2.02	2.27	2.49	2.47	2.41	2.54	2.76	2.94	2.91	2.77	3.02	3.27	3.3	3.63	3.6	3.91
2 pm	0.81	0.84	0.87	0.91	0.99	1.11	1.13	1.21	1.34	1.42	1.34	1.52	1.63	1.65	1.65	1.72	1.96	2.01	2.12	1.87	2.13	2.31	2.47	2.66	2.7	2.69
3 pm	0.35	0.37	0.39	0.41	0.46	0.51	0.54	0.57	0.65	0.69	0.66	0.76	0.8	0.82	0.84	0.9	1.03	1.07	1.08	0.97	1.12	1.23	1.44	1.66	1.67	1.61
4 pm	0.08	0.09	0.1	0.11	0.12	0.14	0.15	0.16	0.19	0.2	0.2	0.24	0.26	0.27	0.28	0.31	0.36	0.4	0.4	0.38	0.43	0.53	0.59	0.72	0.76	0.69
5 pm	0	0	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.03	0.03	0.04	0.04	0.05	0.06	0.07	0.08	0.08	0.11	0.13	0.14	0.19	0.21	0.2
6 pm	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.01	0.01	0.01	0.02	0.02	0.02




Fig. 1. Cape Town: Weekly averages of MED values for different hours of the day.




Fig. 2. Durban: Weekly averages of MED values for different hours of the day.




Fig. 3. Pretoria: Weekly averages of MED values for different hours of the day.

which is of value to dermatologists in advising their patients about ‘safe sun-exposure’. The World Health Organization classifies UVI values >9 as extreme risk, 9 - 7 as high risk, 7 - 4 as moderate risk, <4 as low risk, and <2 as negligible.¹⁶

UVB values across the three cities in South Africa are quite similar, possibly making it acceptable to combine and extrapolate these data for other cities in South Africa. Alternatively, region-specific values can be used, with the Cape provinces using values for Cape Town, Gauteng, Free State, North-West province and Limpopo the values for Pretoria, and KwaZulu-Natal the values for Durban.

We thank Dr H Carman (consultant dermatologist in private practice in Johannesburg) for initially reviewing the paper; Professor G Todd (University of Cape Town) for reviewing the article; and Dr V Akhras (St George’s University Hospital, London) for reviewing and editing the article.

References

1. Gilchrest BA, Rowe JW, Brown RS, et al. Relief of uremic pruritus by UV therapy. *N Engl J Med* 1997;297: 36-138.
2. Mentor A, Korman NJ, Elmets CA, et al. Guidelines of care for phototherapy and photochemotherapy. *J Am Acad Dermatol* 2010;62(1):114-135.
3. Abel JD, Kattan-Byron J. Psoriasis treatment at the Dead Sea: A natural selective ultraviolet phototherapy. *J Am Acad Dermatol* 1985;12:639-643.
4. Abel JD, Kipnis V. Bioclimatology and balneology in dermatology: A Dead Sea perspective. *Clin Dermatol* 1998;16:695-698.
5. Even-Paz Z, Gumon R, Kipnis V, et al. How much Dead Sea sun for psoriasis? *J Dermatolog Treat* 1996;7:17-19.
6. Diffey BL, Larko O, Swanbeck G. UV-B doses received during different outdoor activities and UV-B treatment of psoriasis. *Br J Dermatol* 1982;106:33-41.
7. Kudish AI, Abels D, Harari M. Ultraviolet radiation properties as applied to photodermatology at the Dead Sea. *Int J Dermatol* 2003;42:359-365.
8. Walters IB, Burack LH, Coven TR, Gilleaudeau P, Krueger JG. Suberythemogenic narrow-band UVB is markedly more effective than conventional UVB in treatment of psoriasis vulgaris. *J Am Acad Dermatol* 1999;6:893-900.
9. Zanolli M. Phototherapy arsenal in the treatment of psoriasis. *Dermatol Clin* 2004;22:397-406.
10. Kushelevsky AP, Harari M, Kudish AI, Hristakieva E, Ingber A, Shani J. Safety of solar phototherapy at the Dead Sea. *J Am Acad Dermatol* 1998;3:447-452.
11. Freedberg IM, Eisen AZ, Wolff K, et al., eds. *Fitzpatrick’s Dermatology in General Medicine*, 6th ed. New York: McGraw-Hill, 2003: 1275-1282.
12. Kudish AI, Evseev E, Kushelevsky P. The analysis of ultraviolet radiation in the Dead Sea basin. *Int J Climatol* 1997;17:1697-1704.
13. Larko O, Swanbeck G. Is UVB treatment of psoriasis safe? A study of UVB treated psoriatics compared with a matched control group. *Arch Derm Venereol* 1982;62:507-512.
14. Larko O. Phototherapy of psoriasis: clinical aspects and risk evaluation. *Acta Derm Venereol* 1982;103:1-42.
15. Abel JD, Even-Paz Z, Efron D. Bioclimatology at the Dead Sea. *Clin Dermatol* 1996;14:653-658.
16. Kudish AI, Lyubansky V, Evseev EG, Ianetz A. Inter-comparison of the solar UVB, UVA, and global radiation clearness and UV indices for Beer Sheva and Neve Zohar. *Energy* 2005;30:1623-1641.

Accepted 29 March 2010.