

Tropical Journal of Pharmaceutical Research March 2015; 14 (3): 407-414

ISSN: 1596-5996 (print); 1596-9827 (electronic)

© Pharmacotherapy Group, Faculty of Pharmacy, University of Benin, Benin City, 300001 Nigeria.

All rights reserved.

Available online at <http://www.tjpr.org><http://dx.doi.org/10.4314/tjpr.v14i3.8>

Original Research Article

Enhanced Production of Palmarumycins C₁₂ and C₁₃ in Mycelial Liquid Culture of the Endophytic Fungus *Berkleasmium* sp. Dzf12 with *In situ* Macroporous Resin Adsorption

Yan Mou, Jing Li, Kaiyi Zhou, Ruiting Yu, Dan Xu, Haiyu Luo, Daiwan Lai and Ligang Zhou*

MOA Key Laboratory of Plant Pathology, Department of Plant Pathology, College of Agronomy and Biotechnology, China Agricultural University, Beijing 100193, China

*For correspondence: **Email:** lgzhou@cau.edu.cn; **Tel:** +86-10-62731199; **Fax:** +86-10-62731062

Received: 28 November 2014

Revised accepted: 23 February 2015

Abstract

Purpose: To evaluate *in situ* macroporous resin adsorption for enhancement of palmarumycins C₁₂ and C₁₃ production in mycelial liquid culture of the endophytic fungus *Berkleasmium* sp. Dzf12.

Methods: Ten macroporous adsorption resins (D-101, D1300, HPD-100, X-5, AB-8, DM130, ADS-17, DA-201, NKA-9 and S-8) were tested for adsorption of palmarumycins C₁₂ and C₁₃ in mycelial liquid culture of the endophytic fungus *Berkleasmium* sp. Dzf12.

Results: Among the resins, DA-201 showed the most significant enhancing effect on accumulation of palmarumycins C₁₂ and C₁₃ in mycelial liquid culture of endophytic fungus *Berkleasmium* sp. Dzf12. When resin DA-201 was applied to the medium at a concentration of 4.17 % on day 11 and then harvested on day 15, the maximal yield of palmarumycins C₁₂ and C₁₃ was 149.86 and 55.78 mg/L, which correspond to 70.69- and 1.82-fold higher than for control (2.12 and 30.70 mg/L, respectively). Approximately 95.81 % of palmarumycin C₁₂ and 87.20 % of palmarumycin C₁₃ were distributed in resin DA-201.

Conclusion: The results indicate that *in situ* resin adsorption is an effective strategy for enhancing the production of palmarumycins C₁₂ and C₁₃, and also for facilitating their recovery from the mycelial liquid culture of *Berkleasmium* sp. Dzf12.

Keywords: Endophytic fungus, *Berkleasmium* sp. Dzf12, Spirobisnaphthalene, Palmarumycin, Macroporous adsorption resins, Mycelial liquid culture

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

Endophytic fungi are a special group of microorganisms which spend the whole or part of their lifecycle colonizing inter- and/or intracellularly healthy tissues of the host plant, typically causing no apparent symptoms of disease [1]. They are novel and rich sources of bioactive natural products [2]. Secondary

metabolites such as alkaloids, steroids, phenolics, terpenoids and peptides produced by the endophytes have recently received growing interest from the pharmaceutical industry [3,4]. Spirobisnaphthalenes are a group of naphthoquinone derivatives, consisting of 1, 8-dihydroxynaphthalene-derived spiroketal units linked to a second oxidized naphthalene moiety, that show a great variety of biological activities

such as anti-tumor, anti-bacterial, anti-fungal, anti-leishmanial and enzyme-inhibitory properties [5].

In our previous study, an endophytic fungus isolated from the healthy rhizomes of the medicinal plant *Dioscorea zingiberensis* was identified as *Berkleasium* sp. Dzf12 from which 22 bioactive spirobisnaphthalenes have been successfully obtained [6,7]. Among them, palmarumycins C₁₂ and C₁₃ were found to be the predominant components. Furthermore, these two compounds could be secreted to the extracellular in liquid culture of *Berkleasium* sp. Dzf12. Palmarumycins C₁₂ and C₁₃, which exhibit antibacterial, antifungal and anti-tumor activities as well as inhibitory activity on phospholipase D (PLD), have been isolated from other fungi such as *Coniothyrium* sp., *Cladosporium chlorocephalum* and *Natrassia mangiferae*, [8-12]. With the aim of speeding up application of palmarumycins C₁₂ and C₁₃, considerable efforts have been directed at seeking strategies for improving palmarumycins C₁₂ and C₁₃ production in liquid culture of *Berkleasium* sp. Dzf12. The strategies include optimization of medium and culture conditions [13], utilization of two-phase culture systems [14], application of oligosaccharides and polysaccharides from its host plant *Dioscorea zingiberensis* [15,16], and addition of metal ions [17].

The purpose of this study was to investigate the effects of different types of macro porous adsorption resins including D-101, D1300, HPD-100, X-5, AB-8, DM130, ADS-17, DA-201, NKA-9 and S-8 on palmarumycins C₁₂ and C₁₃ production in liquid culture of the endophytic fungus, *Berkleasium* sp. Dzf12.

EXPERIMENTAL

Endophytic fungus and culture conditions

The endophytic fungus, *Berkleasium* sp. Dzf12 (GenBank accession number EU543255), was isolated from the healthy rhizomes of the medicinal plant, *Dioscorea zingiberensis* C. H. Wright (Dioscoreaceae) in our previous study [6]. It was sub-cultured on potato dextrose agar (PDA) slants at 25 °C in darkness. For preparation of the inoculum, four disks (about 5 mm) with fungal cultures were transferred into each 300-mL Erlenmeyer flask containing 100 mL of potato dextrose broth (PDB). The flasks were placed on a rotary shaker at 150 rpm and 25 °C in darkness for 4 days. For fermentation culture, the seed suspension cultures in 3 % (v/v)

were inoculated in each 150-mL Erlenmeyer flasks containing 30 mL of fermentation medium, which was composed of glucose 40 g/L, peptone 10 g/L, KH₂PO₄ 1.0 g/L, MgSO₄•7H₂O 0.5 g/L and FeSO₄•7H₂O 0.05 g/L. The fermentation cultures were also conducted on the rotary shaker at 150 rpm and 25 °C in darkness. Prior to autoclaving at 121 °C for 20 min, the pH of the medium was adjusted to 6.5.

Macroporous adsorption resins

Ten non-ionic polystyrene resins (D-101, D1300, HPD-100, X-5, AB-8, DM130, ADS-17, DA-201, NKA-9 and S-8) were purchased from Tianjin Haiguang Chemical Company of China (Tianjin, China). They were initially examined as the adsorbents for *in situ* adsorption of palmarumycins C₁₂ and C₁₃ in mycelial liquid culture of *Berkleasium* sp. Dzf12. Their chemical and physical properties, including polarity, particle size, surface area, pore diameter and moisture content are shown in Table 1. Prior to use, the resins were in turn soaked in methanol for 24 h, washed thoroughly with distilled water for 3 - 5 times, immersed in 1 mol/L NaOH solution for 12 h, rinsed by distilled water for 3 - 5 times, immersed in 1 mol/L HCl solution for 12 h, and rinsed with distilled water 3 - 5 times. Each resin was then dried in an oven at 35 - 40 °C to a constant dry weight (dw). The moisture (%) = $[(W_1 - W_2) / W_1] \times 100$, where W₁ is the weight prior to drying, W₂ is the weight after drying.

In situ adsorption culture

The pre-treated dried resin was wrapped with 30 μm-pore nylon cloth (Yanpai Chemical Company, Shanghai, China) into small bags of 0.50 - 2.00 g dry weight (dw). The resin bags were suspended in distilled water and autoclaved at 121 °C for 20 min before being added to the culture broth at the designated time. For screening the optimal resin from the 10 candidates, 1.5 g of each resin in a nylon bag was added to the mycelial culture flask (30 mL medium in 150-mL flask) on day 9 of culture, so the corresponding concentration of resin in the medium was 5.0 % (g/mL). The cultures were harvested on day 15. For determining the suitable concentration of the selected resin DA-201, different quantities (0, 0.50, 0.75, 1.00, 1.25, 1.50, 1.75 and 2.00 g) of resin DA-201 were added in 30 mL medium (their corresponding concentrations were 0 - 6.67 %, g/mL) on day 9 of culture. The cultures were harvested on day 15. For obtaining the optimal addition and incubation time of resin DA-201, 4.17 % (1.25 g in 30 mL medium) of resin was

Table 1: Chemical and physical properties of the macroporous resins employed

Resin	Polarity	Particle size (mm)	Surface area (m ² /g)	Average pore diameter (nm)	Moisture content (%)
D-101	Non-polar	0.30–1.25	480–530	9–11	67.89 ± 0.27
D1300	Non-polar	0.30–1.25	≥600	9–10	68.27 ± 0.57
HPD-100	Non-polar	0.30–1.25	≥650	8–9	68.39 ± 0.48
X-5	Non-polar	0.30–1.25	500–600	29–30	50.13 ± 0.17
AB-8	Weak-polar	0.30–1.25	480–520	13–4	68.33 ± 0.61
DM130	Weak-polar	0.30–1.25	500–550	9–10	66.00 ± 0.32
ADS-17	Middle-polar	0.30–1.25	90–150	25–30	55.16 ± 1.09
DA-201	Polar	0.30–1.25	≥200	10–13	70.64 ± 0.52
NKA-9	Polar	0.30–1.25	250–290	15.5–16.5	70.22 ± 0.23
S-8	Polar	0.30–1.25	100–200	28–30	67.12 ± 0.35

Note: The information was provided by the manufacturers, except for the moisture content

added into the medium on days 0, 3, 6, 9, 10, 11 and 12 of culture, respectively, and the cultures were harvested on day 15. All treatments were performed in triplicate.

Determination of biomass and palmarumycins C₁₂ and C₁₃ yield

The mycelia were harvested by vacuum filtration via a Buchner funnel with a pre-weighed filter paper, and the filtrate volume was measured. The harvested mycelia were washed with deionized water three times to remove medium components, and then dried in an oven at 50 – 55 °C to a constant dry weight (dw). Palmarumycins extraction and determination were carried out as previously described [17]. For analysis of palmarumycins C₁₂ and C₁₃ yield in mycelia, 50.0 mg of dry mycelial powder, was deposited in a vial with 5 mL of methanol-chloroform (9:1, v/v), and then subjected to ultrasonic treatment (three times, 60 min each). After removal of the solid by filtration, the filtrate was evaporated to dryness and re-dissolved in 1 mL of methanol. For analysis of palmarumycins C₁₂ and C₁₃ in broth, 3 mL of the culture broth without mycelia was evaporated to dryness and extracted with 5 mL of methanol-chloroform (9:1, v/v) in an ultrasonic bath, and the liquid extract was then evaporated to dryness and re-dissolved in 1 mL of methanol. For analysis of palmarumycins C₁₂ and C₁₃ in the resins, the nylon cloth sachet with whole resin was extracted with 30 mL of methanol-chloroform (9:1, v/v). The extraction was repeated until the final extract was colorless. After removal of the resin by filtration, the filtrate was evaporated to dryness and re-dissolved in 30 mL of methanol.

Palmarumycins C₁₂ and C₁₃ standards were isolated and identified from the fermentation cultures of the endophytic fungus *Berkleasium* sp. Dzf12 in our previous studies [7,15]. Palmarumycins C₁₂ and C₁₃ yields were analyzed by high performance liquid chromatography

(HPLC) (Shimadzu, Japan), which consisted of two LC-20AT solvent delivery units, an SIL-20A autosampler, an SPD-M20A photodiode array detector, and CBM-20Alite system controller (Shimadzu, Kyoto, Japan), and a reversed-phase Ultimate TM XB C18 column (4.6 × 250 mm, 5 μm, Welch Materials, Inc., Ellicott, MD, USA). The HPLC column was isocratically eluted with MeOH–H₂O (50:50, v/v) in 20 min at a flow rate of 1.0 mL/min. The temperature was maintained at 40 °C, and UV detection at 226 nm. The sample injection volume was 10 μL. The LC-solution multi-PDA workstation was employed to acquire and process chromatographic data. Palmarumycins C₁₂ and C₁₃ were detected and quantified with the standards. The regression equation of palmarumycin C₁₂ was $Y = 121295.5 X + 175236.8$ ($R = 0.9997$), and that of palmarumycin C₁₃ was $Y = 121362.5 X + 256167.5$ ($R = 0.9997$), where Y is the peak area, X is quality (μg) of the sample injected each time, and R the correlation coefficient.

Statistical analysis

All experiments were carried out in triplicate, and the results are presented as mean ± standard deviations (SD). The data were subjected to analysis of variance (one-way ANOVA) to detect significant differences using Proc Anova SAS version 8.2. Significance difference was set at $p \leq 0.05$.

RESULTS

Optimized macroporous resins

The effect of the resins on mycelia biomass, palmarumycins C₁₂ and C₁₃ yields are shown in Figure 1. Compared to the control without resin addition (CK), the mycelial growth was inhibited or promoted by the resins more or less. Resins HPD100, X-5, AB-8, ADS-17 and DA-201 showed slight enhancement, and resins D1300

and NKA-9 exhibited slight inhibition on mycelia growth (Figure 1A).

All the resins exhibited obvious enhancing effects on palmarumycin C₁₂ accumulation (Figure 1B). Among them, resin DA-201 was found to be the most effective in enhancing palmarumycin C₁₂ production. The highest palmarumycin C₁₂ yields (7.99 mg/L and 117.43 mg/L) in mycelia and resin were obtained respectively. The total palmarumycin C₁₂ yield reached 125.43 mg/L, which was 62.71-fold as compared with control (2.00 mg/L). Other effective resins for palmarumycin C₁₂ production included D-101, HPD100, X-5, AB-8, DM130, ADS-17 and S-8.

All the resins exhibited less enhancing effects on palmarumycin C₁₃ production than on palmarumycin C₁₂ production. The palmarumycin C₁₃ yield of the cultures treated with resins D1300, DM130, ADS-17, and NKA-9 was even lower than that of the control without resin addition. Among three resins (HPD100, AB-8 and DA-201) with enhancing effects on palmarumycin C₁₃ production, resin DA-201 was the most effective. When resin DA-201 was applied to the liquid culture of endophyte Dzf12, total palmarumycin C₁₃ yield was increased to 43.44 mg/L, which was 1.41-fold relative to control (30.73 mg/L) (Figure 1C). Overall, resin DA-201 was found to be the most favorable adsorbent for palmarumycins C₁₂ and C₁₃ production in *Berkleasium* sp. Dzf12 liquid culture, and was used in the subsequent experiments.

Effect of resin DA-201 concentrations on production of palmarumycins C₁₂ and C₁₃

The effects of the addition concentrations (1.67, 2.50, 3.33, 4.17, 5.00, 5.83 and 6.67 %) of resin DA-201 on the palmarumycins C₁₂ and C₁₃ production in liquid culture of *Berkleasium* sp. Dzf12 are presented in Table 2. When the concentration of resin DA-201 addition was increased from 0 to 4.17 %, the amount of adsorbed palmarumycins C₁₂ and C₁₃ gradually increased too. The total yields of palmarumycins C₁₂ and C₁₃ reached the highest levels (144.64 and 48.30 mg/L) which were 66.34-fold and 1.59-fold in comparison with those (2.18 and 30.37 mg/L) of the control, respectively, when resin DA-201 was added in the medium at 4.17 % on day 9. There was no significant effect on mycelial growth with resin DA-201 added to the medium at concentrations ranging from 0 to 6.67 % (g/mL). The mycelia biomass was relatively low

when DA-201 resin was added at concentration of 1.67 %.

Effect of resin DA-201 addition and incubation time on production of palmarumycins C₁₂ and C₁₃

The effects of resin DA-201 addition and incubation period on mycelial growth and production of palmarumycins C₁₂ and C₁₃ are shown in Table 3. When the 11-day-old cultures were treated with resin DA-201 at 4.17 %, the total palmarumycins C₁₂ and C₁₃ yields reached the highest values (149.86 and 55.78 mg/L), which were 70.69-fold and 1.82-fold of control yield (2.12 and 30.70 mg/L), respectively. From the above results, day 11 was confirmed as the optimal addition time. It allowed a period of 4 days for adsorption. The percentages of palmarumycins C₁₂ and C₁₃ adsorbed in resin DA-201 were 95.81 and 87.20 %, respectively. Considering production of the two spirobisanthalenes, the highest total yield of palmarumycins C₁₂ and C₁₃ reached 205.64 mg/L, which was 6.25-fold higher than for control (32.82 mg/L), obtained by adding DA-201 resin on day 11 of culture and allowing a period of 4 days for adsorption. As shown in Table 3, if resin DA-201 was added to the cultures early (i.e., the addition time occurred on days 0 and 3), it possibly adsorbed more nutrients and thus caused a decrease of palmarumycins C₁₂ and C₁₃ biosynthesis.

DISCUSSION

The *in situ* product removal by using macroporous resins in fermentation system is an integrated bioprocess of production and separation. It has many benefits for the overall fermentation process by facilitating the product recovery, eliminating feedback inhibition, overcoming product degradation, avoiding product autotoxicity, reducing cost, and improving the product yield efficiently [18,19]. The strategy of employing *in situ* macroporous resin adsorption has been successfully applied in increasing metabolite yields in some fungal culture systems, such as *Fusarium redolens* Dzf2 culture for beauvericin production [20], and *Hyalodendriella* sp. Ponipodef12 culture for botrallin and TMC-264 production [21].

From the results shown in Figure 1B, palmarumycin C₁₂ was undetectable in the broth while it was detectable in the resin and mycelial extracts, which means that palmarumycin C₁₂ can be secreted from the intracellular to the




Figure 1: Effects of the resins on mycelia biomass (A), palmarumycin C₁₂ production (B), and palmarumycin C₁₃ production (C) in liquid culture of *Berkleasium* sp. Dzf12. CK means without addition of the test resin. Different letters (*i.e.*, a, b, c ...) indicate significant differences among the data of each situation at $P \leq 0.05$. The error bars are standard deviations ($n = 3$)

Table 2: Effects of resin DA-201 addition concentrations on mycelial growth, palmarumycins C₁₂ and C₁₃ production in liquid culture of *Berkleasium* sp. Dzf12

Conc. of resin DA-201 (% g/mL)	Mycelia biomass (g dw/L)	C ₁₂ yield in mycelia (mg/L)	C ₁₂ yield in broth (mg/L)	C ₁₂ yield in resin (mg/L)	C ₁₃ yield in mycelia (mg/L)	C ₁₃ yield in broth (mg/L)	C ₁₃ yield in resin (mg/L)	Total C ₁₂ yield (mg/L)	Total C ₁₃ yield (mg/L)	Total C ₁₂ +C ₁₃ yield (mg/L)
0.00	6.42±0.13 ^{ab}	2.18±0.46 ^b	0.00±0.00 ^a	0.00±0.00 ^e	10.50±0.45 ^a	19.87±0.86 ^a	0.00±0.00 ^e	2.18	30.37	32.55
1.67	6.09±0.10 ^b	2.70±1.07 ^b	0.00±0.00 ^a	76.43±5.64 ^d	9.87±1.60 ^{ab}	0.00±0.00 ^b	28.09±1.00 ^d	79.13	37.96	117.09
2.50	6.59±0.18 ^a	3.31±1.02 ^b	0.00±0.00 ^a	92.54±3.18 ^c	10.82±1.28 ^a	0.00±0.00 ^b	32.77±1.24 ^c	95.85	43.59	139.44
3.33	6.86±0.36 ^a	5.34±1.04 ^{ab}	0.00±0.00 ^a	128.70±1.12 ^b	7.86±0.91 ^{bc}	0.00±0.00 ^b	39.03±0.49 ^{ab}	134.04	46.89	180.93
4.17	6.61±0.15 ^a	5.35±2.46 ^{ab}	0.00±0.00 ^a	139.29±4.86 ^a	5.84±0.54 ^{cd}	0.00±0.00 ^b	42.46±2.31 ^a	144.64	48.30	192.94
5.00	6.66±0.13 ^a	7.57±1.04 ^a	0.00±0.00 ^a	136.03±1.67 ^{ab}	4.36±1.04 ^d	0.00±0.00 ^b	39.33±0.87 ^{ab}	143.60	43.69	187.29
5.83	6.44±0.25 ^{ab}	5.05±2.01 ^{ab}	0.00±0.00 ^a	136.86±3.54 ^{ab}	3.33±0.76 ^d	0.00±0.00 ^b	37.60±2.52 ^b	141.91	40.93	182.84
6.67	6.51±0.26 ^{ab}	5.31±2.03 ^{ab}	0.00±0.00 ^a	136.12±4.17 ^{ab}	3.30±1.20 ^d	0.00±0.00 ^b	37.20±0.50 ^b	141.43	40.50	181.93

Notes: The values are expressed as means ± standard deviations (n = 3). Different letters (i.e., a, b, c, ...) indicate significant differences among the treatments in each column at P ≤ 0.05

Table 3: Effects of resin DA-201 addition and incubation time on palmarumycins C₁₂ and C₁₃ production in liquid culture of *Berkleasium* sp. Dzf12

Addition time/Incubation time (d/d)	Mycelia biomass (g dw/L)	C ₁₂ yield in mycelia (mg/L)	C ₁₂ yield in broth (mg/L)	C ₁₂ yield in resin (mg/L)	C ₁₃ yield in mycelia (mg/L)	C ₁₃ yield in broth (mg/L)	C ₁₃ yield in resin (mg/L)	Total C ₁₂ yield (mg/L)	Total C ₁₃ yield (mg/L)	Total C ₁₂ +C ₁₃ yield (mg/L)
CK	6.53±1.08 ^{ab}	2.12±0.31 ^e	0.00±0.00 ^a	0.00±0.00 ^d	11.01±0.53 ^a	19.69±0.59 ^a	0.00±0.00 ^d	2.12	30.70	32.82
0/15	5.14±0.15 ^c	3.29±0.34 ^{de}	0.00±0.00 ^a	77.48±4.76 ^c	3.14±0.05 ^d	0.00±0.00 ^b	10.35±1.51 ^c	80.77	13.49	94.26
3/12	5.47±0.30 ^{bc}	4.85±0.18 ^{cde}	0.00±0.00 ^a	94.46±3.36 ^b	3.10±1.02 ^d	0.00±0.00 ^b	12.94±2.17 ^c	99.31	16.04	115.35
6/9	5.86±0.33 ^{abc}	5.82±1.54 ^{cd}	0.00±0.00 ^a	97.29±8.30 ^b	4.23±0.11 ^{cd}	0.00±0.00 ^b	38.35±1.44 ^b	103.11	42.58	145.69
9/6	6.33±0.38 ^{ab}	5.85±1.66 ^{cd}	0.00±0.00 ^a	133.81±4.10 ^a	6.06±1.31 ^{bc}	0.00±0.00 ^b	41.30±2.02 ^b	139.66	47.36	187.02
10/5	6.62±0.22 ^{ab}	8.03±0.63 ^{bc}	0.00±0.00 ^a	139.04±7.75 ^a	6.46±0.07 ^b	0.00±0.00 ^b	47.79±1.74 ^a	147.08	54.25	193.91
11/4	6.89±0.47 ^a	10.64±2.70 ^{ab}	0.00±0.00 ^a	139.22±3.87 ^a	6.34±1.42 ^b	0.00±0.00 ^b	49.44±2.83 ^a	149.86	55.78	205.64
12/3	6.59±0.12 ^{ab}	13.61±1.09 ^a	0.00±0.00 ^a	131.97±4.32 ^a	6.62±0.61 ^b	0.00±0.00 ^b	47.95±1.14 ^a	145.58	54.57	200.15

Notes: CK, without addition of the test resin. The concentration of resin DA-201 in medium was 4.17% (g/mL). The values are expressed as means ± standard deviations (n = 3). Different letters (i.e., a, b, c, ...) indicate significant differences among the treatments including addition and incubation time in each column at P ≤ 0.05

extracellular and adsorbed by macroporous resins efficiently. The addition of the resin greatly favored palmarumycin C₁₂ production (Tables 2 and 3). 4.17 % of resin DA-201 in the medium is sufficient for achieving the maximal palmarumycins C₁₂ and C₁₃ production from the liquid culture. The yield of palmarumycin C₁₂ was increased about 70 fold by addition of resin DA-201. It is possible that palmarumycin C₁₂ was an intermediate metabolite in the biosynthesis pathway of spirobisanaphthalenes [22]. Addition of resin favored the secretion of palmarumycin C₁₂ and suppressed its metabolism into other compounds.

In this study, we just considered three variables by employing macroporous resin adsorption during fermentation such as the type, concentration and addition time of resin. Other variables such as temperature, pH value, fermentation medium, combination effects of two or more resins, adsorption and desorption kinetics, and bioreactor system for large-scale production by utilizing resin adsorption of palmarumycins C₁₂ and C₁₃ need to be studied in detail [19].

CONCLUSION

This work is the first, to the best of our knowledge, to report the enhancement of palmarumycins C₁₂ and C₁₃ production using solid-phase adsorption in fungal liquid culture. Among the resins, DA-201 showed the greatest enhancement effect on the accumulation of palmarumycins C₁₂ and C₁₃ in mycelial liquid culture of endophytic fungus *Berkleasmium* sp. Dzf12. The results indicate that in situ resin adsorption is an effective strategy for enhancing the production of palmarumycins C₁₂ and C₁₃, and also for facilitating their recovery from mycelial liquid culture of *Berkleasmium* sp. Dzf12.

ACKNOWLEDGEMENT

This work was co-funded by Hi-Tech R&D Program of China (no. 2011AA10A202) and National Natural Science Foundation of China (no. 31071710).

REFERENCES

- Rodriguez RJ, White JF, Arnold AE, Redman RS. Fungal endophytes: diversity and functional roles. *New Phytol* 2009; 182: 314-330.
- Kharwar RN, Mishra A, Gond SK, Stierle A, Stierle D. Anticancer compounds derived from fungal

- endophytes: Their importance and future challenges. *Nat Prod Rep* 2011; 28: 1208-1228.
- Zhao J, Shan T, Mou Y, Zhou L. Plant-derived bioactive compounds produced by endophytic fungi. *Mini-Rev Med Chem* 2011; 11: 159-168.
- Kaul S, Gupta S, Ahmed M, Dhar MK. Endophytic fungi from medicinal plants: a treasure hunt for bioactive metabolites. *Phytochem Rev* 2012; 11: 487-505.
- Zhou L, Zhao J, Shan T, Cai X, Peng Y. Spirobisanaphthalenes from fungi and their biological activities. *Mini-Rev Med Chem* 2010; 10: 977-989.
- Cai X, Shan T, Li P, Huang Y, Xu L, Zhou L, Wang M, Jiang W. Spirobisanaphthalenes from the endophytic fungus Dzf12 of *Dioscorea zingiberensis* and their antimicrobial activities. *Nat Prod Commun* 2009; 4: 1469-1472.
- Shan T, Tian J, Wang X, Mou Y, Mao Z, Lai D, Dai J, Peng Y, Zhou L, Wang M. Bioactive spirobisanaphthalenes from the endophytic fungus *Berkleasmium* sp. *J Nat Prod* 2014; 77: 2151-2160.
- Schlingmann G, West RR, Milne L, Pearce CJ, Carter GT. Diepoxins, novel fungal metabolites with antibiotic activity. *Tetrahedron Lett* 1993; 34: 7225-7228.
- Krohn K, Michel A, Floerke U, Aust HJ, Draeger S, Schulz B. Palmarumycins C1-C16 from *Coniothyrium* sp.: Isolation, structure elucidation, and biological activity. *Liebigs Ann Chem* 1994; 11: 1099-1108.
- Chu M, Truumees I, Patel MG, Gullo VP, Blood C, King I, Pai JK, Puar MS. A novel class of anti-tumor metabolites from the fungus *Nattractia mangiferae*. *Tetrahedron Lett* 1994; 35: 1343-1346.
- Thiergardt R, Hug P, Rihs G, Peter HH. Cladosporine bisepoxide-a novel fungal metabolite structure determination. *Tetrahedron Lett* 1994; 35: 1043-1046.
- Petersen F, Moerker T, Vanzanella F, Peter HH. Production of cladosporine bisepoxide, a new fungal metabolite. *J Antibiot* 1994; 47: 1098-1103.
- Zhao J, Wang X, Sun W, Mou Y, Peng Y, Zhou L. Medium optimization for palmarumycin C₁₃ production in liquid culture of endophytic fungus *Berkleasmium* sp. Dzf12 using response surface methodology. *Electron J Biotechnol* 2013; 16: DOI: 10.2225/vol16-issue6-fulltext-10.
- Zhao J, Li Y, Shan T, Mou Y, Zhou L. Enhancement of diepoxin ζ production with in situ resin adsorption in mycelial liquid culture of the endophytic fungus *Berkleasmium* sp. Dzf12 from *Dioscorea zingiberensis*. *World J Microbiol Biotechnol* 2011; 27: 2753-2758.
- Li Y, Shan T, Mou Y, Li P, Zhao J, Zhao W, Peng Y, Zhou L, Ding C. Enhancement of palmarumycin C₁₂ and C₁₃ production in liquid culture of the endophytic fungus *Berkleasmium* sp. Dzf12 by oligosaccharides from its host plant *Dioscorea zingiberensis*. *Molecules* 2012; 17: 3761-3773.
- Li Y, Li P, Mou Y, Zhao J, Shan T, Ding C, Zhou L. Enhancement of diepoxin ζ production in liquid

- culture of endophytic fungus *Berkleasmium* sp. Dzf12 by polysaccharides from its host plant *Dioscorea zingiberensis*. *World J Microbiol Biotechnol* 2012; 28: 1407-1413.
17. Mou Y, Luo H, Mao Z, Shan T, Sun W, Zhou K, Zhou L. Enhancement of palmarumycins C₁₂ and C₁₃ production in liquid culture of endophytic fungus *Berkleasmium* sp. Dzf12 after treatments with metal ions. *Int J Mol Sci* 2013; 14: 979-998.
18. Schugerl K, Hubbuch J. Integrated bioprocesses. *Curr Opin Microbiol* 2005; 8: 249-300.