

## Minireview

# Enhancing banana weevil (*Cosmopolites sordidus*) resistance by plant genetic modification: A perspective

Andrew Kiggundu<sup>1,2\*</sup>, Michael Pillay<sup>4</sup>, Altus Viljoen<sup>3</sup>, Clifford Gold<sup>4</sup>, Wilberforce Tushemereirwe<sup>1</sup>, Karl Kunert<sup>2</sup>

<sup>1</sup>National Agricultural Research Organization (NARO), Kawanda Agricultural Research Institute (KARI) P. O. Box 7065 Kampala, Uganda.

<sup>2</sup>Department of Botany, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, 74 Lunnon Road, Hillcrest, Pretoria South Africa.

<sup>3</sup>Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, 74 Lunnon Road, Hillcrest, Pretoria South Africa.

<sup>4</sup>International Institute of Tropical Agriculture (IITA), East and Southern Africa Regional Centre (ESARC), P. O. Box 7878, Kampala, Uganda.

Accepted 21 November 2003

**Banana weevil is a serious pest of bananas and plantains in Africa. The development of resistant cultivars is seen as the long term and more sustainable control strategy. The difficulty in conventional breeding of bananas and plantains has prompted efforts towards the use of genetic transformation for banana and plantain improvement. In this review, the current status of banana weevil resistance, sources of resistance and resistance mechanisms is assessed. Further, current efforts and future prospects for identifying resistance genes outside the genus *Musa* with potential to control banana weevil in a transgenic approach are outlined and discussed.**

**Key words:** Banana weevil, host plant resistance, pest resistance genes, transgenic plants.

## BANANA WEEVIL AS AN AFRICAN PEST

The banana weevil is a pest of considerable importance in Africa and significantly affects banana and plantain production (Ostmark, 1974; Gold, 1998; Swennen and Vulysteke, 2001; Gold and Messiaen, 2000; Fogain et al., 2002). The weevil has been associated with rapid decline of banana plantations in east Africa (Gold et al., 1999b) and a phenomenon called "yield decline syndrome" in West Africa (Ortiz and Vulysteke, 1994). The adult weevils are free living, have a nocturnal habit, and rarely fly. Their eggs are deposited at the base of the pseudostem or on exposed corms. Upon hatching, the

larvae tunnel through the corm to feed and develop. The tunnelling damages the corm and weakens the plant reduces, reduces water and mineral uptake, reducing bunch weight (yield) and causing plant toppling during windstorms. In severe weevil infestations, crop losses of up to 100% have been reported (Sengooba, 1986). The establishment of new plantings in infested fields may fail (Price, 1994) and yield loss appears to increase gradually reaching 44% in the fourth ratoon cycle (Rukazambuga et al., 1998).

## CONVENTIONAL BANANA WEEVIL CONTROL

Banana weevil control is currently based on the application of cultural practices, such as the use of clean

\*Corresponding author. Email: [andrew.kiggundu@fabi.up.ac.za](mailto:andrew.kiggundu@fabi.up.ac.za).  
Fax: +27-12-4203960.

**Table I.** Suggested sources of banana weevil resistance in *Musa* spp. taken from field trial cultivar screening experiments.

| Cultivar                | Genome group | Reference   |
|-------------------------|--------------|---|
| Yangambi km-5           | AAA          | Fogain and Price, 1994;<br>Lemaire, 1996;<br>Kiggundu et al., 2003      |
| Sannachenkadali         | AA           | Padmanaban et al., 2001   |
| Sakkali                 | ABB          |   |
| Senkadali               | AAA          |   |
| Elacazha                | BB           |   |
| Njalipoovan             | AB           |   |
| Pisang Awak             | ABB          | Kiggundu et al., 2003   |
| FHIA03                  | AABB         |   |
| TMBx612-74              | IITA hybrid* |   |
| TMB2x6142-1             | IITA hybrid  |   |
| TMB2x8075-7 TMB2x7197-2 | IITA hybrid  |   |
| Long Tavoy              | ABB          | Ortiz et al., 1995  |
| Njeru                   | AA           | Musabyimana et al., 2000  |
| Muraru                  | AA           |   |
| Calcutta-4              | AA           | Fogain and Price, 1994;<br>Ortiz et al., 1995;<br>Kiggundu et al., 2003 |
| Bluggoe                 | ABB          | Fogain and Price, 1994;<br>Kiggundu et al., 2003                        |
| <i>M. balbisiana</i>    | BB           | Fogain and Price, 1994  |

\* IITA – International Institute of Tropical Agriculture

planting material, systematic trapping of adult weevils, in an effort to control population build-up, and field sanitation whereby residues that may form breeding grounds for the weevil are removed (Gold, 2000; Gold and Messiaen, 2000). Although cultural control methods contribute to weevil management, both high labour input and material requirements are often limiting factors for adoption of these practices (Gold, 1998; Gold et al., 2001). The application of effective pesticides is economically not feasible for subsistence producers. In addition, the banana weevil has developed resistance to a range of commonly used pesticides (Collins et al., 1991; Gold et al., 1999a). Consequently, development of resistant plants has been suggested as a potential long-term intervention for weevil control, especially on small-scale farms, as the inclusion of such plants might be part of an integrated pest management (IPM) framework (Seshu-Reddy and Lubega, 1993; Gold et al., 2001).

#### BANANA WEEVIL RESISTANCE AND RESISTANCE MECHANISMS

Considerable work has been done in screening diverse

*Musa* germplasm for weevil resistance in Africa and Asia (Fogain and Price 1994; Pavis and Lemaire, 1997; Kiggundu et al., 1999; Kiggundu et al., 2003). Although plantains and east African highland bananas (EAHB) were found to be the most susceptible banana types, there were a few exceptions. For example, in India, Padmanaban et al. (2001) found that two plantain cultivars (Karumpoovan and Poozhachendu) were resistant to the banana weevil, while Fogain and Price (1994) found that the plantain cultivar Kedong kekang was resistant in Cameroon. Kiggundu et al. (2003) found that some highland banana cultivars including Tereza, Nalukira and Nsowe had intermediate resistance to weevils.

The large variability in weevil response observed in *Musa* germplasm and hybrid testing suggested that useful sources of weevil resistance are available in the *Musa* germplasm (Kiggundu et al., 2003). Candidates for use in conventional breeding for weevil resistance have been selected on the basis of very low levels of weevil damage in the field (Table 1). The AA genome progenitor *Musa accuminata* is more susceptible to weevils than the BB progenitor *Musa balbisiana* (Mesquita et al., 1984), and it is expected that AA

sources of resistance might ultimately produce hybrids with better consumer acceptability. However, at this stage crosses with the more weevil-resistant BB type sources should not be completely excluded for obtaining a weevil-resistant hybrid (Mesquita et al., 1984; E DeLanghe, *personal communication*).

Classical resistance mechanisms (Painter, 1951) have been investigated in *Musa* germplasm, and antibiosis (factors affecting larval performance), rather than antixenosis (attraction), appears to be the most important resistance mechanism in banana (Mesquita et al., 1984; Ortiz et al., 1995; Abera et al., 1999; Kiggundu, 2000; Gold et al., 2001). Although some differences in attracting adult weevils to different cultivars have been identified no direct correlations between weevil resistance and plant damage were found (Buddenburg et al., 1993; Pavis and Minost, 1993; Abera et al., 1999; Kiggundu, 2000; Musabyimana et al., 2000). It was reported that differences in attraction have been due to environmental factors such as soil moisture around a cultivar with high sucker number (Ittyeipe, 1986).

Several banana plant phenological factors seem to contribute to weevil resistance. Corm hardness was the first biophysical factor associated with resistance. Whereas Pavis and Minost (1993) found a negative correlation ( $r = -0.47$ ) between corm hardness and weevil damage, Ortiz et al. (1995) found no relationship between the two parameters in segregating plantain progenies. The latter suggested that other weevil resistance factors such as chemical toxins or anti-feedants might be playing role in weevil resistance. Kiggundu (2000) found that corm dry matter content, resin/sap production and suckering ability were negatively correlated with weevil damage. Whereas corm diameter and resin/sap production were important factors in EAHB accessions, corm dry matter content, corm hardness, resin/sap production and suckering ability (number of suckers) were significant parameters in the resistance response of recently introduced clones in east Africa. In large corms, the weevil larvae can complete their life cycle without borrowing too deep into the corm. This indicates that some form of tolerance exists in cultivars with large corms (Balachowsky, 1963).

High-performance liquid chromatography (HPLC) chromatograms from corm extracts of weevil-resistant AB (Kisubi) and ABB (Pisang awak) cultivars showed peaks that were absent when compared not only to susceptible clones, but also to some resistant clones of the AA (Calcutta 4) and AAA (Yangambi km5) genome group cultivars. This result possibly indicates a type of antibiotic mechanism that may be based on toxic compounds. These compounds are seemingly present in weevil-resistant cultivars with the BB genome suggesting that a different form of resistance may be present in weevil-resistant AA cultivars (Kiggundu, 2000). Furthermore, separation of a methanol extract from Pisang Awak (ABB) allowed the isolation of two highly polar sub-

fractions with high anti-larval activity (Kiggundu, *unpublished data*). In these experiments, unique HPLC peaks were also found in susceptible cultivars, suggesting that the presence of compounds that may be related to weevil susceptibility.

As the identification of resistance mechanisms and weevil resistance genes within *Musa* continues, it is useful to develop biotechnology tools and protocols for the effective transfer of genes into *Musa* cultivars.

## WILL GENETIC MODIFICATION OF BANANA PLAY A ROLE IN WEEVIL RESISTANCE?

The continual need to increase food production necessitates the development and application of novel biotechnologies to enable the provision of improved crop varieties in a timely and cost effective way. Table 2 provides examples of genetically modified (GM) plants that have been produced to date worldwide with increased resistance to different pests using genes from various sources. Although remarkable achievements have already been made in banana transformation, the identification and introduction of useful genes into banana to reduce losses caused by the banana weevil is still a major challenge. A variety of genes are available for genetic engineering for pest resistance (Carozi and Koziel, 1997; Sharma et al., 2000). Among these are proteinase inhibitors. *Bacillus thuringiensis* (Bt) toxins, plant lectins, vegetative insecticidal proteins (VIPs), and alpha-amylase inhibitors (AI).

### Potential of proteinase inhibitors

The two major proteinase classes in the digestive systems of phytophagous insects are the serine and cysteine proteinases. Serine proteinase activity is present in Lepidoptera, Dictyoptera and Hymenoptera, while the cysteine proteinase activity is characteristic for Odoptera and Hemiptera. Coleopteran insects, including the banana weevil, mainly use cysteine proteinases (Gatehouse et al., 1985; Murdock et al., 1987). More recent studies indicate a combination of both serine and cysteine proteinases in the Coleoptera (Gerald et al., 1997). The combination of both cysteine and serine proteinases is considered an advanced digestive strategy. Proteinase inhibitors are expressed naturally as a plant defence against insect attack (Green and Ryan, 1972; Ryan, 1990; Pernas et al., 2000; Ashouri et al., 2001). They operate by disrupting protein digestion in the insect mid-gut via inhibition of proteinases.

The development of genetically modified (GM) banana with weevil resistance is currently the focus of the Uganda National Agricultural Research Organisation (NARO), the Forestry and Agricultural Biotechnology Institute of the University of Pretoria (FABI) and the

**Table 2.** Examples of transgenic plants with increased resistance to insects conferred by genes from various sources. Bt - Cry – *Bacillus thuringiensis* crystal proteins; BAAI - Bean alpha amylase inhibitor; GNA - snow drop lectin; OC-I Oryzacystatin-I cysteine protease inhibitor from rice (*Oryza sativa*); CC-I corn cystatin; CPTi - Cowpea trypsin inhibitor.

| Gene         | Insect-pest  | Transgenic plant | Observed effect                      | References                |
|--------------|--|------------------|--------------------------------------|---------------------------|
| Bt - Cry1Ab  | Rice leaffolder <i>Cnaphalocrocis medinalis</i>                        | Rice             | High stable resistance – no symptoms | Ye et al., 2003           |
| Bt - Cry03Aa | Colorado potato beetle   | Potato           | Increased resistance                 | Willson et al., 1992      |
| Bt - Cry03Aa | Cotton boll weevil   | Cotton           | Increased resistance                 | Willson et al., 1992      |
| CpTi         | Rice stem borers <i>Chilo suppressalis</i> and <i>Sesamia inferens</i> | Rice             | Significant resistance               | Xu et al., 1996           |
| OC-I         | Colorado potato beetle ( <i>Leptinotarsa decemleata</i> )              | Potato           | Increase resistance                  | Lecardonnell et al., 1999 |
| CC-I         | Maize grain weevil ( <i>Sitophilus zeae</i> )                          | Maize            | Increased resistance                 | Irie et al., 1996         |
| BAAI         | Bean bruchids  | Adzuki beans     | Increased resistance                 | Ishimoto et al., 1996     |
| BAAI         | <i>Collasobruchus</i> spp.   | Pea              | Increased resistance                 | Schroeder et al., 1995    |
| GNA          | Peach potato aphid   | Potato           | Increased resistance                 | Gatehouse et al., 1997    |
| GNA          | Sugarcane grub ( <i>Antitrogus consanguineus</i> )                     | Sugarcane        | Increased resistance                 | Nutt et al., 1999         |

International Institute of Tropical Agriculture (IITA) (NARO/FABI/IITA). The objective of this research is to express exogenous cysteine proteinase inhibitors in bananas with the potential to control banana weevil. These inhibitors have already been used for insect control in GM plants (Leple et al., 1995). Cysteine proteinases are not part of the human digestive system, and therefore humans can consume cysteine proteinase inhibitors safely. The NARO/FABI/IITA group by interacting with the University of Laval (D. Michaud) is currently investigating the potential of plant derived cysteine proteinase inhibitors (phytocystatins) from rice and papaya in GM banana. Presently, cysteine proteinase activity has been identified in the mid-gut of the banana weevil and *in-vitro* studies have shown that these cysteine proteinases are strongly inhibited by both a purified recombinant rice (oryzacystatin-I (OC-I)) and papaya cystatin (Kiggundu et al., unpublished data). By using a newly developed bioassay system based on vacuum infiltration of banana stems, it was shown that recombinant papaya cystatin significantly reduced the early growth and development of weevil larvae (Kiggundu et al., unpublished data). In addition, the design of optimised inhibitors using site-directed mutagenesis is currently being carried out to improve the inhibition and stability of the inhibitors for weevil control. GM bananas expressing a modified OC-I gene targeting nematodes have been produced by the John Innes Centre (JIC, Norwich, UK) together with the University of Leeds, UK (P. Vain, John Innes Centre, UK. *personal communication*).

Although the phytocystatin strategy is interesting there are certain stumbling blocks to overcome. Some insects when exposed to a particular proteinase inhibitor

selectively secrete other proteinases that are insensitive to that particular inhibitor. In other cases variation in the gut environment, especially pH, leads to inactivity of the inhibitor thus making the strategy a little more complicated. Strategies for overcoming these and other stumbling blocks in the use of proteinase inhibitors in a transgenic approach have been reviewed by Winterer (2002).

#### Other genes with potential

The expression and biological activity of the *Bacillus thuringiensis* (Bt) toxins has been investigated in GM plants for insect control. Bt gene technology is currently the most widely used system for Lepidopteran control in commercial GM crops (Krettiger, 1997). Bt genes are a group of more than fifty insecticidal crystal proteins. When ingested by an insect feeding on Bt expressing Gm plant, the proteins are solubilized in the alkaline environment of the insect's midgut and become toxic by binding to apical border brush membranes of the columnar cells. This causes lysis of the cells and eventual death of the insect. NARO in collaboration with the Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD) in France, is screening a range of newly isolated Bt toxins for banana weevil toxicity. This work is, however, currently being hampered by the absence of an artificial diet for the banana weevil affecting efficient screening under controlled conditions. The expression of a selected Bt gene for weevil resistance may be a rather long-term strategy since no potential Bt gene with high toxic effects against the banana weevil has been identified as yet. In

earlier screening experiments, 80 Bt strains were tested against the banana weevil at the Katholieke Universiteit Leuven (KUL) without much success (Traore et al., unpublished data). It has been suggested that the acidic environment of the banana weevil gut may be influencing the solubility of the Bt toxin. (J-M.Vassal, CIRAD, personal communication).

Plant lectins confer a protective role against a range of organisms (Sharma et al., 2000). Lectins have been isolated from a wide range of plants including snowdrop, pea, wheat, rice and soybean. Their carbohydrate-binding capability renders them toxic to insects. A lectin from snowdrop, *Galanthus nivalis* agglutinin (GNA), is toxic to several insect pests in the orders Homoptera, Coleoptera and Lepidoptera (Tinjuangjun, 2002). In collaboration with NARO, the effect of GNA and the *Aegopodium podagraria* lectin (APA) on the mortality and reproduction of three nematode species pathogenic to banana is being tested at KUL (Carlens, 2002). Similar work can be extended to banana weevil using *in vivo* assays. A major concern about the use of lectins, however, is that some of them, such as the wheat germ agglutinin (WGA), are toxic to mammals (Jouanin et al., 1998). However, the snowdrop and garlic lectins are toxic only to insects (Boulter, 1993), and are potential candidates for weevil control.

Vegetative insecticidal proteins (VIPs) are similar to Bt toxins. VIP1 and VIP2 are contained in supernatants of *Bacillus cereus* (Warren, 1997) while VIP3 can be found in the supernatants of *B. thuringiensis*. They function by causing gut paralysis followed by lysis of the gut epithelium cells, completely arresting gut function and leading to the death of the insect (Duck and Evola, 1997).

Alpha-amylase inhibitors (AI) and chitinase enzymes might also have a future potential for weevil control. Alpha-amylase inhibitors operate by inhibiting the enzyme alpha-amylase, which breaks down starch to glucose in the insect gut (Le Berre-Anton et al., 1997; Morton et al., 2000). They are divided into two types, AI-1 and AI-2, isolated from common and wild beans (*Phaseolus vulgaris*), respectively. Ishimoto et al. (1996) produced transgenic adzuki beans with enhanced resistance to bean bruchids, which are Coleopteran insects. Alpha-amylase inhibitors may be of interest for banana weevil control in GM banana. Chitinase enzymes are produced as a result of invasion either by fungal pathogens or insects. Transgenic expression of chitinase has shown improved resistance to *Lepidopteran* insect pests in tobacco (Ding et al., 1998). At KUL, a rice chitinase gene has been transferred into bananas with the objective of controlling fungal pathogens (Arinaitwe, 2002).

## THE WAY FORWARD

Plant engineering has, without doubt, the potential to play

a key role in the sustainable production of bananas and plantains in Africa. This has recently been realised by several African governments and international donors by providing, in general, better financial support for research into banana and plantain biotechnology. Such support includes the extensive training of young African scientists in plant biotechnology in both African and overseas research institutions, and the setting up of research networks in plant biotechnology on the African continent.

The current efforts at NARO, IITA and the African Centre for Banana and Plantain (CARBAP) in Cameroon, to develop weevil resistance by conventional means using carefully selected parents and especially improved diploids with resistance should continue to complement the biotechnology efforts. This will provide a two-pronged strategy that would increase the chances of developing resistance against the banana weevil. The development of rapid screening methods for testing weevil resistance in the field and laboratory is an essential step in developing resistant hybrids (K  h   et al., 2000; Messiaen, 2002). A better understanding of resistance mechanisms and genes that control these mechanisms are among the priorities in developing successful resistance for the banana weevil.

Unfortunately, certain barriers exist in Africa that limit the opportunity to take advantage of the benefits plant biotechnology might offer. Affordability is one of the major barriers for the introduction of any costly biotechnology-derived product. Also, what might look exciting in the lab might not necessarily make a good and affordable product when considering the needs of small-scale farmers in Africa. In this respect, one should also consider whether the over-expression of a single selected native plant gene, as is done in many engineering approaches, is not too simplistic and ultimately ineffective in obtaining efficient pest resistance. Natural plant resistance is mostly polygenic. Recent evidence shows, however, that these sophisticated defence mechanisms have been lost during selection for domestication (Carlini and Grossi-de-Sa, 2001). Therefore, one approach would be to optimise a "resistance" gene by protein engineering, or a balanced interaction that involves the simultaneous expression of several protective proteins by using gene pyramiding or multiple resistance engineering (Winterer, 2002). In addition, when a triploid GM banana is produced, it will have to be propagated vegetatively for large-scale multiplication. This raises questions about the genomic integrity of the off-spring plant, such as the occurrence of mutations and epigenetic changes as a result of the plant transformation process (Karp, 1993; Phillips et al., 1994; Labra et al., 2001; Van der Vyver et al., 2003).

Without doubt, any GM banana expressing banana weevil resistance would have great value for the small-scale farmers. However, the current negative public perception of GM crops and concerns of bio-safety has to be addressed. Research on performance of GM bananas

under African field conditions and the long-term response of weevils will be necessary to make a judgement of GM technology on weevil resistance. These concerns are important even in African countries, where crop losses due to stress are most severe and which would, therefore, benefit most from the successful development of GM crops.

The need to strengthen the whole transgenic approach by developing a wide range of technical expertise through training cannot be over emphasised. The benefits would include the development of integrated biotechnology teams with overlapping expertises required for success.

## ACKNOWLEDGEMENTS

We are grateful to the Rockefeller foundation and the Uganda Government for financial support for the work on banana weevil resistance screening and resistance mechanisms in Uganda, and the ongoing work on the potential of cysteine proteinase inhibitors for banana weevil control. We thank, NARO and the International Network for Improvement of Banana and Plantain (INIBAP) for coordinating the projects and the following people for their personal communications Edmond DeLanghe, Philip Vain (JIC, UK) and Jean-Michel Vassal (CIRAD).

## REFERENCES

- Abera AMK, Gold CS, Kyamanywa S (1999). Timing and distribution of attack by the banana weevil (Coleoptera: Curculionidae) in East Africa highland banana (*Musa* spp.). Fla. Entomol. 82:631-641.
- Arinaitwe G, (2002). Genetic improvement of East African Highland Banana (*Musa* spp) via engineering for fungal disease resistance. In: Novel Approaches to the improvement of banana production in Eastern Africa-the application of biotechnological methodologies Project Report (Annexes) INIBAP, Montpellier France
- Ashouri A, Michaud D, Cloutier C (2001). Recombinant and classically selected factors of potato plant resistance to the Colorado potato beetle, *Leptinotarsa decemlineata*, variously affect the potato aphid parasitoid *Aphidius nigripes*. Biocontrol 46:401-18.
- Balachowsky AS (1963). Entomologie appliquée à l'agriculture. Traité I 2:1099-1114.
- Boulter D (1993). Insect pest control by copying nature using genetically engineered crops. Phytochemistry 34:1453-1466.
- Buddenburg WJ, Ndiege IO, Karago FW, Hanson BSS (1993). Behavioural and electrophysiological responses of the banana weevil *Cosmopolites sordidus* to host plant volatiles. J. Chem. Ecol. 19:267-277.
- Carlens K (2002). Nematode control in East African regions by the transgenic banana (*Musa* spp.) expressing different types of lectins. In: Novel Approaches to the improvement of banana production in Eastern Africa-the application of biotechnological methodologies Project Report (Annexes) INIBAP, Montpellier France.
- Carlini CR, Grossi-de-Sa MF (2001). Plant toxic proteins with insecticidal properties. A review on their potentialities as bioinsecticides. Toxicon 40:1515-1539.
- Carozi, N., Koziel M., 1997. Advances in insect control the role of transgenic plants. Taylor and Francis, London.
- Collins PJ, Treverrow NL, Lambkin TM (1991) Organophosphorous insecticide resistance and its management in the banana weevil borer, *Cosmopolites sordidus* (Germar) (Coleoptera: Curculionidae), in Australia. Crop Protect. 10, 215-21
- Ding X, Gopalakrishnan B, Johnson LB, White FF, Wang X, Morgan TD, Kramer KJ, Muthukrishnan S (1998). Insect resistance of transgenic tobacco expressing an insect chitinase gene. Transgenic Res. 7:77-84.
- Duck N, Evola S (1997). Use of transgenes to increase host plant resistance to insects opportunities and challenges In: Carozi, N., Koziel, M., (Eds.), Advances in Insect Control the Role of Transgenic Plants. Taylor and Francis, London, pp. 45-56.
- Fogain R, Price NS (1994). Varietal screening of some *Musa* cultivars for susceptibility to the banana weevil, *Cosmopolites sordidus* (Coleoptera: Curculionidae). Fruits 49:247-251.
- Forgain R, Messiaen S, Foure E (2002). Studies on the banana borer weevil in Cameroon. InfoMusa 11, 1 p. XII
- Gatehouse AMRK, Butler KJ, Fenton KA, Gatehouse JA (1985). Presence and partial characterization of a major proteolytic enzyme in the gut of larval *Callosobruchus maculatus*. Entomol. Exp. Appl. 39:279-286.
- Gatehouse AMR, Davidson GM, Newell CA, Merryweather A, Hamilton WDO, Burgess EPJ, Gilbert RJC, Gatehouse JA (1997). Transgenic potato plants with enhanced resistance to the tomato moth *Lacanobia oleracea*: growth room trials. Mol. Breed. 3:49-63.
- Gerald RR, Kramer KJ, Baker JE, Kanost JF, Behke CA (1997). Proteinase inhibitors and resistance of transgenic plants to insects. In: Carozi, N., Koziel, M., (Eds.), Advances in Insect Control the Role of Transgenic Plants. Taylor and Francis, London, pp. 157-183.
- Gold CS, Bagabe MI, Sendege R (1999a). Banana weevil, *Cosmopolites sordidus* (Germar) (Coleoptera: Curculionidae): tests for suspected resistance to carbofuran and dieldrin in the Masaka District, Uganda. African Entomol. 7:189-196.
- Gold CS, Karamura EB, Kiggundu A, Bagamba F, Abera AMK (1999b). Geographic shifts in highland cooking banana (*Musa* spp. group AAA-EA) production in Uganda. International Journal of Sustainable Development and World Ecology 6: 45-59.
- Gold CS (1998). Banana weevil: ecology pest status and prospects for integrated control with emphasis on East Africa. In: Saini, S.K., (Ed.), Proceedings of a Symposium on Biological Control in Tropical Habitats: Third International Conference on Tropical Entomology. ICIPE, Nairobi, Kenya, pp. 49-74.
- Gold CS (2000). Biology and integrated pest management of banana weevil, *Cosmopolites sordidus* (Germar). In: Molina, A.B., Roa, V.N., Maghuyop, M.A.G., (Eds.), Advancing banana and plantain R&D in Asia and the Pacific vol 10, INIBAP-ASPNET, pp. 28-33.
- Gold CS, Messiaen S (2000). The banana weevil *Cosmopolites sordidus*. *Musa*Pest INIBAP Fact Sheet No 4
- Gold CS, Pena JE, Karamura EB (2001). Biology and integrated pest management for the banana weevil *Cosmopolites sordidus* (Germar) (Coleoptera: curculionidae) Integrated Pest Manage. Rev. 6:79-155.
- Green TR, Ryan CA (1972). Wound-induced proteinase inhibitors in plant leaves: a possible defence mechanism against insects. Science 175:776-777.
- Irie K, Hosoyama H, Takeuchi T, Iwabuchi K, Watanabe H, Abe M, Abe K, Arai S (1996). Transgenic rice established to express corn cystatin exhibits strong inhibitory activity against insect gut proteinases. Plant Mol. Biol. 30:149-157.
- Ishimoto M, Sato T, Chrispeels MJ, Kitamura K (1996). Bruchid resistance of transgenic azuki bean expressing seed  $\alpha$ -amylase inhibitor of common bean. Entomol. Exp. Appl. 79:309-315.
- Ittyeipe K (1986). Studies on host preference of banana weevil borer, *Cosmopolites sordidus* GERM. (Curculionidae-Coleoptera). Fruits 41:375-79.
- Jouanin L, Bonade-Bottino M, Girard C, Morrot G, Giband M (1998). Transgenic plants for insect resistance. Plant Sci. 131:1-11.
- Karp A (1993). Are your plants normal? Genetic instability in regenerated and transgenic plants. Agro-Food-Industry Hi-Tech May/June, pp. 7-12.
- Kéhé M, Toto HC, Gnonhoui P (2000). A method for the rapid detection of sensitivity of bananas to weevil, *Cosmopolites sordidus* Germ. (Coleoptera, Curculionidae): application to research on insecticide efficacy. Acta Hort. 540: 497-503
- Kiggundu A (2000). Host-plant interactions and resistance mechanisms to banana weevil *Cosmopolites sordidus* (Germar) in Ugandan *Musa* germplasm. M.Sc. Thesis. University of the Orange Free State,

- Bloemfontain, South Africa.
- Kiggundu A, Gold CS, Labuschagne M, Vuylsteke D, Louw SVDM (2003). Levels of host plant resistance to banana weevil *Cosmopolites sordidus* (Germar) (Coleoptera: Curculionidae) in African *Musa* germplasm. *Euphytica* 133: 267-277.
- Kiggundu A, Vuylsteke D, Gold CS (1999). Recent advances in host plant resistance to banana weevil, *Cosmopolites sordidus* (Germar) In: Frison, E. A., Gold C.S., Karamura, E. B., Sikora, R. A., (Eds.), Mobilizing IPM for sustainable banana production in Africa Proceedings of a workshop on banana IPM held in Nelspruit, South Africa -23-28<sup>th</sup> November 1998. INIBAP, Montpellier, France, pp. 87-96.
- Krettiger AF (1997). Insect resistance in crops: a case study of *Bacillus thuringiensis* (Bt.) and its transfer to developing countries. International Service for the Acquisition of Agri-biotech Applications. Ithica, USA.
- Labra M, Savini C, Bracale M, Pelucchi N, Colombo L, Bardini M, Sala F (2001). Genomic changes in transgenic rice (*Oryza sativa* L.) plants produced by infecting calli with *Agrobacterium tumefaciens*. *Plant Cell Rep.* 20: 325-330.
- LeBerre-Anton V, Bompard-Gilles C, Payan F, Rouge P (1997). Characterization and functional properties of the alpha-amylase inhibitor (alpha A-1) from kidney bean (*Phaseolus vulgaris*) seeds. *Biochim. Biophys. Acta* 1343:31-40.
- Lecardonnel A, Chauvin L, Jouanin LO, Beaujean A, Prevost G, Sangwan-Novrreel B (1999). Effect of rice cystatin I expression in transgenic potato on Colorado potato beetle larvae. *Plant Sci.* 140:71-79.
- Lemaire L (1996). Les relations sémiocchimiques chez le charançon du bananier *Cosmopolites sordidus* Germar (Coleoptera: Curculionidae) et la resistance de sa plante-hôte, le bananier. Thèse de doctorat, Université de Montpellier II, France.
- Leple JC, Bonadebottino M, Augustin S, Pilate G, Letan VD, Delplanque A, Cornu D, Jouanin L (1995). Toxicity to *Chrysomela tremulae* (Coleoptera, Chrysomelidae) of transgenic poplars expressing a xysteine proteinase inhibitor. *Mol. Breed.* 1:319-328.
- Mesquita ALM, Alvers EJ, Caldas RC (1984). Resistance of banana cultivars to *Cosmopolites sordidus* (GERMAR, 1824). *Fruits* 39:254-257.
- Messiaen S (2002). Components of a strategy for the integrated management of banana weevil *Cosmopolites sordidus* (Germar) (Coleoptera: Curculionidae). Dissertationes de Agriculture No. 450. PhD thesis, Faculty of Agricultural and Applied Biological Sciences, Catholic University of Leuven, Belgium.
- Morton RL, Schoroeder HE, Bateman KS, Chrispeels MJ, Armstrong E, Higgins TJV (2000). Bean  $\alpha$ -amylase inhibitor-I in transgenic peas (*Pisium sativum*) provided complete protection from pea weevil (*Bruchus pisorum*) under field conditions. *Proc. Natl. Acad. Sci.* 97:3820-3825.
- Murdock LL, Brookhart G, Dunn PE, Foard DE, Kelley S, Kitch L, Shade RE, Shuckle RH, Wolfson JL (1987). Cysteine digestive proteinases in Coleoptera. *Comp. Biochem. Physiol.* 87:783-787.
- Musabyimana T, Seshu-Reddy KV, Ngode L (2000). Evaluation of banana cultivars for resistance to the banana weevil *Cosmopolites Sordidus* and nematode complex in western Kenya. *Acta Hort.* 540:233-238.
- Nutt KA, Aillsop PG, McGhie TK, Shepaerd KM, Joyce PA, Taylo GO, McQualter RB, Smith GR, Ogarth DM (1999). Transgenic sugarcane with increased resistance to canegrubs. Proceedings of the 1999 Conference of Australian Society of Sugarcane Technologies, Brisbane, Australia, pp. 171-176.
- Ortiz R, Vuylsteke D, Dumpe B, Ferris RSB (1995). Banana weevil resistance and corm hardness in *Musa* germplasm. *Euphytica* 86:95-102.
- Ortiz R, Vuylsteke D (1994). Plantain breeding at IITA. In Jones (Ed.) The Improvement and Testing of *Musa*: A Global Partnership. INIBAP Montpellier, France, pp. 63-76
- Ostmark HE (1974). Economic insect pests of bananas. *Ann. Rev. Entomol.* 19:161-176
- Padmanaban B, Sundararaju P, Velayudhan KC, Sathiamoorthy S (2001). Evaluation of *Musa* germplasm against banana weevil borers *InfoMusa* 10:26-28
- Painter RH (1951). Insect resistance in crop plants. The Macmillan Co., New York. 520.
- Pavis C, Lemaire L (1997). Resistance of *Musa* germplasm to the banana borer weevil, *Cosmopolites sordidus* Germar (Coleoptera: Curculionidae) *InfoMusa* 5:3-9
- Pavis C, Minost C (1993). Banana resistance to the banana weevil borer *Cosmopolites sordidus* Germar (Coleoptera: Curculionidae); role of pseudostem attractivity and physical properties of the rhizome. In: Proceedings of international symposium on genetic improvement of banana for resistance to diseases and pests, CIRAD-FLHOR, Montpellier France, pp. 129-142.
- Pernas M, Sanchez-Mong R, Salcedo G (2000). Biotic and abiotic stress can induce cystatin expression in chestnut. *FEBS Lett.* 467:206-210.
- Phillips RL, Kaeppler SM, Olhoff P (1994). Genetic instability of plant tissue cultures: Breakdown of normal controls. *Proc. Natl. Acad. Sci.* 91:5222-5226.
- Price NS, (1994). Alternate cropping in the management of *Rodopholus similes* and *Cosmopolites sordidus*, two important pests of banana and plantain. *Int. J. Pest Manage.* 40:237-244.
- Rukazambuga NDTM, Gold CS, Gowen SR (1998). Yield loss in East African Highland Banana (*Musa* spp., AAA-EA group) caused by the banana weevil, *Cosmopolites sordidus* Germar. *Crop Protect.* 17:1-9.
- Ryan CA (1990). Proteinase inhibitors in plants: genes for improving resistance against insects and pathogens. *Ann. Rev. Phytopathol.* 28:425-449.
- Schroeder HE, Gollasch S, Moore A, Tabe LM, Craig S, hardie DC, Chrispeels MJ, Spencer D, Higgins TJV (1995). Bean alpha-amylase inhibitors confers resistance to pea weevil (*Bruchus pisorum*) in trasgenic peas (*Pisum sativum* L.) *Plant Physiol.* 107:1233-1239.
- Sengooba T (1986). Survey of banana pest problem complex in Rakai and Masaka Districts in Uganda. August 1986: Preliminary trip report. Namulonge Research Station, Namulonge, Uganda.
- Seshu-Reddy KV, Lubega MC (1993). Evaluation of banana cultivars for resistance to tolerance of the weevil *Cosmopolites sordidus* Germar, In: Ganry J., (Ed.), Breeding Banana and Plantain for Resistance to Diseases and Pests, CIRAD/INIBAP, Montpellier, France, pp. 143-148.
- Sharma HC, Sharma KK, Seetharama N, Ortiz R (2000). Prospects for using transgenic resistance to insects in crop improvement *EJB Electron. J. Biotechnol.* 3 (2) <http://www.ejb.org/content/vol3/issue2/full/3>
- Swennen R, Vuylsteke D (2001). Banana *Musa* L. In: Raemaekers R.H., (Ed.), Crop Production in Tropical Africa. DGIC Brussels, Belgium, pp. 530-552.
- Tinjuangjun P (2002). Snowdrop lectin gene in transgenic plants: its potential for Asian agriculture [Agbiotech.net.com](http://Agbiotech.net.com) (ABN091)
- Van der Vyver C, Kunert K, Cullis C (2003). Detection and characterization of a labile DNA region in the tobacco genome using representational difference analysis. *Molecular Genetics and Genomics* (submitted).
- Warren GW (1997). Vegetative insecticidal proteins: novel proteins for control of corn pests. In: Carozzi, Koziel, M., (Eds.), Advances in Insect Control: The Role of Transgenic Plants Taylor and Francis, London, pp. 109-121.
- Willson WD, Flint HM, Deaton RW, Fischhoff DA, Perlak FJ, Armstrong TA, Fusch RL, Berberich SA, Parks NJ Stapp BR (1992). Resistance of cotton lines containing *Bacillus thuringiensis* toxin to pink bollworm (Lepidoptera: Gelechiidae) and other insects. *J. Econ. Entomol.* 85:1516-1521.
- Winterer J (2002). The mixed success of proteinase inhibitors to combat insect pests in transgenic crops. *AgBiotechNet* 4, 1-7.
- Xu D, Xue Q, McElroy D, Mawal Y, Hilder V, Wu R (1996). Constitutive expression of a cowpea trypsin inhibitor gene, CpTi, in transgenic rice plants confers resistance to two major rice insect pests. *Mol. Breed.* 2:167-173.
- Ye GY, Yao HW, Shu QY, Cheng X, Hu C, Xia YW, Gao MW, Altosaar I (2003). High levels of stable resistance in transgenic rice with cry1Ab gene from *Bacillus thuringiensis* Berliner to rice leafhopper, *Cnaphalocrocis medilinalis* (guenee) under field conditions. *Crop Protect.* 22:171-178.