

African Journal of Biotechnology Vol. 9(52), pp. 8918-8922, 21 December, 2010  
Available online at <http://www.academicjournals.org/AJB>  
ISSN 1684-5315 ©2010 Academic Journals

## Full Length Research Paper

# Effect of fertilizer treatment on the antimicrobial activity of the leaves of *Ocimum gratissimum* (L.) and *Gongronema latifolium* (Benth)

G. G. E. Osuagwu\* and H. O. Edeoga

Biological Sciences Department, Michael Okpara University of Agriculture, Umudike, PMB 7267, Umuahia, Abia State, Nigeria.

Accepted 5 July, 2010

The effect of fertilizer treatment on the antimicrobial activity of the leaves of *Ocimum gratissimum* (L.) and *Gongronema latifolium* (Benth) was investigated. Cultivated *O. gratissimum* (L.) and *G. latifolium* were applied with NPK (15:15:15) fertilizer at 100, 200, 300, 400 and 500 kg/ha treatment levels in planting buckets derived using the furrow slice method two months after seedling emergence. No fertilizer treatment served as control. Leaves were harvested one month after treatment. The ethanolic extracts of the harvested leaves were used to determine the sensitivity of the extracts on *Klebsiella pneumonia*, *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Streptococcus faecalis*, *Candida albicans* and *Aspergillus niger*. The result obtained showed that the antimicrobial activity of the leaves of *O. gratissimum* and *G. latifolium* was significantly ( $P < 0.05$ ) increased by fertilizer treatment. The inhibition zone increased with increase in the level of fertilizer treatment. The ethanolic extracts of both plants whether treated or not had no antimicrobial effect on *A. niger*. This research revealed that fertilizer treatment might have increased the phytochemical content of the leaves of the plants which in turn enhanced their antimicrobial potential.

**Key words:** Pathogens, NPK fertilizer, ethanolic extracts, inhibition zone.

## INTRODUCTION

The role of plants in the maintenance of good health has been reported (Burkill, 1995; Moerman, 1996). In Nigeria, these indigenous plants contain bioactive compounds that exhibit physiological activities against bacteria and other microorganisms and are also used as precursor for the synthesis of useful drugs (Edeoga et al., 2003; Okwu, 2001; Osuagwu et al., 2007; Osuagwu, 2008; Sofowara, 1993). The antimicrobial activities of these plants and their products such as essential oils are well documented (El-Zaher et al., 2006; Ijeh et al., 2006; Mevy et al., 2007; Sahraoui et al., 2007; Vagionas et al., 2007). Thus, these plants are therefore used in the treatment of many diseases such as rheumatism, diarrhea, malaria, elephantiasis, cold obesity, dysentery, high blood pressure, malnutrition, gonorrhoea and others (Batram, 1998;

Burkill, 1995; Gill, 1992).

The biosynthesis of these bioactive plant chemicals is influenced by various agronomic and environmental factors. Fertilizer treatment is known to determine the concentration of these compounds in plants (Asami et al., 2003; Khalil et al., 2007; Osuagwu and Nwachukwu, 2007; Rasmussen et al., 2008; Saradhi et al., 2007). Water stress (drought) is also reported to influence the concentration of these phytochemicals in plants (Selmar, 2008; Zheng et al., 2006).

The antimicrobial activities of *Ocimum gratissimum* and *Gongronema latifolium* have been investigated and reported. The antimicrobial activities of *O. gratissimum* have been documented (Ezekwesili et al., 2004; Ijeh et al., 2005; Iwalokun et al., 2003; Matasyoh et al., 2007; Mbata and Salkia, 2007; Tchoumboungong et al., 2005) and that of *G. latifolium* reported (Afolabi and Eleyinmi, 2007). The objective of this research is to ascertain the implication of fertilizer treatment on the antimicrobial activity of the

\*Corresponding author. E-mail: [gbekus2002@yahoo.com](mailto:gbekus2002@yahoo.com).

**Table 1.** Physical properties of the soil.

Properties	Values
<b>Particle size distribution</b>	
Sand	70.90%
Silt	15.40%
Clay	13.70%
Texture	5 L
pH (H <sub>2</sub> O)	5.01
Organic carbon (%)	0.75%
Organic matter (%)	1.29%
Available phosphorus (mg/Kg)	46.00
Total nitrogen (%)	0.08%
<b>Exchangeable bases (mg/100g)</b>	
Ca <sup>++</sup>	2.40
Mg <sup>++</sup>	2.00
K <sup>+</sup>	0.07
Na <sup>+</sup>	0.23
Exchangeable acidity (ME/100g)	1.20
Effective cation exchange capacity (ME/100g)	5.90

leaves of *O. gratissimum* and *G. latifolium*.

## MATERIALS AND METHODS

### Plant sample

The seeds of *O. gratissimum* were collected from a homestead garden in Amaogwu village Bende town, Bende Local Government Area of Abia State. The fresh and succulent stem cuttings of *G. latifolium* were obtained from the forest strip of the Forest Department, College of Natural Resources and Environmental Management, Michael Okpara University of Agriculture, Umudike, Umuahia, Abia State. Both plant materials were identified by the taxonomic unit of the Botany section of the Department of Biological Sciences, Michael Okpara University of Agriculture, Umudike, Umuahia, Abia State. The seeds of *O. gratissimum* were raised into seedling in nursery boxes before they were transplanted into planting buckets. Stem cuttings of *G. latifolium* were planted directly into the planting buckets.

Cultivation of the plants was carried out using 24 plastic buckets containing 8 Kg of sterilized soil. The soil used for the research was analyzed to determine the physiochemical properties (Table 1).

Treatments were carried out in four replicates of each treatment. The inorganic fertilizer (NPK 15:15:15) used for the study was obtained from the store of the Abia State Ministry of Agriculture, Umuahia, Abia State. Five levels of fertilizer treatments 100, 200, 300, 400 and 500 Kg/ha derived using the furrow slice method (Brady and Weil, 1999), in four replicates was used. No fertilizer treatment served as control. Treatment occurred two months after seedling emergency. Harvesting of plants leaves for antimicrobial activity investigation was carried out one month after treatment.

### Determination of antimicrobial activity

#### Preparation of plant extract

The ethanolic extracts of the leaves of *O. gratissimum* and *G. latifolium* was prepared using the method of Ijeh et al. (2005). Fifty

grams of the pondered samples was soaked in 200 ml of absolute ethanol and allowed to stand for 24 h. It was filtered using a Whatman (No. 1) filter paper. The filtrate was evaporated to dryness over steam bath. The residue was dissolved in deionized water to obtain the desired plant extract for the antimicrobial tests.

#### Preparation of inocula

*Klebsiella pneumonia*, *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Streptococcus faecalis*, *Candida albicans* and *Aspergillus niger* used in the research were obtained from the stock culture of the Microbiology Laboratory, Federal Medical Centre, Umuahia, Abia State, Nigeria. Viability test of each isolate was carried out by resuscitating the organisms in buffered peptone broth and thereafter, sub-cultured into nutrient agar medium and incubated at 37 °C for 24 h.

#### Antimicrobial test

The sensitivity of the test organisms to the ethanolic extracts of the leaves of *O. gratissimum* and *G. latifolium* was carried out using the diffusion method described by Ebi and Ofoefule (1997).

20 ml of the molten nutrient agar was seeded with 0.2 ml of broth culture of the test organisms in sterile Petri dishes. The Petri-dishes were rotated slowly to ensure a uniform distribution of the organisms. They were left to solidify and in the dish cups of 8.0 mm diameter were made in the agar using a sterile Pasteur pipette. The Petri-dishes were allowed to stand for about 30 min at room temperature to allow for proper diffusion of the extracts to take place. The plates were then incubated at 37 °C for 24 h. The zone of inhibition in millimeter were measured and recorded. The test was carried out in the laboratory of the Department of Biological Sciences, Michael Okpara University of Agriculture, Umudike, Umuahia, Abia State.

#### Statistical analysis

The design for the research was complete randomized design in four replicates of each treatment. Analysis of variance (ANOVA) was used to analyze the data and LSD at 0.05 probability level was used to determine the difference among treatments.

## RESULTS AND DISCUSSION

Fertilizer treatment significantly ( $P < 0.05$ ) affected the antimicrobial activity of the leaves of *O. gratissimum* and *G. latifolium* (Tables 2 – 5).

There was significant increase ( $P < 0.05$ ) in the ability of the leaves of *O. gratissimum* to inhibit the activity of *K. pneumonia*, *E. coli*, *S. aureus*, *P. aeruginosa*, *S. typhi*, *S. faecalis* and *C. albicans* due to fertilizer treatment (Tables 2 and 3). Fertilizer treatment also significantly ( $P < 0.05$ ) increased the ability of the leaves of *G. latifolium* to inhibit the activity of *K. pneumonia*, *E. coli*, *S. aureus*, *P. aeruginosa*, *S. typhi*, *S. faecalis* and *C. albicans* (Tables 4 and 5).

The observed increased ability of the leaves of *O. gratissimum* and *G. latifolium* to inhibit the microbial activity of these pathogens as testified by the inhibition zones in response to fertilizer treatment might be related to the enhanced synthesis and accumulation of phyto-

**Table 2.** Effect of NPK fertilizer treatment of the antimicrobial activity of the leaves of *O. gratissimum* on *K. pneumonia*, *E. coli*, *S. aureus* and *P. aeruginosa*.

Treatment	Zone of inhibition (mm)			
	<i>K. pneumonia</i>	<i>E. coli</i>	<i>S. aureus</i>	<i>P. aeruginosa</i>
Control	20.00 <sup>a</sup>	26.00 <sup>a</sup>	13.00 <sup>a</sup>	16.00 <sup>a</sup>
100 kg/ha	25.00 <sup>b</sup>	28.00 <sup>ab</sup>	21.00 <sup>b</sup>	16.00 <sup>ab</sup>
200 kg/ha	27.00 <sup>bc</sup>	30.00 <sup>bc</sup>	24.00 <sup>bc</sup>	20.00 <sup>bc</sup>
300 kg/ha	29.00 <sup>c</sup>	32.00 <sup>cd</sup>	26.00 <sup>cd</sup>	23.00 <sup>cd</sup>
400 kg/ha	32.00 <sup>d</sup>	34.00 <sup>de</sup>	28.00 <sup>cd</sup>	24.00 <sup>d</sup>
500 kg/ha.	34.00 <sup>d</sup>	36.00 <sup>e</sup>	29.00 <sup>d</sup>	26.00 <sup>a</sup>
LSD (< 0.05)	1.414	1.810	1.863	1.691

Mean of four replicates. Different letters in the superscript in the same row indicates significant difference ( $P < 0.05$ ).

**Table 3.** Effect of NPK fertilizer treatment on the antimicrobial activity of the leaves of *O. gratissimum* on *S. typhi*, *S. faecalis*, *C. albicans* and *A. niger*.

Treatment	Zone of inhibition (mm)			
	<i>S. typhi</i>	<i>S. faecalis</i>	<i>C. albicans</i>	<i>A. niger</i>
Control	17.00 <sup>a</sup>	15.00 <sup>a</sup>	7.00 <sup>a</sup>	0.00
100 kg/ha	20.00 <sup>ab</sup>	18.00 <sup>b</sup>	12.00 <sup>bc</sup>	0.00
200 kg/ha	22.00 <sup>bc</sup>	21.00 <sup>b</sup>	14.00 <sup>c</sup>	0.00
300 kg/ha	25.00 <sup>cd</sup>	22.00 <sup>bc</sup>	10.00 <sup>ab</sup>	0.00
400 kg/ha	27.00 <sup>d</sup>	25.00 <sup>cd</sup>	12.00 <sup>bc</sup>	0.00
500 kg/ha.	28.00 <sup>d</sup>	27.00 <sup>d</sup>	13.00 <sup>bc</sup>	0.00
LSD (< 0.05)	1.599	1.740	1.599	—

Mean of four replicates. Different letters in the superscript in the same row indicates significant difference ( $P < 0.05$ ).

**Table 4.** Effect of NPK fertilizer treatment on the antimicrobial activity of the leaves of *G. latifolium* on *K. pneumonia*, *E. coli*, *S. aureus* and *P. aeruginosa*.

Treatment	Zone of inhibition (mm)			
	<i>K. pneumonia</i>	<i>E. coli</i>	<i>S. aureus</i>	<i>P. aeruginosa</i>
Control	13.00 <sup>a</sup>	15.00 <sup>a</sup>	15.00 <sup>a</sup>	10.00 <sup>a</sup>
100 kg/ha	20.00 <sup>b</sup>	25.00 <sup>b</sup>	23.00 <sup>b</sup>	17.00 <sup>b</sup>
200 kg/ha	24.00 <sup>c</sup>	27.00 <sup>cd</sup>	26.00 <sup>bc</sup>	21.00 <sup>c</sup>
300 kg/ha	26.00 <sup>cd</sup>	30.00 <sup>cd</sup>	25.00 <sup>bc</sup>	20.00 <sup>bc</sup>
400 kg/ha	28.00 <sup>d</sup>	32.00 <sup>cd</sup>	28.00 <sup>cd</sup>	25.00 <sup>d</sup>
500 kg/ha.	32.00 <sup>e</sup>	35.00 <sup>e</sup>	30.00 <sup>d</sup>	28.00 <sup>d</sup>
LSD (< 0.05)	1.546	1.740	1.732	1.683

Mean of four replicates. Different letters in the superscript in the same row indicates significant difference ( $P < 0.05$ ).

chemicals (alkaloids, phenols, saponins, steroids, tannins and other plant chemicals substances) by these plants a consequence of fertilizer treatment. This in turn increased the ability of the extracts of the leaves to inhibit the activity of these pathogens. Enhanced production of phytochemicals, vitamins and other plant chemical substances by plants as a result of fertilizer treatment has been reported (Das et al., 2006; Koloziej, 2007; Mozaffar, 1994; Osuagwu and Nwachukwu, 2007; Osuagwu, 2008;

Omoidbaigi et al., 2008; Polat et al., 2008; Silva et al., 2006; Zheng et al., 2006).

The leaf extracts of *O. gratissimum* and *G. latifolium* had no antimicrobial effect on *A. niger*. The leaves of both treated and untreated plants did not inhibit the microbial activity of *A. niger*. This indicates that *A. niger* is resistant to the chemicals contained in the leaves of the plants. The extracts of the leaves of *O. gratissimum* and *G. latifolium* had more inhibitory effect on the microbial

**Table 5.** Effect of NPK fertilizer treatment on the antimicrobial activity of the leaves of *G. latifolium* on *S. typhi*, *S. faecalis*, *C. albicans* and *A. niger*.

Treatment	Zone of inhibition (mm)			
	<i>S. typhi</i>	<i>S. faecalis</i>	<i>C. albicans</i>	<i>A. niger</i>
Control	12.00 <sup>a</sup>	14.00 <sup>a</sup>	5.00 <sup>a</sup>	0.00
100 kg/ha	24.00 <sup>b</sup>	22.00 <sup>b</sup>	14.00 <sup>b</sup>	0.00
200 kg/ha	26.00 <sup>bc</sup>	25.00 <sup>c</sup>	16.00 <sup>bc</sup>	0.00
300 kg/ha	22.00 <sup>b</sup>	22.00 <sup>b</sup>	16.00 <sup>b</sup>	0.00
400 kg/ha	26.00 <sup>bc</sup>	24.00 <sup>b</sup>	16.00 <sup>bc</sup>	0.00
500 kg/ha.	29.00 <sup>c</sup>	26.00 <sup>bc</sup>	19.00 <sup>c</sup>	0.00
LSD (< 0.05)	1.764	1.856	1.683	—

Mean of four replicates. Different letters in the superscript in the same row indicates significant difference ( $P < 0.05$ ).

activity of bacteria, when compared with the effect on fungi (Tables 2 – 5). The above observation relates to the reactions of different organisms to similar and different environmental conditions.

Furthermore, generally, increasing the level of fertilizer treatment led to corresponding increase in the ability of the leaves of the plants to inhibit the microbial activity of the pathogens, which in turn affect the functions of the quantity of phytochemicals they contain. Wiesler (1997) observed that the impact of nitrogen fertilizer on metabolism of plants depends on factors such as its concentration in the soil and chemical form.

The importance of the finding from this research is that the application of NPK fertilizer at the appropriate levels will enhance the antimicrobial activity of the leaves of *O. gratissimum* and *G. latifolium* and thus, increase their value and efficacy as medicinal plants.

## REFERENCES

- Afolabi F, Eleyinmi A (2007). Chemical composition and antibacterial activity of *Gongronema latifolium*. J. Zheiang Univ. Sci. B. 8(5): 352-358.
- Brady NC, Weil RR (1999). The nature and properties of soils. 12th edition. Prentice Hall, New Jersey. p. 881.
- Burkill HM (1995). The useful plants of West Tropical Africa. families J-L. Royal Botanic Garden, Kew. 3: p. 605.
- Das K, Danga R, Shivananda TN (2006). Effect of major nutrients on biomass production and stevioside content of *Stevia rebaudiana* on pot culture study. Biomedical, 1(1): 22-25.
- Ebi GC, Ofoefule SI (1997). Investigation into the folkloric antimicrobial activities of *Landolphia owerriance*. Phytother. Res. 11: 147-151.
- Edeoga HO, Okwu DE, Mbaebie BO (2003). Mineral and nutritive value of some Nigerian medicinal plants. J. Med. Arom. Plant Sci. 25: 689-694.
- El-Zaher FHA, Fayez M, El-Maksoud HKA, Hosny I (2006). Antimicrobial activity of some Egyptian medicinal and aromatic plant waste extracts. Bull. Nat. Res. Centre, (Cairo), 31(1): 1-20.
- Ezekwesili CN, Obiora KA, Ugwu OP (2004). Evaluation of anti-diarrhoeal property of crude aqueous extracts of *Ocimum gratissimum* L. (Labiata) in rats. Biokemistri, 16(2): 122-131.
- Gill LS (1992). Ethnomedicinal uses of plants in Nigeria. University of Benin press, Benin, Nigeria, p.124.
- Ijeh II, Omodamiro OD, Nwanna IJ (2005). Antimicrobial effects of aqueous and ethanolic fractions of two spices, *Ocimum gratissimum* and *Xylopiia netiopica*. Afr. J. Biotechnol. 4(9): 953-956.
- Iwalokun A, Ghenle GO, Adewole TA, Akinsinde KA (2003). Shigellocidal properties of the Nigerian medicinal plants: *Ocimum gratissimum*, *Terminalia aicennaoides* and *Monmerdica balsamma*. J. Health Popul. Nutr. 19(4): 331-335.
- Khalil MY, Moustafa AA, Naguib MY (2007). Growth, phenolic compounds and antioxidant activity of some medicinal plants grown under organic farming condition. World J. Agric. Sci. 3(4): 451-457.
- Koloziej B (2007). The effect of NPK fertilization on golden rod (*Solidago virgaurea*, L. subsp *Virgaurea*) yield and quality parameters. Herba Polonica, 53(3): 129-134.
- Matasyoh IG, Matasyoh JC, Wachira FN, Kinyua MG, Muigal AW, Muklama TK (2007). Chemical composition and antimicrobial activity of the essential oil of *Ocimum gratissimum* L. grown in Eastern Kenya. Afr. J. Biotechnol. 6(6): 760-765.
- Mbata TI, Salkia A (2007). Antibacterial activity of essential oil from *Ocimum gratissimum* on *Listeria monocytogenes* Int. J. Food Saf. 5(7): 15-19.
- Mevy JP, Bessiere JM, dherrbomez M, Millogo J, Viano J (2007). Chemical composition and some biological activities of the volatile oils of a chemotype of *Lippa chevalier*, Moldenke. Food Chem. 101(2): 682-685.
- Moerman DE (1996). An analysis of the food plants of nature. North Am. J. Ethnopharmacol. 52: 1-22.
- Mozzafar A (1994). Enrichment of some B-vitamins in plants with application of organic fertilizers. Plant Soil, 167(2): 3050-3111.
- Okwu DE (2001). Evaluation of the chemical composition of indigenous species and flavouring agents. Global J. Pure Appl. Sci. 7(3): 455-459.
- Omidbaigi R, Dadman B, Fattahi F (2008). Influence of nitrogen fertilizer on the herb yield, essential oil content and composition of *Tagetes minota* L. J. Essential oil-Bearing Plants. 11(1): 45-52.
- Osuagwu GGE (2008). The effect of rate of application of poultry manure on the phenol, flavonoid and steroid potential of the leaves of *Ocimum gratissimum*. J. Sustain. Agric. Environ. 10(2): 106-111.
- Osuagwu GGE, Nwachukwu CM (2007). Effect of organic fertilizer application on the alkaloids, saponin, tannins and cyanogenic glycosides constituents of *Ocimum gratissimum*. Environ. Ecol. 255(4): 1270-1275.
- Osuagwu GGE, Okwulehie IC, Emenike JO (2007). Phytochemical and Mineral content of the leaves of four Nigerian *Pterocarpus* species. Int. J. Mol. Med. Adv. Sci. 3(1): 6-11.
- Polat E, Demir H, Onys AN (2008). Comparison of some yield and quality criteria in organically and conventionally grown lettuce. Afr. J. Biotechnol. 7(9): 1235-1239.
- Rasmussen S, Parsons AJ, Fraser K, Xue H, Newman JA (2008). Metabolic profiles of *Lolium perenne* as differentially affected by nitrogen supply, carbohydrate content and fungal endophyte infection. Plant Physiol. 146: 1440-1453.
- Sahraoui M, Hellal A, Boutekedjiret C, Bentahar F, Bessiere JM (2007). Antimicrobial activities of essential oils of some Algerian aromatic plants. Int. J. Essential Oil Therapeutica, 1(2): 83-90.
- Saradhi VSP, Salma K, Shivananda BY, Kumar TV, Shivananda TN

- (2007). Effect of NPK fertilizers on chemical constituents of *Aloe vera* leaves. *J. Nat. Rem.* 7(2): 258-262.
- Selmar D (2008). Potential of salt and drought stress to increase pharmaceutical significant of secondary compounds in plants. *Landbauforschung, Kendorf*, 58(112): 139-144.
- Silva FY, Pinto JE, Cardoso MG, Sales JF, Mol DJS, Divino SP (2006). Influence of manure and fertilizer on *Baccharis trimers* (less) d c. growth essential oil yield. *J. Herbs, Spices Med. Plants*, 12(1/2): 1-11.
- Sofowara A (1993). *Medicinal Plants and Traditional Medicine in Africa*. Spectrum Books Limited, Ibadan, Nigeria. p. 289.
- Tchoumboungnong PH, Zollo A, Dongne E, Mekonnen Y (2005). In vivo antimalarial activity of essential oils from *Cymbopogon citratus* and *Ocimum gratissimum* L. on mice infected with *Plasmodium berghei*. *Planta Med.* 71: 20-23.
- Vagionas K, Graikou K, Ngassapa O, Rongyoro D, Chinou I (2007). Composition and antimicrobial activity of the essential oils of three *Satureja* species growing in Tanzania. *Food Chem.* 103(2): 319-324.
- Wiesler F (1997). Agronomical and physiological aspects of ammonium and nitrate nutrition in plants. *Zeltchrt, Fiir, Pflanzenernahrung und Bondenkunde.* 160: 227-238.
- Zheng Y, Dixon M, Saxena (2006). Growing environment and nutrient availability affect the content of some phenolic compounds in *Echinacea purpurea* and *Echinacea angustifolia*. *Planta Med.* 72: 1407-1414.