

Full Length Research Paper

Analysis of diversity in rice (*Oryza sativa* L.) using random amplified polymorphic DNA (RAPD) and simple sequence repeats (SSR) markers

Harsh Bansal¹, Ravindra Kumar^{1*}, Vivek² and Sanjay³

¹Department of Fingerprinting, Sardar Vallabhbhai Patel University of Agriculture And Technology, Modipuram, Meerut, (U.P), India.

²Department of Agronomy, Sardar Vallabhbhai Patel University of Agriculture And Technology, Modipuram, Meerut, (U.P), India.

³Division of Genetics, Indian Agriculture Research Institute, New Delhi, India.

Accepted 17 July, 2013

Molecular markers are useful tool for assessing genetic variations and resolving genotype identity. In the current study, genetic diversity among 20 rice genotypes was assessed using the random amplified polymorphic DNA (RAPD) and simple sequence repeat (SSR). In RAPD analysis, 20 primers generated a total of 116 bands of which 114 were polymorphic. The number of amplification products produced by each primer varied from 4 to 7 with an average of 5.8 bands per primer. Twenty (20) SSR primers generated a total of 65 alleles with an average 3.2 alleles per primer. Genetic diversity of 20 genotypes estimated by polymorphic information content (PIC) value ranged from 0.62 to 0.97 in SSR and 0.33 to 0.88 with RAPD analysis. The cluster dendrogram by SSR revealed two major clusters. Rajeshwari was the only genotype in cluster I. The cluster II further divided into two sub clusters IIA and IIB. II A consisted of 17 genotypes while II B consisted of two genotypes (Apo and Kalakeni). The information generated from this study can be used to maximize selection of diverse parents and broaden the germplasm base for the future rice breeding programs.

Key words: Genetic diversity, molecular markers, rice, markers based estimation of genetic diversity in rice.

INTRODUCTION

Rice (*Oryza* spp.) is one of the most important food crops in the world, being planted on almost 11% of the Earth's cultivated land area over a wide number of ecosystems (Cuevas and Fitzgerald, 2012). Rice belongs to the genus *Oryza* and has two cultivated and 22 wild species.

Rice is an ideal model plant for the study of grass genetics and genome organization due to its diploid genetics, relatively small genome size of 430 Mb (Causse et al., 1994), significant level of genetic polymorphism (McCouch et al., 1998), large amount of well conserved genetically

*Corresponding author. E-mail: kumarrk2000@yahoo.com.

Abbreviations: DNA, Deoxyribonucleic acid; CTAB, cetyl tri-methyl ammonium bromide; PCR, polymerase chain reaction; TE, Tris acetate; EDTA, ethylenediamine tetra acetic acid; RAPD, random amplified Polymorphic DNA, SSRs, simple sequence repeats; dNTPs, deoxyribonucleotide tri phosphate.

diverse material (about 100,000 accessions world wide) and the availability of widely collected compatible wild species. The success of breeding program also depends upon the amount of genetic variability present in the population and extent to which the desirable traits are heritable. The assessment of phenotype may not be a reliable measure of genetic differences. The rapid development of biotechnology allows easy analysis of a large number of loci distributed throughout the genome of plants. Molecular markers have proven to be powerful tool in the assessment of genetic variation and in the elucidation of genetic relationships within and among species (Matin et al., 2012). Molecular markers originate from different parts of the genome including coding and non-coding regions and can cover either the full genome or large genomic segments. Morphological traits are controlled by a relatively small number of loci (Onaga et al., 2013) however; they serve as a valuable guide for effective collection and use of genetic resources. Molecular markers (RAPD) without prior knowledge of DNA sequences are especially useful for unzipping the variations in species with low genetic variability. RAPD markers are considered to be unbiased and neutral markers for genetic mapping applications, in population genetics, taxonomy as well as for genetic diagnostics. Microsatellites are the popular molecular markers in rice for various applications in genetics and breeding. SSR markers are important tool for estimation of genetic variation and identification of germplasm. These markers have some merits like quickness, simplicity, rich polymorphism and stability, thus being widely used in molecular map construction and gene mapping, construction of fingerprints and genetic purity test (Ma et al., 2011), analysis of germ plasm diversity (Jin et al., 2010), utilization of heterosis, especially in identification of species with closer genetic relationship. SSR markers are more popular in rice because they are highly informative, mostly monolocus, co dominant, easily analyzed and cost effective (Prabakaran et al., 2010). Using PCR rapid amplification and gel electrophoresis of high resolving power, we can test SSR length polymorphism rapidly and economically. In the current studies, RAPD, SSR and combined analysis was performed to assess the genetic diversity among 20 genotypes of rice.

MATERIALS AND METHODS

Plant material

Leaf samples of 20 rice genotypes (25 days after transplanting.) were collected from the Genetics Division, Indian Agriculture research institute, Delhi, India from wet season 2010. The details of rice genotypes are presented in Table 1.

DNA extraction

Total DNA was extracted from fresh leaves by the cetyl tri-methyl

ammonium bromide (CTAB) method (Murray and Thompson, 1980). The quality and concentration of extracted DNA were estimated by using a UV-Vis spectrophotometer. The DNA was spooled out, washed twice with 70% ethanol and dissolved in TE (10 mM Tris, 0.1 mM EDTA, pH 8.0) containing 25 µg/ml RNase-A, incubated at 37°C for 30 min and extracted with chloroform:iso-amyl alcohol (24:1 v/v). DNA was re-precipitated and dissolved in TE buffer. DNA was checked for its quality and quantity by 0.8% agarose gel electrophoresis.

PCR analysis and gel electrophoresis

In case of SSR, a set of 20 primers (Chakravarthi et al., 2006) were used. The details of primers used for SSR analysis are presented in Table 2. The PCR reaction was carried out using Taq polymerase (Genei) in 20 ml reaction volume containing 1.5X PCR buffer, 2 mM MgCl₂, 0.02 mM of each dNTPs, 1 mM of forward and reverse primers, 0.5 µl (3 unit) Taq polymerase and 50 ng genomic DNA. Profile used was as follows: an initial hot start and denaturing step at 95°C for 5 min followed by 35 cycles at 94°C for 1 min, appropriate annealing temperature 55°C for 1 min, and primer elongation at 72°C for 2 min. Final extension step at 72°C for 7 min was performed.

RAPD analysis was performed according to Williams et al. (1990) with minor modifications. The primers used for RAPD are given in Table 3. PCR reactions were carried out in 20 µl volume containing 50 ng of total genomic DNA, 10 pmol primer, 200 µM dNTPs, 2 mM MgCl₂, 2.5X PCR buffer and 0.4 µl (3 units) AmpliTaq Polymerase. Twenty (20) 10-mer oligonucleotide random primers were selected for analysis. These primers were obtained from Bangalore Genei, India. Amplification was performed in a Astec thermal cycler with the following profile: 94°C for 4 min (initial denaturation), 94°C for 1 min, 32°C for 1 min, 72°C for 2 min for 42 cycles with a final extension at 72°C for 7 min.

The RAPD-PCR products were analyzed directly on 1.5% agarose gels in TAE buffer while SSR-PCR products were analyzed on 4% agarose gel, visualized by staining with ethidium bromide under short-wave UV light. DNA ladder used in the electrophoresis was of 100 bp. Agrose gel electrophoresis for PCR products of RM-222, SSR marker and OPA-07, RAPD marker is shown in Figure 4 and 5 respectively.

Data analysis

Pair wise comparison of genotypes, based on the presence (1) or absence (0) of unique and shared polymorphic products was used to generate similarity coefficients of Jaccard's coefficient by NTSYS-pc version 2.1 software (Rohlf, 2000). The similarity coefficient was used to construct a dendrogram by the unweighted pair group method with arithmetic averages (UPGMA) according to Rohlf (1993). The polymorphism information content (PIC) value described by Botstein et al. (1980) and modified by Anderson et al. (1993) for self-pollinated species was calculated as follows:

$$PIC_i = 1 - \sum_{j=1}^n P_{ij}^2$$

Where, p_i equals to the frequency of the i th allele and p_j the frequency of the allele. Only data from polymorphic loci were used for analysis.

Table 1. Characteristics of rice genotypes used in investigation.

Name of Variety	Sources/origin	Characteristic features
Apo	IRRI, Philippines	Suitable for aerobic cultivation, larger and fuller grains, tall
Kalakeni	Orissa, India	Semi dwarf, Grain, long slender, tolerant to alkalinity.
Satabdi	West Bengal, India	Semi-dwarf, grains: LS, white, resistant to Sh.B, BB & Sh.R;
Ratna	Orissa, India	Dwarf, long slender and white grain. Moderately resistant to blast, leaf hopper and tolerant to stem borer.
Khandagini	Orrisa, India	Semi dwarf grains- LS, white tolerant to blast, BPH & GM;
IR-36	IRRI, Philippines	Medium duration, irrigated, rain fed upland or lowland areas. semi dwarfs, grain long slender and white.
Swarna	Andhra Pradesh, India	Semi dwarf, grains: LS, resistant to blast, WBPH, moderately resistant to SB
IR-64	IRRI, Philippines	Semi dwarfs, mid/long duration
Safri -17	Chattisgarh, India	Tall, grains –LB, resistant to blast, BPH & BLB
Lalat	Orrisa, India	Semi dwarfs grains – LS, white, resistant to blast, Suitable for aerobic and boro conditions
Khitish	West Bengal, India	Dwarf, grains: LS, moderately resistant to blast & BS,
Naveen	Orissa, India	Medium duration
Suskhasmanat	Bihar, Jharkhand, India	Tall, grains: short bold, white, susceptible to blast, BLB and resistant to drought
Sambhamahsuri	Andhra Pradesh, India	Semi dwarf grains -MS, tolerant to SB,GM & RTV; resistant to blast
Samanta	Andhra Pradesh, India	Semi dwarf short duration lodging & disease resistance
Dandi	Gujrat, India	Semi dwarf, grains-MB, white, moderately resistant to Sh.B, BS,BPH,& BPH
Rajeshwari	Orissa, India	Semi dwarf, grains- MB, white,.. resistant to blast & BLB
Shravani	Jharkhand, India	Grains-LS, white; resistant to blast, Helminthosporium, tolerant to BLB, susceptible to RTV
Abhishek	Jharkhand, India	Medium duration
Udaya	Orissa, India	Semi dwarf grains: long bold, resistant to BPH, blast, root rot and tolerant to RTV;

RESULTS AND DISCUSSION

SSR analysis

In present study, 20 SSR primers distributed from chromosome 7 to 12 were used to estimate genetic diversity among 20 genotypes. A total of 65 alleles were detected among all genotypes. The number of alleles per locus varied from 2 to 7. The mean allele in current studies (3.25 alleles) was comparable with the results of Etemad et al. (2012) as 3.57 in rice. However, it was reported lower by

Prathepha et al. (2012) and Rahman et al. (2012) with an average of 11.85 and 4.18 alleles per locus, respectively.

The overall size of amplified products ranged from 100 (RM264) to 250 bp (RM286). In the current studies, 15 out of 20 SSR primer pairs generated polymorphic bands. PIC values for SSR ranged from 0.62 to 0.97 with mean value of 0.81 (Table 4). The highest PIC value was observed with primer RM-222 and RM-206 (0.97). However, it varied from 0.19 to 0.90 with an average of 0.75 as reported by Borba et al. (2009). About similar

average PIC values (0.78) was reported by Upadhyay et al. (2011). The cluster analysis showed a significant genetic variation among the genotypes with similarity coefficients ranging from 0.63 to 0.85 (Figure 1). The dendrogram revealed two distinct clusters at a similarity coefficient level of 0.63. Cluster II was the largest and included 19 genotypes while, clusters I comprised only one genotype, that is, Rajeshwari. The cluster II further divided into two sub clusters IIA and IIB. II A consisted of 17 genotypes while II B consisted of only two genotypes (Apo and Kalakeni). Apo and

Table 2. List of RAPD Primers used for the analysis of twenty rice genotypes.

Code	Primer sequences 5'-3'	Code	Primer sequences 5'-3'
OPA-04	AATCGGGCTG	OPA-16	AGCCAGCGAA
OPA-07	GAAACGGGTG	OPA-17	GACCGCTTGT
OPA-08	GTGACGTAGG	OPA-18	AGGTGACCGT
OPA-09	GGGTAACGCC	OPA-19	CAAACGTCGG
OPA-10	GTGATCGCAG	OPA-20	GTTGCGATCC
OPA-11	CAATCGCCGT	OPB-06	TGCTCTGCC
OPA-12	TCGGCGATAG	OPB-07	GGTGACGCAG
OPA-13	CAGCACCCAC	OPB-08	GTCCACACGG
OPB-11	GTAGACCCGT	OPB-17	AGGGAACGAG
OPB-13	TTCCCCGCT	OPB-18	CCACAGCAGT

Table 3. List of SSR Primers and their sequences used for the analysis of twenty rice genotypes.

Primer code	Primer sequences 5'-3'	
	Forward	Reverse
RM 21	ACAGT AT TCCGTAGACGG	GCTCCATGAGGG TGGTA AG
RM 536	TCTCTCCTCTTGTGGCTC	ACACACCAACACGACCACAC
RM 206	CCCATGCGTTTAACATTC T	CGTTCCATCGATCCGTATGG
RM 566	ACCCAACACTACGATCATCG	CTCCAGGAACACGCTCTTTC
RM 320	CAACGTGATCGAGGAGAT C	GGATTTGCTTACCACAGCTC
RM 247	TAGTGCCGATCGATGTA ACG	CATATGGTTTTGACAAAGCG
RM 544	TGTGAGCCTGAGCAAT AACG	GAAGCGTGTGATATCGCATG
RM 346	CGAGAGAGCCCATAACTACG	ACAAGACGACGAGGAGGGAC
RM 547	TAGGTTGGCAGACCTTTTCG	GTCAAGATCATTCTCGTAGCG
RM 519	AGAGAGCCCCTAAATTT CCG	AGGTACGCTCACCTGTGGAC
RM 286	GGCTTCATCTTTGGCGAC	CCGGATTCACGAGATAAACTC
RM 581	GAGCTGTTTTGGACTACGGC	GAGTAGCTTTCTCCCACCCC
RM 222	CTTAAATGGGCCACATGCG	CAAAGCTCCGGCCAAAAG
RM 47	ACTCCACTCCACTCCCCA	GTCAGCAGGTCGGACGTC
RM 10	TTGTCAAGAGGAGGCATCG	CAGAAATGGGAAATGGTCC
RM 19	CA AAAACAGAGCAGATGAC	CTCAAGATGGACGCCAAGA
RM 20	ATCTTGTCCTGCAGGTCAT	GAAACAGAGGCACATTTTCATTG
RMR	ACGAGCTCTCGATCAGCCTA	TCGGTCTCCATGTCCCAC
RM 167	GATCCAGCGTGAGGAACACGT	AGTCCGACCACAAGGTGCGTTGTC
RM 264	GTTGCGTCCTACTGCTACTTC	GATCCGTGTGATGATTAGC

Rajeshwari were distantly related at a similarity coefficient of 0.63. Cluster analysis showed that Satabdi and Ratna showed maximum similarity with the similarity coefficient of 0.85. Based on dendrogram, the genotypes Apo, Samantha and Rajeshwari belong to different clusters and are genetically diverse.

RAPD analysis

Using 20 RAPD primers, a total of 116 alleles were detected among the 20 rice genotypes. The number of

alleles per locus varied from 4 to 7. The average number of alleles per locus was 5.8. The overall size of amplified products ranged from 100 (OPA-10) to 1200 bp (OPA-11). All the RAPD primers used for analysis of genetic diversity and relationship generated polymorphic bands among the genotypes. The PIC value for the RAPD ranged from 0.33 to 0.88 with an average of 0.65 (Table 5). Similar results were also reported by Shiva et al. (2013) in rice. The present investigation revealed the effectiveness of RAPD in detecting polymorphism among different genotypes of rice. Nair et al. (2013) also reported

Table 4. Polymorphism Information Content (PIC) of SSR Loci across various genotypes.

Primer	Chromosomal location	Motifs	Molecular weight range (bp)	Total number of alleles	Number of Polymorphic alleles	Number of monomorphic alleles	% Polymorphism	Diversity in value of PIC
RM21	11	(GA)18	130-190	5	5	0	100.00	0.90
RM536	11	(CT)16	150-240	4	4	0	100.00	0.88
RM206	11	(CT)21	200-140	7	7	0	100.00	0.97
RM566	9	(AG)15	250-200	4	4	0	100.00	0.74
RM320	7	(AT)11GTAT(GT13)	250-150	6	6	0	100.00	0.94
RM247	12	(CT)16	250-210	4	4	0	100.00	0.93
RM544	8	(TC)9	180-130	2	2	0	100.00	0.68
RM346	7	(CTT)18	200-100	5	5	0	100.00	0.94
RM547	8	(ATT)19	100-280	5	5	0	100.00	0.90
RM519	12	(AAG)8	150	1	0	1	0.00	0.00
RM286	11	(GA)16	100-150	5	5	0	100.00	0.96
RM581	-	(GA)12	200-150	3	3	0	100.00	0.90
RM222	10	(CT)18	150-250	4	4	0	100.00	0.97
RM47	7	(AG)7(AG)11	200-250	6	6	0	100.00	0.96
RM10	7	(GA)15	160	1	0	1	100.00	0.00
RM19	12	(ATC)10	250	1	1	0	100.00	0.00
RM20	12	(ATT)14	-	0	0	0	0.00	0.00
RMR	-	-	-	0	0	0	0.00	0.00
RM167	11	(GA)16	-	0	0	0	0.00	0.00
RM264	8	(GA)27	100-150	2	2	0	100.00	0.62

that RAPD analysis in *O. sativa* accessions can be successfully used to estimate the genetic diversity in rice.

The highest PIC value (0.88) was observed for primers OPA-17. Cluster analysis revealed two major clusters at a cut-off similarity coefficient of 0.57 (Figure 2). Cluster I was largest and included 14 genotypes while cluster II consisted of six genotypes. Jaccard's coefficient of similarity revealed that 83.8% exist between genotypes

Shravani and Abhishek. Whereas, Apo keeps very low level of similarity with Rajeshwari at similarity coefficient of 0.57%. The major cluster I further divided into two sub clusters IA and IB. The sub cluster IA consist of 10 genotypes namely Khitish, Naveen, Udaya, Sukhasamat, Shravani, Abhishek, Sambhamasuri, Dandi, Samanta and Rajeshwari with the similarity coefficient ranged between 0.62 to 0.83. While sub cluster IB consist of 4 genotypes namely Swarna, IR-64, Safri and Lolat with

similarity coefficient from 0.67 to 0.78.

Combined analysis

The cluster dendrogram with combined analysis revealed two major clusters. Cluster I was the largest and included 14 genotypes, namely Swarna, IR64, Lolat, Safri, Suskhasamanat, Khitis, Naveen, Sambhamasuri, Dandi, Samanta, Rajaewari, Shravani, Abhishek and Udaya with the similarity




Figure 1. Dendrogram showing clustering of 20 genotypes of rice genotypes based on SSR data.




Figure 2. Dendrogram showing clustering of 20 genotypes of rice genotypes based on RAPD data.

Table 5. Polymorphism information content (PIC) of RAPD loci across various genotypes.

Primer	Annealing temperature	Molecular weight range (bp)	Total number of alleles	Number of polymorphic alleles	Number of Monomorphic alleles	% of polymorphism	PIC
OPA-04	30	350-800	5	5	0	100.00	0.64
OPA-07	30	200-700	7	7	0	100.00	0.68
OPA-08	30	200-1100	5	4	1	80.00	0.33
OPA-09	32	100-1000	5	4	1	80.00	0.46
OPA-10	30	100-800	6	6	0	100.00	0.64
OPA-11	30	100-1200	5	5	0	100.00	0.66
OPA-12	30	200-800	7	7	0	100.00	0.72
OPA-13	32	150-700	5	5	0	100.00	0.62
OPA-16	30	300-700	5	5	0	100.00	0.77
OPA-17	30	200-1000	6	6	0	100.00	0.88
OPA-18	30	250-600	5	5	0	100.00	0.82
OPA-19	30	400-1000	7	7	0	100.00	0.66
OPA-20	30	400-900	7	7	0	100.00	0.64
OPB-06	32	300-800	4	4	0	100.00	0.67
OPB-07	32	200-800	5	5	0	100.00	0.69
OPB-08	32	200-550	6	6	0	100.00	0.79
OPB-11	30	300-800	6	6	0	100.00	0.76
OPB-13	32	300-700	6	6	0	100.00	0.78
OPB-17	30	200-900	7	7	0	100.00	0.86
OPB-18	30	300-1000	7	7	0	100.00	0.59

coefficient that ranged between 0.65 to 0.82 while clusters II comprised six genotypes namely Apo, Kalakeni, Satabdi, Ratna, IR36 and Khandagini with the similarity coefficient ranging from 0.69 to 0.82 (Figure 3). Jaccard's coefficient of similarity revealed that high degree of similarity up to the extent of 82.1% existed between IR36 and Ratna. Dendrogram revealed that Apo was distantly located with Udaya with similarity coefficient of 0.62. Dendrogram revealed that genotypes Swarna, Satabdi, and Sukhasmanat, Samanta,

Rajeshwari, Udaya and Apo belong to different clusters and are genetically diverse. SSR markers showed higher PIC value compared to RAPD which indicates that SSR markers are highly informative and are reliable. In the current study, larger range of distinct values for genotypes revealed by microsatellite markers provides greater confidence for the assessments of genetic diversity and relationships, which can be used in future breeding programs. With the aid of microsatellite makers and clustering data, different distantly related rice

genotypes may be combined by intercrossing genotypes, for instance, aromatic rice genotypes with non-aromatic rice genotypes from different clusters to get hybrid varieties with highest heterosis (Sajib et al., 2012). Markers with PIC values of 0.5 or higher are highly informative for genetic diversity studies and can be successfully used to distinguish the polymorphism at a specific locus. The joint use of primer is excellent way of identification of genotypes. A combination of RAPD and SSR help to provide whole genome coverage and




Figure 3. Dendrogram showing clustering of 20 genotypes of rice based on combined (SSR and RAPD) data.




Figure 4. Agrose gel electrophoresis for PCR products of SSR marker (RM-222) of 1 - 20 rice genotypes (As described in Table 1). M=DNA ladder (100 bp).




Figure 5. Agarose gel electrophoresis for PCR products of RAPD marker (OPA-07) of 1 - 20 rice genotypes (as described in Table 1). M=DNA ladder (100 bp).

and reduce the error in estimation of genetic similarity based on any one marker system alone.

ACKNOWLEDGEMENTS

The technical help provided by Mrs. Rekha Satpuri, Technician and Mr. Ashwani Kumar Singh, Senior Research Fellow during the course of investigation is gratefully acknowledged. Financial help provided by SVP&T, Meerut is also gratefully acknowledged.

REFERENCES

- Anderson JA, Churchill GA, Autrique JE, Tanksley SD, Sorrells ME (1993). Optimizing parental selection for genetic linkage maps. *Genom.* 36:181-6.
- Botstein D, White R, Skolnick M, Davis R (1980). Construction of genetic linkage map in man using restriction fragment length polymorphism. *Am. J. Genet.* 32:314-331.
- Causse M, Fulton TM, Cho YG, Ahn SN, Chunwongee J, Wu J, Xiao ZY, Ronald PC (1994). Saturated molecular map of rice genome based on inter specific back cross. *Phyto. pathol.* 85:1251-1274.
- Chakravarthi KB, Rambabu N (2006). SSR marker based DNA fingerprinting and diversity study in rice (*Oryza sativa* L.). *Afr. J. Biotechnol.* 5:684-688.
- Cuevas RP, Fitzgerald MA (2012). Genetic Diversity of Rice Grain Quality and Nutrition Centre, *Int. Rice Res. Inst. Philippines* 15:285-310.
- Etemad A, Maziah M, Daud SK (2012). Determination of genetic relatedness among selected rice (*Oryza sativa*, L.) cultivars using microsatellite markers. *Afr. J. Biotechnol.* 11(28):7158-7165.
- Jin L, Lu Y, Xiao P, Sun M, Corke H, Bao J (2010). Genetic diversity and population structure of a diverse set of rice germplasm for association mapping. *Theor. Appl. Genet.* 121(3):475-487.
- Ma H, Yin Y, Guo ZF, Cheng LJ, Zhang L, Zhong M, Shao GJ (2011). Establishment of DNA fingerprinting of Liaojing series of japonica rice. *Middle-East J. Sci. Res.* 8(2):384-392.
- Matin S, Mohammad A, Islam MM, Sikdar SU, Zobayer N (2012). Molecular marker based (SSR) genetic diversity analysis in deep water rice germplasms of Bangladesh. *Int. J. Biosci.* 10(2):64-72
- McCouch SR, Kochert G, Yu ZH, Wang ZY, Khush GS, Coffman WR, Tanksley SD (1998). Molecular mapping of rice chromosomes. *Theor. Appl. Genet.* 76:815-829.
- Murray MG, Thompson WF (1980). Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Res.* 8:4321-4325.
- Nair GM, Nair AJ, Rajani J, Deepu V (2013). Molecular characterization of selected cultivars of rice, *Oryza sativa* L. using Random Amplified Polymorphic DNA (RAPD) markers. *Int. Food Res. J.* 20(2):919-923.
- Onaga G, Egdane J, Edema R, Abdelbagi I (2013). Morphological and Genetic Diversity Analysis of Rice Accessions (*Oryza sativa* L.) Differing in Iron Toxicity Tolerance. *J. Crop Sci. Biotech.* 16(1):53-62.
- Prabakaran A, Paramasivam K, Rajesh T, Rajarajan D (2010). Molecular characterization of rice land races using SSR markers. *Electron. J. Plant Breed.* 1(4):512-516.
- Prathepha P (2012). Genetic diversity and population structure of wild rice, *Oryza rufipogon* from Northeastern Thailand and Laos. *Australian J. Crop Sci.* 6(4):717-723.
- Rahman MM, Rasaul MG, Hossain MA, Iftekharuddaula KM, Hasegawa H (2012). Molecular Characterization and Genetic Diversity Analysis of Rice (*Oryza sativa* L.) Using SSR Markers. *J. Crop Improv.* 26(2):244-257.
- Rohlf FJ (2000). NTSYS-pc: Numerical taxonomy and multivariate analysis system. Version 2.1. Exeter Publications, New York, USA.
- Rohlf FJ (1993). Numerical taxonomy and multivariate analysis system NTSYS-PC version 1.80 Exeter software, New York.
- Sajib AM, Hossain MM, Mosnaz ATMJ, Hossain H, Islam MM, Md. Shamsher Ali MS, Proshan SH (2012). SSR marker-based molecular characterization and genetic diversity analysis of aromatic landraces of rice (*Oryza sativa* L.). *J. BioSci. Biotech.* 1(2):107-116.
- Siva R, Kumar K, Rajasekaran C (2013). Genetic diversity study of important Indian rice genotypes using biochemical and molecular markers. *Afr. J. Biotechnol.* 12(10):1004-1009.
- Williams K, Kubelik A, Livak K, Rafalski J, Tingey V (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acid Res.* 18:6531-6535.