



Trends in vegetation cover changes in Bonny area of the Niger Delta

AKURO ADOKI

Shell Petroleum Development Company
P.O. Box 263, Port Harcourt, Nigeria
Akuro.adoki@shall.com

ABSTRACT: The main vegetation type in the area is the mangrove forest, which occupies most of the Niger Delta. The other vegetation type is the secondary (re-growth) vegetation that occupies a small area. The secondary vegetation is surrounded by the mangrove swamp forest; a two-layered vegetation. The one-layered forest has mangrove trees of 3-14 m high without undergrowth forming the only layer while the two-layered forest has a top layer made up of *Rhizophora* species (3-32 m high) depending on the height of trees at each site and a ground layer composed mainly of the fern *Acrostichum aureum* and seedlings of the tree species. There is a dearth of epiphytic bryophytes and lichens on the boles and branches of the trees. From satellite imageries of the area, it is evident that the landcover classes changed across the three epochs. The water class covered an area of 111.91km² in 1986, 108.90km² in 1998 and 103.41km² in 2007. Mature forest (Forest I) occupies a total area of 85.64km² in 1986; decreased significantly to 59.68km² in 1998 and 59.30km² in 2007. This could be attributed to man-made and industrial activities. Secondary Forest (Forest II) covered 11.18km² in 1986, but increased to 43.49km² in 1998 and decreased to 23.30km² in 2007. Urban/Industrial/Sand class had a steady increase across the epochs. This class which covered about 10.37km² in 1986 increased to 14.73km² in 1998, and 25.28km² in 2007. This increment is attributed to influx of humans into the area as a result of increase in Oil and Gas industry activities. Mangrove class covered 87.02km² in 1986 but increased to 95.86km² in 1998 and 106.79km² in 2007. Stressed Vegetation occupied 16.38km² in 1986; this class type was only evident during that period. Sparse vegetation which covered an area of 5.23km² in 1986; decreased slightly to 5.07km² in 1998 but increased to 9.07 km² in 2007. Overall, the most dramatic change recorded is with respect to the areal extent of Urban/Industrial/Sand which increased by about 42% between 1986 and 1998 and about 144% between 1986 and 2007. The area covered by mangrove forest consistently declined over all the epochs while the reverse was the case with respect to the area covered by sparse vegetation. The area covered by stressed vegetation in 1986 disappeared during other epochs.

© JASEM

Many, many human actions tend to have indelible imprints on landscape in a short time (Briassoulis, 2000; Goldewijk and Ramankutty, 2004; Fabiyi, 2007). Throughout history, human activities have impacted on the natural ecosystem through the aggressive drive for development (Goldewijk and Ramankutty, 2004). It has been estimated that over the last three centuries, more than 1200 million ha of forests and wood lands have been cleared. Grassland and pastures have diminished by about 560 million ha and cropland areas have increased by about 1200 million ha (Richard and Flint, 1994). Human actions especially those involving biomass fuel consumption, land-use change, and agricultural activities have direct interaction with the land surface and negative consequences on vegetation and environmental qualities (Fabiyyi, 2011). These interactions are rather complex and have attracted research interest in the last four decades (Goldewijk and Ramankutty, 2004).

The Niger Delta region in Nigeria had its share of negative influence of human activities on the natural landscape. The primary vegetation of the delta is fast changing to secondary and derived vegetation due to aggressive incursion of human activities into the seemingly undisturbed ecosystem that characterized the region about a century ago. Human activities including oil exploration and urban development are causing imbalances in the ecosystems of the region with resultant negative consequences on environmental quality and livability (Fabiyyi, 2011). The extent of these environmental alterations has prompted different concerns including political agitations with respect to the social, economic and cultural consequences of the changes that are taking place. The oil multinational companies operating in the Niger Delta area have been fingered as the main change actors by especially the activist and environmentalists. The United Nations Development Programme reported that whereas the Niger Delta has an enormously rich natural

endowment in the form of land, water, forests and fauna, these assets, however, have been subjected to extreme degradation due to oil prospecting. For many people, this loss has been a direct route into poverty, as natural resources have traditionally been primary sources of sustenance" (UNDP, Niger Delta Human Development Report, 2006).

Several forces are responsible for the changes in vegetation quality observed in the Niger Delta. Settlement developments, oil prospecting over the years imprint indelibly on the Niger Delta fragile canvas of the ecosystem. The main visible impact is the change in land use and vegetal cover. Human systems is a part of natural ecosystems but the activities that support human enterprises unfortunately damage the natural landscape of the Niger Delta is in the coastal belt of Nigeria, it is characterized by fragile ecosystems including mangrove, nypa palm, fresh water swamp, sheltered tidal flat and large expanse of vegetated bluff. Niger Delta is home to different fauna and flora species (some of which are in endangered list of IUCN) yet with about 31 million populations in an area of about – square kilometers. The Niger Delta combined the presence of oil rich hydrocarbon deposit with the rich alluvium and the abundance of aquatic life to make the place attractive to rapid expansion. The intense urbanization and industrialization that followed the discovery of oil at Oloibiri in the present day Bayelsa state in 1956, have many consequences on the landscape of the Niger Delta, for example enlargement of natural coastal inlets and dredging of waterways for navigation, port facilities, and oil and gas pipelines have direct impact on the fragile coastal ecosystems. The visible manifestations of these anthropogenic activities include loss of biodiversity and essentially deforestation, environmental degradation, loss in vegetal qualities and soil nutrient loss. A number of large scale and cottage industries sprang up in the last thirty years thereby contributing to environmental degradation of the Niger Delta (Fabiya, 2011).

Other activities such as sand mining, hydrocarbon production like oil and gas, introduction of invasive species (nypa Palm) and engineering constructions such as jetty, seawalls and channelization are few among numerous activities taking place in the region. Apart from the various human induced changes in the ecosystems of the Niger Delta, the global climatic change is another significant change factor in the Delta. These drivers are responsible for long term modification of the coastal

ecosystem including the Niger Delta. Though deforestation and apparent change in vegetal qualities are major land use/ land cover changes occurring in many coastal regions of African countries. The impacts of human activities within the coastal region, and the climate change effects are difficult to separate into different compartments (Fabiya, 2011).

On a continental or global level, climatic fluctuations have been linked to the anthropogenic activities through the release of green house gas into the atmosphere, thus depleting the ozone layer. Effects of climate change have been measured in the continental or regional levels through sea level rise, melting of the icecap, increased rainfall and associated flooding in the low lying areas. In the sub local analyses context, the influence of human activities on the landscape can be directly measured through different approaches such as remote sensing and geographic information system (GIS) techniques (Fabiya, 2011). Recent research discourse focuses on the ways to identify the extent of human influence on the global climate change from the normal perturbation associated with climatic cycle and to what extent will the global warming be reduced with the cutting down of carbon emission by industrialized nations. In the supra local context, it is necessary to separate the impact of the immediate anthropogenic actors on the ecosystems from the changes due to climate changes.

Several methods have been used to measure vegetation cover loss (DeFries et al., 1995). The methods require the definition of the thresholds and classification of the vegetation around the threshold based on some pre-determined schemes. This approach has been fraught with practical inefficiency, especially if there is a need to monitor changes over time. Other methods include the following authors' approaches (Copeland et al., 1996; Bonan, 1999; Houghton, 1999; Postel et al., 1996; Vitousek et al., 1997). Normalized difference of the vegetation index (NDVI) became popular in the last three decades to investigating the quality of vegetal cover. The normalized difference of the vegetation index (NDVI) is a non-linear transformation of the visible (red) and near-infrared bands of satellite information. It is an alternative measure of vegetation amount and condition. It is associated with vegetation canopy characteristics such as biomass, leaf area index and percentage of vegetation cover. NDVI is mathematically defined as:
$$\text{NDVI} = \frac{\text{nir-red} - \text{red}}{\text{nir-red} + \text{red}}$$

red band) Previous studies have used Channels 1 (0.54 to 0.68 m) and 2 (0.73 to 1.10 mm) which are visible and near infrared of the advanced very high resolution radiometer (AVHRR) data (Groten, 1993; Loveland et al, 1991) other works on the use of NDVI to monitor vegetal changes include. Other studies linked NDVI to plant phenology (Defries et al., 1995; Read and Lam, 2002; Mora and Iverson, 1995). Apart from AVHRR NDVI have been calculated from LANDSAT-TM information using bands 3 (0.63 to 0.69 mm) and 4 (0.76 to 0.90 mm). NDVI values range from -1 to +1 for pixel values ranging between 0 to 255. NDVI, which is one of the most successful of many attempts to simply and quickly identify vegetated areas and their "condition" is utilized in this study.

MATERIALS AND METHODS

This project examines the use of Remote Sensing and Geographic information Systems (GIS) in mapping and analyzing Land use/land cover changes within the Bonny area from 1986-2007. Land use /Land cover changes (LULC), which is recognized, as one of the most sensitive indicators of environmental change, reflects the impacts of human activities on the biophysical environment. The impact of activities relating to oil and gas exploration may generate some significant effects on the environment, such as loss of wildlife habitat, changes in surface and subsoil hydrology that may lead to accelerated soil erosion and land degradation, vegetation changes, air pollution and changes in coastline geomorphology.

The area of interest covers an area (see Figure 1) of approximately 322.04km² and it has bounding coordinates of 517899.355E- 540285.491E and 41432.386N-55817.979N in Transverse Mercator Nigerian Mid-belt projection and Minna datum The datasets used for this project were free of negative atmospheric influences such as cloud cover and atmospheric haze.

A number of processing algorithms were used to process the Natural Colour Composite Image, Land Cover Classification and Normalized Difference Vegetation Index. A short description is given of each of the processing method used and interpretation included. Further processing was carried out to delineate areas of vegetative growth and land cover type in the area of interest. This study is expected to reveal or depict the environmental performance of oil and gas facilities over the period in view through land cover change detection analysis.

The study is limited to baseline information generation from field data collection and remote sensing and GIS, based on change analysis of areas lying within the Bonny area. Spatial and satellite data were sourced from the Shell Petroleum Development Company (SPDC).

Landsat TM dataset of 1986, SPOT XI dataset of 1998 and Landsat ETM of 2007 datasets were used. These satellite images were processed and the different landcover types obtained for analysis.

RESULTS AND DISCUSION

Land cover Classification: A Land cover classification was carried out over the Bonny area only, for three epochs using the Landsat TM 1986, SPOT XI 1998, and Land sat ETM of 2007. An unsupervised classification process, which uses an iterative ISO class algorithm to generate a number of classes pre-defined by the processor, was used. This unsupervised classification technique, groups similar pixels based on the spectral characteristics of the image datasets. This method does not require any prior knowledge of the land cover types in the area of interest. The following parameters were used for the unsupervised classification process; 98% unchanged, maximum standard deviation 4.5, minimum distance between class means 3.0. The iterative ISO class processes were carried out on the Landsat TM, SPOT XI, and Land sat ETM datasets. Six main classes were identified using an unsupervised classification technique.

Aerial Environmental Baseline Study: Baseline information was derived from the processing of satellite images and GIS datasets. There is presence of fourteen and nine Oil and Gas Pipelines respectively while there are ten manifolds within the area of interest. Thirty-six wells are found with the area of interest and these are mainly from Bonny, Akikigha, Asaramatoru and Owuanga. Only one Flowstation is found within this area of interest and this is Bonny. Also, within this area could be found the following facilities SHELL Bonny terminal, Nigeria Liquefied Natural Gas (NLNG) and Mobil Terminal.

Land-use : The land use pattern that predominates the area of interest is shown in Figure 2. The Soku-Awoba-Bonny area measures approximately 121.5 x 64 km covering an area of 7782 km². Water and mangroves occupy 45% and 33% of the total area respectively. Undisturbed forest (Forest I) is present as freshwater swamp forest on the northern fringes of the mangrove swamps, riparian forest along

major river channels and on palaeo-beaches along the coastline. Undisturbed forest occupies just over 8 % of the area. Human impact on these natural cover **types result in secondary or disturbed forest** (Forest II) covering 3% and Farmland 5%. Urban and industrial areas occupy just about 1% of the total area.

Landcover types within the pipeline corridor are dominated by mangrove at 75% of the total length. Rivers, creeks and wet mangrove constitute a further 20%. Urban-industrial areas traversed amount to 2.3% of the total length – this is mostly at the south-eastern end at Bonny Terminal but also includes oilfield industrial areas at Ekulama and Cawthorne Channel. The total area of pipeline corridor (given a 1 km width) is 91.00 km² or 1.7% of the total project area (7782 km²).

Vegetation: The main vegetation type in the area is the mangrove forest, which occupies most of the Niger Delta. The other vegetation type is the secondary (re-growth) vegetation that occupies a small area. The secondary vegetation is surrounded by the mangrove swamp forest; a two-layered vegetation. The one-layered forest has mangrove trees of 3-14 m high without undergrowth forming the only layer while the two-layered forest has a top layer made up of *Rhizophora* species (3-32 m high) depending on the height of trees at each site and a ground layer composed mainly of the fern *Acrostichum aureum* and seedlings of the tree species. There is a dearth of epiphytic bryophytes and lichens on the boles and branches of the trees. The vegetation species composition within the area is shown in Table 1.

The detailed tree density per hectare is presented in Table 2. Various plant species make up the vegetation in this area. These include trees, shrubs, fern species, grasses and sedge. The indicator species of the vegetation types are *Rhizophora racemosa*, *Rhizophora harrisonii*, (for the mangrove swamp forest) *Alchornea cordifolia* and *Cocos nucifera* (for the secondary growth forest).

Farmlands with crops were also encountered among the vegetation. The common crops recorded include Cassava (*Manihot esculenta*), Coconut (*Cocos nucifera*), banana (*Musa* sp.), Cocoyam (*Colocasia esculenta*), groundnut (*Arachis hypogea*), Maize (*Zea mays*), Plantain (*Musa sapientum*), Pineapples (*Ananas comosus*), Pepper (*Capsicum* sp.), Okra (*Hibiscus esculentum*), Yam (*Dioscorea* sp.) and Water yam (*Dioscorea alata*).

Mangrove: It is found to be generally concentrated within the northern area, though spreads

Forest II It occurs towards the eastern and western fringes. Within it is found, pockets of forest I.

Water: This comprises mainly the Bonny river flowing south wards in to the ocean and the Andoni River to the north and east of the image. There are quite a few creeks within the area.

Forest I: This type of vegetation is found mainly within the southern area spreading towards the east. The dominant vegetation is mature forest.

This is secondary forest which replaces the primary forest and is in the process of achieving maturity in the successional cycle.. This vegetation type is found in close proximity with the Urban/Industrial and Sparse Vegetation. Small portions of it are found within the mangrove area.

Urban/Industrial/Sand: This is found around the shoreline as sand and also built up areas. Within the areas is a good network of roads connecting the communities.

Sparse Vegetation: This cover the smallest portion of the area and it is found in close proximity with urban/industrial/sand and forest II.

Stressed Vegetation: This type of vegetation appeared near the mangrove in the eastern flange of the area of interest and as pockets around the northern axis. This is type of vegetation is only noticed in the Landsat TM of 1986.

The image representations could be seen in figures 3-5.

Normalized Differential Vegetation Index (NDVI): NDVI was used in highlighting areas showing the different levels of vegetation health. This objective was achieved by applying the vegetation formula to the image. A normalized difference vegetation index is a ratio of Near Infrared (NIR) and Red bands. The formula is presented below

$$DN \text{ (out)} = (\text{band 1} - \text{band 4}) / (\text{band 1} + \text{band 4})$$

An NDVI lookup table was applied to the image. It was observed that

The areas with high NDVI values for the entire epochs are Mangrove, Matured forest (forest I), Secondary forests (forest II), Sparse Vegetation. These are shown in green, yellow and orange colours respectively.

The water, Stressed vegetation and Urban/Industrial are in blue and purple showing low NDVI.

Time-Lapse Change Analysis: Time-lapse analysis is for assessing changes in an area within a period of time. For the case of this study, it was carried out on Bonny Area using the Landsat ETM 2007, SPOT 4 of 1998 and Landsat TM of 1986. This was carried out on the landuse/landcover images (Figures 3 - 15). A breakdown of the various landcover types as a percentage of totals is shown in Tables 6 and 7.

Conclusions: The landcover change detection summary over the area of study between 1986 and 2007 is as follows:

The Time Lapse Analysis revealed that Water class decreased slightly over the years. Water class changed from 111.91km² in 1986 to 108.90km² in 1998, and 103.41 km² in 2007. Since the satellite images were acquired about the same period being dry season, the reason for the decline might be due to the increase in sand deposit along the shoreline.

There was a sharp decline in Mature forest (Forest I) from 85.64km² in 1986, to 59.86km² in 1998. By 2007, it decreased further to 59.30km². The decrease could be attributed to the construction work between 1986 and 1998 which was probably at its peak but reduced by the year 2007. The construction of Mobil facility is one of such activities.

Mangrove which was 87.02 km² in 1986 increased to 95.86 km² in 1998 and 106.79 km² in 2007. The increase might be due to the decline in the area covered by water. In the event of flooding, mangrove area covered by water will be shown as water body on the satellite image. When the water recedes, the vegetation signature can then be mapped from the image.

Stressed Vegetation was only seen in the year 1986 and it had a total coverage of 16.38km². The reason for this is due to the damage caused

to the vegetation by flooding. This is part of the mangrove vegetation. It was separated from mangrove due to the classification done.

There was a sharp increase in secondary forest (Forest II) from 11.18km² in 1986, to 43.49km² in 1998. By 2007, it decreased to 23.30km². The increase between 1986 and 1998 was due to the re-growth that occurred where construction activities did not take place. The decrease between 1998 and 2007 should be due to the forest II being destroyed to make way for activities to go on.

There was a slight decrease in sparse vegetation from 5.23km² in 1986 to 5.07km² in 1998. Between 1998 and 2007, sparse vegetation increased to 9.67km². The decrease between 1986 and 1998 was due to regrowth vegetation while from 1998 to 2007, the increase occurred as a result of the decline in forest II due to construction and human activities.

Urban/Industrial/Sand class increased across the epochs. Urban/Industrial/Sand was 10.37km² in 1986, increased to 14.73km² in 1998, and 25.28km² in 2007. This was due to development and construction activities which brought about the influx of humans into the area.

Overall, the most dramatic change recorded is with respect to the areal extent of Urban/Industrial/Sand which increased by about 42% between 1986 and 1998 and about 144% between 1986 and 2007. The area covered by mangrove forest consistently declined over all the epochs while the reverse was the case with respect to the area covered by sparse vegetation.

REFERENCES

- Bonan GB (1999). Frost followed the plow: impacts of deforestation on the climate of United States. *Ecol. Appl.*, 9(4): 1305-1315.
- Briassoulis H (2000). „Analysis of Land use change: Theoretical and modeling approaches“. The Web book of regional Science. Regional Research Institute, West Virginia University.
- Copeland JH, Pielke RA, Kittel TGF (1996). „Potential climate impacts of vegetation change: A regional modeling study“. *J. Geophys. Res.*, 101: 7409-7418.
- DeFries R, Hansen M, Townshend J (1995). „Global discrimination of land cover types from metrics derived from AVHRR

- Pathfinder data". *Remote Sens. Environ.*, 54: 209-222.
- Fabiyi OO (2007). "Analysis of change- agents in urban land use transition; example from Ibadan city, Nigeria". *J. Environ. Cult.*, 4(2): 23-43.
- Fabiyi, o (2011) Change actors' analysis and vegetation loss from remote sensing data in parts of the Niger Delta region. *J. Ecol. Nat. Environ.*, 3(12), 381-391.
- Fabiyi OO (2011). "Analysis of urban decay from low resolution satellite remote sensing data; Example from organic city in Nigeria". *Int. J. Manage. Dev. Rev.*, 3: 1.
- Goldewijk KK, Ramankutty N (2004). Land cover change over the last three centuries due to human activities: The availability of the new global data sets. *GeoJournal*, 61: 335-344.
- Groten SME (1993). "NDVI-crop monitoring and early yield assessment of Burkina Faso. *Int. J. Remote Sens.*, 14: 1495.
- Houghton RA (1999). The annual net flux of carbon of the atmosphere from changes in land use 1850-1990. *Tellus Ser.*, 51B: 298-313.
- Jensen JR (2004). "Introductory digital image processing: a remote sensing perspective". Prentice Hall, Upper Saddle River. NJ.
- Lu D, Mausel P, Brondizio E, Moran E (2004). Change detection techniques. *Int. J. Remote Sens.*, 25: 2365-2401.
- Millward AA (2011). "Urbanization viewed through a geo-statistical lens applied to remote sensing data" *Area*, 43(1): 53-66.
- Postel SL, Daily GC, Ehrlich PR (1996). Human appropriation of renewable water. *Science*, 271: 758-788.
- Read JM, Lam NS (2002). Spatial methods for characterizing land cover and detecting land cover changes for the tropics. *Int. J. Remote Sens.*, 23: 2457-2474.
- Richard JF, Flint E (1994). "Historic land use and carbon estimates for south and Southeast Asia 1880-1980. Data set NDP 046 CDIA/ORNL. Carbon Dioxide Information Analysis centre". Oak Ridge Natl.Lab., Oak Ridge, Tenn.
- Song C, Woodcock CE, Seto KC, Lenney MP, Macomber SA (2001). Classification and change detection using Landsat TM data. *Remote Sens. Environ.*, 75: 230-244.
- Vitousek PM, Mooney HA, Lubcheno J, Mellilo JM (1997). "Human domination of the Earth ecosystem". *Science*, 277: 494-499.

Trends in vegetation 95

Species	Common Names	Life Form Types	Uses/Economic Importance	Ecological Significance
<i>Acrostichum aureum</i>	Salt water fern	Palm	For making traps to catch fish	Could serve as indicator for deteriorating conditions for growth or <i>Rhizophora racemosa</i> and the rising level of land
<i>Afromomum septum</i>	-	Herb	Medicinal	-
<i>Alchornea cordifolia</i>	Christmas bush	Shrub	Could be used medicinally	Abundant in cleared and disturbed areas
<i>Alstonia booneri</i>	Pattern wood	Woody tree	For carving canoes, furniture Sold as timber	Wetland species
<i>Anacardium occidentale</i>	Cashew apple	Woody tree	Medicinal Fruits edible	Grows in cleared and disturbed areas
<i>Ancistrophyllum secundifolia</i>	Climbing palm	Palm	Stem used as binder in hut construction	Wetland species
<i>Anthocleista vogellii</i>	Gabbage tree	Woody tree	Leaves used for wrapping coconut Root is medicinal	Wetland species
<i>Anthostema aubryanum</i>	Sapling wood	Woody tree	Sold as timber tree For carving canoes and carpentry work	Wetland species
<i>Antidesma laciniatum</i>	-	Woody tree	-	-
<i>Anthonotha macrophylla</i>	-	Woody tree	-	-
<i>Avicenia germanis</i>	White Mangrove	Woody tree	Training/Dye from bark of the tree Leaves, seeds are as curative ointment for ulcers or sores	Trees provide nesting sites for sea and shoreline birds and diverse animal life
<i>Axonopus compressus</i>	Carpet grass	Grass	For making lawns	Species of fallow lands
<i>Borreria ocyroides</i>	Fula's lesser necklace	Herb	Medicinal	Species of fallow lands
<i>Bridelia micranta</i>	-	Woody tree	-	-
<i>Calopogonium mucunoides</i>	Mucuna (climber)	Climber	For Pasture or mulching	Grows in cleared and disturbed areas
<i>Carapa procera</i>	Crab wood	Woody tree	Gum resin Medicinal	Wetland species
<i>Chromolaena odoratum</i>	Awolowo weed	Herb	Medicinal	Species of fallow lands, cleared and disturbed areas
<i>Chrysobalanus orbiculare</i>	Plum of the West Indies	Woody tree	Medicinal	-
<i>Cleistopholis paten</i>	Canoe wood	Woody tree	For canoe making Medicinal Cut and sold as timber	Wetland species
<i>Cocos nucifera</i>	Coconut Palm	Woody tree	Fruits edible/extraction of oil Fibres processed for weaving Coconut fronds for thatching etc.	-
<i>Costus afar</i>	-	Herb	-	Wetland
<i>Crotolaria retusa</i>	-	-	-	-
<i>Cyrtosperma senegalensis</i>	Swamp aerum	Under shrub	Medicinal	Wetland species Grows in marshy lands
<i>Dichospetatum madagascariense</i>	-	Woody tree	-	-
<i>Dalbergia ecastaphyllum</i>	Indian timber	Woody tree	For carving e.g. walking sticks	-
<i>Dictyandra involubrates</i>	Timber	Woody tree	-	-
<i>Diadia serrulata</i>	Star of the ground	Herb	Medicinal	Species of fallow lands
<i>Daniella</i> sp.	-	-	-	-
<i>Dissotis rotundifolia</i>	Rockrose or sheep-eyes	Herb	Medicinal	Species of fallow lands
<i>Drepanocarpus lunatus</i>	-	Tree	-	-
<i>Dracaenia arborea</i>	Mottled leaf	Tree sp.	Ornamental	-
<i>Elaeis guineensis</i>	Oil palm tree	Tree	Source of red palm oil Fronds used for weaving baskets and for thatched houses	-
<i>Emelia praetermissa</i>	Bush bulk's ear	Herb	Medicinal and eating as salad	Species of fallow lands
<i>Erafrotis tenella</i>	Stink grass	Grass	Fodder for cattle	Species of fallow lands
<i>Ficus capensis</i>	Fig tree (sand paper tree)	Woody tree	Plants have medicinal value	-
<i>Fimbristylis</i> sp.	Sedge	Sedge	Used for making mats	-

Trends in vegetation 96

			Medicinal	
<i>Fuirena ciliaris</i>	Sedge	Sedge	Burnt to make salt	Wetland species
<i>Gilbertiodetifron brachystegoides</i>	-	-	-	-
<i>Hibiscus suratensis</i>	Wild sour	Herb	Medicinal	Species of fallow lands
<i>Irvingia gabonensis</i>	Bush mango	-	Fruit edible	-
<i>Indigofera</i> sp.	Indigo sp.	Under shrub	Medicinal	-
<i>Killingia robusta</i>	Sedge	Sedge	Used for fumigant Medicinal	-
<i>Klainedoxa gabonensis</i>	-	-	-	-
<i>Laguncularia racemosa</i>	Black Mangrove	Woody trees	Mangrove wood is a source of fuel, timber, electric poles and scaffolding materials in building construction	Stilt roots bind the tidal mudflats together, hence ensuring the stabilization of coasts and protection of shoreline from erosional processes
<i>Lophora lanceolata</i>	-	Woody tree	-	-
<i>Lycopodium cernuum</i>	-	Woody tree	-	-
<i>Mariscus ligularis</i>	Sedge	Sedge	-	Abundant in cleared and disturbed areas
<i>Mangifera indica</i>	Mango tree	Woody tree	Leaf – medicinal Fruit edible	-
<i>Mitragyna ciliata</i>	Abura (Timber)	Woody tree	Construction of light canoes Making of household utensils (e.g. furniture) Cut and sold for timber	Wetland species
<i>Milletia arboemis</i>	-	-	-	-
<i>Musanga cecropiodes</i>	Umbrella tree (corkwood)	Woody tree	Soft wood tree for household utensils and sold as timbers Cut and used for buoyancy	Wetland species
<i>Nauclea diderrichii</i>	Opepe (sapling tree)	Woody tree	Valuable commercial/industrial wood Transmission poles/piling timber	-
<i>Nephrolepis biserrata</i>	Climbing fern	Fern	-	Wetland species
<i>Nypa fruticans</i>	Salt water coconut (Nipa palm)	Palm	Palm for making thatch houses/basket weaving	Stabilization of coasts and protection of shoreline from erosional processes
<i>Newbouldia laevis</i>	-	-	-	Possess rapid colonizing potentials and can displace mangrove.
<i>Palisota</i> sp.	Break hoe or knee cap	Undershrub	Medicinal	Species of fallow lands
<i>Pandanus candelebrum</i>	Panadana luster (ornamental screw pine)	Palm	Leaves used for household uses	Wetland species
<i>Panicum</i> sp.	Elephant grass/guinea grass	Grass	For fodder and roofing thatch houses	Species of fallow lands
<i>Paspalum vaginatum</i>	Salt water sedge	Grass	Used as fodder	Grows in marshy areas Can tolerate high degree of salinity and oil contamination
<i>Phoenix reclinata</i>	-	-	-	-
<i>Pennisetum hordeoides</i>	Shura grass	Grass	Used as fodder grass	Grows in cleared and disturbed areas
<i>Pentacletra macrophylla</i>	-	Woody tree	Timber tree	Wetland species
<i>Psychotria vogeliana</i>	Woodcock	Woody tree	Medicinal	-
<i>Raffia hookeri</i>	Raffia palm	Woody tree	Tree tapped for its wine Fronds used as rooting sheets and woven into baskets Fruits used as bait for animals	Wetland species
<i>Rubiaceae</i>	-	-	-	-
<i>Rhizophora harrisonii</i>	Red Mangrove	Woody trees	Mangrove wood is a source of fuel, timber, electric poles and scaffolding materials in building construction	Tidal mudflats, stilt roots/rootlets serve as critical nursery for fishes, prawn and habitat for crabs and molluses
<i>Rhizophora racemosa</i>	Red Mangrove	Woody trees	Mangrove wood is a source of fuel, timber, electric poles and scaffolding materials in building construction	Tidal mudflats, stilt roots/rootlets serve as critical nursery for fishes, prawn and habitat for crabs and molluses

Trends in vegetation 97

<i>Selaginella myosurus</i>	Club plant	Twining climber	Medicinal	Wetland species
<i>Setaria</i> sp.	Foxtail grass	Grass	For pasture	-
<i>Sida acuta</i>	Hornbeam leaf	Herb	Medicinal	Species of fallow lands
<i>Sporobolus pyramidalis</i>	Rat's tail grass	Grass	For fodder	Species of fallow lands
<i>Symphonia globulifera</i>	Hog Gum tree	Woody tree	For making canoes Household utensils Sold as timber/planks	Wetland species
<i>Syzygium guineensis</i>	Berry of Kizza	Shrub	Medicinal	Species of fallow lands
<i>Tridax procumbens</i>	Snail's salia	Herb	Medicinal	Species of fallow lands
<i>Vernonia cinera</i>	-	Herb	Medicinal	-
<i>Zanthoxylum gillettii</i>	-	-	-	-

Table 1: Species Composition of vegetation in the Soku/Awoba-Bonny Area

Key: - not known

Table 2: Tree Density per hectare within the Soku/Awoba-Bonny Area.

Plant species	Mean Density/Hectare
<i>Achrosticum aureum</i>	850
<i>Alchornea cordifolia</i>	600
<i>Anthocleista vogelli</i>	450
<i>Anthonotha macrophylla</i>	300
<i>Antidesma laciniatum</i>	100
<i>Avicennia germinans</i>	500
<i>Bridelia micrantha</i>	200
<i>Cocos nucifera</i>	330
<i>Dalbergia ecastaphyllum</i>	300
<i>Daniella</i> sp	200
<i>Dichopetalum madagascariense</i>	200
<i>Dictyandra involucreta</i>	600
<i>Drepanocarpus lunatus</i>	300
<i>Elaeis guineensis</i>	400
<i>Ficus capensis</i>	600
<i>Gilbertiodetifron brachystegoides</i>	160
<i>Irvingia gabonensis</i>	100
<i>Klainedoxa gabonensis</i>	100
<i>Laguncularia racemosa</i>	530
<i>Mangifera indica</i>	500
<i>Milletia arboensis</i>	100
<i>Musanga cecropioides</i>	100
<i>Newbouldia laevis</i>	100
<i>Nypa fruticans</i>	550
<i>Pandanus candelabrum</i>	100
<i>Phoenix reclinata</i>	1300
<i>Rhizophora harrisonii</i>	1386
<i>Rhizophora racemosa</i>	1389
<i>Zanthoxylum gillettii</i>	200

Table 3: Landcover summary for 2007

Landcover Types	sq km	%
Water	103.41	31.55
Mangrove	106.79	32.58
Forest I	59.30	18.09
Forest II	23.30	7.11
Sparse Vegetation	9.67	2.95
Urban/Industrial/Sand	25.28	7.71
Total	327.74	100

Table 4: Landcover summary for 1998

Landcover Types	sq km	%
Water	108.90	33.23
Mangrove	95.86	29.25
Forest I	59.68	18.21
Forest II	43.49	13.27
Sparse Vegetation	5.07	1.55
Urban/Industrial/Sand	14.73	4.50
Total	327.74	100

Table 5: Landcover summary for 1986

Landcover Types	sq km	%
Water	111.91	34.15
Mangrove	87.02	26.55
Forest I	85.64	26.13
Forest II	11.18	3.41
Sparse Vegetation	5.23	1.60
Stressed Vegetation	16.38	5.00
Urban/Industrial/Sand	10.37	3.16
Total	327.74	100

Table 6: Land cover summary for the three Epochs

Landcover Types	1986 (%)	1998 (%)	2007 (%)	1986 (sq km)	1998 (sq km)	2007 (sq km)
Water	34.15	33.23	31.55	111.91	108.90	103.41
Mangrove	26.55	29.25	32.58	87.02	95.86	106.79
Forest I	26.13	18.21	18.09	85.64	59.68	59.30
Forest II	3.41	13.27	7.11	11.18	43.49	23.30
Sparse Vegetation	1.60	1.55	2.95	5.23	5.07	9.67
Stressed Vegetation	5.00	0.00	0.00	16.38	0.00	0.00
Urban/Industrial/Sand	3.16	4.50	7.71	10.37	14.73	25.28





Fig 1: Satellite image of Niger delta major towns and study area






Fig 3: Landcover map of Bonny Area derived from Landsat ETM 2007




Fig 4: Landcover map of Bonny Area derived from SPOT 4 1998




Fig 5: Landcover map of Bonny Area derived from Landsat TM 1986




Fig 6: Pictorial representation of the landcover types in 2007 Landcover types

Trends in vegetation 100




Fig. 7: Pictorial representation of the Land cover types in 1998




Fig. 8: Pictorial representation of the Land cover types in 1986




Fig 9: Normalized Difference Vegetation Index Map of Landsat ETM for 2007




Fig 10: Normalized Difference Vegetation Index Map of SPOT 4 for 1998




Fig 11: Normalized Difference Vegetation Index Map of Landsat TM for 1986




Fig 13: Landsat ETM of 2007













Fig 14: SPOT 4 of 1998












Fig 15: Landsat TM of 1986

Trends in vegetation 102

Table 7: Land use and Land cover Profile changes

Changes	Date 1 1986	Date 2 1998	Date 3 2007
Change 1	 <p data-bbox="352 574 758 602">Absence of LNG Staff Quarters near Finima</p>	 <p data-bbox="842 574 1142 602">LNG Staff Quarters in existence</p>	 <p data-bbox="1339 574 1692 602">Completion of the LNG Staff Quarters</p>
Change 2	 <p data-bbox="338 878 743 906">Construction at the LNG site had just begun</p>	 <p data-bbox="856 873 1199 901">Construction reaches advanced stage</p>	 <p data-bbox="1346 889 1751 938">Construction completed and activities in full throttle</p>
Change 3	 <p data-bbox="386 1192 667 1240">Presence of appreciable Sand deposition along the Shoreline</p>	 <p data-bbox="848 1192 1276 1219">Presence of Spit at the entrance of Bonny river</p>	 <p data-bbox="1352 1192 1696 1240">Spit disappears from the entrance of Bonny river due to erosion</p>

Trends in vegetation 103

<p>Change 4</p>	 <p>Absence of Oil facility near Finima</p>	 <p>Construction of the Oil Facility site reaches advanced stage</p>	 <p>Construction probably completed</p>
<p>Change 5</p>	 <p>Presence of Sand deposit</p>	 <p>Presence of Settlement</p>	 <p>Settlement grows due human influx</p>
<p>Change 6</p>	 <p>Stressed Vegetation within the Mangrove vegetation</p>	 <p>Absence of Stressed Vegetation within the Mangrove vegetation</p>	 <p>Absence of Stressed Vegetation within the Mangrove vegetation</p>