

Bull. Chem. Soc. Ethiop. 2021, 35(1), 33-42.
 © 2021 Chemical Society of Ethiopia and The Authors
 DOI: <https://dx.doi.org/10.4314/bcse.v35i1.3>

ISSN 1011-3924
 Printed in Ethiopia
 Online ISSN 1726-801X

SYNTHESIS, CHARACTERIZATION AND ANTIMICROBIAL ACTIVITY OF COPPER(II) SCHIFF BASE ADDUCTS OF SOME *p*-SUBSTITUTED ANILINE SCHIFF BASES

Abdullahi Owolabi Sobola^{1,2*}, Gareth Mostyn Watkins², Rafiu Olanrewaju Shaibu³, Sheriff Adewuyi⁴ and Saliu Alao Amolegbe⁴

¹Department of Chemistry, Lagos State University, P.O. Box 0001, Ojo, Nigeria

²Department of Chemistry, Rhodes University, Grahamstown 6140, South Africa

³Department of Chemistry, University of Lagos, Akoka, Nigeria

⁴Department of Chemistry, College of Physical Sciences, Federal University of Agriculture, Abeokuta, P.M.B 2240, Nigeria

(Received March 31, 2019; Revised January 1, 2021; Accepted January 20, 2021)

ABSTRACT. The synthesis, characterization and antimicrobial activity of Cu(II) complexes of some *p*-substituted aniline Schiff base ligands have been carried out. The Schiff bases were obtained from salicylaldehyde and *o*-vanillin. The Cu(II) complexes have been characterized by elemental analysis, conductivity measurement, infrared and electronic spectral data. The complexes were obtained either as metal chelates [Cu(L)₂] or Schiff base adducts (CuCl₂·2LH).xH₂O. The metal chelates were non-electrolytes while the Schiff base adducts exhibited 1:1 or 2:1 electrolytes in methanol. The Cu(II) complexes exhibited slight antimicrobial activity against *Escherichia coli* ATCC® 8739TM*, *Staphylococcus aureus subsp. aureus* ATCC® 6538TM*, *Bacillus subtilis subsp. spizizenii* ATCC® 6633TM* and *Candida albicans* ATCC® 2091TM*. The complexes exhibited significant antifungal activity.

KEY WORDS: Metal Chelates, Schiff bases, Adducts, Cu(II) complexes, Salicylaldimines

INTRODUCTION

Metal complexes have received overwhelming attention in main group and transition metal chemistry due to the ease of their preparation and structural variability. Schiff base metal complexes exist as metal chelates or as Schiff base adducts. Schiff base ligands especially salicylaldimines have been extensively reported to form metal chelates readily with metal ions [1-9]. In most cases, the *ortho*-hydroxyl Schiff base ligands coordinate as bidentate monobasic [5-8] via the imine nitrogen and the deprotonated phenolic oxygen, Figure 1.




Figure 1. Schiff base metal chelate.

On the other hand, reaction of salicylaldimine (HSal) type ligands with metal salts could as well result in the formation of Schiff base adducts [10-14] containing the intact ligand moieties,

*Corresponding author. E-mail: abdullahi.sobola@lasu.edu.ng

This work is licensed under the Creative Commons Attribution 4.0 International License

MA_n.mLH. This is an uncommon occurrence and has been scarcely reported. The ability of the HSal ligand to retain the acidic hydrogen atoms when reacting with Lewis acids to form Schiff base adducts depends upon structural peculiarities, the nature of the coordinate bond and reaction conditions [14]. Specifically, compounds exhibiting strong intra-molecular hydrogen bonding, such as salicylaldimine type ligands, have high tendency to form adducts with Lewis acids. Also, most reported cases [10-14] of Schiff base adducts involve the use of metal chloride (MCl₂.xH₂O) salt. Bamfield [12] attributed the formation of Schiff base adducts to the interaction between the metal chloride salt and the keto-tautomer of the Schiff base ligands. A variety of diverse coordination modes, as shown in Figure 2, have been proposed for Schiff base adducts of salicylaldimine type ligand [14]. Although, the assignment of coordination mode for the Schiff base adducts could be somewhat ambiguous, the structure of an organotin Schiff base adduct, Me₂SnCl₂.SalenH₂, has been resolved by X-ray single crystal refinement [15]. The SalenH₂ ligand coordinates to the tin atom through the oxygen atom [11]. Therefore, formation of Schiff base adducts provides another structural variability to the chemistry of Schiff base metal complexes.




Figure 2. Structural types of salicylaldimine Schiff base adducts.

This study presents the synthesis and characterization of some Cu(II) metal chelates and Schiff base adducts. The Schiff base ligands were prepared from salicylaldehyde or *o*-vanillin and *p*-substituted aniline. The Schiff base ligands and the Cu(II) complexes have been evaluated for their antimicrobial activity.

EXPERIMENTAL

All the chemicals and reagents were of reagent grade and used without further purification. The ¹H- and ¹³C-NMR spectra for the Schiff base ligands were recorded on Bruker Avance NMR equipment operating at 400 MHz using deuterated chloroform with TMS as internal standard. The FT-IR absorption spectra and the electronic spectra for the compounds were recorded on Perkin Elmer Spectrum 100 FT-IR and Perkin Elmer Lambda 25 spectrophotometer, respectively. The FT-IR spectrometer was equipped with a universal attenuated total reflectance (ATR) accessory. The elemental analysis, CHN, was done on Vario MICRO V1.6.2 elemental

analysen system GmbH, while the melting points (uncorrected) of the compounds were determined using the Galenkemp melting point apparatus. All the micro-organisms used for the antimicrobial study were obtained from Microbiologics, Cape Town.

Table 1. $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ Spectral data for the Schiff base ligands.

| S/N | Ligands | $^1\text{H-NMR}$ (δ) ppm | $^{13}\text{C-NMR}$ (δ) ppm |
|-----|---------|---|--|
| 1 | L1 | 13.26 (1H, s, -OH); 8.66 (1H, s, HC=N); 7.44 (3H, m); 7.32 (3H, m); 7.07 (1H, d, J = 8.10); 6.98 (1H, t, J = 7.96). | 163.08 (Ar-OH); 161.60 (HC=N); 149.00 (Ar-C=N); 133.53, 132.66, 122.98, 127.27, 121.56, 119.66, 119.44, 117.68 (Ar-C). |
| 2 | L2 | 13.01 (1H, s, -OH); 8.63 (1H, s, HC=N); 7.40 (4H, m); 7.30 (2H, d, J = 8.59); 7.10 (1H, d, J = 8.57); 7.00 (1H, t, J = 7.48). | 163.39 (Ar-OH); 161.56 (HC=N); 147.53 (Ar-N=C); 133.83 (Ar-Cl); 132.93, 132.78, 129.94, 122.83, 119.58, 117.74 (Ar-C). |
| 3 | L3 | 12.97 (1H, s, -OH); 8.63 (1H, s, HC=N); 7.58 (2H, d, J = 8.64); 7.43 (2H, m); 7.19 (2H, d, J = 8.64); 7.06 (1H, d, J = 8.59); 6.98 (1H, t, J = 7.47). | 163.45 (Ar-OH); 161.57 (HC=N); 148.03 (Ar-N=C); 133.88, 132.91, 132.80, 123.20, 120.78, 119.59, 117.75 (Ar-C). |
| 4 | L4 | 13.37 (1H, s, -OH); 8.66 (1H, s, HC=N); 7.40 (2H, m); 7.22 (2H, d); 7.20 (2H, d); 7.10 (1H, d, J = 8.16); 7.0 (1H, m); 2.48 (3H, s, -CH ₃). | 162.11 (Ar-OH); 161.56 (HC=N); 146.37 (Ar-N=C); 137.29 (Ar-CH ₃); 133.29, 132.50, 130.40, 121.39, 119.71, 117.63 (Ar-C); 21.42 (-CH ₃). |
| 5 | L5 | 13.41 (1H, s, -OH); 8.64 (1H, s, HC=N); 7.39 (2H, m); 7.32 (1H, t); 7.30 (1H, q); 7.05 (1H, m); 7.0 (1H, t); 6.98 (1H, t); 6.95 (1H, dd); 3.88 (3H, s, -OCH ₃). | 161.44 (Ar-OH); 160.88 (HC=N); 159.30 (Ar-OCH ₃); 141.81 (Ar-N=C); 133.07, 132.34, 122.68, 119.82, 119.36, 117.57, 115.07 (Ar-C), 55.95 (-OCH ₃). |
| 6 | L6 | 12.54 (1H, s, -OH); 8.63 (1H, s, HC=N); 8.30 (2H, d, J = 8.91); 7.5 (2H, m); 7.4 (2H, d, J = 8.91); 7.10 (1H, d, J = 8.15); 7.0 (1H, t, J = 7.49). | 165.84 (Ar-OH); 161.74 (HC=N); 154.64 (Ar-N=C); 146.50 (Ar-NO ₂); 134.96, 133.45, 125.69, 122.35, 120.02, 119.14, 117.98 (Ar-C). |
| 7 | L7 | 13.68 (1H, s, -OH); 8.67 (1H, s, HC=N); 7.46 (2H, t, J = 7.75); 7.31 (3H, m); 7.06 (2H, m); 6.94 (1H, t, J = 7.88Hz); 3.98 (3H, s, -OCH ₃). | 163.02 (HC=N); 151.99 (Ar-N=C); 148.97 (Ar-OCH ₃); 148.66 (Ar-OH); 129.82, 127.37, 121.56, 118.92, 115.38, (Ar-C); 56.63 (-OCH ₃). |
| 8 | L8 | 13.36 (1H, s, -OH); 8.64 (1H, s, HC=N); 6.93 (1H, t, J = 7.88); 7.10 (2H, m); 7.19 (2H, d, J = 8.69); 7.57 (2H, d, J = 8.69); 3.97 (3H, s, -OCH ₃). | 163.44 (HC=N); 151.85 (Ar-OCH ₃); 148.96 (Ar-N=C); 147.73 (Ar-OH); 132.93 (Ar-Cl), 123.21, 120.88, 119.44, 119.13, 115.67 (Ar-C); 56.68 (-OCH ₃). |
| 9 | L9 | 13.39 (1H, s, -OH); 8.63 (1H, s, HC=N); 7.10 (2H, dd, J = 7.22); 7.30 (2H, d, J = 8.51); 7.40 (2H, d, J = 8.54); 6.90 (1H, t, J = 7.86); 3.97 (3H, s, -OCH ₃). | 163.38 (HC=N); 151.84 (Ar-OCH ₃); 148.95 (Ar-N=C); 147.23 (Ar-OH); 133.01 (Ar-Br); 129.95, 122.85, 119.11, 115.63, (Ar-C), 56.68 (-OCH ₃). |
| 10 | L10 | 13.82 (1H, s, -OH); 8.66 (1H, s, HC=N); 7.04 (2H, m); 7.25 (4H, m); 6.91 (1H, t, J = 7.87); 3.97 (3H, s, -OCH ₃) and 2.42 (3H, s, -CH ₃). | 162.03 (HC=N); 151.99 (Ar-OCH ₃); 148.96 (Ar-OH); 146.01 (Ar-N=C); 137.40 (Ar-CH ₃); 130.42, 121.39, 119.66, 118.82, 115.19 (Ar-C); 56.66 (-OCH ₃); 21.42 (-CH ₃). |
| 11 | L11 | 12.86 (1H, s, -OH); 8.68 (1H, s, HC=N); 8.30 (2H, d, J = 8.98); 7.40 (2H, d, J = 8.98); 7.10 (2H, d, J = 8.16); 7.0 (1H, m); 3.98 (3H, s, -OCH ₃). | 165.80 (HC=N); 154.43 (Ar-N=C); 152.04 (Ar-OCH ₃); 149.04 (Ar-OH); 146.64 (Ar-NO ₂); 125.62, 124.73, 122.29, 119.54, 119.16, 116.47 (Ar-C); 56.74 (-OCH ₃). |
| 12 | L12 | 13.58 (1H, s, -OH); 9.62 (1H, s, -OH); 8.88 (1H, s, HC=N); 7.31 (2H, d); 7.18 (1H, d); 7.08 (1H, d); 6.89 (1H, d); 6.85 (2H, d) and 3.82 (3H, s, -OCH ₃). | 161.15 (HC=N); 157.83 (Ar-OH); 151.32 (Ar-OH); 148.74 (Ar-C=N); 139.90, 124.56, 123.47, 120.23, 119.29, 116.87, 116.03 (Ar-C), 56.78 (-OCH ₃). |

Note: L¹ = Sal-H; L² = Sal-Cl; L³ = Sal-Br; L⁴ = Sal-Me; L⁵ = Sal-OMe; L⁶ = Sal-NO₂; L⁷ = Ovan-H, L⁸ = Ovan-Cl, L⁹ = Ovan-Br, L¹⁰ = Ovan-Me, L¹¹ = Ovan-NO₂, L¹² = Ovan-OH.

Synthesis of the Cu(II) complexes

The Cu(II) complexes were prepared by reacting a series of *p*-substituted aniline derived Schiff bases with Cu(II) chloride dihydrate, CuCl₂·2H₂O in hot ethanolic solution under stirring condition. The Schiff bases, L¹–L¹² were obtained from salicylaldehyde, *o*-vanillin and *p*-substituted aniline (X = H, Cl, Br, Me, OMe, NO₂ and OH). The ¹H- and ¹³C-NMR spectral data for the Schiff base ligands are presented in Table 1.

Hot ethanolic solution of CuCl₂·2H₂O was gradually added to a hot ethanolic solution of the ligand in a ratio 1:2. The resulting solution was stirred for 30 min with slight heating to obtain the Cu(II) complexes. Diethyl ether was used to induce precipitation in cases where the precipitate did not form on stirring. The precipitate was filtered under suction, washed with ethanol and dried in a vacuum desiccator over silica gel. The analytical and the spectroscopic data are presented in Tables 2 and 3, respectively.

Table 2. Physical and analytical data for the complexes.

| Complex | Colour | %Yield | M.P. (°C) | Molar mass (g/mol) | % Found (calculated) | | | | *Λ _M |
|--|--------|--------|--------------|-----------------------|----------------------|----------------|------------------|------------------|-----------------|
| | | | | | C | H | N | Cu | |
| Cu(L ¹) ₂ Cl ₂ ·2½H ₂ O | Brown | 69 | 128-130 | 572.05 | 54.59 (54.69) | 4.41 (4.26) | 4.90 (5.91) | 11.11 (11.62) | 183.00 (2:1) |
| Cu(L ²) ₂ Cl ₂ ·2H ₂ O | Brown | 55 | 160-162 | 633.95 | 49.26 (49.61) | 3.82 (3.33) | 4.42 (4.72) | 11.02 (10.15) | 138.00 (2:1) |
| Cu(L ³) ₂ Cl ₂ ·H ₂ O | Brown | 75 | 171-172 | 704.77 | 44.31 (44.56) | 3.15 (2.82) | 3.97 (4.07) | 9.02 (8.72) | 126.50 (2:1) |
| Cu(L ⁴) ₂ Cl ₂ ·H ₂ O | Brown | 57 | 156-158 | 575.01 | 58.49 (58.68) | 4.91 (4.71) | 4.85 (5.00) | 11.79 (11.05) | 83.00 (1:1) |
| Cu(L ⁵) ₂ | Red | 77 | 170-172 | 516.07 | 65.08 (65.16) | 4.92 (4.69) | 5.34 (5.43) | 11.98 (12.31) | 1.60 (DMF) |
| Cu(L ⁶) ₂ ·½H ₂ O | Orange | 90 | >250 | 555.03 | 56.60 (56.26) | 3.48 (3.48) | 10.13 (10.09) | 11.12 (11.45) | 6.39 (DMF) |
| Cu(L ⁷) ₂ Cl ₂ ·H ₂ O·½EtOH | Brown | 88 | 124-126 | 629.42 | 55.34 (55.52) | 4.97 (4.69) | 4.45 (4.44) | 10.10 (9.80) | 134.2 (2:1) |
| Cu(L ⁸) ₂ | Green | 84 | 194-196 | 584.85 | 57.30 (57.50) | 4.05 (3.79) | 4.76 (4.79) | 11.01 (10.87) | 1.57 (DMF) |
| Cu(L ⁹) ₂ Cl ₂ ·H ₂ O | Brown | 56 | 179-180 | 764.93 | 43.97 (43.47) | 3.43 (3.16) | 3.66 (3.61) | 8.31 (7.72) | 99.80 (1:1) |
| Cu(L ¹⁰) ₂ Cl ₂ ·2½H ₂ O | Brown | 60 | 168-170 | 671.19 | 53.69 (54.41) | 5.41 (5.33) | 4.17 (4.23) | 9.47 (9.60) | 82.70 (1:1) |
| Cu(L ¹¹) ₂ ·2EtOH | Brown | 78 | >250 | 698.09 | 55.17 (55.06) | 4.91 (4.38) | 8.03 (8.00) | 9.10 (8.79) | 6.61 (DMF) |
| Cu(L ¹²) ₂ Cl ₂ ·H ₂ O | Brown | 67 | 178-180 | 637.09 | 52.79 (52.44) | 4.11 (4.31) | 4.40 (5.03) | 9.98 (9.77) | 131.00 (2:1) |

*Λ_M (Ω⁻¹cm²mol⁻¹). Note: L¹ = Sal-H; L² = Sal-Cl; L³ = Sal-Br; L⁴ = Sal-Me; L⁵ = Sal-OMe; L⁶ = Sal-NO₂; L⁷ = Ovan-H; L⁸ = Ovan-Cl; L⁹ = Ovan-Br; L¹⁰ = Ovan-Me; L¹¹ = Ovan-NO₂; L¹² = Ovan-OH.

RESULTS AND DISCUSSION*Elemental analysis and conductivity measurement*

The physical and the analytical data for the complexes are presented in Table 2. The Cu(II) complexes were obtained in high yield and the micro analytical data indicate the complexes are relatively pure. Analysis of the CHN results indicates the isolation of the Cu(II) complexes as inner chelates for ligands L⁵, L⁶, L⁸ and L¹¹ while ligands L¹–L⁴, L⁷, L⁹, L¹⁰ and L¹² complexes

were obtained as Schiff base adducts in 1:2 stoichiometric ratio comprising two molecules of the ligands, two atoms of chlorine and one of copper. The metal chelates were non-electrolytes with molar conductivity values of $6.61\text{--}1.57\ \Omega^{-1}\text{cm}^2\text{mol}^{-1}$ while the Schiff base adducts were either 1:1 ($99.8\text{--}82.7\ \Omega^{-1}\text{cm}^2\text{mol}^{-1}$) or 2:1 ($183\text{--}126.5\ \Omega^{-1}\text{cm}^2\text{mol}^{-1}$) electrolytes in methanol [16]. The complexes are, therefore, proposed to be of the forms $[\text{CuL}_2]\cdot x\text{H}_2\text{O}$ for the metal chelates and $[\text{Cu}(\text{LH})_2\text{Cl}]\text{Cl}\cdot x\text{H}_2\text{O}$ or $[\text{Cu}(\text{LH})_2]\text{Cl}_2\cdot x\text{H}_2\text{O}$ for the Schiff base adducts (Figure 3).




Figure 3. Proposed structures for the Schiff base adducts.

Infrared study

Some selected infrared spectral data for the Schiff base complexes are presented in Table 3. The phenolic hydroxyl stretching vibration for the free Schiff base ligands were observed as a broad and weak band at $3100\text{--}2200\ \text{cm}^{-1}$ due to strong intramolecular hydrogen bonding which is characteristic of ortho-hydroxyl Schiff base ligands [17]. In addition, the vibrational absorption band for the azomethine group, $\text{C}=\text{N}$, appeared at $1616\text{--}1598\ \text{cm}^{-1}$ [18-21]. The broad hydroxyl band of the OH group disappeared in the Cu(II) chelates of L^5 , L^6 , L^8 & L^{11} , indicating coordination of the ligands via the deprotonated oxygen atom [22-23]. This was corroborated by the bathochromic shift in the phenolic C-O stretch of the free ligands at $1280\text{--}1270\ \text{cm}^{-1}$ to $1330\text{--}1318\ \text{cm}^{-1}$. Also, the azomethine absorption band underwent negative shift in the spectra of the metal chelates by $-20\ \text{cm}^{-1}$ to $-8\ \text{cm}^{-1}$, suggesting coordination via the imine nitrogen [2, 6, 17, 18]. A new strong band was, however, observed in the spectral of the Schiff base adducts at $1641\text{--}1628\ \text{cm}^{-1}$. This band was absent in the metal chelates and it indicates the presence of a quinoid system in the Schiff base adducts [12] as shown in Figure 3. In addition, the presence of the N-H group is indicated by the appearance of a weak to medium band at $3500\text{--}3100\ \text{cm}^{-1}$ [6] in the spectra of the Schiff base adducts. The OH stretching vibration of the lattice water also absorbed in this region [6, 22]. The coordination mode of the Schiff base ligands was further substantiated by the assignment of metal-ligand vibration bands in the far-infrared spectra of the metal complexes. The bands corresponding to the $\nu_{\text{Cu-N}}$, $\nu_{\text{Cu-O}}$ and $\nu_{\text{Cu-Cl}}$ bonds were observed at $592\text{--}534\ \text{cm}^{-1}$, $515\text{--}424\ \text{cm}^{-1}$ and $387\text{--}346\ \text{cm}^{-1}$, respectively [20, 24].

Table 3. Selected infrared and electronic spectral data for the complexes.

| S/N | Compound | $\nu_{\text{NH}}/\nu_{\text{OH}}$ | $\nu_{\text{C=N}}$ | ν_{CO} | $\nu_{\text{Cu-N}}$ | $\nu_{\text{Cu-O}}$ | $\nu_{\text{Cu-Cl}}$ | λ (nm) |
|-----|--|-----------------------------------|--------------------|-------------------|---------------------|---------------------|----------------------|------------------------------|
| 1 | Cu(L ¹) ₂ Cl ₂ . 2½H ₂ O | 3261, 3215 | 1606, 1634 | 1315 | 534, 564 | 497, 515 | 387 | 230, 280, 383, 713 |
| 2 | Cu(L ²) ₂ Cl ₂ . 2H ₂ O | 3322, 3241, 3205 | 1606, 1628 | 1309 | 589 | 426, 463 | 346 | 240, 282, 386, 810 |
| 3 | Cu(L ³) ₂ Cl ₂ . H ₂ O | 3119 | 1603, 1633 | 1309 | 586 | 442, 512 | 368 | 223, 240, 282, 385, 804 |
| 4 | Cu(L ⁴) ₂ Cl ₂ . H ₂ O | 3251, 3210 | 1606, 1638 | 1316 | 585 | 465, 503 | 364 | 282, 383, 750 |
| 5 | Cu(L ⁵) ₂ | - | 1600 | 1325 | 591 | 426, 496 | - | 230, 273, 299, 384, 501, 674 |
| 6 | Cu(L ⁶) ₂ . ½H ₂ O | - | 1610 | 1320 | 587 | 491, 424 | - | 374, 423, |
| 7 | Cu(L ⁷) ₂ Cl ₂ . H ₂ O. ½EtOH | 3494, 3438 | 1608, 1633 | 1309 | 540 | 464, 424 | 355 | 231, 287, 391, 748 |
| 8 | Cu(L ⁸) ₂ | - | 1598 | 1318 | 584, 562 | 492, 441 | - | 234, 293, 396, 405, 517 |
| 9 | Cu(L ⁹) ₂ Cl ₂ .H ₂ O | 3114 | 1613, 1637 | 1308 | 543, 528 | 492, 436 | 372 | 284, 394, 756, 819 |
| 10 | Cu(L ¹⁰) ₂ Cl ₂ .2½H ₂ O | 3251, 3207 | 1611, 1641 | 1305 | 595, 537 | 495, 442 | 347 | 231, 288, 388, 794 |
| 11 | Cu(L ¹¹) ₂ . 2EtOH | 3210 | 1603 | 1330 | 540, 573 | 466, 442 | - | 307, 412, 770 |
| 12 | Cu(L ¹²) ₂ Cl ₂ .H ₂ O | 3448, 3180 | 1608, 1641 | 1311 | 592, 538 | 492, 448 | 387 | 280, 313, 389, 502 |

Note: L¹ = Sal-H; L² = Sal-Cl; L³ = Sal-Br; L⁴ = Sal-Me; L⁵ = Sal-OMe; L⁶ = Sal-NO₂; L⁷ = Ovan-H; L⁸ = Ovan-Cl; L⁹ = Ovan-Br; L¹⁰ = Ovan-Me; L¹¹ = Ovan-NO₂; L¹² = Ovan-OH.

Electronic spectra study

The electronic spectral data for the Cu(II) complexes exhibited three different absorption bands due to transitions arising from the Schiff base ligands. The high energy bands at 234–223 nm and 293–273 nm correspond to the $\pi \rightarrow \pi^*$ transition of the benzene rings [25] and the azomethine group respectively [20, 26] while the third band at 412–341 nm was assigned to the $n \rightarrow \pi^*$ of the azomethine group, C=N [19]. In addition, a broad band corresponding to the d–d transition of the Cu(II) ion was observed at 674–423 nm and 819–713 nm in the spectra of the Cu(II) chelates and the Schiff base adducts, respectively [27].

Biological study

The Cu(II) complexes and the free ligands were evaluated for their antimicrobial activity against *Escherichia coli* ATCC® 8739™*, *Staphylococcus aureus subsp. aureus* ATCC® 6538™*, *Bacillus subtilis subsp. spizizenii* ATCC® 6633™* and *Candida albicans* ATCC® 2091™* using disc diffusion technique and two-fold serial dilution methods.

Disc diffusion technique

The qualitative antimicrobial susceptibility testing of the compounds was evaluated using the disc diffusion technique [28]. The micro-organisms were cultured on nutrient agar at 37 °C for 24 h to obtain the primary culture. About 10 mL bacterial suspension containing 10⁶-10⁸ CFUs was prepared from the primary culture using saline water. Sterile Mueller Hinton plates were then inoculated with 0.1 mL of the bacterial suspension and discs soaked with the test compounds were firmly placed on it. The plates were incubated at 37 °C for 16 h and the zone of inhibition was measured as millimetres diameter. Ampicillin and dimethylformamide (DMF)

were used as standard antibacterial drug and control solvent respectively. The test was repeated two more times for those compounds that showed activity of more than 6.5 mm and their activity was recorded as average zone of inhibition as presented in Figures 4 and 5. Similar procedure was repeated for the antifungal susceptibility testing of the compounds at 28 °C for 48 h using potato disc assay and ketoconazole in place of Mueller Hinton agar and penicillin, respectively.




Figure 4. Disc diffusion result of the salicylaldehyde-based complexes.




Figure 5. Disc diffusion result of the *o*-vanillin Schiff base complexes.

Minimum inhibitory concentration (MIC)

The quantitative antimicrobial activity of the test compounds was evaluated using micro dilution broth method according to Clinical and Laboratory Standard Institute (formally NCCLS) [28]. Two-fold serial dilutions of the compounds were prepared in 96 micro wells plates using sterile nutrient broth as diluent. The plates were inoculated with 5 μ L bacterial suspensions containing

10^6 – 10^8 CFUs and incubated at 37 °C for 16–18 h. The MIC value was defined as the lowest concentration of the compounds giving complete inhibition of visible growth. The MIC values for the compounds varied from 3.125 mg/mL to 0.0122 mg/mL and the result is presented in Table 4.

Nearly all the compounds exhibited significant antifungal activity against *Candida albicans* and were more potent than the standard antifungal agent, ketoconazole (Keto). They were, however, not as effective as ampicillin (Amp). The *Bacillus subtilis* and *Escherichia coli* strains exhibited the most and least antimicrobial susceptibility respectively. *E. coli* is a gram negative bacterium with a thick peptidoglycan cell wall which reduces the permeation of the test compounds into the microorganisms. Furthermore, the *o*-vanillin derived compounds possess higher activity than the salicylaldehyde analogues. The presence of the *ortho*-methoxy group could be responsible for the enhancement of the antimicrobial activity of the Schiff base ligands in the *o*-vanillin derived compounds. The biological activity of Schiff base compounds is often associated [20] with the presence of the of the azomethine functional group, HC=N, and the basicity of the nitrogen atom in this group depends on the nature of the substituents on either or both of the aldehyde and amine moiety of the Schiff base compounds. The inductive effect of the methoxyl group at the *ortho* position reduces the basicity of the azomethine nitrogen and may thus be responsible for the enhanced antimicrobial activity of the *o*-vanillin based compounds over the salicylaldehyde analogues. Similarly, the aniline based compounds exhibited higher antimicrobial activity than the *p*-substituted analogues.

Lastly, the antimicrobial activity of the Cu(II) complexes did not follow any regular pattern. It varied from positive to no effect and even slight reduction in the antimicrobial activity of the free ligands. The complexes of ligands L1, L4, L5, L7, L11 and L10 exhibited higher activity than the free ligands while the ligand L12 complex showed lower potency than the free ligand. The antimicrobial activity of the remaining complexes did not differ significantly from the free Schiff base ligands.

Table 4. MIC values for the Cu(II) complexes (1×10^{-1} mg/mL).

| Compounds | Gram-positive | | Gram-negative |
|-------------------|------------------|--------------------|----------------|
| | <i>S. aureus</i> | <i>B. subtilis</i> | <i>E. coli</i> |
| L ¹ | - | 7.8125 | - |
| CuL ¹ | 3.9062 | 3.9062 | 7.8125 |
| L ² | - | 3.9062 | - |
| CuL ² | - | 3.9062 | - |
| L ⁴ | - | 7.8125 | - |
| CuL ⁴ | 0.9766 | 7.8125 | - |
| CuL ⁵ | 31.25 | 7.8125 | - |
| L ⁷ | 3.9062 | 3.9062 | 7.8125 |
| CuL ⁷ | 0.2441 | 1.9531 | 0.3125 |
| L ⁸ | 7.8125 | 7.8125 | - |
| CuL ⁸ | *ns | *ns | *ns |
| L ⁹ | - | 31.25 | - |
| CuL ⁹ | - | 31.25 | - |
| L ¹⁰ | - | 3.9062 | - |
| CuL ¹⁰ | 0.1220 | 7.812 | 31.250 |
| L ¹¹ | - | 3.9062 | - |
| L ¹¹ | 15.625 | 3.9062 | 3.9062 |
| L ¹² | 0.4883 | 0.9766 | 3.9062 |
| CuL ¹² | 7.8125 | 0.9766 | 1.9531 |

*ns: Not soluble.

CONCLUSION

The reaction of Cu(II) chloride salt, $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$, with a series of *p*-substituted salicylaldimine Schiff bases yielded both the prevalent inner metal chelates $[\text{CuL}_2]$ and the partially resolved Cu(II) Schiff base adducts $(\text{LH})_2 \cdot \text{CuCl}_2$. The metal chelates were non-electrolytes while the Schiff base adducts were either 1:1 or 2:1 electrolytes in dimethylformamide. The Cu(II) complexes exhibited significant antifungal activity against *Candida albicans*.

ACKNOWLEDGEMENTS

A.O. Sobola is grateful to Rhodes University for providing facilities for this research.

REFERENCES

- Jiang, S.; Ni, H.; Liu, F.; Gu, S.; Yu, P.; Gou, Y. Binuclear Schiff base copper(II) complexes: Syntheses, crystal structures, HSA interaction and anti-cancer properties. *Inorg. Chim. Acta* **2020**, 499, 119186.
- Salehi, M.; Rahimifar, F.; Kubicki, M.; Asadi, A. Structural, spectroscopic, electrochemical and antibacterial studies of some new nickel(II) Schiff base complexes. *Inorg. Chim. Acta* **2016**, 443, 28-35.
- Liu, X.; Hamon, J.R. Recent developments in penta-, hexa- and heptadentate Schiff base ligands and their metal complexes. **2019**. *Coord. Chem. Rev.* 389, 94-118.
- Gaber, M.; El-Wakiel, N.; El-Baradie, K.; Hafez, S. Chromone Schiff base complexes: Synthesis, structural elucidation, molecular modeling, antitumor, antimicrobial, and DNA studies of Co(II), Ni(II), and Cu(II) complexes. *J. Iran. Chem. Soc.* **2019**, 16, 169-182.
- Bhunora, S.; Mugo, J.; Bhaw-Luximon A.; Mapolie, S.; Van Wyk J.; Darkwa, J.; Nordlander, E. The use of Cu and Zn salicylaldimine complexes as catalyst precursors in ring opening polymerization of lactides: Ligand effects on polymer characteristics. *Appl. Organomet. Chem.* **2011**, 25, 133-145.
- Maity, D.; Drew, M.G.B.; Godsell, J.F.; Roy, S.; Mukhopadhyay, G. Synthesis and characterization of Cu(II) complexes of tetradentate and tridentate symmetrical Schiff base ligands involving *o*-phenylenediamine, salicylaldehyde and diacetyl monoxime. *Transition Met. Chem.* **2010**, 35, 197-204.
- Chen, W.-H.; Wei, H.-H.; Lee, G. -H.; Wang, Y. Dimeric iron(III) complexes with salicylaldimine ligands: Structures, magnetic properties, and activity of H_2O_2 disproportionation. *Polyhedron* **2001**, 20, 515-521.
- Percy, G.C.; Thornton, D.A. N-Aryl salicylaldimine complexes. Infrared and PMR spectra of the ligands and vibrational frequencies of their metal(II) chelates. *J. Inorg. Nucl. Chem.* **1972**, 34, 3357-3367.
- Sacconi, L.; Ciampolini, M.; Nardi, N. Investigation on the occurrence of tetrahedral forms of substituted bis(N-alkylsalicylaldimino)nickel(II) complexes. *J. Am. Chem. Soc.* **1964**, 86, 819-823.
- Bowden, F.L.; Ferguson, D. Redox reactions in the formation of titanium complexes with aromatic Schiff bases. *J. Chem. Soc., Dalton Trans.* **1974**, 460-462.
- Randaccio, L. The polymeric structure of the adduct of dimethyltin dichloride with bis (salicylaldehyde) ethylenediamine. *J. Organometal. Chem.* **1973**, 55, c58-c60.
- Bamfield, P. The reaction of cobalt halides with N-arylsalicylideneimines. *J. Chem. Soc. A. Inorg. Phys. Theor.* **1967**, 804-808.

13. Kwiatkowski, M.; Kwiatkowski, E.; Olechnowicz, A.; Kołczyński, B.Z. Cadmium complexes with the tridentate amino-ketoenamides derived from pentane-2, 4-dione and aliphatic diamines. *Polyhedron* **1991**, *10*, 945-947.
14. Garnovskii, A.D.; Nivorozhkin, A.L.; Minkin, V.I. Ligand environment and the structure of Schiff base adducts and tetracoordinated metal-chelates. *Coord. Chem. Rev.* **1993**, *126*, 1-69.
15. Calligaris, M.; Nardin, G.; Randaccio, L. Structural aspects of metal complexes with some tetradentate Schiff bases. *Coord. Chem. Rev.* **1972**, *7*, 385-403.
16. Geary, W.J. Use of conductivity measurements in organic solvents for the characterization of coordination compounds. *Coord. Chem. Rev.* **1971**, *7*, 81-122.
17. Kovacic, J.E. The carbon:nitrogen stretching frequency in the infrared spectra of Schiff's base complexes. I. Copper complexes of salicylidene anilines. *Spectrochim. Acta* **1967**, *23A*, 183-187.
18. Percy, G.C.; Thornton, D.A. Infrared spectra of N-aryl Schiff base complexes. *Inorg. Nucl. Chem. Lett.* **1971**, *7*, 599-604.
19. De, A.; Ray, H.P.; Jain, P.; Kaur, H.; Singh, N. Synthesis, characterization, molecular docking and DNA cleavage study of transition metal complexes of *o*-vanillin and glycine derived Schiff base ligand. *J. Mol. Struct.* **2020**, *1199*, 126901.
20. Bahron, H.; Khaidir, S.S.; Tajuddin, A.M.; Ramasamy, K.; Yamin, B.M. Synthesis, characterization and anticancer activity of mono- and dinuclear Ni(II) and Co(II) complexes of a Schiff base derived from *o*-vanillin. *Polyhedron* **2019**, *161*, 84-92.
21. Iftikhar, B.; Javed, K.; Khan, M.S.U.; Akhter, Z.; Mirza, B.; McKee, V. Synthesis, characterization and biological assay of salicylaldehyde Schiff base Cu(II) complexes and their precursors. *J. Mol. Struct.* **2018**, *1155*, 337-348.
22. Sakthi, M.; Ramu, A. Synthesis, structure, DNA/BSA binding and antibacterial studies of NNO tridentate Schiff base metal complexes. *J. Mol. Struct.* **2017**, *1149*, 727-735.
23. Kalitha, M.; Bhattacharjee, T.; Gogoi, P.; Barman, P.; Kalita, R.D.; Sarma, B.; Karmakar, S. Synthesis, characterization, crystal structure and bioactivities of a new potential tridentate (ONS) Schiff base ligand N-[2-(benzylthio) phenyl] salicylaldehyde and its Ni(II), Cu(II) and Co(II) complexes. *Polyhedron* **2013**, *60*, 47-53.
24. Zhou, Y.; Liu, L.; Yang, M.; Lu, R.; Jin, Y.; Chen, W. Synthesis, crystal structure and biological property of a novel phenolato-bridged trinuclear copper(II) complex derived from bis-Schiff base ligand. *Inorg. Nano-Met. Chem.* **2019**, *48*, 291-295.
25. Zhou, Y.; Ye, X.; Xin, F.; Xin, X. Solid state self-assembly synthesis of cobalt(II), nickel(II), copper(II) and zinc(II) complexes with a bis-Schiff base. *Transition Met. Chem.* **1999**, *24*, 118-120.
26. Turkyilmaz, M.; Onder, A.; Baran, Y. Synthesis, spectroscopic, and thermal properties of some azomethine complexes of Cu(II), Ni(II), and Pt(II). *J. Therm. Anal. Calorim.* **2012**, *109*, 991-998.
27. Samanta, B.; Chakraborty, J.; Choudhury, C.R.; Dey, S.K.; Dey, D.K.; Batten, S.R.; Jensen, P.; Yap, G.P.A.; Mitra, S. New Cu(II) complexes with polydentate chelating Schiff base ligands: Synthesis, structures, characterisations and biochemical activity studies. *Struct. Chem.* **2007**, *18*, 33-41.
28. Schalbe, R.; Steele-Moore, L.; Goodwin, A.C. (Eds). *Antimicrobial Susceptibility Testing Protocols*, CRC Press: Boca Raton, Florida; **2007**; p 53.