



Herbal Medicines in Kenya: A Review of the Toxicity and Quality Control Issues

Maina N¹, Kagira JM², Achila O¹, Karanja SM¹, Ngotho M²

1. Department of Biochemistry, Jomo Kenyatta University of Agriculture and Technology, PO Box 62000-00200, Nairobi, Kenya.
2. Institute of Primate Research, PO Box 24481-00502, Nairobi, Kenya;

***Corresponding author:** Dr. Naomi Maina, Department of Biochemistry, Jomo Kenyatta University of Agriculture and Technology, PO Box 62000-00200, Nairobi, Kenya. Email: nmaina@hotmail.com; Tel: +254727726785

Financial support: National Council of Science and Technology

SUMMARY

In sub-Saharan Africa, it is estimated that 80% of the population depends on indigenous medicines for primary health-care. These herbs often contain highly active pharmacological compounds whose pharmacotherapeutic and toxicity profiles have not been well characterized. Toxicity may be related to several intrinsic and extrinsic factors. Most of the available reports related to the toxic effects of herbal medicines cite hepatotoxicity as the most frequently experienced toxicity. However, noxious effects involving kidneys, the nervous system, skin, blood, the cardiovascular system, mutagenicity and carcinogenicity have also been published. This article presents a systematic review on safety and toxicity of herbal medicines used in Kenya.

Keywords: Herbal medicine, herbal preparations, toxicity, Kenya, regulations, complementary and alternative medicine.

[*Afr J Health Sci.* 2013; 24:29-47]

Introduction

Herbal medicine is an important part of the culture and traditions of most African communities. Kenya in particular has a rich and diverse repertoire of flora that has been used by various ethnic communities as medicines for centuries. Ethnobotanical information and usage of traditional herbal medicine by different ethnic communities in Kenya has been investigated by

several researchers (Jeruto *et al.*, 2006; Bussman, 2006; Nguta *et al.*, 2010; Njoroge *et al.*, 2010; Kareru *et al.*, 2008; Nagata *et al.*, 2011; Nanyingi *et al.*, 2008). According to some estimates, traditional healers provide the first line of care for 80-90% of the population (Lambert *et al.*, 2011). While herbal usage, especially in rural communities is high, an estimated 23% of the sick do not seek conventional health care (Njoroge *et al.*, 2010; Muthaura *et al.*, 2011). The



importance of herbal medicines in Kenya is further evidenced by the fact that traditional herbal medical practitioners (THMPs) far outnumber conventional or allopathic providers. Given the estimated 40,000 THMPs and assuming a population of 38 million Kenyans, there is a THMP – patient ratio of 1 to 950 (Lambert *et al.*, 2011). Currently, the conventional doctor to patient ratio stands at 1 to 33 000 (Ibid).

The growing popularity of herbal medicines has been attributed to a number of factors including accessibility, affordability, aggressive marketing by THMP and the perception that herbal medicines are natural hence safe (Sahoo *et al.*, 2010). Parallel with the growing popularity is an increasing interest in the scientific rationale for plant usage. Towards this end, several research groups in Kenya have investigated the efficacy of specific herbs. Crude or fractionated extracts of medicinal herbs have been screened in *in vitro* and *in vivo* systems for antibacterial (Kareru *et al.*, 2008; Matasyoh *et al.*, 2008); antiviral (Kanyara *et al.*, 2005) and antiplasmodial activity (Kigundu *et al.*, 2011; Wagate *et al.*, 2008). Still, little is known about their mechanisms of actions. Another aspect, which is largely under investigated is the toxicity associated with the use of herbs (Huxtable, 1990). This should raise some concern given the recent warnings by biomedical health practitioners on the potential adverse effects of herbal remedies. In this paper, we evaluate the safety and toxicity herbal medicines used in the country.

Herbal Medicine in Kenya


Current regulatory status

Herbal medicine practitioners in Kenya are registered by the Ministry of Culture and Social Services. Registration is non-obligatory and is subject to the requirement that herbal medicines used should be submitted to national laboratories, namely: Kenya Medical Research Institute (KEMRI), Nairobi University's Mitishamba Drug Research Unit (MDRU) and Government Chemists (GC) for evaluation of safety. However, toxicity tests on the herbal preparations presented are mostly not done. These institutions mainly rely on literature search to predict the efficacy and safety of the herbs. The literature review has some important shortcomings, for instance: most of these studies are directed at the toxicological properties of single plant formulations. However, the labels on most herbal preparations, especially those sold as panaceas for common chronic diseases; indicate that most contain multiple herbs.

For mass produced herbal products, registration is done by the Pharmacy and Poisons Board of Kenya (PPBK) under the Ministry of Health Services. However, PPBK relies, in part, on the same laboratories mentioned earlier and is thus subject to the same challenges highlighted above. In addition, the regulatory scheme by PPBK covers only Good Manufacturing Practices (GMP) and Good Supply Practices (GSP) but not Good Agricultural and Collection Practices (GACP). Figure 1 shows the circulation of herbal medicines in Kenya and how these regulatory regimes apply.



Figure 1: Flow diagram of the circulation of herbal medicines in Kenya.



Red flow lines represent non regulated channels of the flow of herbal medicines or materials; The blue flow lines represent part of the channel regulated by the Ministry of Culture. However, most herbal medicine practitioners are not registered by the Ministry of Culture. Therefore the blue line represents partial regulation. The black flow lines represent part of the channel regulated by the Poisons and Pharmacy Board (PPB).

Safety concerns and challenges in the regulation of herbal preparations are many and include:

- Some practitioners combine herbal treatment with spirituality making regulation difficult (Kokwaro, 1993).
- There are unscrupulous traders/healers who typically use misleading adverts to market their products. This has prompted to the establishment of guidelines for advertisement and promotion of herbal

products (Robison, 2011). However, the enforcement of these guidelines remains a challenge.

- There is a general decline in authentic knowledge of traditional remedies; this issue has health implications since sourcing of herbs by non-experts can result in misidentification which may lead to the inadvertent use of toxic herbs.
- There are no forums for exchange of information/knowledge among the herbal medical practitioners (THMPs), and conventional medical practitioners (Lambert *et al.*, 2011).
- Further, the emerging non-traditional use of herbal preparation (e.g., use of sliming products and herbal teas) may lead to prolonged use. Such prolonged usages enhance the likelihood of cumulative toxic consequences (Aronson, 2009).
- Pharmacovigilance for herbal medicines is nonexistent in Kenya (Onyambu, 2011).



Risks associated with herbal medicines.

This may be attributed to a number of quality related issues (Zhang *et al.*, 2011). From cultivation of medicinal herbs to the final product, several factors (internal and external) can influence quality.

External Quality Issues

External quality issues include: contamination, adulteration, substitution and misidentification.

Contamination and substitution of herbal medicines

Contamination of herbal medicines with heavy metals (Lead (Ld), Cadmium (Cd), Mercury (Hg), Arsenic (Ag), misidentified plants, pesticides, excessive levels of microorganisms or their toxins and other noxious organic materials are possible sources of danger in non-regulated use of herbal medicines (Mobusz *et al.*, 2006; Aronson, 2009). Contamination can occur at any stage from harvesting through packaging and storage (WHO, 2005). Experts believe that toxicities associated with herbal medicines are mostly due to these contaminants and clinical reports have vindicated this belief (Robinson *et al.*, 2011). A recent study in Kenya, found that 67% of the herbal samples had bacterial count in the range of 6.00×10^5 to 1.56×10^{10} cfu/ ml and fungal count in the range of 5.30×10^4 to 1.56×10^9 cfu/ ml. Analysis confirmed the presence of microbes including, *Escherichia coli*, *Shigella* spp, *Klebsiella pneumonia* (Onyambu, 2011).

Adulteration of herbal medicines

Another area of concern is the adulteration of herbal medicines with conventional drugs or other plant

materials (WHO, 2005). For instance, a study spanning 1990 to 2001 which evaluated 41 products out of 3320 Chinese Proprietary Medicines (CPM) sold in Singapore reported that all these products were contaminated with nineteen synthetic drugs (Sahoo *et al.*, 2010). When 243 proprietary herbal formulations in California were tested, it was revealed that 7% contained undeclared pharmaceuticals (non-steroidal anti-inflammatory drugs (NSAIDs) and antihistamines, among others) (Phua and Heard, 2009). Since some herbs sold in the Country are imported from China, it can be asserted that Kenyans are exposed to the dangers posed by the presence of undeclared pharmaceuticals in TCM. In fact, reports published recently by the PPB indicate that some Chinese herbal contraceptives sold in the country contain high levels of conventional contraceptive (levonorgestrel and quinestrol) (PPB, 2012). Children born to mothers using these concoctions presented with several complications including preconscious sexual maturity, edema, muscular pain and impaired speech (PPB, 2012; Nzioki, 2011; Okwemba 2010).

Internal quality issues

Internal quality challenges are related to the pharmacologically active phytochemicals present in a formulation (Zhang *et al.*, 2011).

Dosage of the active compounds

The concentration of bioactive phytochemicals is determined by several factors including geographical (Matasyoh *et al.*, 2008) and environmental factors (Zhang *et al.*, 2011), species differences, part of the plant used, diurnal and seasonal variations (Saad *et*



al., 2001; Snodgrass, 2001; Farhat *et al.*, 2001), soil composition and its contaminants and length of storage (Stickel *et al.*, 2005). This means that without proper standardization, the actual dose of bioactive compounds being consumed is often variable, unpredictable or simply unknown. Indeed, batch-to-batch variation of pharmacologically active constituents of herbal formulations presents a major problem when it comes to standardization of herbal medicines (Mobusz *et al.*, 2006). While testing the antimicrobial activity of hydro – distilled volatile oils from the leaves of *Ocimum gratissimum* L. (Lamiaceae) of 13 populations of different silvicultural zones in Kenya; a recent analytical study found significant variation in the antibacterial activity of the essential oils sourced from different parts of the country (Matasyoh *et al.*, 2008). Plants sourced from the Meru region had the greatest strength. Such variation in dosage may have greater effects in special group including children, geriatrics, invalids, and malnourished/undernourished, drug users,

HIV/AIDS patients, patients with liver or kidney complications and those on long term medications (Snodgrass, 2001).

Toxicity of some commonly used herbs in Kenya

Literature on toxicity of herbal medicines used by THMPs in Kenya is generally scant. Therefore, many questions regarding the safety and toxicity of specific herbs used in Kenya cannot be answered definitely by the available scientific data. However, the fact that some of the herbs used by THMPs are toxic cannot be discounted. This is underscored by the fact that some plants used in the country like *Aristolochia sp.* are listed by the European Commission's Committee for Proprietary Medicinal Products (CPMP) as plants which pose serious health risks (Kokwaro, 1993; Hamill *et al.*, 2000). In Table 1 below, we present a list of the toxicological profiles of some commonly used medicinal plants.



Table1: Toxicity of some commonly used herbs

Family	Species	Common Name	Part of plant used	Ethnic Group	Indications and Parts used	Toxicity	Compound	Reference
Amaranthaceae	<i>Achyranthes aspera</i>	Chaff Flower	RT, LV:	Nandi(1); Meru,Mbeere(7); Luo(10); Maasai (6);Kamba and Luyha (10)	Amoebiasis, Bleeding wounds and stomach ache. Tooth ache, cough	Antiandrogenic and abortifacant property. elevation of thyroid hormone levels†	Unknown	Anuja <i>et al.</i> , 2010 Tahiliani and Kar., 2000
Apocynaceae	<i>Acokanthera longiflora and Acokanthera schimperi</i>	Common poison bush	RT:	Massai (10)	Syphilis	Cardio toxicity	Cardenolide s: (ouabain, acolongifl oside	Donald., 2008
	<i>Thevetia peruviana</i>	Yellow Oleander	LV	Luo (8), Kikuyu (10)	Skin rash, cuts, burns, Blocked nose	Subendocardial and perivascular hemorrhage, focal	Thevetia A and B, thevetoxin, thevetin	Donald., 2008 Bandara <i>et al.</i> , 2009
	<i>Nerium Oleander</i>	Rose Laurel		Luo(8);Samburu (10)		Cardiovascular effects†	Cardiac glycosides	Bandara <i>et al.</i> , 2009
Aristolochiaceae	<i>Aristolochia bracteata</i>		LV	Turkana (10)	chest pain	Nephropathy and Carcinogenicity	Aristolochic acid	Worbs <i>et al.</i> , 2011
	<i>Aristolochia petersiana</i>		RT	Swahili (10)	Cough	Nephropathy and Carcinogenicity	Aristolochic acid	Worbs <i>et al.</i> , 2011



Aspidiaceae	<i>Pteridium aquilinum</i>	Bracken	LV	Nandi (1)	Skin diseases	Immunotoxic material Carcinogenic	(cooked safe), thiaminase	Ptaquiloside,	Latorre <i>et al.</i> , 2011
Balanitaceae	<i>Balanites aegyptiaca</i>	Balanites	LV, BK, FR	Luo (8); Nandi (1)	Itch, bone Problem. Diarrhea, chest pain, cough	Cardiovascular effect †	Unknown		Mohamed <i>et al.</i> , 1999
Bignoniaceae	<i>Kigelia africana</i>	Sausage Tree	BK,FR	Luo (8); Kikuyu (10); Meru (7); Taita and Luyha (10)	Stomach ache, anemia, cold, flu	Potentially mutagenicity	Unknown		Fennell <i>et al.</i> , 2004
Brassicaceae	<i>Heliotropium subulatum</i>	Heliotrope	RT	Turkana (10)	Post-parturition disease in women	Hepatotoxic and carcinogenic		Pyrrolizidine alkaloids	Donald., 2008
Capparidaceae	<i>Warbugia salutaris</i>	Fever Tree	BK	Luhya, Kikuyu, Maasai,	Muscle pain, fever, cough	GIT irritant	Unknown		Donald., 2008
	<i>Capparis tomentosa</i>	African Caper	LV, BK	Kamba and Luo (10)	Asthma and abdominal	Human fatalities reported following root	Stachydrine		Duke <i>et al.</i> , 2002
	<i>Maerua edulis</i>		RT	Taita (10)	Syphilis and gonorrhoea	Deaths reported	Unknown		Kokwaro 1993
Caesalpinaceae	<i>Senna occidentalis</i> (formerly	Coffee Senna	RT	Nandi (1).	Stomach ache, diarrhea	Liver and muscle degeneration especially in animals	Unknown		Yadav <i>et al.</i> , 2010 Barbosa <i>et al.</i> ,2005 Ferreiro <i>et al.</i> , 2009



	<i>Cassia occidentalis</i>)								
Celastraceae	<i>Catha edulis</i>	Khat	RT	Used nationally	As a recreational drug. RT: Amebas, gonorrhoea, sexually transmitted infections	Cardiovascular effects	Cathinone	Getahun <i>et al.</i> , 2010 Duke <i>et al.</i> , 2002 Aronson <i>et al.</i> , 2009	
	<i>Tithonia diversifolia</i>	Tithonia	LV, RT	Luo (7); Kikuyu, Meru, Mbeere (7). Eastern	Stomach ache, headache.	Hematological and acute toxicity of liver and kidney †	Unknown	Elifioye <i>et al.</i> , 2009 Duke <i>et al.</i> , 2002	
	<i>Bidens pilosa</i>	Spanish Needles	LV	Luo (1), Meru and . Mbeere	Stomach ache,	Genotoxicity	Unknown	Elifioye <i>et al.</i> , 2009	
	<i>Senecio sp. (S. discifolius, S. lyratipartitus, S. syringifolius)</i>		RT, LV	Meru, Luo, Digo (10)	Emetic, cold, stomach ache in pregnant women	Genotoxicity (veno-occlusive disease)	Pyrrrolizidine alkaloids	Ernest and Pittler 2003	
Euphorbiaceae	<i>Euphorbia tirucalli</i>	Pencil Tree	LV	Luo (8); Kalenjin (2); Marakwet (10)	(Chira) Unexplained weight loss.	Kerato – conjunctivitis , uveitis carcinogenic	Ingenane – and tigliane – tvne	Donald., 2008	
	<i>Europhobia candelabrum</i>	Candelabra Cactus	ST	Luo, Kipsigis, Maasai (10)	Pregnant women	Kerato – conjunctivitis, dermatitis, uveitis	Diterpene hydrocarbon	Donald., 2008	



	<i>Croton macrostachyus</i>		LV	Luyha, Meru, Kikuyu and Kamba (10)	LV: Anti-helminthic, Malaria and	Phorbol esters	Diterpene hydrocarbons	Kokwaro, 1993
	<i>Jatropha circus</i>	Bubble Bush or Puraina	LV, ST	Giriama (10)	Kidney disease	Amnesia, convulsions, delirium, diarrhea, nausea, vertigo and	Curcin	Donald., 2008
	<i>Ricinus communis</i>	Castor	LV	South Coast (4); Kikuyu ; Akamba (5); Samburu (10); Meru and Luo (10)	Malaria, Fever	Ribosome inactivation, hypersensitivity reactions , Genotoxic†	Ricin (mostly in seeds)	Aronson <i>et al.</i> , 2009 Fennell <i>et al.</i> , 2004
Fabaceae	<i>Abrus precatorius</i>	Jequirity	LV, BK, RT	Luo (8); Eastern Kenya (5)	Cough, cold, Gonorrhoea, stomach ache	Ribosome inhibitor†	Abrin (mostly in seeds)	Ferreiro <i>et al.</i> , 2009
Meliaceae	<i>Azadirachta indica</i>	Neem	LV	Luo (8); Akamba (5); Meru, Mbeere (7), Kikuyu (10), Maasai (6)	Malaria, skin, rashes, Rash	Nausea, diarrhea, vomiting, drowsiness, tachypnea†	Azadirachtin compounds	Boeke <i>et al.</i> , 2004
	<i>Melia azedarach</i>	Chinaberry	LV	Luo (8); Kikuyu (10)	Measles,	Muscle, contractions, weakness, ataxia, paresis, abdominal distress,	Tetranortriterpe ne limonoid compounds(mel	Ferreiro <i>et al.</i> , 2009



	<i>Trichilia emetica</i>	Trichilia	BK, SP	Meru and Mbeere (7)	Kidney problems, skin rashes	Genotoxicity	Unknown	Fennell <i>et al.</i> , 2004 Duke <i>et al.</i> , 2002
Myrtaceae	<i>Psidium guajava</i>	Guava	LV	Kikuyu(10); Luo (8)	ENT infections, Diarrhea, gingivitis	Hypersensitivity reaction. Should not be used for a period exceeding 30 days†	Hydrosable tannins, essential oils, terpenes and Estragole	WHO 1999
Lamiaceae	<i>Ocimum basilicum</i>	Sweet Basil	LV	Kikuyu(10); Akamba(5); Maasai (3); Miji Kenda (4)	ENT infections (Bronchial Cattarh), Malaria	Potentially carcinogenic	Estragole	Duke <i>et al.</i> , 2002
Leguminosae	<i>Cassia didimobotrya</i>		LV,ST, RT	Kamba, Luyha, Meru, Luo (10)	purgative	GIT irritation	Sennosides	Donald., 2008
Liliaceae	<i>Aloe vera</i>	Aloe	LV	Used nationally	Swelling, boils, diarrhea, burns. Forms part of a large number of formulations sold nationally.	Overdose is associated with diarrhea †. Interaction with other drugs is also suspected	Anthraquinone compounds	Rodriguez-Fragoso <i>et al.</i> , 2008
Poaceae	<i>Cymbopogon</i>	Lemon	LV	Luo (8); Kikuyu	Headache,	Associated with GIT	Unknown	Fandohan <i>et al.</i> , 2008



	<i>citratus</i>	Grass		(10)	purify blood, clean pores	irritation, potentially neurotoxic †		
Rhamnaceae	<i>Rahmnus prinoides</i>			Kikuyu (10)	Malaria	Potentially genotoxic	Unknown	Fennell <i>et al.</i> , 2004
Rosaceae	<i>Prunus africana</i>	Pygeum	BK	Meru and Mbere (6); Maasai (6)	Cancer. Typhoid	Potentially Genotoxic	Docosanol, β -sitosterol,	WHO 2002
	<i>Hagenia abyssinica</i>	Kouso	LV	Nandi (1); Meru (7); Kikuyu (10)	Eye infections	Retinotoxicity	Unknown	Arbo <i>et al.</i> , 2009
Solanaceae	<i>Datura stramonium</i>	Jimsonweed	ST	Kikuyu (10). Swahili (10)	Tonsillitis	Acute psychosis. May cause peripheral Anticholinergic toxicity (decreased salivation,	Tropane belladonna alkaloids	Phua <i>et al.</i> , 2009
	<i>Withania somnifera (Physalis somnifera)</i>	Ashwagandha	DR	Kikuyu (10); Masaai (6).	ENT infections, Malaria	May potentiate barbiturates, berries may cause severe GIT pain. †	Withanolides I–XI, Withasomniferols A–C, Alkaloids	Duke <i>et al.</i> , 2002
Verbenaceae	<i>Lantana camara</i>	Wild Sage	LV, RT	Kikuyu(10);South coast (4); Luo (8); Maasai (6).	Measles, sore throat, cough, Malaria	Cholestasis, hepatotoxicity and photosensitization†	Lantadene A, B	Donald., 2008 Jordan <i>et al.</i> , 2011

(a) Ethno medical information was obtained from Jeruto *et al.*, 2006¹; Okello *et al.*, 2010²; Bussman *et al.*, 2006³; Nguta *et al.*, 2010⁴; Njoroge *et al.*, 2010⁵; Kigondu *et al.*, 2011⁶; Kirira *et al.*, 2006⁷; Nagata *et al.*, 2011⁸; Kokwaro, 1993¹⁰.

(b). LV: leaves; RT: root; WP: Whole plant; BK: bark; SE: seed; ST: stem; FR: fruit; SP: sap; ENT, Ear nose and throat

(c). † No health hazards known at proper dosage levels.



Hepatotoxicosis

Hepatotoxicity is one of the biggest concerns raised by the indiscriminate use of herbal medicines. Ephedra and pyrrolizidine containing plants including *Crotalaria*, *Heliotropium* and *Senecio* spp. are commonly used in Kenya and have been associated with hepatotoxicity (Huxtable, 1990; Stickel *et al.*, 2005; Willet *et al.*, 2004; Ernst and Pittler., 2003; Seeff *et al.*, 2001). The reported liver pathologies range from mild elevation of liver enzymes to fulminant liver failure requiring liver transplantation (Pak *et al.*, 2004). According to Steenkamp and colleagues, the excessive exposure to pyrrolizidine alkaloids may be responsible for the high rates of liver cancer, veno-occlusive disease (VOD) and cirrhosis seen in Africa (Steenkamp, *et al.*, 2000). In Kenya, *Senecio* species used by the Luo (Kokwaro, 1993) Meru, Mbeere (Kareru *et al.*, 2008), Nandi (Jeruto *et al.*, 2006) and the Miji Kenda (Nguta *et al.*, 2010) are known to contain pyrrolizidine alkaloids. Research in animal's studies indicates that unbound pyrroles are highly reactive hepatocarcinogens (Stickel *et al.*, 2005). Another commonly used plant which may cause liver toxicity is *Lantana camara* which grows in most of the agro-ecological zones in Kenya. While the exact mechanism of toxicity is unknown, animal studies have demonstrated that extracts from the plant can cause widespread changes in mitochondrial, microsomal, and membrane enzymes resulting in cholestasis, hepatotoxicity, and photosensitization (Bevilacqua *et al.*, 2011; Ernst and Pittler, 2003). Human fatalities

have been attributed to ingestion of green berries (Duke *et al.*, 2002).

Gastro-intestinal tract toxicities

Several herbs including *Aloe sp.*, *Cinnamomum caphora*, *Cassia didimobotrya*, *Warbugia salutaris*, *Melia azedarach*, *Turraea mombassana*, *Cocculus hirsutus*, *Olinia rochetiana*, *Pittosporum viridiflorum* are known to contain toxins which are detrimental to the gastrointestinal tract (Duke *et al.*, 2002). Toxalbumins, which are toxic to gastro-intestinal tract, have been isolated from plants such as *Ricinus communis*, *Abrus precatorius*, *Jatropha curcas* and *Jatropha multifida* (Levin *et al.*, 2000). Some studies have shown that administration of the fruit of *Jatropha curcas* increases fetal resorptions when administered soon after implantation (Donald, 2008). Other toxins such as ricin and abrin which are derived from *Ricinus communis* and *Abrus precatorius*, respectively, feature highly on the list of substances which might be used in bioterrorism (Fennell *et al.*, 2004). The compounds are highly toxic, heterodimeric type II ribosome inactivating proteins. Ingestion of meliatoxins, a toxin from ripe berries of *Melia azedarach*, has also been associated with vomiting and diarrhea in children (Ferreiro *et al.*, 2010). *Aloe Vera*, a herb which has found widespread commercial application, contains several biologically active constituents including emodin and anthrone (Duke *et al.*, 2002). Emodin anthrones act as GI tract irritant. Therefore, it is generally contraindicated for patients with inflammatory intestinal diseases such as Crohn disease, ulcerative colitis, and irritable bowel



syndrome or to children less than ten years of age (Ibid).

Cardiovascular system

A number of cardenolide-containing garden plants are used in traditional herbal formularies in Kenya. Over 400 cardiac glycosides have been isolated from the plant kingdom. In Kenya, commonly used plants containing cardiac glycoside include *Thevetia peruviana* and *Nerium oleander* (Nagata *et al.*, 2011; Nanyingi *et al.*, 2008; Kirira *et al.*, 2006). High concentrations of cardiac glycosides are found in seed, stems, and roots, followed by the fruit (Donald, 2008). Overdosage of *Nerium Oleander* may cause arrhythmia, bradycardia, cardiodepression, confusion, cyanosis, diarrhea, headache, hyperkalemia, nausea, neurodepression, stupor and vomiting (Duke *et al.*, 2002). Another herb that is grown in Kenya and used as stimulant is *Catha edulis* (Khat). Habitual consumption of khat may enhance the risk of acute myocardial infarction and hypertension (Getahun *et al.*, 2010). Myocardial necrosis in rabbits fed on *Senna occidentalis* has also been reported (Yadav *et al.*, 2010).

Genotoxicity

A large proportion of plants which are used in traditional medicine in Kenya have *in vitro* mutagenic effects (Elgorashi *et al.*, 2003; Fennell *et al.*, 2004). A recent study in South Africa, which screened more than 50 plants sourced from different parts of the continent for genotoxicity showed that both polar and apolar extracts of some plants maybe genotoxic.

Plants used locally which were shown to be to be highly toxic (caused both DNA damage and chromosomal aberrations) are *Prunus africana*, *Rhamnus prinoides* L'Hér, *Ricinus communis* and *Trichelia emetica* (Fennell *et al.*, 2004). Extracts derived from *Euphorbiaceae sp.* have also been linked to tumorigenesis (Norhanom and Yadav, 1995). According to some scholars, the high prevalence of Burkitt lymphoma in East Africa maybe associated with the commonplace use of these plants in folk medicine (Ibid). The observation is premised on the findings that extracts from some *Euphorbiaceae sp.* have Epstein Barr virus (EBV) – activating capacity (Ibid). Other locally used plants which have been linked to carcinogenesis include anthrone – containing laxatives from *Aloe spp.* While *Aloe* has been identified as a possible risk factor in colorectal cancer, the claim is still controversial since studies in rats suggest that *Aloe* can have antitumorigenic effects (Duke *et al.*, 2002).

Toxicities associated with herb-conventional drug interactions

The concurrent use of herbal preparations alongside conventional drugs raises concerns about possible adverse reactions due to drug-herbal interactions. Some constituents of herbal medicines can induce or inhibit enzymes involved in drug metabolism (Sahoo *et al.*, 2010). Evaluating how phytochemicals and cytochrome P450 interact has been proposed as a possible way of predicting both the therapeutic outcome of herbal medicines and possible side effects (Ibid). Competitive inhibition at the P450 active site



has been proposed as the putative point of herb–drug interaction in these reactions (Ulbright *et al.*, 2008).

At present, the number of studies on this subject is scant. However, several warnings have been issued on the strength of these studies. In Kenya, studies have shown that African potato (*Hypoxis hemerocallidea*) extracts which are sold to HIV patients may inhibit HAART drug metabolism and transport (Mills *et al.*, 2006). Therefore, some experts are of the opinion that the concomitant use African potato extracts and HAART by HIV patients should be discouraged (Izzo, 2004). In addition, studies have shown that use of *Allium sativum* (garlic) supplements may reduce the plasma concentrations of the antiretrovirals (indinavir and saquinavir) (Williamson *et al.*, 2009). Other studies have suggested that *Aloe vera* extracts, (Aloe gel) might have additive effects when used with anti–diabetes drugs (Anuja *et al.*, 2010). Preliminary evidence suggests that aloe gel might lower blood glucose levels and may interact with conventional anti–diabetic drugs (Rodríguez–Fragoso *et al.*, 2008) and increase the risk of hypoglycemia in some patients. Other studies suggest that excessive use of *Aloe* and cardiac glycoside drugs might potentiate diuretic induced potassium loss, increasing the risk of hypokalaemia. Since *Aloe* inhibits thromboxane A2 and prostaglandins; concomitant use with anticoagulants might prolong bleeding time (Ibid).

Conclusion

The toxicity associated with the use of herbs is largely under–investigated and only a few studies have looked at toxicity of some herbs used in Kenya. At the same

time, very few studies have focused on chronic toxicity or interactions with concurrently administered drugs. Therefore, toxicological investigations remain a high priority area. In recognition of this, we suggest that evaluation of toxicity should be included in the registration process. In addition, research has shown that the dangers associated with the use of herbal medicines are not necessarily inherent in the herbs themselves. Contamination (deliberate or otherwise), adulteration with conventional drugs, substitution, misidentification and concomitant herb–drug usage have also been cited as possible sources of danger. In view of this, enforcement of regulation including GMP, GACP and GSP should be prioritized. In addition, continuous monitoring of the quality of herbal medicines sold in the local market should be established and the public should be sensitized to the dangers of indiscriminate use of herbal medicines.

Reference

1. Anuja, M., Nithya, R., Rajamanickam, C. and Madambath, I.; 2010. Spermatotoxicity of a protein isolated from the root of *Achyranthes aspera*: a comparative study with gossypol. *Contraception*. **82**: 385–390.
2. Aronson, J.; 2009. *Meyler's Side Effects of Herbal Medicines*. Amsterdam: Elsevier.
3. Arbo, M., Schmitt, G., Limberger, F., Charão M., Moro, A., Ribeiro, G., Dallegrove, E., Garcia, S., Leal, M. and Limberger, R.; 2009. Subchronic toxicity of *Citrus aurantium* L. (Rutaceae) extract and p–



- synephrine in mice. *Regulatory Toxicology and Pharmacology*. **54**: 114–117.
4. Bandara, V., Weinstein, S., White, J. and Eddleston, M.; 2009. A review of the natural history, toxicology, diagnosis and clinical management of *Nerium oleander* (common oleander) and *Thevetia peruviana* (yellow oleander) poisoning, *Toxicol.* **56**: 273–281.
5. Barbosa – Ferreira, M., Dagli, M., Maiorka, P. and Gorniak, S.; 2005. Sub-acute intoxication by *Senna occidentalis* seeds in rats, *Food and Chemical Toxicology* **43**: 497–503.
6. Barnes, J., Anderson, L. and Phillipson, J.; 2007. *Herbal Medicines*. 3rd ed. London: pharmaceutical press.
7. Bevilacqua, A., Suffredini, I., Romoff, P., Lago, J. and Bernard, I.M.; 2011. Toxicity of apolar and polar *Lantana camara* L. crude extracts in mice, *Research in Veterinary Science*. **90**: 106–115.
8. Boeke, S., Boersma, M., Alink, G., Joop, J., Huis A., Dicke, M. and Ivonne, M.; 2004. Safety evaluation of neem (*Azadirachta indica*) derived pesticides, *Journal of Ethnopharmacology* **94**: 25–41.
9. Bussmann, R.; 2006. Ethnobotany of the Samburu of Mt. Nyiru, South Turkana, Kenya. *Journal of Ethnobiology and Ethnomedicine*. **2**:35.
10. Calapai, G., Firenzuoli, F., Saitta, A., Squadrito, F., Arlotta, M., Costantino, G. and Inferrera G.; 1999. Antiobesity and cardiovascular toxic effects of *Citrus aurantium* extracts in the rat: a preliminary report, *Fitoterapia*. **70**: 586–592.
11. Costa, R., Diniz, A., Mantovani, M. and Jorda, O.; 2008. *In vitro* study of mutagenic potential of *Bidens pilosa* Linne' and *Mikania glomerata* Sprengel using the comet and micronucleus assays, *Journal of Ethnopharmacology*. **118**: 86–93.
12. De smet, P.; 2006 Clinical risk management of herb–drug interaction. *British Journal of Clinical Pharmacology*. **63**: 258–267.
13. Donald, G.; 2008. *Medical Toxicology of Natural substances: Foods, Fungi, Medicinal Herbs, Plants, and Venomous Animals*, New Jersey: A John Wiley and Sons.
14. Duke, J., Bogenschutz–Godwin, M. and Duke, P.; 2002. *Handbook of Medicinal Herbs* 2nd Ed. London: CRC Press.
15. Elifioye, T., Alatise, O., Fakoya, F., Agbedahunsi, J. and Houghton, P.; 2009. Toxicity studies of *Tithonia diversifolia* A. Gray (Asteraceae) in rats. *Journal of Ethno pharmacology* **122**: 410–415.
16. Elgorashi, E., Taylor, J., Maes, J., De Kimpe, N. and Verschaeve.; 2003. Screening of medicinal plants used in South African traditional medicine for genotoxic effects. *Toxicology Letters*. **143**: 195–207.
17. Elvin – Lewis, M.; 2005. Safety issues associated with herbal ingredients. *Advances in Food and Nutrition Research*. **50**: 220–300.
18. Ernst, E. and Pittler, M.; 2003. Systematic review: hepatotoxic events associated with herbal medicinal products' *Alimentary Pharmacology and Therapeutics* **18**: 451–471.



19. Ernst, E.; 2002. Toxic heavy metals and undeclared drugs in Asian herbal medicines. *Trends Pharmacological Science*. **23**:136–9.
20. Fandohan, P., Gnonlonfin, B., Laleye, A., Gbenou, J., Darboux, R. and Moudachirou, M.; 2008. Toxicity and gastric tolerance of essential oils from *Cymbopogon citratus*, *Ocimum grattisum* and *Ocimum basilicum* in Wistar rats, *Food and Chemical Toxicology*. **46**: 2493–2497.
21. Farhat, G., Affara, N. and Gali-Muhtasib, H.; 2001. Seasonal changes in the composition of the essential oil extract of East Mediterranean sage (*Salvia libanotica*) and its toxicity in mice. *Toxicol* **39**: 1601–1605.
22. Ferreiro, D., Orozco, J., Miron, C., Real, T., Harnandez –Moreno, D., Soler, F. and Perez–Lopez, M.; 2010. Chinaberry Tree (*Melia azedarach*) Poisoning in Dog: A case report, *Topics in Companion Animal Medicine*. **25**: 64–67.
23. Fennell, C., Lindsey, K., McGraw, L., Sparg, S., Stafford, G., Elgorashi, E., Grace, O. and Van Staden, J.; 2004. Assessing African medicinal plants for efficacy and safety pharmacological screening and toxicology, *Journal of Ethnopharmacology*. **94**: 205–217.
24. Getahun, W., Gedif, T. and Tesfaye, F.; 2010. Regular Khat (*Catha edulis*) chewing is associated with elevated diastolic blood pressure among adults in Butajira, Ethiopia: A comparative study. *BMC Public Health*. **10**: 1–8.
25. Hamill, F., Apio, S., Mubiru, K., Mosango, M., Bukenya–Ziraba, R. and Maganyi, O.; 2000. Traditional herbal drugs of southern Uganda. *Journal of Ethnopharmacology* **70**: 281–300.
26. Huxtable, R.; 1990. The harmful potential of herbal and other plants products. *Drug Safety* **5**: 126–136.
27. Izzo, A.; 2004. Herb–drug interactions: an overview of the clinical evidence. *Fundamental of Clinical Pharmacology* .**19**: 1–16.
28. Jeruto, P., Lukhoba, C., Ouma, G., Otieno, D. and Mutai, C.; 2006. Herbal treatments in Aldai and Kaptumo divisions in Nandi district, Rift valley Province, Kenya. *African Journal of Complementary and Alternative Medicines*. **5**: 103–105.
29. Jordan, S., Cunningham, D. and Marles, R.; 2010. Assessment of herbal medicinal products: Challenges, and opportunities to increase the knowledge base for safety assessment. *Toxicology and Applied Pharmacology*. **243**: 198–216.
30. Kanyara, J. and Njagi ,N.; 2005. Anti-HIV–1 activities in extracts from some medicinal plants as assessed in an in vitro biochemical HIV–1 reverse transcriptase assay. *Phytotherapy Research* **19**: 287–290.
31. Kareru, P., Keriko, J., Gachanja, A. and Kenji, G.; 2008. Direct detection of Triperpenoid in medicinal plants. *Traditional Complementary and Alternative Medicines*. **5**: 56–60.
32. Kigondu, E., Rukunga, G., Gathirwa, J., Irungu, B., Mwikabe, N., Amalemba, G., Omar, S. and Kirira, P.; 2011. Antiplasmodial and cytotoxicity activities of some selected plants used by the Maasai community, Kenya. *South African Journal of Botany*. **77**: 725–729.



33. Kirira, P., Rukunga, G., Wanyonyi, A., Gathirwa, J., Mathaura, C., Omar, S., Tolo, F., Mungai, G. and Ndiege, I.; 2006. Anti-plasmodial activity and toxicity of extracts of plants used in traditional malaria therapy in Meru and Kilifi Districts in Kenya. *Journal of Ethnopharmacology*. **106**: 403–407.
34. Kokwaro, J.; 1993. *Medicinal Plants of East Africa*. Nairobi: Kenya Literature Bureau.
35. Lambert, J., Leonard, K., Mungai, G., Omindi-Ogaja, E., Gatheru, G., Mirangi, T., Owara, J., Herbst, C., Romana, G. and Lemiere, C.; 2011. *The Contribution of Traditional Herbal Medicine Practitioners to Kenyan Health Care Delivery: Results from community health-seeking behavior vignettes and a traditional herbal medicine practitioner survey*. Washington: World Bank.
36. Latorre, A., Caniceiro, B., Wysocki, H., Haraguchi, M., Gardner, D. and Gorniak, S.; 2011. Selenium reverses *Pteridium aquilinum*-induced Immunotoxic effects. *Food and Chemical Toxicology*. **49**: 464–470.
37. Levin, Y., Sherer, Y., Bibi, H., Schlesinger, M. and Hay, E.; 2000. Rare *Jatropha multiday* intoxication in two children, *Journal Emergence Medicine*. **16** : 173–175.
38. Li, Y., Robert, C. and Martin, I.I.; 2011. Herbal medicine and hepatocellular carcinoma: Applications and Challenges. *Evidence Based Complementary and Alternative Medicine* 1–14
39. Matasyoh, L., Matastoh, J., Wachira, F., Kinyua, M., Anne, W., Muigai, T. and Mukiama, T.; 2008. Antimicrobial activity of essential oils of *Ocimum gratissimum* L. From Different populations of Kenya. *African Journal of Traditional, Complementary and Alternative Medicines*. **5** : 187–193.
40. Mattocks, A.; 1986. Chemistry and toxicology of pyrrolizidine alkaloids. London: Academic Press.
41. Mendonca-Filho, R.; 2006. Bioactive Phytocompounds: New approaches in the phytosciences. Edit. Ahmad, I, Aqil, F., Owais, M. Turning Medicinal plants into drugs. *Modern Phytomedicine* .
42. Mills, E., Dugoua, J-J., Perri, D. and Koren, G.; 2006. Herbal Medicine in Pregnancy and lactation. London: Taylor and Francis Group. Mohamed, A., Eltahir, K., Ali, M., Galal, M., Ayeed, I., Adam, S. and Hamid, O.; 1999 Some pharmacological and toxicological studies on *Balanites aegyptiaca* bark, *Phytotherapy Research* **13**: 439–441.
43. Mobusz, M.; 2006. Forensic Science Handbook of Analytical Separations, Edit. Bogusz, M. J. and Al-Tufail. Toxicological aspects of herbal remedies **6**: 608–610.
44. Muthaura, C., Keriko, J., Derese, S., Yenesew, A. and Rukunga, G.; 2011. Investigation of some medicinal plants traditionally used for treatment of malaria in Kenya as potential sources of antimalarial drugs. *Experimental Parasitology*. **127**: 609–626.
45. Nagata, J., Jew, A., Kimeu, J., Salmen, C., Bukusi E. and Cohen, C.; 2011. Medical pluralism on Mfangano Island: Use of medicinal plants among persons living with HIV/AIDS in Suba District, Kenya. *Journal of Ethnopharmacology*. **135**: 501–509.
46. Nanyingi, M., Mbaria, M., Lanyasunya, A., Wagate, C., Koros, B., Kaburia, H., Munenge, R. and Ogara, W.; 2008. Ethnopharmacological survey of



- Samburu District, Kenya. *Journal of Ethnobiology and Ethnomedicine*. 4:14.
47. Nguta, J., Mbaria, J., Gakuya, D., Gathumbi, P. and Kiama, S.; 2010. Traditional antimalarial phytotherapy remedies used by the South Coast community, Kenya. *Journal of Ethnopharmacology*. **131**; 256–267.
48. Njoroge, N., Kaibui, M., Njenga, P. and Odhiambo, O.; 2010. Utilisation of priority traditional medicinal plants and local people's knowledge on their conservation status in arid lands of Kenya (Mwingi District), *Journal of Ethnobiology and Ethnomedicine*. **6**: 2–8.
49. Norhanom, A. and Yadav, M.; 1995 Tumour promoter activity in Malaysian Euphorbiaceae, *British Journal of Cancer*. **71**: 776–779.
50. Nzioki, P.; 2011 Beware of herbal remedies, warn experts. *Daily Nation, Kenya*. 2011, April, 21.
51. Onyambu, M.; 2011. *Identification and characterization of microbial contaminants of herbal medicines in Kenya*. Research Thesis. Nairobi: Nairobi University.
52. Okello, S., Nyunja, R., Netondo, G. and Onyango, J.; 2010. Ethnobotanical study of medicinal plants used by Saboats of Mt. Elgon. *African Journal of Traditional and Complementary Medicine*. **7**: 1 – 10.
53. Okwemba, A.; 2010. Be warned, herbal prescriptions could be harmful, *Daily Nation* 2010, Feb 7.
54. Pharmacy and Poisons Board.; 2011. www.pharmacyboardkenya.org/ accessed on 10th Oct 2011
55. Pharmacy and Poisons Board.; 2012. www.pharmacyboardkenya.org/ accessed on 10th May 2012.
56. Phua, D. and Heard, Z.; 2009. Dietary supplements and herbal medicine toxicities—when to anticipate them and how to manage them, *International Journal of Emergency Medicine*. **2**: 69–76.
57. Robinson, M. and Zhang, X.; 2011. *The world medicines situation 2011 – Traditional Medicines: Global Situation, Issues and Challenges*. Geneva: WHO.
58. Rodriguez–Fragoso, L., Reyes–Esparza, J., Burchiel S., Herrera–Ruiz, D., Torres, E.; 2008. Risks and benefits of commonly used herbal medicines in Mexico, *Toxicology and Applied Pharmacology*. **227**: 125–135.
59. Reedman, L., Shih R., Hung, O.; 2008. Survival after an Intentional Ingestion of Crushed *Abrus Seeds*. *Western Journal of Emergency Medicine* IX **3**: 157–159.
60. Saad, B., Azaizeh, H., Abu–Hijleh, H. and Said, O.; 2006. Safety of traditional Arab herbal medicine, *eComplementary and Alternative Medicine*. **3**: 433–439.
61. Sahoo, N., Manchikanti, H. and Dey, S.; 2010. Herbal drugs: standards and regulation, *Fitoterapia*. **81**: 462–471.
62. Seeff, L., Lindsay, K., Bruce, R., Thomas, F. and Hoofnagle, J.; 2001. Complementary and alternative medicine in chronic liver disease. *Journal of Hepatology* **34**: 595–602.



63. Snodgrass, W.; 2001 Herbal Products: Risks and benefits of use in children. *Current Therapeutics Research* **62**: 724–737.
64. Steenkamp, V., Stewart, M. and Zuckerman, M.; 2000. Clinical and analytical aspects of pyrrolizidine poisoning caused by South African traditional medicines. *Therapeutic Drug Monitoring*. **22**: 302–306.
65. Stickel, F., Patsenker, E. and Schuppan, D.; 2005. Herbal Hepatotoxicity, *Journal of Hepatology*. **43**: 901–910.
66. Tahiliani, P. and Kar, A.; 2000. *Achyranthes aspera* elevates thyroid hormone levels and decreases hepatic lipid peroxidation in male rats, *Journal of Ethnopharmacology*. **71**: 527–532.
67. Ulbright, C., Chao, W., Costa, D., Rusie-Seamon, E., Weissner, W. and Woods, J.; 2008. Clinical evidence of herb–drug interactions: a systematic review by the natural standard research collaboration. *Current Drug Metabolism*. **9**: 1063–1120
68. Vanherweghem, J., Depierreux, M. and Tielemans, C.; 1993. Rapidly progressive interstitial renal fibrosis in young women: Association with slimming regimen including Chinese herbs. *Lancet*. **341**: 387–391.
69. Wagate, G., Gakuya, W., Nanyingi, M., Njonge, F. and Mbaria, J.; 2008. Antibacterial and Cytotoxic Activity of Kenyan Medicinal Plants, *Mem Inst Oswaldo Cruz* **103**. 650–652.
70. Wanyoike, G., Chhabra, S., Langat-Thoruwa, C. and Omar, S.; 2004. Brine shrimp toxicity and antiplasmodial activity of five Kenyan medicinal plants. *Journal of Ethnopharmacology*. **90**: 129–133.
71. Willet, K., Roth, R. and Walker, L.; 2004. Workshop overview: Hepatotoxicity assessments for botanical dietary supplements, *Oxford Journals Life Sciences and Medicine Toxicological Studies*. **71**: 4–9.
72. Williamson, E., Diver, S. and Baxter, K.; 2009. *Stockleys's Herbal Medicines Interactions*. London: Pharmaceutical Press.
73. Worbs, S., Kohler, K., Pauly, D., Anondet, M–A., Schaer, M., Dorner, M. and Dorner, B.; 2011. *Ricinus Communis* Intoxication in Human and Veterinary Medicine – A Summary of Real Cases. *Toxins* **3**:1332–1372
74. World Health Organisation; 2005. *National policy on traditional medicine and regulation of herbal medicines*: Report of a WHO global survey, ISBN 92 4 159323 7
75. World Health Organisation; 1999. *WHO Monographs on Selected Medicinal Plants*. World Health Organisation, Geneva 1.
76. World Health Organisation; 2002. *WHO monographs on selected medicinal plants*. World Health Organisation, Geneva 2.
77. Yadav, J., Arya, V., Yadav, S., Panghal, M., Kumar, S. and Dhankhar, S.; 2010. *Cassia occidentalis* L.: A review on its ethnobotany, phytochemical and pharmacological profile, *Fitoterapia*. **81**: 223–230.
78. Zhang, J., Wider, B., Shang, H., Li, X. and Ernst E.; 2011. Quality of herbal medicines: Challenges and Solutions. *Complementary Therapeutic Medicine*. **1100**: 1–70.