

<http://dx.doi.org/10.4314/ajtcam.v11i3.19>ANTIBACTERIAL AND ANTI-BIOFILM ACTIVITY OF FLAVONOIDS AND TRITERPENES ISOLATED FROM THE EXTRACTS OF *FICUS SANSIBARICA* WARB. SUBSP. *SANSIBARICA* (MORACEAE) EXTRACTSGbonjubola V. Awolola<sup>1</sup>, Neil A. Koorbanally<sup>1\*</sup>, Hafizah Chenia<sup>2</sup>, Francis O. Shode<sup>3</sup> and Himansu Baijnath<sup>2</sup><sup>1</sup>School of Chemistry, University of KwaZulu-Natal, Durban 4000, South Africa; <sup>2</sup>School of Life Sciences, University of KwaZulu-Natal, Durban 4000, South Africa; <sup>3</sup>Department of Chemistry, University of Zululand, Kwadlangezwa, 3886, South Africa.\*E-mail: [Koorbanally@ukzn.ac.za](mailto:Koorbanally@ukzn.ac.za)

## Abstract

**Background:** *Ficus* species are used in African traditional medicine in the treatment of a wide variety of ailments and diseases such as convulsive disorder, wound healing, gonorrhoea, tuberculosis, diabetes, diarrhoeal infections, dysentery, malaria and HIV. The aim of this study was to isolate the phytochemical constituents in the plant and test them for their antibacterial activity.

**Materials and methods:** The fruits, leaves and stem bark were extracted with organic solvents and the compounds in the extracts separated and purified by column chromatography before being identified by NMR spectroscopy and by comparison of the NMR data against values reported in the literature. The antibacterial activity of the pure compounds and extracts were tested using the disk diffusion method.

**Results:** Three triterpenes and three flavonoids: lupeol acetate (**1**); cycloart-23-ene-3,25-diol (**2**);  $\beta$ -sitosterol (**3**); 5,7,4'-trihydroxyflavan-3-ol (**4**); epicatechin (**5**); and isovitexin (**6**) were isolated in this study. Antimicrobial activity was observed at 8 mg mL<sup>-1</sup> for *Staphylococcus aureus* ATCC 29213 with four of the six isolated compounds, with no activity being observed at 1 – 4 mg mL<sup>-1</sup> against *Escherichia coli* ATCC 25922, *E. coli* ATCC 35218 and *S. aureus* ATCC 43300. Epicatechin (**5**) was found to decrease adhesion of *E. coli* ATCC 25922 and *S. aureus* ATCC 29213. Decreased adhesion of *S. aureus* ATCC 29213 was also observed with 5,7,4'-trihydroxyflavan-3-ol (**4**) and isovitexin (**6**).

**Conclusions:** The results of this study provide baseline information on *F. sansibarica*'s potential validity in the treatment of infections associated with Gram-positive microorganisms.

**Key words:** Moraceae, *Ficus sansibarica*, antimicrobial, anti-adhesion properties, flavonoids, triterpenoids.

## Introduction

Primary healthcare in Africa still remain inaccessible to the vast majority of its populace due to inadequate transport systems to and from hospitals and clinics; the cost of drugs and the high rates charged for in-hospital treatment. In most cases, ~80% of the world's population, especially those living in the rural areas of African countries, rely on traditional medicine for the healing and treatment of various ailments from coughs and colds to HIV, malaria and tuberculosis (WHO, 1996; Street and Prinsloo, 2013). Research into pharmacologically active African medicinal plants is being carried out with the aim of providing scientific evidence for the use of the plants in traditional medicine, commercialization and pharmaceutical application of the active components. This has primarily centered on antimicrobial activity but is now being targeted towards microbial virulence mechanisms such as biofilm formation and/or quorum sensing. Microbial biofilm formation of microorganisms is associated with persistent tissue and foreign body infections which are resistant to treatment with antimicrobial agents. Up to 80% of human bacterial infections are biofilm-associated; with infections most frequently being caused by *Staphylococcus aureus*, *S. epidermidis*, *Pseudomonas aeruginosa*, and *Escherichia coli* (Römling and Balsalobre, 2012).

In tropical and subtropical regions of the world, the genus *Ficus* (Moraceae), of the Mulberry family constitutes one of the largest genera of angiosperms with over 900 species consisting of trees, shrubs and epiphytes (Lansky and Paavilainen, 2011). Among these, about 110 *Ficus* species are indigenous to Africa (Berg and Wiebes, 1992; Singh *et al.*, 2011). Many *Ficus* species have been used in African traditional medicine either in the treatment or management of a variety of ailments and diseases such as convulsive disorder, wound healing, gonorrhoea, tuberculosis, diabetes, diarrhoeal infections, dysentery, malaria and HIV (Wakeel *et al.*, 2004; Yadav *et al.*, 2006; Parekh and Chanda, 2007; Annan and Houghton, 2008; Abdulla *et al.*, 2010; Lamorde *et al.*, 2010; Nadembega *et al.*, 2011). *Ficus* species have also been used in Indian ayurvedic and traditional medicine (Joseph and Justin Raj, 2010). Extracts of *Ficus* species have also been documented to have antioxidant, anti-diabetes, antibacterial, antifungal, antiviral, anti-protozoal, anticancer, cytotoxic, anti-ulcer, anti-inflammatory, anti-hyperglycemic, anti-diarrhoea, hepato-protective, muco-protective and gastro-protective activity (Kuete *et al.*, 2009). Phytochemical studies on species from the genus *Ficus* have shown these plants to be rich sources of flavonoids, lignans, terpenoids, alkaloids, coumarins steroids, and ceramides (Ragasa *et al.*, 1999; Kuete *et al.*, 2008; Ramadan *et al.*, 2009; Ueda *et al.*, 2009; Chen *et al.*, 2010; Kiem *et al.*, 2011; Naressi *et al.*, 2012). However, although there are many phytochemical and pharmacological reports on species within the genus *Ficus*, there are many species that have not been studied and whose ethno-botanical relevance is yet to be investigated. *Ficus sansibarica* Warb. subsp. *sansibarica* is a large tree with a spreading crown which is up to 20 × 30 m and is commonly known as the Zanzibar fig, Angola fig or knobbly fig. It is usually without aerial roots and contains milky latex. The knobby fig is distributed in South Africa (Mpumalanga and Northern Province), Swaziland, Mozambique, Zimbabwe, Zambia, Malawi, Tanzania and Kenya. It has also been visually recorded in Zululand (Burrows and Burrows, 2003). In Botswana, both adults and children eat the fresh ripe fruits (Daniel and Topo, 2012). The ripe fruits are also consumed by birds, fruit bats, baboons, monkeys and other animals as well as insectivorous birds which feed upon the wasps that pollinate the figs. The leaves are browsed by nyala, kudu, elephant and giraffe (Venter and Venter, 1996; Boon, 2010). The only biological study reported so far was the analysis of the fresh ripe fruits for its nutrients, total phenols and antioxidant activity (Daniel and Topo, 2012). This prompted us to investigate the phytochemical composition of *Ficus sansibarica* which to the best of our knowledge has not been studied previously, and to test the isolated compounds for antimicrobial and anti-biofilm activity with the aim of providing a scientific rationale for the use of the plant in the treatment of some bacterial infections.

## Materials and Methods

### Collection of plant materials

The fruits, leaves and stem bark of *Ficus sansibarica* were collected in 2011, in KwaZulu-Natal, South Africa. The plant was identified and a voucher specimen (G. V Awolola & H. Baijnath 1), was deposited in the Ward herbarium at the University of KwaZulu-Natal.



<http://dx.doi.org/10.4314/ajtcam.v11i3.19>

### Antimicrobial susceptibility testing

Antimicrobial susceptibility of the crude MeOH leaf and stem bark extracts and compounds **1-6** were determined using the disk-diffusion method. Crude extracts and isolated compounds were dissolved in chloroform or DMSO to a final concentration of 100 mg mL<sup>-1</sup>. Blank discs (6 mm; MAST, UK), were impregnated with 10 µL (1 mg), 20 µL (2 mg) and 40 µL (4 mg) and allowed to dry.

Bacterial isolates (Gram-negative: *Escherichia coli* ATCC 25922 and *E. coli* ATCC 35218; Gram-positive: *Staphylococcus aureus* ATCC 29213 and *S. aureus* ATCC 43300), were grown overnight on tryptic soy agar (TSA) plates. Cell suspensions were prepared with sterile de-ionized water and the turbidity adjusted equivalent to that of a 0.5 McFarland standard. These were used to inoculate Mueller-Hinton (MH) agar plates by streaking swabs over the entire agar surface followed by the application of the respective phytochemical extract discs (CLSI, 2007). Plates were then incubated for 24 h at 37 °C. Testing was done in duplicate with tetracycline (TE30; 30 µg mL<sup>-1</sup>), and ampicillin (AMP10; 10 µg mL<sup>-1</sup>) discs being used as standard antimicrobial agent controls, in addition to DMSO and chloroform.

Following the preliminary disk-diffusion assays, the agar-well diffusion assay was also used for *S. aureus* ATCC 29213, in order to accommodate a greater concentration of the isolated compounds. Instead of using impregnated discs, agar wells with a diameter of ~6 mm were cut into MH agar plates swabbed with *S. aureus* ATCC 29213. Wells were filled with 80 µL of 100 mg mL<sup>-1</sup> stock solutions of compounds **1, 2, 4 and 5** (8 mg mL<sup>-1</sup>), and 80 µL of 20 mg mL<sup>-1</sup> stock solution of compound **6** (1.6 mg mL<sup>-1</sup>) (Table 2). Plates were then incubated for 24 hrs at 37 °C. Testing was done in duplicate and tetracycline (TE30) and ampicillin (AMP10), discs were used as standard antimicrobial agent controls, in addition to DMSO and chloroform as negative controls. Zone diameters were determined and averaged. The following zone diameter criteria were used to assign susceptibility or resistance to the phytochemicals tested: Susceptible (S) ≥ 15 mm, Intermediate (I) = 11 – 14 mm, and Resistant (R) ≤ 10 mm (Okoth *et al.*, 2013). Criteria for assigning susceptibility or resistance to AMP10 was as follows: (S) ≥ 17 mm, (I) = 14 – 16 mm, (R) ≤ 13 mm, while those for TE30 were: (S) ≥ 19 mm, (I) 15 – 18 mm, (R) ≤ 14 mm (CLSI, 2007).

### Biofilm microtitre plate assays

In order to ascertain the anti-biofilm potential of the isolated compounds, *E. coli* ATCC 29922, *E. coli* ATCC 35218, *S. aureus* ATCC 29213 and *S. aureus* ATCC 43300 were exposed to lupeol acetate **1** (100 mg mL<sup>-1</sup> stock; 2, 5, 10 and 15 mg mL<sup>-1</sup>); 5,7,4'-trihydroxyflavan-3-ol **4** (100 mg mL<sup>-1</sup> stock; 1 and 2.5 mg mL<sup>-1</sup>); epicatechin **5** (100 mg mL<sup>-1</sup> stock; 1, 2.5, 5 and 7.5 mg mL<sup>-1</sup>); and isovitexin **6** (20 mg mL<sup>-1</sup> stock; 0.2 and 0.5 mg mL<sup>-1</sup>) using microtitre plate assays (Basson *et al.*, 2008). The purified cycloart-23-ene-3,25-diol **2** could not be tested due to unavailability of sufficient material. All isolates were cultured overnight in tryptic soy broth (TSB), and centrifuged for 2 min at 12000 rpm. Cell pellets were washed and re-suspended to a turbidity equivalent to a 0.5 McFarland standard. Each well of the sterile 96-well U-bottomed polystyrene micro-titre plates contained 90 µL of TSB medium and 10 µL of culture after standardization. To investigate the effect on adherence, respective volumes of extracts were added to the TSB medium at the time of inoculation in the micro-titre plate assays and volumes were standardized to 200 µL with sterile distilled water. Positive control wells contained TSB broth and respective cultures without the tested compounds whilst the negative control wells contained un-inoculated TSB broth only to test for sterility and non-specific binding of media. Plates were incubated aerobically for 24 h with agitation on an Orbit P4 micro-titre plate shaker (Labnet), at 37 °C to allow cell attachment and biofilm formation (Basson *et al.*, 2008).

Following incubation, the crystal violet assay was performed. The contents of each well were aspirated. Micro-titre plates were washed three times with sterile de-ionized water to remove loosely attached cells. The remaining cells were fixed with 200 µL of MeOH for 15 min and air-dried. Following drying, the wells were stained with 150 µL of 2% Hucker's crystal violet for 5 min. Thereafter, wells were rinsed gently with water, air-dried and 150 µL of 33% (v/v) of glacial acetic acid was added for re-solubilization of dye bound to adherent cells. Plates were then read using the GloMax®-Multi+ Detection System (Promega) at an optical density of 600 nm. Tests were done in triplicate on two separate occasions and the results were averaged (Basson *et al.*, 2008). OD<sub>600 nm</sub> values of treated cells were compared to untreated cells to investigate the increase or decrease in adherence as a result of phytochemical exposure. Treated and untreated samples were compared statistically using Paired *t*-tests (SigmaStat V3.5, Systat Software, Inc).

## Results and Discussion

The phytochemical investigation of the leaves, stem bark and fruits of *F. sansibarica* yielded six compounds: three triterpenoids and three flavonoids. The non-polar hexane extract of both leaves (**1** and **2**) and stem bark (**3**) contained triterpenoids while the flavonoids were isolated from the more polar EtOAc extract of both the stem bark (**4** and **5**), and the fruits (**6**). The isolated compounds were identified based on their <sup>1</sup>H and <sup>13</sup>C NMR spectra and 2D NMR spectra and compared with literature values. Thus, they were identified as lupeol acetate **1** (Mahato and Kundu, 1994), cycloart-23-ene-3,25-diol **2** (Abdel-Monem *et al.*, 2008; Escobedo-Martínez *et al.*, 2012; Khan *et al.*, 2006), sitosterol **3** (Chaturvedula and Prakash, 2012), 5,7,4'-trihydroxyflavan-3-ol **4** (Dafalla, 2005; Iida *et al.*, 2007), epicatechin **5** (Hubert *et al.*, 2011; Markham and Ternai, 1976) and isovitexin **6** (Lin *et al.*, 2009).

Epicatechin **5** has been isolated from several *Ficus* species like *F. microcarpa* and *F. spragueana* (Kiem *et al.*, 2011; Ragab *et al.*, 2013). In contrast, isovitexin **6** was only reportedly isolated from *F. microcarpa* (Kiem *et al.*, 2011), while 5,7,4'-trihydroxyflavan-3-ol **4** was isolated from *F. capensis* and *F. spragueana* (Owolabi *et al.*, 2009; Ragab *et al.*, 2013). Lupeol acetate **1** and sitosterol **3** are ubiquitous to plants. To the best of our knowledge, this is the first report of cycloart-23-ene-3,25-diol **2**, a cycloartane type triterpenoid with molecular formula C<sub>30</sub>H<sub>50</sub>O<sub>2</sub> in the genus *Ficus*. Furthermore, a comparison of the NMR data obtained for **2** with that published in the literature (Abdel-Monem *et al.*, 2008; Escobedo-Martínez *et al.*, 2012; Khan *et al.*, 2006) matched the NMR data of the cycloartane backbone only and there were discrepancies in the NMR data of the side chain, which we have unambiguously assigned from HMBC data. The *cis* isomer of **2** is also reported (Gherraf *et al.*, 2010), and is also included here for comparison. We speculate that the reports of **2** in Abdel-Monem *et al.* (2008), Escobedo-Martínez *et al.* (2012), and Khan *et al.* (2006), are also that of the *cis* isomer as the H-23 and H-24 resonances were always reported as overlapping multi-plets at δ<sub>H</sub> 5.60. In our spectra, we were able to clearly define the *trans* nature of the δ<sup>23</sup> double bond by clear coupling constants. A full NMR structural elucidation is given below and a comparison with that in the literature given in Table 1.

The <sup>1</sup>H-NMR spectrum of compound **2** displayed two pairs of doublets with resonances at δ<sub>H</sub> 0.55 (1H, d, *J* = 3.4 Hz, H-19<sub>a</sub>) and at 0.30 (1H, d, *J* = 3.4 Hz, H-19<sub>b</sub>), characteristic of cycloartane-type triterpenes. In addition, there were the presence of seven methyl groups at δ 0.82 (s, CH<sub>3</sub>-29), 0.89 (d, *J* = 6.5 Hz, CH<sub>3</sub>-21), 0.90 (s, CH<sub>3</sub>-28), 0.99 (s, CH<sub>3</sub>-30, CH<sub>3</sub>-18), 1.36 (s, CH<sub>3</sub>-26/27), the typical H-3 resonance for triterpenoids at δ 3.26, a doublet of doublet of doublets (ddd) at δ<sub>H</sub> 5.71 (1H, *J* = 15.7, 8.5, 6.1 Hz, H-23), a doublet at δ<sub>H</sub> 5.53 (1H, *J* = 15.7 Hz, H-24) and one proton of the H-22 methylene group at δ 2.26 which could clearly be distinguished. The <sup>13</sup>C NMR spectrum was able to be

<http://dx.doi.org/10.4314/ajtcam.v11i3.19>

resolved into 7 methyl, 10 methylene, 7 methine and 6 singlet carbon resonances, including two olefinic methine resonances for the double bond in the side chain, an oxygenated singlet resonance at  $\delta$  82.3 and an oxygenated methine resonance at  $\delta$  78.8, indicating two hydroxy groups in the molecule.

**Table 1:**  $^1\text{H}$  and  $^{13}\text{C}$  NMR data of cycloart-23-ene-3,25-diol **2** (600 MHz,  $\text{CDCl}_3$ )

Pos.	$^1\text{H}^a$	$^1\text{H}^{bc}$	$^1\text{H}$ (2)	$^{13}\text{C}^a$	$^{13}\text{C}^b$	$^{13}\text{C}$ (2)
1	1.56 (m), 1.24 (m)	1.30 (m), 0.90 (m)	1.58 (m), 1.26 (m)	31.98	31.93	31.96
2	1.76 (m), 1.56 (m)	1.40 (m), 1.23 (m)	1.78 (m), 1.58 (m)	30.40	30.33	30.38
3	3.24 (m)	3.10 (m)	3.26 (m)	78.86	78.80	78.84
4	-	-	-	40.50	40.45	40.48
5	1.30 (m)	0.91 (m)	1.30 (m)	47.11	47.10	47.09
6	1.59 (m), 0.79 (m)	1.26 (m), 0.45 (dd, 12.5, 8)	1.60 (m), 0.80 (m)	21.12	21.08	21.09
7	1.33 (m), 1.08 (m)	1.65 (m), 1.40 (m)	1.35 (m), 1.10 (m)	26.02	28.04	25.99
8	1.51 (m)	1.16 (dd, 5, 13)	1.51 (m)	47.98	48.00	47.94
9	-	-	-	20.02	20.00	19.98
10	-	-	-	26.11	25.96	26.10
11	1.99 (m), 1.20 (m)	0.77 (m, 2H)	2.00 (m), 1.15 (m)	26.46	26.07	26.45
12	1.61 (m, 2H)	0.93 (m, 2H)	1.63 (m, 2H)	32.80	35.55	32.80
13	-	-	-	45.32	45.29	45.33
14	-	-	-	48.83	48.80	48.83
15	1.30 (m, 2H)	1.27 (m, 2H)	1.30 (m, 2H)	35.59	32.76	35.56
16	1.91 (m), 1.30 (m)	1.60 (m, 2H)	1.90 (m), 1.30 (m)	28.09	26.42	28.08
17	1.57 (m)	1.24 (m)	1.58 (m)	52.02	52.00	52.07
18	0.97 (s, 3H)	0.62 (s, 3H)	0.99 (s, 3H)	18.10	18.04	18.07
19	0.57 (d, 4.2), 0.34 (d, 4.2)	0.22 (d, 4.5), 0.01 (d, 4.5)	0.57 (d, 3.9), 0.35 (d, 3.9)	29.90	30.00	29.86
20	1.46 (m)	1.10 (m)	1.50 (m)	36.40	36.66	36.29
21	0.86 (d, 6.5, 3H)	0.53 (d, 3.5, 3H)	0.89 (d, 6.5, 3H)	18.30	18.25	18.22
22	2.18 (m), 1.74 (m)	1.88 (m)	2.26 (m), 1.81 (m)	39.05	39.00	39.36
23	5.59 (m)	5.26 (d, 8.0)	5.71 (ddd, 15.7, 8.5, 6.06)	125.63	139.31	130.74
24	5.60 (m)	5.26 (d, 8.0)	5.53 (d, 15.7)	139.36	125.57	134.45
25	-	-	-	70.77	70.75	82.30
26	1.31 (s, 3H)	0.98 (s, 3H)	1.36 (s, 3H)	30.00	29.66	24.40
27	1.32 (s, 3H)	0.98 (s, 3H)	1.36 (s, 3H)	29.90	29.83	24.34
28	0.89 (s, 3H)	0.56 (s, 3H)	0.89 (s, 3H)	19.31	19.26	19.30
29	0.81 (s, 3H)	0.42 (s, 3H)	0.82 (s, 3H)	14.02	13.9	13.99
30	0.97 (s, 3H)	0.62 (s, 3H)	0.98 (s, 3H)	25.45	25.41	25.43

<sup>a</sup> Escobedo-Martinez et al. (2012); <sup>b</sup> Gherraf et al. (2010),  $\delta^{23}$  *cis* isomer; <sup>c</sup> It is highly likely that the  $^1\text{H}$  NMR spectrum in Gherraf et al. (2010) is offset by approximately  $\delta$  0.33; coupling constants are given in parenthesis.

The splitting pattern of the two olefinic protons, which appeared as a ddd at  $\delta_{\text{H}}$  5.71 ( $J = 15.7, 8.5, 6.1$  Hz, H-23) and doublet at  $\delta_{\text{H}}$  5.53 ( $J = 15.7$  Hz, H-24) arose as a result of H-23 coupling with H-24, and H-23 coupling with the two diastereotopic H-22 methylene protons. These coupling constants clearly indicated that the double bond at  $\delta^{23}$  was *trans* because of the large  $J_{23,24}$  value of 15.7 Hz, typical of *trans* olefinic protons. Coupling in the COSY spectrum between H-23 and H-24 and between H-23 and the two H-22 proton resonances at  $\delta$  2.26 and 1.81, both multiplets was also present. The corresponding carbon resonances of C-23 and C-24 were identified at  $\delta$  130.74 and 134.45 respectively. The C-23 carbon resonance was used to identify the 3H-26/27 methyl resonances through a HMBC correlation with  $\delta$  1.36 and in turn the C-25 oxygenated singlet resonance at  $\delta$  82.30 through a HMBC correlation between  $\delta$  1.36 and  $\delta$  82.30. The H-23 proton resonance also showed a HMBC correlation to C-25. The doublet of 3H-21 at  $\delta$  0.89 showed HMBC correlations to C-17, C-22 and C-20 at  $\delta$  52.07, 39.36 and 36.29 respectively, completing the side chain assignment and linking it to the core skeleton.

The rest of the proton and carbon resonances were confirmed by COSY and HMBC correlations and matched well with that in Escobedo-Martinez et al. (2012).

### Antimicrobial susceptibility testing

The isolated compounds and extracts were subjected to antimicrobial screening and the results are presented in Table 2. Initial disk-diffusion assays with 1 – 4 mg mL<sup>-1</sup> of the six isolated compounds, as well as the crude methanolic leaf and stem-bark extracts proved ineffective against the four bacterial strains (Gram-negative:  $\beta$ -lactam susceptible *E. coli* ATCC 25922 and  $\beta$ -lactam resistant *E. coli* ATCC 35218; Gram-positive: methicillin-susceptible *S. aureus* ATCC 29213 and methicillin-resistant *S. aureus* ATCC 43300) tested, with no zones of inhibition being observed (Table 2). However, *S. aureus* ATCC 29213 demonstrated intermediate susceptibility to 8 mg mL<sup>-1</sup> for four of the compounds: lupeol acetate **1**, cycloart-23-ene-3,25-diol **2**, sitosterol **3** and epicatechin **5**, using the agar-well diffusion assay (Table 2). However, the flavonoids, 5,7,4'-trihydroxyflavan-3-ol **4** and isovitexin **6** were not active against the bacteria being tested (Table 2). Most phytochemicals work best against Gram-positive bacteria and there are very limited small-molecule plant-derived antimicrobial compounds with high activity against Gram-negative species (Lewis, 2013).

The limited inhibition observed for the crude MeOH extracts is not surprising. Reports on the antimicrobial activity of extracts from other *Ficus* species have been variable, with differences related to the *Ficus* species tested, the part of the plant sampled, the extraction solvent and the method of antimicrobial testing (Salem et al., 2013). Lawal et al. (2012) observed *F. exasperata* root bark MeOH extract antimicrobial activity against three *S. aureus* strains at  $\geq 12.5$  mg mL<sup>-1</sup>, while activity against *E. coli* strains required  $\geq 25$  mg mL<sup>-1</sup>. They also noted the potency of the MeOH extract compared to the EtOAc extract. Murti and Kumar (2011) demonstrated that the *F. racemosa* ethanolic root extract

<http://dx.doi.org/10.4314/ajtcam.v11i3.19>

(25, 50 and 75 mg mL<sup>-1</sup>) showed maximum inhibition against *S. aureus* when compared with *F. benghalensis* ethanolic root extract. Crude extracts and compounds from *Ficus conraui* stem barks also demonstrated selective antimicrobial activity varying from weak to moderate (Kengap et al., 2011). In contrast, MICs of  $\geq 625 \mu\text{g mL}^{-1}$  were obtained with *F. ovata* stem bark MeOH extracts from *S. aureus* and *E. coli*. (Kueté et al., 2009), while *F. polita* MeOH root extract and euphol-3-*O*-cinnamate and (*E*)-3,5,4'-trihydroxystilbene-3,5-*O*- $\beta$ - $\delta$ -diglucopyranoside MICs ranged from 32 -  $\geq 512 \mu\text{g mL}^{-1}$  against Gram-positive and Gram-negative bacteria (Kueté et al., 2011).

The antimicrobial potencies of the two classes (triterpenes and flavonoids), of compounds isolated have previously been established (Cowan, 1999). Akiyama et al. (2001), obtained epicatechin MICs of 8 mg mL<sup>-1</sup> with 18 clinical *S. aureus* strains, since it causes limited damage to bacterial plasma membranes, while Mahmoud et al. (2013) observed no antimicrobial effect against methicillin-susceptible and -resistant *S. aureus* strains with 32 - 1024  $\mu\text{g mL}^{-1}$  of epicatechin. The reported weak activity of some isolated cycloartane-type triterpenoids from the genus *Aphanamixis* against *S. aureus* (Wang et al., 2013) was consistent with the weak activity observed for cycloart-23-ene-3,25-diol (**2**) in this study.

**Table 2:** Antimicrobial susceptibility profile of *Staphylococcus aureus* ATCC 29213 following exposure to six compounds isolated from *Ficus sansibarica* and two crude methanol extracts

Compound/Extract	Concentration mg mL <sup>-1</sup>	Zone diameter (mm) 40 $\mu\text{L}$ <sup>*</sup>	Zone diameter (mm) 80 $\mu\text{L}$ <sup>#</sup>
Lupeol acetate ( <b>1</b> )	8	0	13.5 (I)
Cycloart-23-ene-3,25-diol ( <b>2</b> )	8	0	11.5 (I)
Sitosterol ( <b>3</b> )	8	0	13 (I)
5,7,4'-trihydroxyflavan-3-ol ( <b>4</b> )	8	0	0 (R)
Epicatechin ( <b>5</b> )	8	0	13.5 (I)
Isovitexin ( <b>6</b> )	1.6	0	0 (R)
MeOH leaf extract	8	0	0 (R)
MeOH stem bark extract	8	0	9 (R)
Ampicillin	10	25 (S)	25 (S)
Tetracycline	30	28 (S)	28 (S)
Chloroform		0	0 (R)
DMSO		0	0 (R)

\* Tested using disk-diffusion assay;

# Tested using agar-well diffusion assay; and


S = susceptible, I = intermediate susceptibility, and R = resistant (CLSI, 2007).

### Biofilm microtitre plate assays

In the face of increasing incidence of antimicrobial resistance and recalcitrance of bacteria to current antimicrobial therapy, drug discovery research is focusing on limiting the pathogenicity mechanisms demonstrated by bacteria. Given the focus on biofilms and their role in microbial pathogenicity, screening for the anti-biofilm activity of phytochemicals is imperative in identifying alternative therapeutic options. The anti-biofilm activity of lupeol acetate **1**; 5,7,4'-trihydroxyflavan-3-ol **4**; epicatechin **5** and isovitexin **6** against *E. coli* ATCC 29922, *E. coli* ATCC 35218, *S. aureus* ATCC 29213 and *S. aureus* ATCC 43300 was variable. Lupeol acetate (triterpene), exposures resulted in statistically significant increased adhesion for all strains (Fig. 2a) at all concentrations tested (2 - 15 mg/ml).

Epicatechin **5** exposures decreased adhesion of antimicrobial-susceptible *E. coli* ATCC 25922 and *S. aureus* ATCC 29213 at all concentrations tested (1 - 7.5 mg mL<sup>-1</sup>), with 5 mg mL<sup>-1</sup> being the most effective (Fig. 2b) indicating that epicatechin has the ability to interfere with the adhesion ability of these microorganisms to polystyrene surfaces. El-Adawi (2012) obtained a 55-66% decrease in *Streptococcus mutans* biofilm formation upon exposure to 2-15% epicatechin. Previous reports have suggested that epicatechin was unable to inhibit biofilm formation as observed for *Eikenella corrodens*, a periodontopathogenic bacterium (Matsunaga et al., 2010) and for *P. aeruginosa* PAO1 biofilm formation after 18 h of growth (Vandeputte et al., 2010). Similarly, Nyila et al. (2012) observed that epicatechin from *Acacia karroo* did not reduce *Listeria monocytogenes* biofilms. The observed anti-biofilm activity of epicatechin was, however, strain-specific as adhesion of  $\beta$ -lactam resistant *E. coli* ATCC 35218 and methicillin-resistant *S. aureus* ATCC 43300 was increased at all concentrations tested (Fig. 2b). Biofilm formation of *P. aeruginosa* PAO1 was enhanced 3- to 7-fold under the action of epicatechin. Epicatechin caused an up to 5-fold enhancement of the *P. aeruginosa* PAO1 biofilm formation at sub-inhibitory concentrations up to 750  $\mu\text{g mL}^{-1}$  (Plyuta et al., 2013).

574',-Trihydroxyflavan-3-ol **4** (1 - 2.5 mg mL<sup>-1</sup>) and isovitexin **6** (0.2 - 0.5 mg mL<sup>-1</sup>) exposures also decreased adhesion of methicillin-susceptible *S. aureus* ATCC 29213 (Figs. 2c - 2d). Exposure to 2.5 mg mL<sup>-1</sup> of **4** almost completely inhibited adhesion of *S. aureus* ATCC 29213 and decreased adhesion of  $\beta$ -lactam susceptible *E. coli* ATCC 25922 (Fig. 2c). No previous studies have examined the effect of lupeol acetate, 5,7,4'-trihydroxyflavan-3-ol **4** and isovitexin **6** on microbial adhesion. The anti-adhesion activity of **4**, **5** and **6** was strain-specific, being most effective against *S. aureus* ATCC 29213, while lupeol acetate (**1**) increased adhesion of all *E. coli* and *S. aureus* strains tested.



**Figure 2:** Anti-adhesion activity of 1 – 15 mg mL<sup>-1</sup> of compounds **1** and **4-6** on biofilm formation by *E. coli* and *S. aureus* strains: (a) lupeol acetate (**1**); (b) epicatechin (**5**); (c) 5,7,4'-trihydroxyflavan-3-ol (**4**); (d) isovitexin (**6**).

## Conclusion

The phytochemical investigation of *Ficus sansibarica* resulted in the isolation of three triterpenes and three flavonoids. The antimicrobial activity of the crude extracts and the isolated compounds were weak with a narrow-spectrum activity against Gram-positive bacteria. Most phytochemicals work best against Gram-positive bacteria and there are very limited small-molecule plant-derived antimicrobial compounds with high activity against Gram-negative species. The anti-biofilm activity demonstrated by the flavonoids 574',-trihydroxyflavan-3-ol, epicatechin and isovitexin (**4-6**), is most significant for *S. aureus* ATCC 29213. The results of this study provide baseline information on *F. sansibarica* potential validity in the treatment of Gram-positive biofilm-associated infections, despite its limited antimicrobial potential.

## Acknowledgments

Research funding from UKZN and the National Research Foundation is gratefully acknowledged. Staff supplementation (study leave) from the University of Ilorin, Ilorin, Kwara State, Nigeria is also acknowledged for Gbonjubola V. Awolola.

## References

1. Abdel-Monem, A. R., Abdel-Sattar, E., Harraz, F. M and Petereit, F. (2008). Chemical investigation of *Euphorbia schimperi* C. Presl. Rec. Nat. Prod. **2**(2): 39-45.
2. Abdulla, M. A., Ahmed, K. A., Abu-Luhood, F. M. and Muhanid, M. (2010). Role of *Ficus deltoidea* extract in the enhancement of wound healing in experimental rats. Biomed. Res. **21**(3): 241-245.

<http://dx.doi.org/10.4314/ajtcam.v11i3.19>

3. Akiyama, H., Fujii, K., Yamasaki, O., Oono, T., and Iwatsuki, K. (2001). Antibacterial action of several tannins against *Staphylococcus aureus*. J. Antimicrob. Chemother. **48**: 487-491.
4. Annan, K. and Houghton, P. J. (2008). Antibacterial, antioxidant and fibroblast growth stimulation of aqueous extracts of *Ficus asperifolia* Miq. and *Gossypium arboreum* L., wound healing plants of Ghana. J. Ethnopharmacol. **119**: 141-144.
5. Basson, A., Flemming, L. A. and Chenia, H. Y. (2008). Evaluation of adherence, hydrophobicity, aggregation and biofilm development of *Flavobacterium johnsoniae*-like isolates. Microb. Ecol. **55**: 1-14.
6. Berg, C. C. and Wiebes, J. T. (1992). African fig trees and fig wasps. Royal Netherlands Academy of Arts and Sciences Department of Physics treatises, 2<sup>nd</sup> series, Part 89. North-Holland, Amsterdam.
7. Boon, R. (2010). Pooley's Trees of Eastern South Africa, Flora and Fauna Publications Trust, South Africa.
8. Burrows, J. and Burrows S. (2003). Figs of Southern and South-Central Africa. Umdaws Press, South Africa.
9. Chaturvedula, V. S. P., and Prakash, I. (2012). Isolation of stigmaterol and  $\beta$ -sitosterol from the dichloromethane extract of *Rubus suavisissimus*. Int. Curr. Pharm. J. **1**: 239-242.
10. Chen, L-W., Cheng, M-J., Peng, C-F. and Chen, I-S. (2010). Secondary metabolites and antimycobacterial activities from the roots of *Ficus nervosa*. Chem. Biodiversity. **7**: 1814-1821.
11. Clinical and Laboratory Standards Institute (CLSI) (2007) M100-S17. Performance Standards for Antimicrobial Susceptibility Testing, 17th informational supplement, vol. 27, No. 1, Wayne, PA.
12. Cowan, M. M. (1999). Plant products as antimicrobial agents. Clin. Microbiol. Rev. **12** (4): 564-582.
13. Dafalla, H. A. A. (2005). Studies on the constituents of *Ficus capensis* (Thunb). Pak. J. Social Sci. **3**: 751-754.
14. Daniel, M. and Topo, E. (2012). Analysis of nutrients, total phenols and antioxidant activity of *Ficus sansibarica* warb. fruits from Eastern Botswana. J. Drug Delivery Ther. **2** (6): 1-5.
15. El-Adawi, H. (2012). Inhibitory effect of grape seed extract (GSE) on cariogenic bacteria. J. Med. Plants Res. **6**: 4883-4891.
16. Escobedo-Martínez, C., Lozada, M. C., Hernández-Ortega, S., Villarreal, M. L., Gnecco, D., Enríquez, R. G., and Reynolds, W. (2012). <sup>1</sup>H and <sup>13</sup>C NMR characterization of new cycloartane triterpenes from *Mangnifera indica*. Magn. Reson. Chem. **50**: 52-57.
17. Gherraf, N., Zellagui, A., Mohamed, N. S., Hussien, T. A., Mohamed, T. A., Hegazy, M-E. F., Rhouati, S., Moustafa, M. F. M., El-Sayed, M. A. and Mohamed, A. E-H. (2010). Triterpenes from *Euphorbia rigida*. Pharmacognosy Res. **2** (3): 159-162.
18. Hubert, D. J., Dawe, A., Florence, N. T., Gilbert, K. D. W. F., Angele, T. N., Buonocore, D., Finzi, P. V., Vidari, G., Bonaventure, N. T., Marzatico, F., and Paul, M. F. (2011). *In vitro* hepatoprotective and antioxidant activities of crude extract and isolated compounds from *Ficus gnaphalocarpa*. Inflammopharmacol. **19**: 35-43.
19. Iida, N., Inatomi, Y., Murata, H., Inada, A., Murata, J., Lang, F.A., Matsuura, N and Nakanishi, T. (2007). A new flavone xyloside and two new flavan-3-ol glucosides from *Juniperus communis* var. depressa. Chem. Biodiversity. **4**: 32-42.
20. Joseph, B., and Justin Raj, S. (2010). Phytopharmacological and phytochemical properties of three *Ficus* species - An overview. Int. J. Pharma Bio Sci. **1** (4): 246-253.
21. Kengap, R.T., Kapche, G. D. W. F., Dzoyem, J-P., Simo, I. K., Ambassa, P., Sandjo, L. P., Abegaz, B. M and Ngadjui, B. T. (2011). Isoprenoids and flavonoids with antimicrobial activity from *Ficus conraui* Warburg (Moraceae). Helv. Chim. Acta. **94**: 2231 - 2238.
22. Khan, M. T. H., Khan, S. B., and Ather, A. (2006). Tyrosinase inhibitory cycloartane type triterpenoids from the methanol extract of the whole plant of *Amberboa ramosa* Jafri and their structure-activity relationship. Bioorg. Med. Chem. **14**: 938 - 943.
23. Kiem, P. V., Cuong, N. X., Nhiem, N. X., Thu, V. K., Ban, N. K., Minh, C. V., Tai, B. H., Hai, T. N., Lee, S. H., Jang, H. D., and Kim, Y. H. (2011). Antioxidant activity of a new C-glycosylflavone from the leaves of *Ficus microcarpa*. Bioorg. Med. Chem. Lett. **21**: 633-637.
24. Kuete, V., Kamga, J., Sandjo, L. P., Ngameni, B., Poumale, H. M. P., Ambassa, P., and Ngadjui, B. T. (2011). Antimicrobial activities of the methanol extract, fractions and compounds from *Ficus polita* Vahl. (Moraceae). BMC Complementary Altern. Med. **11**: 1-6.
25. Kuete, V., Nana, F., Ngameni, B., Mbaveng, A. T., Keumedjio, F., and Ngadjui, B. T. (2009). Antimicrobial activity of the crude extract, fractions and compounds from stem bark of *Ficus ovata* (Moraceae). J. Ethnopharmacol. **124**: 556-561.
26. Kuete, V., Ngameni, B., Fotso Simo, C. C., Tankeu, R. K., Ngadjui, B. T., Meyer, J. J. M., Lall, N., and Kuete, J. R. (2008). Antimicrobial activity of the crude extracts and compounds from *Ficus chlamydocarpa* and *Ficus cordata* (Moraceae). J. Ethnopharmacol. **120**: 17-24.
27. Lamorde, M., Tabuti, J. R. S., Obua, C., Kukunda-Byobona, C., Lanyero, H., Byakika-Kibwika, P., Bbosa, G. S., Lubega, A., Ogwali-Okeng, J., Ryan, M., Waako, P. J., and Merry, C. (2010). Medicinal plants used by traditional medicine practitioners for the treatment of HIV/AIDS and related conditions in Uganda. J. Ethnopharmacol. **130**: 43-53.
28. Lansky, E. P., and Paavilainen, H. M. (2011). Traditional Herbal medicines for Modern Times: Figs: The Genus *Ficus*. CRC Press .Taylor and Francis Group LLC. USA. pp. 13-50.
29. Lawal, I. O., Borokini, T. I., Oyeleye, A., Williams, O. A., and Olayemi, J. O. (2012). Evaluation of extract of *Ficus exasperata* Vahl root bark for antimicrobial activities against some strains of clinical isolates of bacterial and fungi. Int. J. Mod. Bot. **2**: 6-12.
30. Lewis, K. (2013). Platforms for antibiotic discovery. Nat. Rev. Drug Discovery **12**: 371-387.
31. Lin, Y-P., Chen, T-Y., Tseng, H-W., Lee, M-H., and Chen, S-T. (2009). Neural cell protective compounds isolated from *Phoenix hanceana* var. formosana. Phytochemistry **70**: 1173-1181.
32. Mahato, S.B and Kundu, A.P. (1994). <sup>13</sup>C NMR Spectra of pentacyclic triterpenoids-A compilation and some salient features. Phytochemistry **37** (6): 1517-1575.
33. Mahmoud, M., Alkhaleefah, F., and Sherif, D. M. (2013). Antimicrobial effects of epigallocatechin gallate and epicatechins of green tea on planktonic and biofilm forms of *Staphylococcus aureus*, including MRSA. Nat. Sci. **11** (6): 70-79.
34. Markham, K. R., and Ternai, B. (1976). <sup>13</sup>C NMR of Flavonoids-II: Flavonoids other than flavone and flavonol aglycones. Tetrahedron. **32**: 2607-2612.
35. Matsunaga, T., Nakahara, A., Minnatul, K. M., Noiri, Y., Ebisu, S., Kato, A., and Azakami, H. (2010). The inhibitory effects of catechins on biofilm formation by the periodontopathogenic bacterium, *Eikenella corrodens*. Biosci. Biotechnol. Biochem. **74**: 2445-2450.
36. Murti, K., and Kumar, U. (2011). Antimicrobial activity of *Ficus benghalensis* and *Ficus racemosa* roots L. Am. J. Microbiol. **2**: 21-24.
37. Nadembega, P., Boussim, J. I., Nikiema, J. B., Poli, F., and Antognoni, F. (2011). Medicinal plants in Baskoure, Kourittenga Province, Burkina Faso: An ethnobotanical study. J. Ethnopharmacol. **133**: 378-395.
38. Naressi, M. A., dos Santos Ribeiro, M. A., Bersani-Amado, C. A., Zamuner, M. L. M., da Costa, W. F., Abe Tanaka, C. M., and Sarragiotto, M. H. (2012). Chemical composition, anti-inflammatory, molluscicidal and free-radical scavenging activities of the leaves of *Ficus radicans* 'Variegata' (Moraceae). Nat. Prod. Res. **26** (4): 323-330.
39. Nyila, M. A., Leonard, C. M., Hussein, A. A., and Lall, N. (2012). Activity of South African medicinal plants against *Listeria monocytogenes* biofilms, and isolation of active compounds from *Acacia karroo*. S. Afr. J. Bot. **78**: 220-227.
40. Okoth, D. A., Chenia, H. Y., and Koorbanally, N. A. (2013). Antibacterial and antioxidant activities of flavonoids from *Lannea alata* (Engl.) Engl. (*Anacardiaceae*). Phytochem. Lett. **6**: 476-481.

<http://dx.doi.org/10.4314/ajtcam.v11i3.19>

41. Owolabi, O. J., Nworgu, Z. A., Falodun, A., Ayinde, B.A., and Nwako, C. N. (2009). Evaluation of tocolytic activity of ethanol extract of the stem bark of *Ficus capensis* thunb. (Moraceae). *Acta Poloniae Pharm.- Drug Res.* **66** (3): 293-296.
42. Parekh, J., and Chanda, S. V. (2007). *In vitro* antimicrobial activity and phytochemical analysis of some Indian medicinal plants. *Turk. J. Biol.* **31**: 53-58.
43. Plyuta, V., Zaitseva, J., Lobakova, E., Zagorskina, N., Kuznetsov, A., and Khmel, I. (2013). Effect of plant phenolic compounds on biofilm formation by *Pseudomonas aeruginosa*. *APMIS*. doi: 10.1111/apm.12083.
44. Ragab, E.A., Mohammed, A. E-S. I., Abbass, H. S., and Kotb, S. I. (2013). A new flavan-3-ol dimer from *Ficus spragueana* leaves and its cytotoxic activity. *Pharmacognosy Mag.* **9** (34): 144-148.
45. Ragasa, C. Y., Juan, E., and Rideout, J. A. (1999). A triterpene from *Ficus pumila*. *J. Asian Nat. Prod. Res.* **1** (4): 269-275.
46. Ramadan, M. A., Ahmad, A. S., Nafady, A. M., and Mansour, A. I. (2009) Chemical composition of the stem bark and leaves of *Ficus pandurata* Hance. *Nat. Prod. Res.* **23** (13): 1218-1230.
47. Römmling, U., and Balsalobre, C. (2012). Biofilm infections, their resilience to therapy and innovative treatment strategies. *J. Intern. Med.* **272**: 541-561.
48. Salem, M. Z. M., Salem, A. Z. M., Camacho, L. M., and Ali, H. M. (2013). Antimicrobial activities and phytochemical composition of extracts of *Ficus* species: An overview. *Afr. J. Microbiol. Res.* **7**: 4207-4219.
49. Singh, D., Singh, B., and Goel, R. K. (2011). Traditional uses, phytochemistry and pharmacology of *Ficus religiosa*: A review. *J. Ethnopharmacol.* **134**: 565-583.
50. Street, R. A., and Prinsloo, G. (2013). Commercially important medicinal plants of South Africa: A review. *J. Chem.* **2013**: 1-16 (Article ID 205048).
51. Ueda, J-y., Takagi, M., and Shin-ya, K. (2009). Aminocarpophenone- and pyrrolidine-type alkaloids from the leaves of *Ficus septica*. *J. Nat. Prod.* **72**: 2181-2183.
52. Vandeputte, O. M., Kiendrebeogo, M., Rajaonson, S., Diallo, B., Mol, A., El Jaziri, M., and Baucher, M. (2010). Identification of catechin as one of the flavonoids from *Combretum albiflorum* bark extract that reduces the production of quorum-sensing-controlled virulence factors in *Pseudomonas aeruginosa* PAO1. *Appl. Environ. Microbiol.* **76**: 243-53.
53. Venter, F., and Venter, J.-A. (1996). Making the most of indigenous trees. Briza Publications, Pretoria.
54. Wakeel, O. K., Aziba, P. I., Ashorobi, R. B., Umukoro, S., Aderibigbe, A.O., and Awe, E. O. (2004). Neuropharmacological activities of *Ficus platyphylla* stem bark in mice. *Afr. J. Biomed. Res.* **7**: 75-78.
55. Wang, X-Y., Tang, G-H., Yuan, C-M., Zhang, Y., Zou, T., Yu, C., Zhao, Q., Hao, X-J., and He, H. (2013). Aphagrandinoids A-D, cycloartane triterpenoids with antibacterial activities from *Aphanamixis grandifolia*. *Fitoterapia* **85**: 64-68.
56. World Health Organisation (1996). WHO Guideline for the Assessment of herbal medicines. WHO Expert Committee on specification for pharmaceutical preparation. Technical Report series no. 863, Geneva.
57. Yadav, J.P., Kumar, S., and Siwach, P. (2006). Folk medicine used in gynaecological and other related problems by rural population of Haryana. *Indian J. Traditional Knowledge.* **5** (3): 323-326.