

Available online/abstracted at <http://www.biolineinternational.org.br/njps>; www.ajol.info/journals/njps; www.cas.org

OSTEOCALCIN AND BONE-SPECIFIC ALKALINE PHOSPHATASE IN SICKLE CELL HAEMOGLOBINOPATHIES

E. C. AZINGE and D. M. BOLARIN*,

Department of Clinical Pathology, College of Medicine, University of Lagos, Lagos, Nigeria.

*Department of Chemical Pathology, College of Medical Sciences,
University of Calabar, Calabar, Nigeria.

Summary: Osteocalcin or bone gamma-carboxyglutamic acid (gla) protein and Bone-specific alkaline phosphatase (b-AP) total protein levels were evaluated as indicators of bone turnover in twenty patients with sickle cell haemoglobinopathies and in twenty normal healthy individuals. The serum bone-specific alkaline phosphatase total protein level was measured by immunoradiometric (IRMA) method. The concentrations of serum bone-specific alkaline phosphatase total protein were higher in the study group than in the control group ($p < 0.05$). The serum osteocalcin (BGP) showed no significant difference with the control healthy subjects. There was no correlation between the serum osteocalcin and serum bone-specific alkaline phosphatase total protein in the patient group. In conclusion, serum bone-specific alkaline phosphatase total protein determined or measured by IRMA can be considered a sensitive marker of bone turnover and could be especially useful as valuable non-invasive biochemical marker for identifying sickle cell patients with bone complications.

Key Words: *Bone-specific alkaline phosphatase (b-AP); Osteocalcin; sickle cell haemoglobinopathies.*

Introduction

Sickle cell haemoglobinopathies are probably the commonest known hereditary blood disorders in Nigeria (David *et al*, 1993; Ballas, 1995). The sickle cell anaemia (SS) is the homozygous state in which the sickle gene is inherited from the father and the mother. Sickle cell disease may also occur in a heterozygous form in conjunction with other haemoglobin beta chain abnormalities. The common heterozygous forms include sickle haemoglobin C disease (HbSC), and sickle cell haemoglobin thalassaemia (HbS β ⁺thal. or HbS β ⁰thal.) (David *et al*, 1993; Ballas, 1995; Ballas 2001; Smith, 1996; Aster, 2004; Iwegbu and Fleming 1985).

Sickle cell disorder may be classified as an uncompensated haemolytic anaemia in which the life span of the erythrocytes is shortened due to increased rate of destruction and this is insufficiently balanced by increase in erythropoiesis. Vascular occlusion plays a major role in the clinical course of sickle cell disorders (Lee *et al*, 1981; Embury *et al*, 1994; Francis and Johnson, 1991; Hebbel, 1991; Powars, 1990). Vascular occlusion may occur in both the microcirculation and the

macrocirculation. When vascular occlusion occurs in the microcirculation it leads to acute sickle cell painful episodes, which is the hallmark of the sickle cell disorder. Vascular occlusion in the macrocirculation is associated organ failure (Lee *et al*, 1981; Embury *et al*, 1994; Francis and Johnson, 1991; Hebbel, 1991; Powars, 1990).

Patients with sickle cell disorders often suffer from both chronic haemolytic anaemia, which causes bone marrow hyperplasia (Lee *et al*, 1981; Embury *et al*, 1994; Francis and Johnson, 1991; Hebbel, 1991; Powars, 1990) and repeated episodes of vaso-occlusion (Lee *et al*, 1981; Embury *et al*, 1994; Francis and Johnson, 1991; Hebbel, 1991; Powars, 1990). The results of infection and infarction, which lead to orthopaedic complications, have been well documented by several authors (Lee *et al*, 1981; Embury *et al*, 1994; Francis and Johnson, 1991; Hebbel, 1991; Powars, 1990).

Avascular necrosis is the commonest and a major chronic complication of sickle cell disorders (David *et al*, 1993; Ballas, 1995; Ballas 2001; Smith, 1996; Mitchell *et al*, 1986; Rao *et al*, 1985; Rao *et al*, 1989; Amundsen *et al*, 1984; Milner *et al*, 1993; Mitchell *et al*

1987). The sickle cell disease is a cause of hip deformities in the patients (David *et al*, 1993; Ballas, 1995; Ballas 2001; Smith, 1996; Lee, *et al*, 1981; Embury *et al*, 1994; Francis and Johnson, 1991; Hebbel, 1991; Powars, 1990). Involvement of the humeral head is also very common (Rao *et al*, 1985; Rao *et al*, 1989; Amundsen *et al*, 1984; Milner *et al*, 1993; Mitchell *et al*, 1987). More than 30 percent of patients with sickle cell disorders end up with complications of avascular necrosis of the bone (David *et al*, 1993; Ballas, 1995; Ballas 2001; Smith, 1996).

The disappointing results of treatment of avascular necrosis after bony collapse have led to the suggestion by many workers that future efforts should be directed to the early detection or diagnosis of this bone disease before the onset of the skeletal or structural failure in patients with sickle cell haemoglobinopathies. Bone scan with technetium ^{99m}Tc and magnetic resonance imaging (MRI) can detect disease but these tests are very expensive and difficult to use routinely in a developing country like Nigeria (David *et al*, 1993; Ballas, 1995; Ballas 2001; Smith, 1996; Rao *et al*, 1985; Rao *et al*, 1989; Amundsen *et al*, 1984; Milner *et al*, 1993; Mitchell *et al*, 1987). Recently there has been introduction of new biochemical markers of bone metabolism such as osteocalcin and others (Bettica and Moro, 1995; Kleerekoper 1997; De la Piedra *et al*, 1996; Bolarin, 1996).

a. Osteocalcin

Osteocalcin or bone gamma-carboxyglutamic acid (gla) protein (BGP) is the most abundant non-collagenous protein found in bone (Bolarin, 1996). Osteocalcin has a molecular weight of 5.7 kD. It is synthesized exclusively by osteoblasts and by its dental form, the odontoblasts (Lian and Grundberg 1988; Hauschka *et al*, 1989). Osteocalcin represents the activity of osteoblasts, since it is synthesized and released into circulation by the osteoblasts. The protein is eliminated by glomerular filtration and degraded by the renal tubule (Bolarin, 1996; Lian and Grundberg 1988; Hauschka *et al*, 1989; Garnero *et al*, 1994).

Measurement of osteocalcin is by immunoassays using monoclonal or polyclonal antibodies. It is advised that blood samples be collected in ice and plasma or serum is stored at -20 °C or -70 °C. It should be thawed once to prevent degradation of the protein (Bolarin, 1996; Lian and Grundberg 1988; Hauschka *et al*, 1989; Garnero *et al*, 1994).

b. Alkaline phosphatase (EC. 3.1.3.1)

Bone-specific alkaline phosphatase (b-AP) is a plasma membrane enzyme. For many years this enzyme has been the most commonly used biochemical marker to assess osteoblast function (Bolarin, 1996). Bone-specific alkaline phosphatase activity correlates with bone formation rate in metabolic disease of the bone (Bolarin, 1996; Grundberg, 1993; Bolarin, 2001) and during normal physiological bone growth (Bolarin, 1996; Grundberg, 1993; Bolarin, 2001). The serum alkaline phosphatase activity is present in tissues other than bone. The isoenzymes can be divided into four groups (placental, carcinoplacental, intestinal, and hepatic, renal, skeletal group). The major fraction of the total serum is from the liver and bone isoforms or isoenzymes, which are both products of the same genetic loci (Bolarin, 1996; Grundberg, 1993; Bolarin, 2001). It is often difficult to distinguish between the two isoenzymes. But distinction is necessary for clinical use (Bolarin, 1996).

There are several methods for the estimation of serum bone-specific alkaline phosphatase. These methods include inactivation some of the isoenzymes by heat or using chemical compounds, precipitation with wheat germ lectin and separation by electrophoresis. These techniques, however, do not seem to provide the sensitivity, specificity and reliability required for routine clinical use (Bolarin, 1996; Grundberg, 1993; Bolarin, 2001). A two-site immunoradiometric assay that preferentially recognizes bone-specific alkaline phosphatase is generally being used in most clinical or research laboratories (Bolarin, 1996; Grundberg, 1993; Bolarin, 2001). These new assay methods have shown improved diagnostic accuracy with regards to bone diseases and their therapeutic monitoring (Bolarin, 1996; Grundberg, 1993; Bolarin, 2001).

In the present preliminary study, serum osteocalcin is assayed by enzyme immunoassay using monoclonal antibody. Serum bone-specific alkaline phosphatase protein (b-AP) was also measured by immunoradiometric assay (IRMA). Both biochemical markers were used to investigate bone turnover in patients with sickle cell haemoglobinopathies.

Materials And Methods

Study Population

Twenty patients consisted of 9 men and 11 women with sickle cell haemoglobinopathies. Mean age \pm standard deviation was 31.6 ± 11.5 years (median 30

years, range 18 - 51 years). A reference control group of 20 apparently healthy subjects (9 males and 11 females), age range 20 - 52 years (median 30 years). None of the persons had a history serious disease or was taking drugs known to affect bone metabolism or the blood vessels. The control had no history of fractures nor any nutritional, metabolic, liver, renal or bone disease. None of the females were on contraceptives prior to obtaining blood samples. All control subjects had haemoglobin AA. The control groups were individuals of African-American ancestry just like the patients used in the study. The diagnosis of sickle cell disease was based on the haemoglobin pattern on cellulose acetate electrophoresis. Family studies were consistent where available. The genotypes of the patients were divided as follows:

- a. SS - sickle cell anemia 14
- b. S β - sickle cell thalassaemia 4
- c. SC-sickle cell hemoglobin C disease 2

Five patients were identified clinically and by physical diagnostic methods (such as radiography, etc.) to have skeletal complications due to sickle cell disease. All the other patients were asymptomatic for skeletal complications at the time blood samples were taken for the biochemical assays. The clinical history and routine liver function tests and urinalysis carried out on the experimental subjects including the control group showed that none of the patients or control subjects had liver, renal, or inflammatory disease during the time blood and urine samples were taken for the assays. The patients were attendees at sickle cell disease out-patient clinic.

Blood samples:

Blood samples were collected at the same time in the morning during clinic. Sera were separated from blood by centrifugation at 4 °C and were stored frozen at -60 °C until assayed. Assays were carried out within 1 to 2 weeks of storage.

Assay of Biochemical markers of bone metabolism

a. *Determination of serum osteocalcin (S-BGP):* Serum levels of osteocalcin (S-BGP) were determined in duplicate samples using an

IRMA kit from CIS bio International, (GifSurYvette, France). The tracer is a ¹²⁵I-mono-clonal antibody against human BGP. The mono-clonal antibody used in this study is raised against human osteocalcin and therefore measured total serum osteocalcin protein levels. The serum concentration of osteocalcin was expressed as $\mu\text{g/L}$.

b. *Determination of serum bone-specific alkaline phosphatase (S-b-AP):* Serum b-AP protein level was determined with an IRMA (Tandem-R-OSTASE, Hybritech, San Diego, CA), using two mono-clonal antibodies, specific for b-AP over liver alkaline phosphatase. Briefly, 100 μl of standard or serum and 100 μl tracer (100 nCi [¹²⁵I] anti-b-AP mouse mono-clonal antibody) were added to beads coated with mouse mono-clonal antibody directed to human b-AP. After incubation for 19 ± 2 h at 4 °C, the solid beads were washed three times with detergent solution. Total and bound radioactivity was counted in a gamma-counter. The level of b-AP protein is expressed as μg per liter.

Statistical Analysis

The results are presented as means \pm SD. Median values and range are included. The mean \pm SD are compared, as applicable, by Student's t-test.

Results

There were no statistically significant differences in age between the groups. The results of the biochemical markers of bone formation, serum osteocalcin (S-BGP) and serum bone-specific alkaline phosphatase (b-AP) are summarized in Table 1. The mean level of serum osteocalcin was not statistically ($p < 0.05$) different from the control group. The concentration of bone-specific alkaline phosphatase protein was significantly higher in the patients with sickle cell disease than in control group ($p < 0.05$). The upper limit of the reference range (i.e. mean + 2 standard deviation, SD) of the control for the bone-specific alkaline phosphatase was 25.5 $\mu\text{g/l}$. The upper limit of the reference ranges for serum osteocalcin (S-BGP) were exceeded by 2/20 patients or 10 per cent and 11/20 patients or 55 per cent of all the sickle cell disease patients studied respectively.

Table 1: Summary of the biochemical markers in control group and patients with sickle cell Haemoglobinopathies

Biochemical markers	Control Group			Sickle Cell Disease			P Value
	Mean \pm SD*	(Median)	Range	Mean \pm SD	(Median)	Range	
Serum bone-specific alkaline phosphatase (S-b-sAP, μ g/l)	10.5 \pm 7.5	(10)	6.0 – 15	16.9 \pm 22.1	(17.0)	9.0 – 27.0	< 0.05
Serum Osteocalcin (BGP, μ g/l)	20.3 \pm 8.7	(16.6)	11.5– 37.3	19.9 \pm 12.7	(16.2)	9.3 – 60.1	NS

*SD** = Standard Deviation *NS* =Not Significant

Discussion

Serum alkaline phosphatase is used extensively for the assessment of bone turnover in patients with various metabolic bone disorders, but its clinical use is often impaired by serum changes occurring in non-bone alkaline phosphatase. In this study, we used the two-site immunoradiometric assay (IRMA) developed by Hybritech (Tandem-R Ostase) to determine the bone isoenzyme in sera of normal subjects and in patients with sickle cell haemoglobinopathies. With this IRMA, serum bone-specific alkaline phosphatase protein increased significantly in patients with sickle cell disease when compared with those in the normal healthy subject group. The results of bone-specific alkaline phosphatase total protein levels in this study agree with data obtained in patients with metabolic bone disease, showing increased serum bone-specific alkaline phosphatase (Duda *et al*, 1988). Our data shows that serum bone-specific alkaline phosphatase total protein determination may be useful in early diagnosis of bone changes in sickle cell haemoglobinopathies, without any sign of hepatic disorder (Bolarin, 1983). There was no difference in the mean values of serum osteocalcin in both control group and patients with sickle cell haemoglobinopathies. The value of osteocalcin as a biochemical marker of bone matrix formation is unclear as fluctuation in circulating osteocalcin may only reflect changes in the equilibrium between bone matrix and bone (Bolarin, 1996; Duggan, 2001). Osteocalcin is unstable in circulation and subject to variable vitamin K-dependent carboxylation. Because of the heterogeneity of circulating osteocalcin, different assays give differing results (Bolarin, 1996; Duggan, 2001).

In conclusion, we have shown that IRMA assay of serum bone-specific alkaline

phosphatase total protein was more sensitive and specific than the determination of serum osteocalcin (S-BGP) to detect the increase in bone turnover due to bone remodeling in sickle cell haemoglobinopathies. The pilot data presented here also form the basis for further confirmatory studies leading to development of noninvasive biochemical markers to monitor skeletal complications in sickle cell disease. It may be useful to monitor the course of therapy of these complications in these patients, since it is characterized by increased sensitivity and organ specificity (Behr and Barnett, 1986).

Acknowledgments

This study was carried out while one of the authors, Dr. Bolarin was working in Wayne State University, Detroit, Michigan, USA and it was supported by the University Interdisciplinary Grant 1444981. The authors are grateful to Dr P. Swerdlow, Division of Haematology for the patients.

References

- Amundsen, T. R., Siegel, M. J., Siegel, B. A. (1984). Osteomyelitis and infarction in sickle cell haemoglobinopathies: Differentiation by combined technetium and gallium Scintigraphy. *Radiology* 159: 807-813.
- Aster, J. C. (2004). Red blood cell and bleeding disorders. In Robbins and Cotrans Pathologic Basis of Disease. Kumar V, Abbas AK and Nelson F. Eds. 7th Edition. *Elsevier Saunders, Philadelphia, Pennsylvania, USA*. pp 619 – 659.
- Ballas, S. K. (1995). The sickle cell painful crisis in adults: phases and objective signs. *Haemoglobin* 19: 323-333.

- Ballas, S. K. (2001). Sick cell: Current clinical management. *Semin. Hematol.* 38: 307–315.
- Behr, W., Barnett, J. (1986). Quantitation of bone alkaline phosphatase in serum by precipitation with wheat-germ lectin: A simplified method and its plausibility. *Clin. Chem.* 32: 1960-1966.
- Bettica, P., Moro, L. (1995). Biochemical markers of bone metabolism in the assessment of osteoporosis. *J. Int. Fed. Clin. Chem.* 7: 16-22.
- Bolarin, D. M. (1996). Biochemical Markers for the assessment of skeletal growth in children. *Nig. Quart. J. Hosp. Med.* 6: 256-261.
- Bolarin, D. M. (2001). Bone-specific alkaline phosphatase protein, total alkaline phosphatase activity and lactate dehydrogenase in sera of patients with sickle cell disease. *Haematologica* 31: 51-56.
- Bolarin, D. M. (1983). Liver function and the diagnostic significance of conjugated cholic acid and chenodeoxycholic acid in serum of African patients with sickle cell disease. *Acta Tropica.* 40: 59-64.
- David, H. G., Bridgman, S. A, Davies, S. C. (1993). The shoulder in sickle cell disease. *J. Bone Joint Surg.* [Br.] 75-B: 538 - 545.
- De la Piedra, C., Rapado, A., Diego, D. E. M., Martin, D. M.A., Aguirre, C., Gavilanes, L. E., Curiel, D. M. (1996). Variables efficacy of bone remodeling biochemical markers in the management of patients with Paget's disease of bone treated with tiludronate. *Calcif. Tissue Int.* 59: 95-99.
- Duda, R. J., O'Brien, J. F., Katzmann, J. A., (1988). Concurrent assays of circulating bone Gla protein and bone alkaline phosphatase: effects of sex, age and metabolic bone disease. *J. Clin. Endocrinol. Metabol.* 66: 951-957.
- Duggan, R. (2001). Biochemical markers of bone and Ca⁺⁺ metabolism. *Clin. Lab. Int.* 24: 14.
- Embury, S. H., Hebbel, R. P., Mohandas, N., Steinberg, M. H. Eds. (1994). Sick cell Disease. *Basic Principles and Clinical Practice.* New York, Raven Press.
- Francis, R. B., Johnson, C. S. (1991). Vascular occlusion in sickle cell disease. Current concepts and unanswered questions. *Blood* 77: 1404-1405.
- Garnero, P., Grimaux, M., Seguin, P., Delmas, P. D. (1994). Characterization of immunoreactive forms of human osteocalcin generated in vivo and in vitro. *J. Bone Miner. Res.* 9: 255-264.
- Gundberg, C. M. (1993). Alkaline phosphatase and osteocalcin. In: Primer on the metabolic bone diseases and disorders of mineral metabolism. Favus M. J. Ed. New York: *American Society of Bone and Mineral Research*, 2nd edition. pp. 74-76.
- Hauschka, P. V., Lian, J. B., Cole, D. E. C., Grundberg, C. M. (1989). Osteocalcin and matrix gla protein: vitamin K dependent proteins in bone. *Phys. Rev.* 69: 990-1047
- Hebbel, R. P. (1991). Beyond hemoglobin polymerization. The red blood cell membrane and sickle cell disease pathophysiology. *Blood* 77: 214-217.
- Iwegbu, C. G., Fleming, A. F. (1985). Avascular necrosis of the femoral head in sickle cell disease: a series from the Guinea savannah of Nigeria. *J. Bone Joint Surg.* [Br] 67-B: 29-32.
- Kleerekoper, M. (1997). Evaluating and managing osteoporosis. The emerging role of biochemical bone markers. *Clinical Laboratory News* 23: 6-7.
- Lee, R. E. J., Golding, J. S. R., Serjeant, G. R. ((1981). The radiological features of avascular necrosis of the femoral head homozygous sickle cell disease. *Clin. Radiol.* 32: 205-214.
- Lian, J. B., Grundberg, C. M. (1988). Osteocalcin: Biochemical consideration and clinical application. *Clin. Orthop.* 226: 267-291.
- Milner, P. F., Kraus, A. P., Sebes, J. I. (1993). Osteogenesis of the humeral head in sickle cell disease. *Clin. Orthop. Related Res.* 289: 136-143.
- Mitchell, D. G., Rao, V. M., Dlinka, M. K. (1987). Femoral head avascular necrosis: correlation of MR imaging, radiographic staging, radionuclide imaging and clinical findings. *Radiology* 162: 709-715.
- Powars, D. R. (1990). Sick cell anaemia and major organ failure. *Haemoglobin* 14: 573-598.
- Rao, S., Solomon, N., Miller, S. (1985). Scintigraph differentiation of bone infarction from osteomyelitis in children sickle cell disease. *J. Pediat.* 107: 685-689.
- Rao, V. M., Mitchell, D. G., Rifkin, M. D. (1989). Marrow infarction in sickle cell anaemia: correlation with marrow type and distribution by MRI. *Magn. Reson. Imaging* 7: 39-44.
- Smith, J. A. (1996). Bone disorders in sickle cell disease. *Hematol./Oncol. Clin. N. Am.* 10: 1345 - 1356.

Received: 15/9/06

Accepted: 16/11/06