

Journal of Science and Technology, Vol. 35, No. 3 (2015), pp25-33 25

© 2015 Kwame Nkrumah University of Science and Technology (KNUST)

<http://dx.doi.org/10.4314/just.v35i3.3>

RESEARCH PAPER

A MULTI-PERIOD MARKOV MODEL FOR MONTHLY RAINFALL IN LAGOS, NIGERIA

K. O. Aiyesimoju and A. O. Busari

Department of Civil and Environmental Engineering

University of Lagos, Akoka-Yaba, Lagos, Nigeria

Corresponding author: k_simoj@yahoo.com

ABSTRACT

Long periods of historical hydrological data such as rainfall and streamflow which are necessary for planning and design of water resources projects, are often not available and have to be forecasted. Many models available for this were developed and tested in developed countries in temperate climates and so their application in tropical climates is questionable. A twelve-period Markov model has been developed for the monthly rainfall data for Lagos, along the coast of South Western Nigeria. The goodness of fit of the model was assessed by estimating the autocorrelations of the residuals of the historical data (from January 1924 to December 1983) for lags one to sixty. A 95% confidence band was also established for the autocorrelations. The results show that all but two of the autocorrelations fall within the 95% confidence band confirming that the residuals are indeed white noise. This indicates that the model is very adequate.

Keywords: *Markov, multi-period, rainfall model*

INTRODUCTION

The problem of inadequacy or sometimes a total lack of stream flow data if not resolved would definitely lead to an underestimation of water resources facilities or their overestimation. This is why the first attempt to solve the problem of short record length was as far back as the beginning of the 20th century. Hazen (1914) suggested combining records from several stations whilst Sudler (1927) on the other hand wrote historic records on cards and randomly picked the cards, with replacement, to

generate 1000 years record. These approaches have the shortcoming of being unable to generate flows outside the actual historical range. This is serious because the potential for large and complex variation over time of hydrological data (see Koutsoyinnannis and Montanori, 2007) may result in future flows well outside of the historically observed range.

In reservoir size estimation, the required capacity of a reservoir depends on the sequence of inflows into it versus the sequence of required

withdrawals from it. In this case, it is not so important to know the exact sequence of future inflows. All that is needed is to generate future inflows with the same stochastic properties as the historical record. This is why stochastic models were introduced in hydrology to generate inflows that have equal probability of occurrence as the historical values.

RELATED WORKS

Stochastic models were pioneered by Barnes (1954) and Thomas and Fiering (1962). Box and Jenkins (1972) generalized existing models and further developed them. The resulting Box-Jenkins models; namely Autoregressive (AR), Moving Average (MA), and their combinations greatly popularized stochastic modeling despite their shortcomings which include short range dependence, parsimony problems and a lack of physical meaning especially for the higher order models. These shortcomings led to the development of other models such as Fractional Gaussian Noise Processes (Mandelbrot, 1965), Broken Line Processes (Mejia *et al.*, 1972), Disaggregation Models (Valencia and Schaake, 1973), etc. More recently Artificial neural Network (ANN) models (Abrahart *et al.*, 2007; Birikundavyl *et al.*, 2002; Corzo and Solomatine, 2005; De Vos and Rietjes, 2007, Kant *et al.* 2013, Borga *et al.*, 2011, Chen *et al.*, 2015) which are connectionist in nature, have been developed. Also, advances in non-linear dynamical systems (Langousis and Koutsoyiannis, 2006, Leedal *et al.*, 2013) have led to the development of models which possess a deterministic basis. These latest models are however data driven and of a black box nature thus providing no process insight. Worse still, they do not provide tools for Monte Carlo simulation which is a serious problem as the whole point of stochastic simulation is to extend the record. Several other authors focused on the effect of uncertainty on the predictions of hydrological phenomena e.g. Weerts *et al.* (2011) and Zappa *et al.* (2011). More recently, Ursu and Perea (2015) utilized a periodic autoregressive process to model river flow; Deo *et al.* (2015) utilized an adaptive regression

spline model to estimate monthly evaporative loss; Pinto *et. al* (2015) compared models for forecasting monthly streamflows; but all of them were developed in a different environment from tropical Africa.

Selection of a suitable model among the numerous options available may not be clear but factors that will influence the suitability of a model are the nature of the physical processes involved, the quality of the data available, environmental factors such as climate, and, the use to be made of the model. The aim of this work is to develop a multi-period Markov model for monthly rainfall forecasting using monthly rainfall data for Lagos. The model will be assessed by checking if the autocorrelations at different lags of the residuals of the model when applied to the historical data are statistically insignificant. This will confirm the suitability of such a model for stochastic simulation of monthly rainfall data in our tropical and coastal environment. Monthly rainfall is well known to exhibit a seasonality of twelve months and thus twelve periods would be the natural choice and is adopted here.

Periodic Markov Model

The standard Markov Model is based on the premise that the variate at a particular time is linearly dependent on that at the immediately preceding time plus a random component as follows:

$$X_i = rX_{i-1} + a_i \quad (1)$$

where X_i is the variate value at time i , a_i is random deviate at time i and r is the lag one autocorrelation coefficient for series X_i . Taking the expectation and variance of Equation (1) implies

$$E(X_i) = rE(X_{i-1}) + E(a_i) \quad (2)$$

$$\begin{aligned} \text{Var}(X_i) &= \text{Var}(rX_{i-1}) + \text{Var}(a_i) \\ &= r^2\text{Var}(X_{i-1}) + \text{Var}(a_i) \end{aligned} \quad (3)$$

Assuming stationarity, then $E(X_i) = E(X_{i-1}) = \mu_x$

$Var(X_i) = Var(X_{i-1}) = \sigma_x^2$ and let $E(a_i) = \mu_a$ or

and $Var(a_i) = \sigma_a^2$

Then from Equations (2) and (3)

$$\mu_a = \mu_x(1-r) \tag{4}$$

$$\sigma_a^2 = (1-r^2)\sigma_x^2 \tag{5}$$

Assuming our variates are normally distributed and if t_i represents a standard normal random deviate at time i , then

$$a_i = \mu_a + t_i\sigma_a = \mu_x(1-r) + t_i\sigma_x\sqrt{(1-r^2)} \tag{6}$$

Dropping the subscript X , Equation (1) can be written in terms of the parameters of the original variate X as

$$X_i = rX_{i-1} + \mu(1-r) + t_i\sigma\sqrt{(1-r^2)} \tag{7}$$

which can be written as

$$\frac{X_i - \mu}{\sigma} = r\frac{X_{i-1} - \mu}{\sigma} + t_i\sqrt{(1-r^2)} \tag{8}$$

Equation (8) in effect shows that prior normalization of the deviates simplifies the model.

Hydrological data however exhibit an annual period. Thus for monthly rainfall that is of interest here, this translates to a period of twelve months. In Equation (8), the variate for a time i depends on only the variate for the previous time $i-1$ and a random component. However, seasonality implies that the dependence of the variates belonging to period $j+1$ on their immediately preceding values (period j) is different from the dependence of the variates belonging to period j on their immediately preceding values (period $j-1$). Let $j(i)$ be the period to which time i belongs, a multi-period Markov model can thus be written as

$$\frac{X_i - \mu_{j(i)}}{\sigma_{j(i)}} = r_{j(i)}\frac{X_{i-1} - \mu_{j(i-1)}}{\sigma_{j(i-1)}} + t_i\sqrt{(1-r_{j(i)}^2)} \tag{9a}$$

$$x_i = r_{j(i)}x_{i-1} + t_i\sqrt{(1-r_{j(i)}^2)} \tag{9b}$$

where x_i is the normalized X_i as follows

$$x_i = \frac{X_i - \mu_{j(i)}}{\sigma_{j(i)}}$$

Note above, r_j is the autocorrelation of the normalized variates belonging to period j to those immediately preceding them (period $j-1$). Index i is consecutively numbered from 1 to total no of data whilst index j will vary from 1 to number of periods, which is twelve here. For example, j for all January values will be 1 whilst that for all December values will be 12. μ_j and σ_j are ideally the population mean and standard deviation respectively for period j values. Let $i \in j$ represent only i values belonging to period j , the sample estimates used were as follows

$$\mu_j = \frac{\sum_{i \in j} X_i}{T_j} \tag{10}$$

$$\sigma_j = \sqrt{\frac{\sum_{i \in j} (X_i - \mu_j)^2}{T_j}} \tag{11}$$

$$r_j = \frac{\frac{1}{T_j} \sum_{i \in j} (X_i - \mu_j)(X_{i-1} - \mu_{j-1})}{\sigma_j\sigma_{j-1}} \tag{12a}$$

or

$$r_j = \frac{\sum_{i \in j} x_i x_{i-1}}{T_j} \tag{12b}$$

where T_j is the number of values in the summation belonging to period j . Equations (10), (11) and (12a) are used to estimate the parameters of the model.

GOODNESS OF FIT MODEL

After the parameters of the model have been determined, application of the model to the

historical data will not show a perfect fit. The resulting errors are termed as residuals and they represent the random components corresponding to the historical data. For a good model, the residuals should be white noise in which case, their autocorrelation ρ_j at any lag j should be zero. However, since only a sample is available here, the $100(1-\alpha)$ % confidence interval for the sample autocorrelation ρ_j at any lag j , is (Yevjevich, 1972)

$$\frac{-t(T-2, \alpha/2)}{(T-2+t^2(T-2, \alpha/2))^{1/2}} \leq \rho_j \leq \frac{t(T-2, \alpha/2)}{(T-2+t^2(T-2, \alpha/2))^{1/2}} \quad (13)$$

where T is the sample size and $t(N, \alpha)$ is a student-t random deviate with N degrees of freedom and a probability of exceedance of α . The residuals \hat{t}_i of the historical data are obtained by simply replacing the random component in Equation (9a) or (9b) with the residual \hat{t}_i to give

$$\frac{X_i - \mu_{j(i)}}{\sigma_{j(i)}} = r_{j(i)} \frac{X_{i-1} - \mu_{j(i-1)}}{\sigma_{j(i-1)}} + \hat{t}_i \sqrt{(1 - r_{j(i)}^2)}$$

This implies

$$\hat{t}_i = \frac{\frac{X_i - \mu_{j(i)}}{\sigma_{j(i)}} - r_{j(i)} \frac{X_{i-1} - \mu_{j(i-1)}}{\sigma_{j(i-1)}}}{\sqrt{(1 - r_{j(i)}^2)}} \quad (14a)$$

or

$$\hat{t}_i = \frac{x_i - r_{j(i)} x_{i-1}}{\sqrt{(1 - r_{j(i)}^2)}} \quad (14b)$$

RESULTS AND DISCUSSION

Monthly rainfall data from January 1924 to

South Western Nigeria (Table 1) were obtained, making a total of 720 data points. The parameters of the model estimated from the historical data using Equations (10), (11) and (12a) are as shown in Table 2. To test the goodness of fit, the residuals (model errors) for the historical data were obtained using Equation (14a). The autocorrelation values of these residuals for lags one to sixty are plotted in Fig. 1. The 95% confidence band for these from Equation (13) is also plotted. This shows that all but two fall within the 95% confidence band which is very good. Furthermore, 1000 years of data were generated with the model (using Equation (9a)). The parameters obtained for them are shown in Table 3 and they compare very well with Table 2, further validating the model.

CONCLUSION

A twelve-period Markov stochastic model has been developed for the monthly rainfall data for Lagos along the coast of South Western Nigeria. The goodness of fit of the model was assessed by estimating the autocorrelations of the residuals of the historical data (from January 1924 to December 1983) for lags one to sixty. A 95% confidence band was also established for the autocorrelations. The results show that all but two of the autocorrelations fall within the 95% confidence band, confirming that the residuals are indeed white noise. This indicates that the model is very adequate. This confirms the suitability of such a model for stochastic simulation of monthly rainfall data in our tropical and coastal environment. An important limitation of the model is the fact that theoretically, negative values are possible which is practically meaningless. Thus in practice, negative values are treated as zeros.

Table 1: Rainfall data (mm) for Lagos from 1924 to 1983

Year	JAN	FEB	MAR	APR	MAY	JUNE	JULY	AUG	SEP	OCT	NOV	DEC
1924	49.28	28.45	134.11	191.77	87.63	140.46	62.99	2.54	104.14	396.75	21.08	48.77
1925	38.1	10.16	167.89	177.8	308.86	518.16	386.59	32.51	151.89	75.69	72.9	0
1926	0	76.45	69.6	324.1	347.73	331.72	255.78	6.6	280.67	96.67	138.94	1.78
1927	63.25	56.69	70.61	85.6	208.03	179.83	217.63	6.35	77.22	338.58	60.45	29.72
1928	44.96	56.39	208.28	176.78	389.38	534.67	64.26	52.04	142.24	321.82	13.72	3.3
1929	0.51	37.08	26.92	178.82	288.04	629.67	506.22	20.57	78.99	153.16	104.14	152.91
1930	33.05	56.13	83.06	127.25	218.69	337.31	467.36	16.76	67.82	316.48	47.75	42.93
1931	23.87	37.33	149.6	181.86	225.29	450.34	452.36	53.34	318.51	149	56.09	23.62
1932	0.5	11.17	66.29	96.52	288.03	358.14	21.84	76.7	104.34	131.06	6.8	0
1933	125.22	52.07	118.61	100.33	167.89	377.44	495.04	38.35	139.44	154.94	134.87	24.63
1934	5.08	0	120.65	145.56	136.65	398.27	368.04	200.91	109.72	341.63	29.71	20.14
1935	0	31.49	212.64	159.25	335.34	537.97	408.68	10.66	58.52	154.43	10.92	166.6
1936	23.62	39.11	146.81	62.27	306.82	373.38	4.06	28.44	44.95	352.55	135.38	61.46
1937	0	70.35	128.27	122.17	254.4	509.01	458.97	36.06	316.23	138.4	9.65	32
1938	21.84	78.23	42.16	153.16	276.09	292.6	88.39	40.89	115.57	218.94	158.49	4.06
1939	8.63	45.21	75.94	95.5	315.72	399.42	315.21	141.22	17.01	91.94	13.71	61.72
1940	20.06	4.52	83.05	158.24	208.53	584.96	193.29	38.1	193.23	313.18	58.92	0
1941	32	43.68	50.03	238.76	401.32	166.59	275.33	154.17	186.43	26.87	186.18	3.04
1942	54.1	40.64	170.94	122	428.75	552.9	122.17	11.43	224.02	159.76	92.51	82.29
1943	152.14	23.36	58.42	128.63	254.5	615.44	86.86	24.38	160.52	425.95	92.93	24.89
1944	33.26	65.53	134.87	104.18	197.1	563.11	341.12	73.91	35.56	112.77	64.51	0
1945	1.78	4.32	124.2	65.02	151.13	219.71	392.27	21.84	101.85	309.37	101.09	0
1946	23.87	29.97	446.73	59.43	416.56	556.26	0.13	0.25	28.95	93.47	60.19	18.24
1947	1.01	21.59	88.13	251.46	247.65	385.06	785.7	130.04	80.51	151.13	43.18	25.9
1948	0	140.2	33.78	278.13	342.39	305.05	19.55	45.72	164.84	82.04	19.18	0
1949	0	2.79	99.56	57.4	294.13	400.81	148.33	104.9	39.62	159	113.79	0
1950	0	26.16	28.44	227.07	244.85	434.08	195.32	17.78	217.42	123.69	64	0.5
1951	97.02	0.25	150.62	55.37	265.4	355.09	383.54	124.96	91.44	367.79	111.25	0
1952	58.42	64.77	178.05	169.92	298.45	259.69	141.72	7.87	171.7	166.62	88.13	60.19
1953	0.25	131.57	65.27	89.66	359.65	536.19	247.14	22.6	395.47	207.77	127.5	2.79
1954	51.05	144.52	120.23	88.13	255.77	617.47	40.13	39.11	170.94	290.32	42.67	16.51
1955	80.01	9.9	139.44	144.01	175.26	629.15	129.15	105.15	140.2	354.58	22.09	17.78

Table 1 (contd): Rainfall data (mm) for Lagos from 1924 to 1983

Year	JAN	FEB	MAR	APR	MAY	JUNE	JULY	AUG	SEP	OCT	NOV	DEC
1956	0.25	25.4	118.11	144.01	252.22	322.07	230.12	9.39	157.98	74.67	87.37	87.37
1957	7.11	13.2	113.03	115.06	350.52	405.63	613.15	194.56	268.73	197.61	78.74	39.37
1958	51.05	47.24	151.38	83.05	513.08	818.38	3.3	11.68	170.94	107.69	119	35.3
1959	63.24	28.19	105.51	167.89	422.91	327.4	422.14	100.07	124.96	191.26	45.21	0
1960	53.34	76.7	38.06	293.11	183.38	489.45	66.54	105.41	191.13	199.13	69.08	15.03
1961	3	0.3	152.7	219.5	230.6	673.4	319	1.8	130.6	96.9	38.9	49.3
1962	23.4	12.7	143.3	157.7	179.8	833.6	273.8	157	38.1	167.7	12.04	20.6
1963	1.8	18	46.5	222	301.2	303.3	493	370.3	295.9	123.7	86.9	1.5
1964	2.5	19.8	104.4	156.5	218	468.8	48.3	58.2	77	177.5	39.6	4.6
1965	4.1	56.4	100.6	216.4	260.4	434.8	648.2	138.9	299.5	133.9	24.6	1.3
1966	13.2	20.1	69.6	198.9	194.6	272.3	447.3	50.3	134.4	219.2	79.8	11.7
1967	51.6	19.3	106.9	115.1	128.5	405.4	441.2	4.8	133.1	220.7	66.3	10.4
1968	12.7	52.6	21.3	181.9	172	597.2	926.6	527.6	534.9	189.7	31.2	18
1969	31.2	1.8	217.4	162.6	368.6	527.1	283	134.4	67.6	221	4.3	52.1
1970	126.7	7.6	34.5	101.6	301.5	587.2	476.5	34.8	327.8	305.5	67.6	0
1971	8.4	95.3	26.2	57.7	213.6	356.6	255.6	272.5	219.7	58.9	47.5	17.8
1972	35.6	37.3	108.7	89.9	220.5	271.3	77.7	272.5	238.8	194.2	40.6	8.6
1973	82.1	25.9	23.6	157.2	196.9	468.4	0	57.7	116.6	74.2	20.1	15.5
1974	79.2	25.7	112	84.3	324.1	393.7	359.7	185.2	278.9	78.5	16.3	0
1975	0	79.2	13.5	210.6	159	297.7	647.4	22.4	62	110.2	40.6	109.5
1976	0	140.2	224	206.2	145.5	588.8	119.9	23	54.9	119.9	42.2	14.5
1977	103.9	0.4	27.5	108.4	222.9	417.7	211.3	24.1	86.8	147.1	6.8	9.4
1978	18.3	60.9	100.8	292.3	293.7	189.4	268.8	268.7	283.9	275.4	100.6	26.7
1979	2.9	10.1	60.4	135.7	377.9	364.2	467.8	399.7	239.4	161.2	120	4.1
1980	133.1	45.7	51.8	75.1	267	341.8	141.9	97.4	200.5	122.2	170.9	0.3
1981	6.7	2.7	51.8	169.1	255.7	449.1	96.3	84.3	240.6	115	22.2	0.9
1982	0	16.7	24.7	215	288.2	665.9	279.9	5	64.3	109.1	31.5	0
1983	0	0	29.6	60.8	423.7	603.5	86.8	13.6	175.2	34.7	47.6	0

Table 2: Estimated parameters of model from observed data

Lag j	1	2	3	4	5	6	7	8	9	10	11	12
μ_j	3.264E+0	3.966E+0	1.502E+0	2.694E+0	4.404E+0	2.797E+0	8.852E+	1.624E+0	1.834E+0	6.434E+0	2.466E+0	
	1	1	1.029E+02	2	2	2	01	2	2	2	1	1
	3.880E+0	3.615E+0	6.556E+0	8.727E+0	1.520E+0	2.080E+0	1.074E+	1.019E+0	9.771E+0	4.494E+0	3.519E+0	
σ_j	1	1	7.103E+01	1	1	2	2	02	2	1	1	1
	1.906E-02	-1.177E-	-2.021E-	-8.829E-	1.675E-	-7.414E-	4.100E-	-8.015E-	4.765E-01	1.595E-	-7.134E-	
	01	01	-4.073E-02	01	02	01	01	01	02	02	02	02
r_j	1.906E-02	-1.096E-	-1.865E-	2.917E-	-1.014E-	2.116E-	-7.687E-	7.335E-	-5.587E-			
	01	01	-8.002E-02	01	01	01	01	01	02	03	02	02
b_j	2.102E-02											




Fig. 1: Auto correlations of the residuals from the observed data for the various Lags

Table 3: Estimated parameters of 1000 years of forecasted data

Lag, j	1	2	3	4	5	6	7	8	9	10	11	12
μ_j	3.247E+0	3.871E+0	1.043E+0	1.478E+0	2.685E+0	4.405E+0	2.857E+0	8.847E+0	1.579E+0	1.831E+0	6.290E+0	2.527E+0
	1	1	2	2	2	2	2	01	02	2	1	1
	3.730E+0	3.731E+0	7.254E+0	6.370E+0	8.574E+0	1.551E+0	2.084E+0	1.067E+0	1.042E+0	9.636E+0	4.564E+0	3.628E+0
σ_j	1	1	1	1	1	2	2	02	02	1	1	1
	-1.516E-	-1.216E-	-6.820E-	-1.499E-	-6.987E-	2.218E-	-9.670E-	4.309E-	4.394E-	-8.304E-	-6.743E-	-5.153E-
	02	01	02	01	02	01	02	01	01	02	03	02
r_j	-1.558E-	-1.217E-	-1.326E-	-1.316E-	-9.405E-	4.011E-	-1.300E-	2.206E-	4.292E-	-7.678E-	-3.194E-	-4.097E-
	02	01	01	01	02	01	01	01	01	02	03	02
b_j												

REFERENCES

Abrahart, R. J., Heppenstall, A. J. and See, L. N. (2007). Timing error correction procedure applied to neural network rainfall-runoff modeling. *Hydrological Sciences Journal*, 52(3): 414 – 431.

Barnes, F. B. (1954). Storage required for a city water supply. *Journal of Institution of Engineers, Australia*. 26(9): 198–203.

Birikundavyl, S., Labid, R., Trung, H. and Rousselle, J. (2002). Performance of Neural Networks in Daily Streamflow Forecasting. *Journal of Hydrologic Engineering*, 7(5):392–398.

Borga, M., Anagnostou, E. N., Blöschl, G., Creutin, J. D. (2011). Flash flood forecasting, warning and risk management: the HYDRATE project, *Environmental Science and Policy*, 14(7):834–844

Box, G. E. P. and Jenkins, G. M. (1972). Time series analysis, forecasting and control. Holden Day Inc., San Francisco

Chen, X. Y., Chau, K. W. and Busari A. O. (2015). A comparative study of population-based optimization algorithms for downstream river flow forecasting by a hybrid neural network model. *Engineering Applications of Artificial Intelligence*, 46(A): 258–268

Corzo, G. S. and Solomatine, D. (2007). Base-flow separation technique for modular artificial neural network modeling in flow forecasting. *Hydrological Sciences Journal*, 52 (3): 491–507.

De Vos, N. J. and Rietjes, T. H. M. (2007). Multi-objective performance comparison of an artificial neural network and a conceptual rainfall-runoff model. *Hydrological Sciences Journal*, 52(3): 397–413.

Deo, R. C., Samui, P. and Kim, D. (2015). Esti-

- mation of monthly evaporative loss using . relevance vector machine, extreme learning machine and multivariate adaptive regression spline models. *Stochastic Environmental Research and Risk Assessment*, DOI 10.1007/s00477-015-1153-y. , pp. 1-16
- Hazen, A. (1914). Storage to Be Provided in Impounding Reservoirs for Municipal Water Supply. *Trans. ASCE*, .77: 1539-1669
- Kant, A., Suman, P. K., Giri, B. K., Tiwari, M. K., Chatterjee, C., Nayak, P. C. and Kumar, S. (2013). Comparison of multi-objective evolutionary neural network, adaptive neuro-fuzzy inference system and bootstrap-based neural network for flood forecasting, *Neural Computing and Applications*,23(Suppl 1):S231-S246
- Langousis, A. and Koutsoyinnanis, D. (2006). A stochastic methodology for generation of seasonal time series reproducing overheard scaling. *Journal of Hydrology*, 322(48):138–154.
- Leedal, D., Weerts, A. H., Smith, P. J. and Beven, K. J. (2013). Application of data-based mechanistic modelling for flood forecasting at multiple locations in the Eden catchment in the National Flood Forecasting System (England and Wales), *Hydrology and Earth System Sciences*, 17:177-185
- Koutsoyinnanis, D. and Montanori, A. (2007). Statistical analysis of hydroclimatic time series: uncertainty and insights. *Water Resources Research*, 43(5): 5429:1–9.
- Mandelbrot, B. B. (1965). Une Class de processus stochastiques qui sont homogènes à soi-même. Application à la loi climatologique. *Comptes Rendus, Academic Science*. 260:3284–3277.
- Mejia, J. M., Rodriguez-Iturbe, I. and Dawdy, D. R. (1972). Streamflow Simulation; The Broken Line Process as a Potential Model for Hydrologic Simulation. *Water Resources Research*, 8(4):931-941
- Pinto, W. P., Lima, G. B. and Zanetti, J. B. (2015). Comparative Analysis of Models for Times to Series Modeling and Forecasting of Scheme of Average Monthly Streamflow of the Doce River, Colatina Espirito Santo, Brazil. *Ciência e Natura*, 37(4): 1-11
- Sudler, C. E. (1927). Storage Required for the Regulation of Streamflow. *Trans. ASCE*, 91: 622-704
- Thomas, H. A. S. and Fiering, M. B. (1962). Mathematical Synthesis of streamflow sequences for the analysis of river basins by simulation. In *Design of Water Resources Systems*, by A Maass, et al. Harvard University Press, Cambridge, Mass.
- Ursu, E. and Perea J. C. (2015). Application of periodic autoregressive process to the modeling of the Garonne river flows. *Stochastic Environmental Research and Risk Assessment*. DOI 10.1007/s00477-015-1193-3., pp. 1-11
- Valencia, D. and Schaake, Jr. J. C. (1976). Disaggregation process in stochastic hydrology. *Water Resources Research*, 9:580–585
- Weerts, A. H., Winsemius, H. C. and Verkade, J. S. (2011). Estimation of predictive hydrological uncertainty using quantile regression: examples from the National Flood Forecasting System (England and Wales), *Hydrology and Earth System Science*, 15(1):255-265
- Yevjevich, V. (1972). *Probability and Statistics in Hydrology*. Water Resources Publications, Fort Collins, Colorado
- Zappa, M., Jaun, S., Germann, U., Walser, A. and Fundel, F. (2011). Superposition of three sources of uncertainties in operational flood forecasting chains, *Atmospheric Research*, 100(2&3):246–262