

Original Article

HEPATITIS B AND C: AN ASSESSMENT OF RISK EXPOSURE AND PREVALENCE AMONG PRECLINICAL MEDICAL STUDENTS IN NORTHWESTERN NIGERIA.

Musa Abubakar Kana, Nanben Victoria Omole, Awawu Grace Nmadu, Istifanus A. Joshua,
Zainab Kwaru Muhammad-Idris .
Department of Community Medicine, Kaduna State University, Kaduna, Nigeria.

ABSTRACT**Background**

The risk of transmission of hepatitis B and C (HBV and HCV) infection is of particular concern among medical students training in high endemic regions. In this study, we investigated the risk and prevalence of hepatitis B and C viruses among preclinical medical students in a government-owned medical school in northwestern Nigeria.

Methods

We employed a cross-sectional study targeting the whole population of preclinical medical students of Kaduna State University, Kaduna State northwestern Nigeria. Data were collected using a **self-administered, semi-structured questionnaire and** blood testing for HBsAg and HCV antibodies using rapid screening kits. **Data were analyzed using SPSS version 22.**

Results

A total of 133 students participated, and 79.7% of them had a previous risk exposure to injuries by sharps. The seroprevalence for HBV and HCV was 0.8% for both infections. Complete HBV vaccination uptake (3 doses) before enrolment in medical school was 18.8%. The knowledge that HBV infection is preventable was a significant determinant for the previous testing for hepatitis B and C.

Conclusion

Risk exposure to injuries by sharps was high in preclinical students. But their seroprevalence for both hepatitis B and C were low.

Keywords: Hepatitis B and C viruses; prevalence; medical students; sharp injuries

NigerJmed 2020; 62-68
© 2020. Nigerian Journal of Medicine

INTRODUCTION

Hepatitis B and C viruses (HBV and HCV) cause infections that affect the liver. Worldwide HBV is the most common cause of chronic liver disease, including cirrhosis and hepatocellular cancer.¹ Globally, there are about 2 billion people infected with HBV and an estimated 360 million with chronic infections. Annually, the disease also accounts for the death of about 1 million people.^{3,4} Nigeria is one of the countries in which HBV is highly endemic with an estimated 12% of the population being chronic carriers, although studies among selected populations report varying prevalence rates.^{5,6} HBV is spread between people usually through contact with the blood or other body fluids of infected persons, among other modes of transmission. Healthcare workers and medical students are at risks of infection through occupational exposure to blood and body fluids.^{7,8} HCV remains a large healthcare burden to the world, and it

has been estimated to have a worldwide prevalence of about 3% with the virus affecting about 170 million people.⁹ The prevalence of HCV infection is greatest in the low-income countries of sub-Saharan Africa and south-east Asia, where about 8-10% of the population are chronic carriers. The disease is transmitted primarily through parenteral exposure (bloodborne). Sexual transmission almost certainly occurs, but its public health importance is not well quantified. HCV-infected persons can remain asymptomatic for many years following initial infection; however, the virus is a major cause of acute hepatitis and chronic liver disease; including cirrhosis, liver cancer and decompensated liver failure.

Medical students are at a high risk of acquiring blood-borne infections including HBV and HCV, as, during the course of their clinical work, they are in direct contact with patients, blood, injections and surgical instruments. They are also at risk of workplace accidents that occur during the period of their practical training. The lack of professional skills and experience on the part of the students increases their risk exposure in the course of invasive medical procedures. Previous studies have shown that medical and para-medical students lack adequate knowledge about hepatitis B and C.¹⁰⁻¹³ Other studies showed prevalence ranges of between 4.2 to 12.5% for HBV and 1.5 to 3.6% for HCV and a vaccination rate of 19.2% for hepatitis B amongst medical students.¹⁴⁻¹⁷ Vaccination is the best means

Correspondence to :

Dr Musa Abubakar Kana,
Department of Community Medicine,
Kaduna State University, Kaduna, Nigeria.
E-mail: musakana77@yahoo.com

by which one can arm oneself against hepatitis B infection. The transmission rate is as high as 30% in those who are not immune but rare in those who have been immunized.¹⁸

Medical students come into contact with patients immediately they commence the clinical phase of medical school. It is relevant that sufficient consideration be given to prevention measures like vaccination and health education. These measures will be most effective if it targets the medical students prior to their first contact with patients. The vaccination against hepatitis B is recommended to minimize the risk of this occupational infection. Although the exposure to the clinical environment of the hospital will increase the risk of exposure to bloodborne viral hepatitis for medical students, the availability of information about their seroprevalence status during the preclinical phase of their training will provide evidence for the appropriate timing for the hepatitis B vaccination. Therefore, the objective of this study was to determine the prevalence and risk exposure of hepatitis B and C virus infections of pre-clinical medical students in a government-owned medical school.

Methods

Study Design

A cross-sectional study was conducted between January-March 2013 among pre-clinical, medical students of Kaduna State University (KASU), Kaduna, Nigeria.

Study Location and Population

Kaduna State University is sited in Kaduna, which is the capital of Kaduna State, located in northwestern Nigeria. The medical school of this university was established in 2007 with an initial intake of 46 students in 2009. At the time of conducting the study in 2013, the number of students had grown to 148, distributed across three levels of study. The study population included all registered pre-clinical medical students of the university.

Sampling Method

A whole population sampling method was used for the study, and all medical (pre-clinical) students of the university who were duly registered at the time of the study and gave informed consent were included in the study. Thus, a total sample population of 133 respondents (representing 89.9% of the student population) participated in the study.

Data collection

A self-administered, semi-structured questionnaire was used to collect information about: 1) Respondents' socio-demographic characteristics including age, gender, year of study in medical school, previous enrolment in other medical school(s) and previous employment as a health worker. 2) Previous risk exposure with regard to sharp injuries, including needles. 3) Previous testing and outcome for hepatitis B and Hepatitis C.

Before commencement of data collection members of the

research team composed of three public health physicians and two laboratory scientists were trained on the questionnaire and its mode of administration. The principal investigator supervised the overall data collection activities. Informed consent was obtained from all medical students that consented, but three students completed the questionnaire but declined to have their blood samples taken for undisclosed reasons. The questionnaire administration and taking of blood samples were done during afternoon breaks on consecutive days until all the students that consented were reached. Blood samples for testing were obtained from participants through a finger prick with a lancet under sterile conditions. Thereafter, laboratory testing for hepatitis B and C viral infections was performed on the samples.

Laboratory Testing:

For the diagnosis of hepatitis B virus and hepatitis C virus infections, finger prick specimens of whole blood were obtained from the participants with a lancet after cleaning the area to be lanced with an alcohol swab. Testing of each participant's whole blood sample for HBsAg and HCV antibodies were done using Micro-point HBsAg and HCV antibodies Gold Rapid Screening kits according to the manufacturer's instruction. Each test kit is a chromatographic immunoassay (CIA) for the direct qualitative detection of HBsAg and HCV antibodies. Similar rapid screening tests were also used in a study to assess the prevalence of hepatitis C in Makurdi, north-central Nigeria and the prevalence of hepatitis B and C and co-infection with HIV in Malawi.^{19,20}

Data Analysis:

The questionnaires were checked for completeness and consistency. Data were entered into data entry templates and analyzed using Statistical Package for Social Sciences (SPSS), version 23.0 (IBM Corp., New York, USA). Main outcome variable was hepatitis B and hepatitis C serostatus while the independent variables were socio-demographic characteristics and risk factors for hepatitis B and C. Descriptive statistics such as mean and standard deviation were generated for variables including frequencies and percentages for categorical variables. Bivariate analysis was carried out, and the Chi-square (χ^2) test was used to test associations between variables. A p-value of less than 0.05 was used as the cut off level for statistical significance.

Ethical Considerations:

The Ethics Research Committee of the Kaduna State Ministry of Health approved the research protocol. Permission was also obtained from the management of the Kaduna State University prior to conducting this study. Written informed consent was obtained from all students that consented to participate. Confidentiality was assured for all information provided and for the results of the investigations. Subsequently, administration of hepatitis B vaccine to HBV-negative individuals was done. Those who tested positive were linked with the health care system for further management.

Results

A total of 133 preclinical students participated in the study. This consists of 85 (63.9%) males and 48 (36.1%) females. Risk exposure and awareness of the consequences of the same were over 75% among participants (Table 1). Only 30.5% of the respondents were previously tested for HBV, out of which 2 (5.0%) tested positive for HBV. The prevalence of those that previously tested for HCV was 33.6% and the seropositivity of 4.5% (Table 2). Only one respondent tested positive for each antigen (with a seroprevalence of 0.8% for each antigen) during current testing; the majority of participants (96.2%) tested negative for both HBV and HCV. Almost one-third (29.0%) of the participants had previously been vaccinated against HBV, 65.8% of whom had received 3 doses of the vaccine; while the majority of same (73.7%) had a fairly recent (<5 years) history of HBV vaccination.

Among the evaluated socio-demographic factors, only knowledge that viral hepatitis is preventable is positively associated with the previous testing of both hepatitis Band C (Table 3). Meanwhile, gender, age, year of study in medical school, previous enrolment in medical school or employment as a health worker, and knowledge of hepatitis being preventable of the medical student were not associated with previous vaccination against hepatitis B (Table 4).

Discussion

The health risks associated with patient care are a recognized occupational hazard in medical and paramedical professions globally; as accidental transmissions of infectious diseases have been reported in developing and developed countries.²¹ The developing countries are of particular concern because over 90% of such occupational exposures among health workers occur in this setting.²² Fundamentally, the risk of these occupational mishaps have been observed to be more among novices in the profession such as students, interns and other categories of trainees¹². This has been attributed to several factors such as inadequate knowledge, lack of experience and professional expertise, fatigue, stress, anxiety, lack of supervision and tutorial support from older professionals etc.^{11-12,22}

Invasive medical procedures involving the use of sharp instruments particularly with ungloved hands exposes healthcare workers and students to the blood and other body fluids of patients, which may be infectious in nature.^{7,23-24,25} These exposures increase the risk of blood-borne infections like hepatitis B and C viruses.²⁶ Our study finding indicates a high prevalence of previous risk exposure due to injury by sharps among pre-clinical medical students. This is plausible considering the common use of sharp objects for personal (shaving, barbing and nail cutting), domestic and occupational purposes as well as cultural practices like scarification marks in the study setting. However, our result contrasts with lower prevalence as reported in other studies conducted among medical students as well as students and

interns of other paramedical and health-related professions in developing countries, including Nigeria.²⁸⁻³¹ A much lower prevalence has been in the same category of students in developed nations.^{32,33} Generally, developing countries have prevalence level similar to the global prevalence of needle stick injuries among medical students, which is lower than our finding.³⁴ Nevertheless, exceptionally higher prevalence has been reported in some developing nations.^{22,35}

It is remarkable that only a negligible number of the participants of this study had either been enrolled in a medical school or served as health workers in the past. Hence, the high prevalence of previous exposure to injury by sharps obtained in this study is rather alarming for preclinical students because most needle stick injuries have been documented to occur during clinical training.³⁶ Further investigation will be required to assess the risk exposure to sharp injuries in pre-clinical years, particularly during anatomy dissection session or Physiology practical classes. This will provide evidence about the suitability of instituting pre-exposure prophylactic vaccination for medical students at the onset of medical school rather than wait until the beginning of clinical training.³⁰

The knowledge of the potential dangers and risks posed by such injuries by sharps, including needles for infection with hepatitis B and C was high among our study participants. This finding compares with studies among medical students in other parts of Nigeria.^{28,17} On the contrary, there are studies that showed limited or inadequate knowledge regarding sharps injuries among medical and dental students in Scotland and India.^{32,37} Precisely, a lack of awareness among medical students entering into the profession and a significant difference between knowledge of pre-clinical and clinical students has been observed.^{11,13} In spite of this relatively good level of awareness among the cited Nigerian studies, the continuous teaching and training of medical students on the risks and hazards associated with injuries with sharps. Additionally, the measures to be taken to avert them as well as manage them in the event of their occurrence should be encouraged as the mainstay of prevention of occupationally-acquired infections.

An estimated hepatitis B seroprevalence of one-tenth the adult population has been reported for sub-Saharan Africa, including Nigeria.²⁷ Thus, the laboratory testing to ascertain the hepatitis B and C status of individuals is key to prevention and control of viral hepatitis. In this study, only one-third of the participants had previously been tested for both hepatitis B and C. This low proportion of those that had previously tested for hepatitis B and C among the participants may be a reflection of the poor practice of routine or voluntary medical examinations in developing countries.³⁸ Our finding corroborates other Nigerian studies for low previous testing of hepatitis B and C even though a higher proportion had been screened for HIV at least once.³⁹

The voluntary submission to routine medical examinations is uncommon among Nigerians as well as

nationals of many other developing nations. In this study, the knowledge that hepatitis B is a preventable disease was significantly associated with the previous testing for hepatitis B and C. Contrary to expectation the proportion of those that had previously tested for hepatitis C were more than for hepatitis B. Generally, there is a higher level of awareness and knowledge in the larger population, of HBV infection as a disease entity, as compared to HCV.⁴⁰ Likewise, the World Health Organization reports that prevalence data and studies on HBV abound as compared with HCV, which has a relative paucity of information in circulation; and therefore, a lower level of awareness and understanding in the general population.⁹ Furthermore; this situation is also reported to have engendered some confusion between HBV and HCV within certain populations.⁴¹ This might have contributed to the resultant outcome reflected in this study.

Majority of our study participants had a negative outcome for current testing, with only a participant that tested positive for either hepatitis B or C. It is documented that with a seroprevalence of over 10% reported from various studies and surveys, Nigeria is classified as a high endemic region.^{5,6,42,43} The seroprevalence for HBV obtained in this study is extremely lower than that reported by various authors working among undergraduate students in different parts of Nigeria.^{15,44-47} The low seroprevalence we found may be due to the fact that the entire study population are pre-clinical students, who are yet to commence direct interactions with patients or the clinical environment with its attendant risks.

Pre and immediate post-exposure prophylactic vaccination with hepatitis B vaccine are globally recognized and advocated measures for the prevention and control of HBV infection. In this study, only about one-third of the participants had previously been vaccinated against HBV, and only 66% of them had received the recommended 3 doses of the vaccine.⁴⁸ The observed inadequate or sub-optimal vaccination is comparable with studies among medical students in Nigeria and other parts of sub-Saharan Africa.^{17,30,28,49} These sampled studies reported values lower than among medical students and personnel in developed countries.^{7,8} In the United States, for instance, the Centre for Disease Control

and Prevention (CDC) has since set a goal of achieving 98% hepatitis B vaccination coverage among health workers, with the intention of eliminating occupationally-acquired HBV infection in that country.⁵⁰ Therefore, the goal of forestalling the huge morbidity and mortality consequences on human resource for health should be enhanced by advocacy for pre-exposure vaccination and instituted as mandatory for all medical students and personnel.

Finally, the majority of our participants had recent (1-4 years) history of HBV vaccination, which corresponded with the period of their admission to medical school. Whether this is merely a coincidence or an initiative taken up by these respondents as a prophylactic measure, in view of their enrollment into medical school was not further explored because of its non-inclusion in the scope of this study. Personal motivation could explain this behaviour, but none of the socio-demographic factors was found to significantly determine previous vaccination against HBV. The knowledge that HBV is preventable was positively associated with previous testing for the disease and not for vaccination. This might suggest other barriers to HBV vaccine uptake among medical students. These factors include limited or no opportunity, lack of time (busy or tight schedule), financial constraints (out-of-pocket expenses either for the vaccine or the consumables used, or both), information gaps (such as not aware a vaccine exists, fears and concerns regarding possible vaccine side effects etc.), indifference and procrastination.^{28,30} As valid and genuine as some of these factors might be, all efforts need to be put in place to ensure higher HBV vaccine uptakes for medical students and health care workers in moderate to high endemic areas.

Conclusion

The risk exposure to injuries by sharps, including needlestick, was observed to be high among preclinical students. Fortunately, the seroprevalences recorded for both hepatitis B and C were low. However, the low hepatitis B vaccine uptake in the context of a high endemic adult population constitutes a major public health problem. Consequently, we recommend regular training on universal precaution and hepatitis B vaccination early in medical training.

Table 1: Previous exposure and testing for hepatitis B and C viruses among preclinical medical students

Variable	Frequency (%)
Previous needle or sharp injury	106 (80.9)
Aware of the consequences of sharp injuries	102 (77.9)
Previously tested for hepatitis B	40 (30.5)
Previously tested positive for hepatitis B	2 (5.0)
Previously tested for hepatitis C	44 (33.6)
Previously tested positive for hepatitis C	2 (4.5)
Tested positive for only HBV in current testing*	1 (0.8)
Tested positive for only HCV in current testing	1 (0.8)
Tested positive for both HBV and HCV in current test	0

*Declined consent for current testing=3 (2.3%); Some variables have missing records

Table 2. Prevalence of previous vaccination against hepatitis B virus among preclinical medical students

Variable	Frequency (%)
Previously vaccinated (n=131)	
Yes	38 (29.0)
Number of Doses Given (n=38)	
1	4 (10.5)
2	9 (23.7)
3	25 (65.8)
Time interval since previous vaccination in years (n = 38)	
1-2	8 (21.1)
3-4	20 (52.6)
5	10 (26.3)

Some variables have missing records

Table 3: Association between socio-demographic characteristics and previous testing for hepatitis B and C viruses among preclinical medical students

Characteristic (n=131)	Previous testing for hepatitis B		P-value	Previous testing for hepatitis C		P-value
	Yes (n=40)	No (n=91)		Yes (n=44)	No (n=87)	
Gender						
Female	14 (35.0)	31 (34.1)	1.000	11 (25.0)	34 (39.1)	0.123
Male	26 (65.0)	60 (65.9)		33 (75.0)	53 (60.9)	
Age in years						
<20	8 (20.0)	18 (20.0)	0.165	10 (22.7)	16 (18.6)	0.211
20-24	23 (57.5)	63 (70.0)		25 (56.8)	61 (70.9)	
25	9 (22.5)	9 (10.0)		9 (20.5)	9 (10.5)	
Year of study in medical school						
1st	7 (17.5)	20 (26.4)	0.341	11 (25.0)	20 (23.0)	0.499
2nd	13 (32.5)	22 (24.2)		15 (34.1)	20 (23.0)	
3rd	15 (37.5)	26 (28.6)		11 (25.0)	30 (34.5)	
4th	5 (12.5)	19 (20.9)		7 (15.9)	17 (19.5)	
Ever been previously enrolled in a medical school						
Yes	3 (7.5)	3 (3.3)	0.262*	3 (6.8)	3 (3.4)	0.322*
Previous employment as health worker						
Yes	2 (5.0)	3 (3.3)	0.485*	2 (4.5)	3 (3.4)	0.547*
Knowledge of hepatitis B is preventable						
Yes	36 (90.0)	33 (36.3)	0.003*	40 (11.8)	54 (62.1)	<0.05*

*Fischer's exact test; Some variables have missing records

Table 4: Association between socio-demographic characteristics and previous vaccination against Hepatitis B among preclinical medical students

Characteristic (n=133)	Previous vaccination against Hepatitis B		P-value
	Yes (n=38)	No (n=93)	
Gender			
Female	12 (31.6)	33 (35.5)	0.669
Male	26 (68.4)	60 (64.5)	
Age in years			
<20	8 (21.1)	18 (19.6)	0.093
20-24	21 (55.3)	65 (70.7)	
25	9 (23.7)	9 (9.8)	
Year of study in medical school			
1st	6 (15.8)	25 (26.9)	0.280*
2nd	13 (34.2)	22 (23.7)	
3rd	14 (36.8)	27 (29.0)	
4th	5 (13.2)	19 (20.4)	
Ever enrolled in medical school			
Yes	3 (7.9)	3 (3.2)	0.246*
Previous employment as health worker			
Yes	2 (5.3)	3 (3.2)	0.581
Knowledge of hepatitis B is preventable			
Yes	33 (86.8)	61 (65.6)	0.014*

*Fischer's exact test; Some variables have missing records

References

- Lai CL, Ratzu V, Yeun MF, Poynard T. Viral hepatitis B. *Lancet* 2003;362(9401):2089-94.
- Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: global burden of disease study. *Lancet* 1997;349(9061):1269-76.
- Drosten C, Nipprasche T, C Manegold, H Mensel and V Brixner. Prevalence of HBV DNA in anti-HBV positive/HBsAg-negative sera-correlate with HCV but not HIV serostatus. *J of Virol* 2004; 29:59- 68: 1-4.
- Ziraka AK, Bwogi J, Namale A, Wainaina CW and Mayanja-Kizza H. Sero-prevalence and risk factors for hepatitis B virus infection among healthcare workers in tertiary hospital in Uganda. *BMC infectious Dis*2010; 10:191-6.
- Mbaawuga EM, Enebeaku MNO, Okopi JA and Damen JG. Hepatitis B virus (HBV) infection among pregnant women in Makurdi, Nigeria. *Afric J of Biomed Res* 2008; 11:155-9.
- Alao EE, Okwori EE, Egwu E and Audu F. Sero-prevalence of hepatitis B surface antigen among prospective blood donors in urban area in Benue State. *Internet J of Haematol* 2009; vol5.
- Deisenhammer S, Radon K, Nowak D and Reichert J. Needlestick injuries during medical training. *J Hosp Infect* 2006; 63: 263-67.
- Simard EP, Miller JT, George PA, Wasley A, Alter MJ, Bell BP and Finelli L. Hepatitis B vaccination coverage levels among healthcare workers in the United States 2002-2003. *Infect Control Hosp Epidemiol* 2007; 28: 783-90.
- World Health Organisation (WHO). Global surveillance and control of hepatitis C. Report of a WHO Consultation organized in collaboration with the viral hepatitis prevention board, Antwerp, Belgium. *J Viral Hepat* 1999; 6:35-47.
- Al-faleh FZ and Fachar ZT. Hepatitis B infection in Saudi Arabia. *Ann Saudi Med* 1988; 8:474-80.
- Anjum Q, Siddiqui H, Ahmed Y, Rizvi SR and Usman Y. Knowledge of students regarding hepatitis and HIV/AIDS of a private medical university in Karachi. *J Pak Med Assoc* 2005; 55(7):285-8.
- Osborn EH, Papadakis MA, and Gerberding JL. Occupational exposures to body fluids among medical students: a seven-year longitudinal study. *Ann of internal med* 1999; 130(1):45-51.
- Eisa ZM, Eifan SA and Al-sum BA. Awareness of viral hepatitis B and C infection among first-year medical science students in Jazan University. *Pub Health Res* 2012; 2(5):161-73.
- Pido M and Kagimu M. Prevalence of hepatitis B virus infection among Makerere University medical students. *Afric Health Sci* 2005; 5(2):92-8
- Pennap GR, Nwachukwu O, Ishaleku D and Ombugadu RJ. Hepatitis B virus carriage among students of a Nigerian tertiary institution: a cohort of eligible blood donors. *Res J of Med Sci* 2011; 5(2):90-3.
- Ugbabor O, Aigbiriori M, Osazuwa F, Enabudoso and Zabayo D. The prevalence of hepatitis viral infections among pregnant women. *North Am J of Med Sci* 2011; 3(5):238-41
- Peterside B, Duru CO, Adeyemi OO, Kunle-Olowu OE, Kunle-Olowu AO and Akinbami FO. Hepatitis B vaccination rate among medical students at the Niger Delta University Teaching Hospital, Bayelsa State, Nigeria. *Global Adv Res J of Med & Med Sci* 2012; 1(10): 280-85.
- Centres for Disease Control and Prevention (CDC). Recommendations for preventing transmission of human immunodeficiency virus and hepatitis B virus to patients during exposure-prone invasive procedures. *MMWR Recomm Rep* 1991; 40:1-9.
- Achingi GI, Malu AO, Mbaave PT, Bitto TT, Shaahu VN, Mohammed H and Misauno MA . Prevalence of hepatitis C in Makurdi, north-central Nigeria. *IOSR J of Dent & Med Sci (IOSR-JDMS)*2013; 7(5):06-10
- Nyirenda M, Beadsworth MJB, Stephany P, Hart CA, Hart IJ, Munthali C, Beeching NJ and Zijlstra EE. Prevalence of infection with hepatitis B and C virus and co-infection with HIV in medical in-patients in Malawi. *J of Infect*2008; 57(1):72-7
- Taylor R, Sladden T, Levy S, Gust I, Macaskill P, Rushworth L and Gaxibarich G: A seroepidemiological study of hepatitis B amongst Fiji health care workers. *SE Asian J Trop Med Pub Health* 1991; 22(4):567-76.
- World Health Report. Reducing Risks, promoting healthy life. WHO: Geneva (CH); 2002. Available from: http://www.who.int/whr/2002/en/whr02_en.pdf?ua=1. (Last accessed 13th November 2017).
- Pruss-Ustun A, Rapiti E and Hutin Y: Estimation of the global burden of disease attributable to contaminated sharps injuries among health-care workers. *Am J Ind Med* 2005; 48(6):482-90.
- Askew SM. Occupational exposures to blood and body fluids: a study of medical students and health professions students in Virginia. *Workplace and Health Safety J* 2007; 5 5 (9) : 3 6 1 - 7 1 (D O I : <https://doi.org/10.1177/216507990705500904>)
- Ghasemzadeh I, Kazerooni M, Davoodian P, Hamed Y and Sadeghi P. Sharp injuries among medical students. *Glob J Health Sci* 2015; 7(5):320-25.
- Tarantola A, Abiteboul D and Rachline A. Infection risks following accidental exposure to blood or body fluids in health care workers: a review of pathogens transmitted in published cases. *Am J Infect Control* 2006; 34:367-75
- World Health Organization (WHO). Hepatitis B. 2002, Geneva, WHO, Department of Communicable Disease Surveillance and Response
- Okeke E, Ladepe NG, Agaba E and Malu AO. Hepatitis B vaccination status and needle stick injuries among medical students in a Nigerian university. *Niger J Med* 2008; 17(3):330-2
- Osazuwa-Peters N, Obarisiagbon A, Azodo CC, Ehizele AO and Obuekwe ON. Occupational exposure to sharp injuries among medical and dental house officers in Nigeria. *Int'l J of Occup Med and Env Health* 2013; 26(2):283-90
- Noubiap JJN, Nansseu JRN, Kengne KK, Ndoula ST and Agyingi LA. Occupational exposure to blood, hepatitis B vaccine knowledge and uptake among medical students in Cameroon. *BMC Med Educ* 2013; 13: 148
- Díaz ML and Cadena ALP. Risk of hepatitis B infection in Peruvian medical students following occupational exposure to blood and body fluids. (2002). *Rev Gastroenterol Peru* 2003; 23(2):107-10
- Varsou O, Lemon JS and DickFD. Sharps injuries among medical students. *Occup Med* 2009; 59(7):509-11
- Schmid K, Schwager C and Drexler H. Needlestick injuries and other occupational exposures to body fluids amongst employees and medical students of a German university: incidence and follow-up. *J of Hosp Infec* 2007; 65(2):124-30
- Sharma GK, Gilson MM, Nathan H and Makary M. Needlestick injuries among medical students: incidence and implications. *Acad Med* 2009; 84(12):1815-21
- Liyanaage IK, Caldera T, Rwna R, Liyange CK, De Silva P and Karunathilake IM. Sharps injuries among medical students in the faculty of medicine, Colombo, Sri Lanka. *Int'l J of Occup Med and Env Health* 2012; 25(3):275-80
- Siegmann S, Muth T, Kluth W, Hofbauer U, Angerer P and Schwarze S. Needlestick Injuries to medical students.

- Gesundheitswesen* 2016;78(1):22-7.
37. Kumar S. Knowledge, attitude and practices regarding needlestick injuries among dental students. *Asian J of Pharm and Clin Res* 2016; 9(4):312-15
 38. World Health Organisation (WHO). Hepatitis B. Bulletin of the World Health Organization 2013; 91:242-43.
 39. Oladeinde BH, Ekejindu IM, Omoregie R, and Ikpomwonosa O. Uptake of HIV, HBV and HCV testing services among medical laboratory scientists in Nigeria. *New Zealand J of Med Lab Sci* 2014; 68(2):54-9
 40. Gautier A, Jestin C and Jauffret-Roustide M. Perception and knowledge of viral hepatitis: results of the Nicolle Survey, France, 2006. *Bull Epidemiol Hebd* 2009; 20-21:208-11
 41. Brouard C, Gautier A, Saboni L, Jestin C, Semaille C and Beltzer N. Hepatitis B knowledge, perceptions and practices in the French general population: the room for improvement. *BMC Pub Health* 2013; 13:576-85
 42. Bukbuk DN, Bassi AP and Mangoro ZM. Sero-prevalence of hepatitis B surface antigen among primary school pupils in rural Hawal valley, Borno State, Nigeria. *J Comm Med PHC*. 2005; 17(1):20-3
 43. McMahon BJ. Natural history of chronic hepatitis B-clinical implications. *Medscape J Med* 2008; 10:91-101
 44. Mbotto CI and Edet EJ. Prevalence and risk factors of hepatitis B virus infection among students in University of Uyo. *Int'l J Modern Biol Med* 2012; 2(2): 101-11
 45. Imarenezor EPK, Brown STC, Yakubu OE and Soken DC. Survey of hepatitis B and C among students of Federal University Wukari, Taraba State, Nigeria. *Int Res J Med Med Sci* 2016; 4(3):31-3
 46. Musa BM, Bussell S, Borodo MM, Samaila AA and Femi OL. Prevalence of hepatitis B virus infection in Nigeria, 2000-2013: a systematic review and meta-analysis. *Nig J of Clin Pract* 2015; (18)2:163-72
 47. Chhabra P, Grover VL and Agrawal K. Do our medical students have enough knowledge of Hepatitis B? A Delhi based study. *J Commun Dis* 2002;34:221-5
 48. Centres for Disease Control and Prevention. Hepatitis B. In Hamborsky J, Kroger A, Wolfe S, (Editors). *Epidemiology and Prevention of Vaccine-Preventable Diseases*. 13th edition 2015. Public Health Foundation. Washington D.C
 49. Odusanya OO, Meurice FP and Hoet B. Nigerian medical students are at risk for hepatitis B infection. *Trans R Soc Trop Med Hyg* 2007;101(5):465-8.
 50. US Department of Health and Human Services. *Healthy People 2010 with Understanding and Improving Health and Objectives for Improving Health*. 2 vols. 2nd ed. Washington, DC: US Government Printing Office; 2002