

Taxonomic diversity of benthic macroinvertebrates along the Oum Er Rbia River (Morocco): implications for water quality bio-monitoring using indicator species

F. Souilmi^{*1,2}, K. Ghedda¹, A. Fahde², F. Z. EL Fihri², S. Tahraoui³, F. Elasri² and M. Malki²

¹ National Office for Electricity and drinking Water (ONEE), Rabat, Morocco

² Department of Biology, University Hassan II, Faculté des Sciences Ain Chock, Casablanca, Morocco

³ Department of Biology, Université Chouaib Doukkali faculté des sciences El jadida, Morocco

*Corresponding Author: fatimasouilmi@gmail.com

Abstract

The macroinvertebrates of the Oum Er Rbia River were studied from samples collected seasonally from September 2015 to September 2016 at 10 sampling sites. The macroinvertebrates found during the sampling period were distributed into twelve orders. The most abundant order was diptera, having 9618 individuals, followed by the order Ephemeroptera with 2985 individuals. Coleoptera, odonates and crustaceans represent only a small fraction of the total fauna. Hydropsyche, Chironomidae sp. and Simuliidae are numerically more inventoried. The composition and distribution of the species were directly or indirectly affected by the physicochemical variables and the quality of the habitat. Correspondence analysis results showed that habitat quality and quality of water represented species distribution patterns and species can be used as indicators to assess the quality of the Oum Er Rbia River system. Habitat management along the Oum Er Rbia river should be aimed at preserving native species, especially during the summer, when the biotope requirements are optimal. The results obtained in this study showed an alarming situation of the water quality of the Oum Er Rbia River and particularly in downstream segment. To solve this problem, we recommend the development of the wastewater discharge of Khenifra and Kasba Tadla and the purification of wastewater before it is discharged into the river.

Introduction

The identification of environmental factors influencing the composition and abundance of stream biological communities is an important point in a stream restoration (Richards *et al.*, 1993). Water chemistry and physical characteristics of stream have an impact on river communities (Meyer *et al.*, 1988, Resh and Rosenberg, 1984) and the characteristics of the riparian zone affecting invertebrate communities (Gregory *et al.*, 1991). Obviously, the distribution of benthic macroinvertebrates is impacted by environmental factors (Benbow *et al.*, 2003) as well as chemicals and physical factors produced by human activities (Prat and Ward 1994). The substrate, flow regime, geomorphology and temperature are the physical factors that affect river macro-invertebrate communities (Minshall, 1984; Poff & Ward, 1989, Huryn & Wallace 1987; Stannz, Gore and Resh, 1988, Vannote & Sweeney, 1980, Ward & Stanford,

1982;). Macroinvertebrate communities change spatially and temporally (Townsend & Hildrew 1994) and the combination of local environmental characteristics and large-scale geographical factors bring occurrence of a species (Townsend *et al.*, 2003; Mykra, Heino & Muotka, 2007). Knowledge about distribution, diversity and composition of macro invertebrate assemblages in gravel-water interfaces are very important, because they are highly dependent on sediment particle size (Beauger *et al.*, 2006). Besides, stream bed structures, such as clusters and gravel pits that create stable habitats for benthic communities, affect the growth of macro invertebrates and biodiversity (Duan *et al.*, 2009). Urbanization, deforestation, construction, irrigation, drainage and pollution affect aquatic habitats. (Dudgeon, 2008). The organisms which live in the water respond to all environmental stresses (Veroli *et al.*, 2010), due to all these traits. Benthic macroinvertebrates have been

widely used as indicators of water quality in river management. (Hellawell 1986, Metcalfe 1989, Jeffries and Mills 1990, Ro-senberg and Resh 1993), but the way functional diversity of macroinvertebrate assemblages influence processes in freshwater ecosystems needs to be broadened (De'nes *et al.*, 2016).

The use of macroinvertebrates in bioassessments has increased significantly (Cairns & Pratt, 1993 and Lenat & Resh, 2001). Macroinvertebrates represent a key indicator group for monitoring environmental change (Neville & Yen, 2007), but there is still much debate about the level of taxonomic identification and the sample size needed to determine anthropogenic impacts (Resh and McElravy, 1993) though the results depend on the particular objectives of the study (Daniel F. Buss and Anderson S. Vitorino, 2010). Generally, literature suggests that family classification is sufficient to detect impacts from point sources and other gross impacts in freshwater and marine systems (Wright *et al.* 1995, Vanderklift, Ward & Jacoby, 1996). Many studies have shown that the lowest level taxonomic resolution (genus) usually does not improve detection of impairment because of its reduced community variability when decreasing taxonomic resolution to the family level (Bowman and Bailey, 1997). Yet, the genre gave the same description of community models as family or order.

Although identification of species have been more successful at classifying streams than the family, Furse *et al.* (1984) suggested that the family was sufficient to detect significant site-related environmental gradients. They also suggested that family identification could offer substantial savings in time and money over species identification. (Marchant, Barmuta and Chessman, 1995) found that the community pattern among 40 streams along an altitudinal gradient was preserved when quantitative species data were converted to quantitative or qualitative family data.

The main objective of the study was to establish as complete an inventory as possible of the different taxa that may be encountered in the aquatic system of the Oum Er Rbia River with the view to:

- enriching the list of Moroccan biodiversity,
- determining the taxonomic level (genus, family),
- detecting the gradient of pollution in Oum Er Rbia River and
- comparing the biodiversity of different freshwater ecosystems.

Materials and methods

Study area

The Oum Er Rbia River, one of the most important in Morocco , is located in the Mid-West of Morocco (Figure. 1). The River




Figure 1: Location of the sampling station

originates from the Middle Atlas Mountains at 1800 m and flows into the Atlantic Ocean at Azemmour city. The Middle Atlas is characterized by a humid cold climate and is classified as Mediterranean mountain climate (Martin, 1981). Rainfall in the Oum Er Rbia basin varies between 1100 mm on the Middle Atlas and 300 mm in the downstream sections, with an average of 550 mm (USAID, 2010). Many dams and reservoirs have been constructed on the Oum Er Rbia River to generate hydroelectric power and to provide water for irrigation.

Sampling sites

The sampling sites (S1, S2, S3, S4, S5, S6, S7, S8, S9, S10) were marked out alongside the Oum Er Rbia Rivers (Figure 1). The samples were taken seasonally from September 2015 to September 2016. At each site, we made qualitative samplings of macroinvertebrates, using a Surber sampler (catching area: 0.025 m²) (Berrahou *et al.*, 2000). At each site, eight samples were taken at different habitats. All captured organisms were placed in plastic bottles and preserved in 10% formaldehyde. The benthic macroinvertebrate identification was done to the lowest possible taxonomic level in the laboratory based on keys presented

by Tachet *et al.*, (2000); Tachet *et al.* (2010). Physico-chemical data and heavy metals were obtained from the National Office for Electricity and drinking Water (ONEE) of Morocco. For each sampling site, data on pH, temperature, electrical conductivity (EC), dissolved oxygen (DO), chemical oxygen demand (COD), biochemical oxygen demand (BOD5) were collected. Total suspended solids (TSS), ammonium (NH₄⁺) and total phosphorus (P) were obtained.

Data analysis

Correlations between biological indices and chemical variables were computed using the Principal Component Factor Analysis of the Correlation Matrix. All statistical analyses were performed using the basis of Ward's method (H.Øyvind *et al.*, 2001).

Results and discussion

The physic-chemical factor values

The physical and chemical characteristics of the sites are presented in Table.1. However, except for Electrical Conductivity and COD, all measured parameters did not show any significant difference among the sampling occasions (Table 1). The measured temperature

TABLE 1
Mean values of physicochemical variables measured in the Oum Er Rbia River

Sites	(COD)	T°	pH	(NH ₄ ⁺) mg N/L	(EC) (µs/cm)	TSS mg/L	BOD5	(DO) mg O ₂ /L
S1	9.25±18.5	18.9±4.6	8.5±0.10	0.04±0.07	3450±685.3	99.8± 119.3	6.75 ± 5.9	6.4±0.6
S2	9.75±19.5	20.65±3.8	8.4±0.03	0.30±0.47	2340±105.5	98.6± 107.9	6.25 ± 4.9	6.9±0.5
S3	9.5±19.0	20.25±3.6	8.4±0.05	0.27±0.42	2390±150.1	106.3± 110.6	6 ± 5.0	6.9±0.4
S4	0±0.0	20.6±3.7	8.4±0.10	0.03±0.06	2405±147.3	1295.3±2470.2	4.5 ± 5.3	6.4±0.5
S5	4.75±9.5	20.55±3.0	8.4±0.11	0.26±0.16	2347.5±111.5	1394.8±2670.6	5.75 ± 7.6	6.5±1.1
S6	98.5±181.4	20.1±3.6	8.4±0.15	0.39±0.21	2310±202.2	1447.8±2768.6	34.5 ± 50.4	6.2±1.5
S7	0±0.0	17.67±2.0	8.4±0.15	0.30±0.13	2220±249.1	34.9±30.2	3.75 ± 4.8	6.2±0.7
S8	0± 0.0	16. 7±2.3	8.2±0.10	0.38±0.28	1987.5±130.0	34.3±28.2	0 ± 0.0	6.1±0.5
S9	0±0.0	20.8±4.8	8.3±0.13	0.45±0.29	2017.5±88.1	23.3±29.9	14.25 ± 7.8	6.0±0.3
S10	0± 0.0	24.55±1.8	8.3±0.09	0.02±0.04	1235±20.8	24.5±30.4	0 ± 0.0	7.3±0.3

Order (cont.)	Family	Code	Sta. 1	Sta. 2	Sta. 3	Sta. 4	Sta. 5	Sta. 6	Sta. 7	Sta. 8	Sta. 9	Sta. 10
Odonata	F/Gomphidae <i>Mesogomphus</i>	Mes	+	+			+	+				
	F/ Calopterygidae <i>Calopteryx sp</i>	Cal	+	+	+	+						
	F/Leptophlebiidae <i>Choroterpes</i>	Cho	+	+	+	+						
	F/Coenagrionidae <i>Coenagrion sp</i>	Coe	+	+	+	+						
	F/ Chironimidae <i>Chironomus sp</i> <i>Tanytarsus sp.</i>	Chi Tan	+	+	+	+	+	+	+	+	+	+
Diptera	F/Simuliidae <i>Eusimulium</i> <i>Simulium</i>	Eus Sim	+	+	+	+	+	+	+	+	+	+
	F/Ceratopogonidae <i>Culicoides sp</i>	Cul	+	+	+	+						+
	F/Tabanidae <i>Tabanus sp.</i>	Tab	+	+	+	+	+	+	+			+
	F/Anthomyidae	Ant	+		+					+	+	
	F/Empididae	Emp	+	+	+	+				+		
	F/Rhagionidae	Rha	+	+	+							
	F/Tipulidae	Tip	+	+	+							
	F/ Athericidae	Ath	+	+	+							
	F/Stratiomyidae <i>Stratiomys sp.</i>	Str	+	+	+	+						
	F/Scatophagidae	Sca	+	+	+	+				+		
	F/Psychodidae	Psy	+	+	+	+				+		
	F/Ephydriidae	Eph	+	+	+							
	F/Limoniidae	Lim	+	+	+							
	F/Syrphidae	Syr	+	+	+							
	F/Dixidae	Dix	+	+	+							
	F/Ptychoptera <i>Ptychoptera sp.</i>	pty	+	+	+							
Trichoptera	F/Hydropsychidae <i>Hydropsyche sp</i>	Hyd	+	+	+	+	+	+	+	+	+	
	F/ Hydroptilidae <i>Orthotrichia sp</i>	Ort	+	+	+	+	+			+		+
	F/philopotamidae <i>Chiamarra</i>	Chi	+	+	+							
	F/polycentropodidae	Pol	+	+	+							
	F/Leptoceridae <i>Setodes</i>	Set	+	+	+							
	F/Limnephilidae	Lim	+	+	+							
	F/Sericostomatidae	Ser	+	+	+							
Pulmonata	F/Physidae <i>Physa acuta</i>	Phy	+	+	+	+	+	+	+	+	+	+
Architaenioglosse	F/Viviparoidea Ss-F/viviparidae <i>Viviparus sp.</i>	Viv	+	+	+							+

Order (cont.)	Family	Code	Sta. 1	Sta. 2	Sta. 3	Sta. 4	Sta. 5	Sta. 6	Sta. 7	Sta. 8	Sta. 9	Sta. 10
Haplotaxida	F/ Haplotaxidae <i>Haplotaxis sp</i>	Hap	+	+	+	+	+	+	+	+	+	+
Coleoptera	F/Dytiscidae <i>Agabus sp.</i>	Aga	+	+	+	+			+			
	<i>Ilybius sp.</i>	Ily	+	+	+							
	<i>Hydrovatus sp</i>	Hyv	+	+	+							
	F/ Elmidae <i>Elmis sp</i>	Elm	+	+	+	+						+
	F/Gyrinidae <i>Gyrinus sp</i>	Gyr	+	+	+	+		+				+
	F/Chrysomelidae	Chr	+	+	+							
	F/ Hydrophilidae <i>Enochrus sp.</i>	Eno	+		+							
	<i>Hydrochara sp.</i>	Hyr	+	+	+							
	<i>Coelostoma sp.</i>	Coe	+	+	+							
	<i>Anacaena sp.</i>	Ana	+	+	+							
Amphipoda	F/ Gammaridae <i>Gammarus sp.</i>	Gam	+	+	+	+						+
Heteroptera	F/ Notonectidea <i>Notonecta maculate</i>	Not	+	+	+	+				+		
	F/Corixidae <i>Arctocorixa sp.</i>	Arc	+	+	+	+			+			
	<i>Corixa sp</i>	Cor	+	+		+						
	F/Aphelocheiridae <i>Aphelocheirus sp</i>	Aph	+									
	F/Nepidae <i>Nepa rubra</i>	Nep	+									
	F/Hydrometridae <i>Hydrometra</i>	Hym	+									
	F/Hydroptilidae <i>Orthotrichia sp</i>	Hyo	+	+		+						
	F/Hydraenidae	Hya	+									
Mecoptera	F/Boreidae	Bor	+	+		+				+		
Rhynchobdellida	F/ Glossiphoniidae <i>Glossiphonia complanata</i>	Glo	+		+		+	+	+	+	+	+

system and thus make it possible to enrich the list of Moroccan biodiversity. The faunistic inventory established groups and the distribution of stands in the different sampling stations (Table 2).

The macroinvertebrates found during the sampling period were distributed into twelve orders (Table 2). The most abundant order was *diptera*, having 9618 individuals (around 70% of the total), followed by the order *Ephemeroptera* with 2985 individuals (22%).

The abundance of the order diptera was seen at all stations (Table 2) and during the whole sampling period, while the other orders were present occasionally. Station S3 had the highest relative abundance (Table 3), with a total of 2749 individuals reported (20%), followed by S6 with 2161 (16%), S7, 1757 (13%), S1, 1690 (12.1%) and S2, 1698 individuals (11.9%). Analysis of the entire population during the study period shows that the insects are numerically the most inventoried and

TABLE 3
Average values of the indices calculated on the 10 study stations

Sites	°/° EPT	PE	Pchiro	P hyd	PINS	IBGN	IHF	Sha	Marg	Indi
S1	24.5	13,1	4	2,7	97,5	14	61	3.208	7.4	1690
S2	21,3	9,8	6	3,6	90	13.25	57	2.896	6.32	1698
S3	19,6	8,1	5	4,0	94,4	13.5	58	2.547	5.683	2749
S4	9	6,5	20	0,5	85,0	10.75	54	2.648	4.045	483
S5	6,5	3,27	21	5,3	53,8	6.5	50	1.81	1.375	695
S6	6,5	4,9	16	18,1	79,0	5.5	37	1.883	1.302	2161
S7	8,19	6,5	17	7,3	83,0	6	54	1.963	1.472	1757
S8	9,83	6,5	30	1,3	91,2	10.5	58	1.596	1.979	1182
S9	4,91	4,91	23	0,5	92,3	7.5	42	0.9978	1.143	1093
S10	8,1	6,5	4	0	61,7	11.5	62.7	2.392	2.458	198

%d'ephemeroptera :PE
Individuals :indi

% d'insectes : PINS
Shannon_H :Sha

°/° Hydropsychidae : P hyd
Margalef : Marg

%Chironomes : Pchiro

represent the highest percentage in all sites in Oum Er Rbia.

Table 3 shows the average values of the indices of diversity and biotic indices according to the stations. The highest mean values in ETP (*Ephemeres*, *Trichoptera* and *Plecoptera*) (19.6 to 24.5 taxa) are located at the stations where the species richness is the most important (S1, S2, S3). The stations disturbed (S5, S6, S9) had very low values, (6.5, 4.91). Also, the *Gammaridae* is more collected in upstream stations S1, S2, S3 (Table 2) represented by a single genus (*Gammarus*). We noticed that the taxonomic richness in Diptera was more diversified upstream than downstream, we had inventoried only five genera in medium stations. These spatial variations may be due to the various influences on the environment and also the nature of the different habitats. In fact, poor water quality can create favourable conditions for certain organisms which are more tolerant of pollution (polluoresistants). Relative percentages of *chironomids* are lower at Oum Er Rbia source and bass stations (S1, S2, S3 and S10) than medium stations (Table 2). The medium stations were characterized

by the lowest biotic indices and diversity indices : Shannon :1.02 to 1.95, Margalef: close to 3.95, IBGN (Global Standard Biological Index) : less than 5.5. This is not surprising because pollution-sensitive taxa, e.g. *Ephemeroptera*, *Trichoptera* and *Odonata* disappeared at these sites, as a result of lower biotope quality at these sites compared to other sites of the river . This could be related to natural and / or anthropogenic factors influencing this watercourse. Indeed, the discharge of domestic untreated wastewater in Kasba Tadla and Khenifra, undoubtedly, contribute considerably to the occurrence of unfavorable conditions for benthic population. The polluted stations are characterized by the lowest biotic indices and diversity indices confirming the poor quality of these environments: Shannon: 0.9 to 1.96, Margalef: close to 1.14, IBGN (Global Standard Biological Index) : less than 5.5. The loss of indicator faunal groups, such as the *Plecoptera* Perlodidae and the *Trichoptera* Philopotamidae, explains the low values of the IBGN at these stations. These indices of diversity and biotic indices have a significant




Figure 2: Correspondence analysis of stations , Macroinvertebrate families and environmental variables




Figure 3: Correspondence analysis of stations and Macroinvertebrate families

linear relationship with the Oum Er Rbia stations (Fig. 2 and 4).

Figure 2 shows the relationship between the biotic indices and diversity indices and the IHF. This relationship is identical to that presented in Barbour *et al.* (1999). The condition of macroinvertebrate communities decreases with decreasing habitat quality. The River Habitat Index IHF shows a good relationship with the IBGN. These two indices, the IHF and the IBGN, have positive correlation respectively with the Shannon and Margalef (Fig. 4).

Using a variety of multivariate analysis,

family and genus showed most of the same correlations to biotic indices and diversity indices (Fig. 4). Although the response to diversity indices was different, species were grouped in the center of the CCA ordinal triplate (Fig. 3). This suggests that no environmental variables explained species composition and distribution between sites. However, DO was negatively correlated with Chironomidae (Fig. 3).

According to diversity indices (Shannon , Taxa_S and Margalef), higher richness of benthic macroinvertebrates were recorded at the S1 ,S2, S3 with 10 genus restricted to




Figure 4: Correspondence analysis of stations and indices calculated on the 10 study stations

these sites, probably due to optimal habitat conditions, these sites are located upstream from major polluting source and can be considered as a local reference sites. In fact, the River Habitat Index IHF showed that these sites have good habitat quality (61 to 54) (Table 2). Riparian vegetation can also play a role in protecting benthic fauna from the direct effects of UV radiation (Quinn *et al.* 2004). In this study, although habitat quality was important across the sites (particularly at Headwaters), the habitat characteristics were positively correlated with benthic macroinvertebrates. Also, we noticed that percentages of ephemeroptera (PEPH) are higher at S1, S2, S3. However, most families belonging to the PEPH group are highly sensitive (Dickens and Graham 2002). Populations of benthic macroinvertebrate species responded differently to disturbance regimes and many species have different habitat requirements at each stage of their development cycle (Simaika and Samways 2011).

Conclusion

In summary, Oum Er Rbia River has a regime

characterized by the irregularity of the flows and by brutal hydrological manifestations. The regime is marked by a poor flow in summer and autumn, and by a high flow in winter and spring. An inventory of the fauna carried out in the present study constitutes a first important database. The fauna studied is characterized by a variable taxonomic diversity according to the degree of water pollution. The fauna identified in this work consists of 52 families and belong to 3 faunistic groups (annelids, molluscs and arthropods). The benthic population showed that Diptera, Ephemeroptera and Trichoptera were dominant. Gammaridae, Heteroptera, Coleoptera and Odonata were only a small fraction of the harvested fauna. Hydropsyche, Chironomidae sp. and Simuliidae are the most numerically inventoried. This is related to the relative resistance of these invertebrates to pollution and the fact that Hydropsyche larvae are a common component of water benthos (Verneaux *et al.*, 1976). The biological indices (IBGN, Shannon and Equitability) used for the characterization of the waters of the Oum Er Rbia River, made the study stations quite distinctive based on their level of pollution. The specific richness of the Oum Er Rbia

River depends on the ecological conditions at each station; it is even higher when the biotope is heterogeneous and less influenced by anthropogenic activities. Oum Er Rbia River can be considered as poor in wildlife (as to the presence / absence of taxa and their abundance); compared to the fauna inventoried by Chahlaoui in 1996 (Chahlaoui *et al.* 1996) and Karrouch in 2010 (Karrouch *et al.* 2010) in Oued Boufekrane.

In conclusion, we suggest that for the next studies in the Oum Er Rbia River should focus on the selection of the indicator species and monitor them as substitutes for the entire community. However, it is not possible to develop a macroinvertebrate index that can be applied directly and without any adaptation or modification to all Morocco rivers. Some grouping of rivers based on ecological similarities which include both physical characteristics and information on macroinvertebrate, can allow equivalent macroinvertebrate indices (IBGN) to be developed for similar river sections or in rivers in other parts of the country.

References

- Arab, A., Lek, S., Lounaci, A. and Park, Y.S.** (2004). Spatial and temporal patterns of benthic communities in an intermittent river (North Africa). *Annales de Limnologie International Journal of Limnology*, 40:317-327.
- Beauger, A., Lair, N., Marchant, P.R. and Peiry, J. L.** (2006). The distribution of macroinvertebrate assemblages in a reach of the River Allier (France) in relation to riverbed characteristics. *Hydrobiologia*, 571:63-76.
- Benbow, M.E., Burky A.J. and Way C.M.** (2003). Life cycle of a torrenticolous Hawaiian chironomid (*Telmatogeton torrenticola*): stream flow and microhabitat effects. *Annales de Limnologie International Journal of Limnology*, 39:103-114.
- Bowman, M. F. and R. C. Bailey.** (1997). Does taxonomic resolution affect the multivariate description of the structure of freshwater benthic macroinvertebrate communities. *Canadian Journal of Fisheries and Aquatic Sciences*, 54:1802-1807.
- Cairns, J. Jr. and Pratt, J.R.** (1993). A history of biological monitoring using benthic macroinvertebrates. *Freshwater Biomonitoring and Benthic Macroinvertebrates*, 10-27.
- Carl, R., George, E.H. and John, W.A.** (1993). Identification of predominant environmental factors structuring stream macroinvertebrate communities within a large agricultural catchment. *Freshwater Biology*, 29: 285-294.
- Chahlaoui, A.** (1996). Hydrobiological study of Oued Boufekrane - Impact on the environment and health, Thesis in Biology. Université Moulay Ismail Faculty of Science, 256pp.
- Chessman, B.C.** (1986). Diatomora of an Australian river system: Spatial patterns and environmental relationships. *Freshwater Biology*, 16:805-819.
- Daniel, F.B. and Anderson, S.V.** (2010). Rapid Bioassessment Protocols using benthic macroinvertebrates in Brazil: evaluation of taxonomic sufficiency. *The North American Benthological Society*, 29:562-571.
- De'nes, S, Jani, H., Ja'nos, P, Tibor and E., Sylvain, D.** (2017). Functional diversity: a review of methodology and current knowledge in freshwater macroinvertebrate research. *Hydrobiologia*, 787:27-44.

- Duan, X.H., Wang, Z.Y., Xu, M. Z. and Zhang ,K.**(2009).Effect of streambed sediment on benthic ecology. *International Journal of Sediment Research*, 24: 325–338.
- Dudgeon, D.**(2008).*Tropical Streams Ecology*. Elsevier,USA,343pp.
- Elliot, J.M.**(1971).Some Methods for the Statistical Analysis of Samples of Benthic Invertebrates. Scientific Publication . *Fresh water Biology Association*,59:302-302.
- Faith, D.P. and Norris, R.H.** (1989). Correlation of environmental variables with patterns of distribution of common and rare freshwater macroinvertebrates. *Biological Conservation*,50:77-98.
- Furse, M.T., Moss ,D., Wright, J.F. and Armitage P.D.** (1984). The influence of seasonal and taxonomic factors on the ordination and classification of running-water sites in Great Britain and on the prediction of their macroinvertebrate communities. *Freshwater Biology*, 14:257–280.
- Gregory, S.V.,Swanson ,F.J., McKee, W,A. and Cummins, K.W.** (1991).An ecosystem perspective of riparian zones. *Bioscience Journal*, 11:540-551.
- Hellawell, J.** (1986). Biological indicators of freshwater pollution and environmental management. *Elsevier Applied Science Publishers*,14,188.
- Huryn, A.D. and Wallace, J.B.** (1987). Local geomorphology as a determinant of macrofaunal production in a mountain stream. *Ecology*, 68:1932-194.
- Karrouch, L.** (2010).Bio-évaluation de la qualité des eaux courantes de la région Meknès (Centre-Sud, Maroc) – Impact sur l’environnement et la sante. Thèse doctorat es sciences en biologie. *Faculté des sciences. Meknès*, 216pp.
- Learner, M.A. and Densem, J.W. and Iles T.C.** (1983).A comparison of some classification methods used to determine benthic macroinvertebrate species associations in river survey work based on data obtained from the river Ely (South Wales). *Freshwater Biology*,13:13-36.
- Lenat, D.R. and Resh, V.H.**(2001). Taxonomy and stream ecology – the benefits of genus- and species-level identifications. *Journal of the North American Benthological Society*, 20:287–298.
- Maranda,Y. and Sasseville, J.L.**(1999). Efficacité de l'assainissement des eaux usées sur le bassin de la rivière Chaudière (Québec, Canada). *Revue des sciences de l'eau*. 12:485-507.
- Marchant, R., Barmuta, L.A. and Chessman, B.C.** (1995).Influence of sample quantification and taxonomic resolution on the ordination of macroinvertebrate communities from running waters in Victoria, Australia. *Marine and Freshwater Research*, 46:501–506.
- Meyer, J.L., McDowell, W.H., Bott, T.L., Elwood J.W., Ishizaki C, Melack J.M., Peckarsky B.L, Peterson, B.J. and Rublee P.A.** (1988).Elemental dynamics in streams. *Journal of the North American Benthological Society*, 7:410-432.
- Michelle, F. B. and Robert, C.B.** (1997).Does taxonomic resolution affect the multivariate description of the structure of freshwater benthic macroinvertebrate communities. *Canadian Journal of Fisheries and Aquatic Sciences*, 54:1802-1807.
- Moroccan Standards for surface water Quality** .(2002) ,Official Bulletin n° 5062.
- Mykra, H., Heino, J. and Muotka, T.** (2007).Scale-related patterns in the spatial and environmental components of stream

- macroinvertebrate assemblage variation. *Global Ecology and Biogeography*, 16:149–159.
- Neville, P.J., Yen, A.L.** (2007): Standardising terrestrial invertebrate biomonitoring techniques across natural and agricultural systems. *Australian Journal of Experimental Agriculture*, 47:384–391.
- Ouyang, Y.** (2005) :Evaluation of river water quality monitoring stations by Principal Component Analysis. *Water Research*, 39:2621–2635.
- Plafkin, J., Barbou, M.T., Porter, K.D., Gross, S.K. and Hughes, R.M.** (1989).Rapid bioassessment protocols for use in streams and rivers: benthic macroinvertebrates and fish. EPA/444/4-89/001. Once of Water Regulations and Standards, U. S. Environmental Protection Agency, Washington, DC.
- Poff, N.L. and Ward, J.V.** (1989).Implications of streamflow variability and predictability for lotic community structure: a regional analysis of streamflow patterns. *Canadian Journal of Fisheries and Aquatic Sciences*, 46:1805-1818.
- Prat, N. and Ward, J.V.**(1994).The tamed river. In R. Margalef (editor). *Limnology now: A paradigm of planetary problems. Elsevier Science B. V.*, 219-236.
- Resh, V.C. and Mcelravy, E.P.** (1993). Contemporary quantitative approaches to biomonitoring using benthic macroinvertebrates. In: *Freshwater Biomonitoring and Benthic Macroinvertebrates* (Eds D.M. Rosenberg and V.H. Resh), pp. 159–194.
- Resh, V.H. and Rosenberg, D.M.** (1984).The ecology of aquatic insects. Praeger, NYY. *Limnology and Oceanography*, 29:1350–1351.
- Statzner, B., Gore, J.A. and Resh ,V.H.** (1988).Hydraulic stream ecology: observed patterns and potential applications. *Journal of the North American Benthological Society*, 7: 307-360.
- Thorne, R. St. J. and Williams, W.P.** (1997). The response of benthic macroinvertebrates to pollution in developing countries: a multimetric system of bioassessment. *Freshwater Biology*. 37: 671-686.
- Townsend., C.R., Doledec, S., Norris, R., Peacock, K. and Arbuckle. C .** (2003). The influence of scale and geography on relationships between stream community composition and landscape variables: description and prediction. *Freshwater Biology*, 48:768–785.
- Townsend, C.R., Scarsbrook, M.R. and Dolédec, S.** (1997).Quantifying disturbance in streams: alternative measures of disturbance in relation to macroinvertebrate species traits and species richness. *Journal of the North American Benthological Society*, 16:531-544.
- Vanderklift, M.A., Ward, T.J. and Jacoby. C.A.** (1996).Effect of reducing taxonomic resolution on ordinations to detect pollution-induced gradients in microbenthic. *Freshwater Biology*, 49:474-489.
- Vannote, R.L. and Sweeney, B.A .** (1980). Geographic analysis of thermal equilibria: A conceptual model for evaluating the effect of natural and modified thermal regimes on aquatic insect communities. *American Naturalist*, 115:667-695.
- Veroli, A.D.,Selvaggi, R., Pellegrino, R.M. and Goretti, E.**(2010). Sediment toxicity and deformities of Chironomid larvae in Lake Piediluco. *Chemosphere*,79:33–39.
- Verneaux, J., and Faessel, B.** (1976).Larves du genre Hydropsyche (Trichoptères

- Hydropsychidae). Taxonomie, données biologiques et écologiques. *Annales de Limnologie*, 12:7–16.
- Ward, J.V.** (1986). Altitudinal zonation in a Rocky Mountain stream. *Archiv für Hydrobiologie, Supplement*, 74 (2):133-199.
- Wright, I.A., Chessman, B.C., Fairweather, P.G. and Benson, L.J.** (1995). Measuring the impact of sewage effluent on the macroinvertebrate community of an upland stream: The effect of different levels of taxonomic resolution and quantification. *Australian Journal of Ecology*, 20: 142–149.
- Øyvind, H, David Harper, A.T. and Paul, D.P.** (2001). Past: paleontological statistics software package for education and data analysis. *Palaeontol Electronica*, 4:1 – 9.