

African Research Review

International Multi-Disciplinary Journal, Ethiopia

Vol. 5 (1), Serial No. 18, January, 2011

ISSN 1994-9057 (Print)

ISSN 2070-0083 (Online)

Comparative Studies on the Larvicidal Action of Novaluron (Mosquiron® 100EC) and Moringa Oliefera (LAM) Seed Oil against Aedes Aegypti (Diptera: Culicidae) Larvae

(Pp. 424-437)

Nwankwo, E. N. - Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria

Okonkwo, N. J. - Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria

Ozumba, N. A. - Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria

Okafor, E. G. - Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria

Abstract

Inhibition of emergence (IE) by Novaluron (Mosquiron 100EC) and Moringa oliefera Lam.(Moringaceae) seed oil (MOSO) was assayed with first and fourth instar larvae of Aedes aegypti (Diptera: Culicidae) at ambient laboratory conditions. graded concentration ranging from 3.125- 200µg/ml including a control were tested. Four replicates of each concentration in RCBD were variously challenged with cohorts of 20 active larvae of a given instar. Mortality resulting from eclosion inhibition was recorded at 3-hourly intervals for 12 hours, and data analysed using log/probit transformation. Results indicated dosage-related mortality differences. High concentration of Novaluron gave 95% mortality of both 1st and 4th instars, while sublethal dosages resulted in 45 and 35% mortality of 1st and 4th instars, respectively.

*Exposure to Moringa oil resulted in 90 and 100% mortality of 1st and 4th instars, respectively, and sublethal dosages gave 30 and 25% mortality, respectively. Both toxicants showed significant ($P < 0.05$) inhibition of emergence following post-treatment culturing. The IE_{50} values for 1st instar larvae were 4.546 μ g/ml and 15.44 μ g/ml of Novaluron and Moringa oil, respectively; and 8.028 μ g/ml and 9.977 μ g/ml, respectively, for 4th instar. Although the 1st instar larvae were more susceptible, Novaluron was about twice more potent. Nevertheless, Moringa oil has shown promise as biopesticide for *Ae. aegypti* larvae control.*

Key Words: Novaluron, *Moringa oliefera*, *Aedes aegypti*, Toxicity, Nigeria.

Introduction

Mosquitoes are vectors of many human diseases. According to Clements, 1992, Reinert, 2000) there are over 3000 different species of mosquitoes throughout the world, about 1,900 species occur in the humid tropics and subtropics where the climatic conditions are favourable for rapid immature stages development and adult survival. Global efforts to reduce the number of mosquitoes usually are due to the deadly diseases they transmit to man and animals. The important mosquito transmitted disease with high public health importance include malaria, yellow fever, Dengue haemorrhagic fever and filariasis, these are transmitted by various species of Anopheles, *Aedes* and *Culex* mosquitoes.

Early attempts to control these vectors have relied heavily on the use of synthetic insecticides resulting in several problems, including environment pollution, insect resistance, unacceptable levels of residues, escalating cost of production among others (Otto, 1992).

Science and Technology have recognized the need to reduce synthetic pesticide usage in order to achieve a cleaner environment, even though this would certainly not result in reduced vector control. This conflict in the goal to reduce synthetic pesticide usage and elimination of vectors of diseases, provides a strong impetus for the development of cost-effective and ecologically more benign alternatives.

Emphasis is on the use of botanical products, ecological controls and others such as insect growth regulators. Synthetic pesticides are used only when necessary and truly justifiable and when acceptable alternatives are lacking (Lim *et al*, 1997). Although, successful control of mosquito vectors with Malathion has been reported (Fradin *et al* 2002), many pyrethroid

insecticides and insect growth regulators (IGRS) are now being directed towards mosquito control. Prominent among the IGRS is Novaluron, a novel acylurea acting as a chitin synthesis inhibitor (Mulla *et al* 2003) affecting the moulting in insect development. It causes mortality in larvae on acdysis after ingestion or contact and causes abnormal endocuticular deposition and abortive moulting (Stephen, 2005). Emergence prevention is the primary goal of IGR larvicidal treatment and inhibition of emergence (IE) results in death. Mulla *et al* (2003) reported that inhibition of larval emergence was 100% at 0.25 to 1.0g/l of Novaluron® under laboratory condition while in the field condition using clay jars, there was 86-96% inhibition of emergence (IE) for about 190 days. Novaluron is registered as insecticide for both food crop and ornamentals in many countries. World Health Organization assessed its mosquito Larviciding activity for *Aedes species* and recommended a dosage application of 0.05mg/l (WHO, WHOPEs, 2005).

On the other hand, plants are by far the most efficient “factory” of chemical compounds, synthesizing many products that are used in the defense against many pests and vectors (Schoonhoven *et al*, 1998). Thus, the use of plant extract in agro-ecosystem is now emerging as one of the primary means to protect the environment from toxic synthetic pesticides pollution (Schumuterer; 1995). Jany *et al.* (2002) reported the larvicidal activity of leguminous seeds and grains against *Aedes aegypti* and *Culex pipens pallens* while Bassole *et al* (2003), investigated the ovicidal and larvicidal activities of leaves of three plants naturally growing in Burkina Faso.

Many researchers are now investigating Moringa plant for its use as pesticides. Donli and Dauda (2002) reported the use of aqueous Moringa seed extracts as seed dressing bio-fungicide for groundnuts in Nigeria. Adandonon *et al.* (2006) combined a biocontrol with *M. oleifera* seed extract for integrated control of Sclerotium damping-off of cowpea stem and root. Ajayi (2007) compared efficacy of *Moringa oleifera* seed oil with *Sesamum indicum* and *Olea europaea* against *T. castaneum*. Aqueous extracts of leaves of *Moringa oleifera* Lam, *Vernonia amygdalina* and *Annona muricata* were evaluated for the control of *Collectotrichum destructivum* on seeds of cowpea (*Vigna uniguculata*) (Akinbode and Ikotun, 2008). Badruddoza and Rahman (2008) reported the larvicidal action of *M. oleifera* root together with other nineteen Indian plants against *Ae. albopictus* and *C. quinquefasciatus*.

Horse radish, *Moringa oleifera* Lam. is an abundant invasive plant found throughout Nigeria and many other African countries which has generated a

lot of interests among Scientists in many African countries for its medicinal and other uses (Ozumba, 2005). Its potentials as bio-pesticide need to be further evaluated. In the present study investigations were aimed at evaluating the larvicidal action/ inhibition of emergence by an insect growth regulator, Novaluron (Mosquirom® 100EC) and botanical plant *Moringa oleifera* seed oil extract for the control of mosquito, *A. aegypti* larvae.

Materials and Methods

These investigations were carried out in the Research Laboratory of the Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka, Nigeria.

Sources of Mosquito Larvae and Rearing

Eggs of *Aedes aegypti* were obtained from at the Federal Ministry of Health, Department of Public Health National Arbovirus and Vector Research Center, Enugu, Nigeria. They were hatched and reared in the laboratory at 26±2°C, 80±5% r.h. The eggs were placed in transparent plastic containers containing 500mls of distilled water and allowed to hatch to 1st instar larvae and kept to reach the 4th instar. The larvae were fed on crumbs of biscuit brand, Yale Fortune Cabin Sweetened Biscuit; the feed was applied on alternate days for normal development.

Moringa Seed Collection and Oil Extraction

The *Moringa oleifera* seed was collected from the moringa plantation in Enugu, Nigeria owned by the third author. The seeds were dried under shed for 7 days and decorticated to obtain the white kernel. The dried seeds were pulverized using an electric blender to obtain fine powder. Thereafter, the oil extraction was carried out using petroleum ether in Soxhlet Extractor for 3 hours. About 40g of the pulverized sample was wrapped in a double layer of Whatman No. 1 filter paper and placed inside the Soxhlet thimble. Measured volume of 250ml of petroleum ether was poured through a funnel, by-passing the thimble, containing the sample into the round bottom flask system of the Soxhlet apparatus. This was heated about 5cm above a hot electric plate while cold water was allowed to reflux in and out of the Extractor to cool the condenser compartment. After many refluxes, the petroleum ether was gradually evaporated from the oil extract. The process was repeated six times to allow the solvent distill off at about 75°C, leaving the light yellow oil.

Preparation of the Oil Extracts

Serial dilutions of the *M. oleifera* seed oil (MOSO) were prepared in acetone. The pure oil extract was regarded as 100% concentrate and subsequently

diluted serially to 20%, 10%, 5%, 2.5%, 1.25%, 0.625% and 0.3125% in acetone.

Novaluron (Mosquiron® 100EC) Preparation

The Novaluron (Mosquiron®100EC) was obtained from Dizengoff West Africa Ltd, Nigeria. Based on the WHO (2005), recommended dosage of 10% ai, a measured volume of Mosquiron® 100EC was first diluted down 10% ai equivalent to (10EC) subsequently serial dilutions were made in acetone to obtain 200µg/ml, 100µg/ml, 50µg/ml, 25µg/ml, 12.5µg/ml, 6.25µg/ml, and 3.125µg/ml.

Bioassay Tests

The method and standard procedures used were those of Mulla *et al.* (1986) and WHO (1981). The different concentrations of the above mentioned preparation obtained from diluting in acetone were used. Aliquots of 1ml of each dilution were separately placed in plastic cups containing 160ml distilled water. Each container was challenged with batches of 20 active fourth or first instar larvae of *Aedes aegypti* accordingly. Controls were also included.

Each treatment and control was replicated five times. The bioassay was repeated two times. The tests were carried out at ambient laboratory temperature of $26 \pm 2^\circ\text{C}$, $80 \pm 5\%$ relative humidity and photoperiod 12:12 light and dark hours. Mortality/ inhibition of emergence assessments were made at 3-hourly intervals for 24 hrs. Dead larvae were counted and recorded. However, exuviae or shed cuticle started appearing in control treatments after about 2 hrs but none became pupae or adult. Both moribund larvae, that is, those unable to wriggle were counted as dead.

Data Analysis

Mortality data obtained were corrected using Abbot Formula (1925). Log-probit regression analysis was carried out (Finney, 1971) for determining LD_{50} or level of inhibition of emergence ($1E_{50}$). Analysis of variance (ANOVA) was further carried out also performed on the mortality data and means separated using least significant difference (LSD) effectiveness at the different time intervals and concentration levels.

Results

The mortality values of 4th instar larvae of *Ae aegypti* exposed to Novaluron (Mosquiron 100EC), an IGR and *Moringa oleifera* seed oil extract at 3 hourly intervals are presented in Table 1 and 2 respectively. The results show

that novaluron and moringa oil exhibited significant ($P < 0.05$) levels of toxicity to *Ae. Aegypti* larvae. Mortality increased with increasing concentration of the IGR and oil and also with increasing period of exposure. However, there were much greater responses at higher concentrations of toxicant (no mortality was observed in the control). Generally, dose-time related mortality responses were noted in all treatments with higher ranges of the toxicant causing almost 100% death or inhibition of emergence in the larvae, while in lower concentrations, larval mortality was incomplete. However the time of exposure had more significant impact than dosage. The log-probit regression analysis revealed that after 12 hours of exposure, the LD_{50} (IE_{50}) of the novaluron treated 4th instar larvae was 8.028 μ g/ml and moringa is 9.977 μ g/ml (Fig 1).

Tables 3 and 4 respectively showed that mortality of 1st instar larvae of *Ae. aegypti* exposed to novaluron and moringa oil increased significantly ($P < 0.05$) with increasing levels of concentration and also with increasing period of exposure to the toxicant. The LD_{50} (IE_{50}) value of novaluron on 1st instar larvae after 12 hours was 4.546 μ g/ml and that moringa was 15.44 μ g/ml (Fig 2). Mortality in the second 3- hours for both 4th and 1st instar larvae was significantly ($P < 0.05$) higher than that of the first 3- hours. This trend was observed in all the treatment irrespective of the dosage. However, at highest concentration, 200 μ g/ml, the percentage mortality caused by the IGR on the 4th and 1st instar were 95% respectively (Tables 1 and 3). A similar trend was observed in the moringa treated larvae (Tables 2 and 4). The 1st instar was significantly ($P < 0.05$) more susceptible to the two toxicant than the 4th instar larvae.

Discussion

The larvicidal action of Novaluron (Mosquiron® 100EC) and *Moringa oliefera* seed oil (MOSO) on both 1st and 4th instar larvae of *Ae. aegypti* was studied in the laboratory. The studies are important to understand how these two insecticides of both synthetic and plant origin can be used as independent control method or as adjunct to other control measures in reducing the impact of mosquitoes. In the present study the general performance of Novaluron in killing the larvae of *Ae. aegypti* was observed in the relatively high percentage larval mortality recorded in both 1st and 4th instar larvae, high percentage inhibition of emergence and LD_{50} values. This may explain why WHO recommended its use as a mosquito larvicide (WHO, 2005).

There was increase in the percentage mortality with increased concentration, as the time progressed. The 100% inhibition of emergence (IE %) in both 1st and 4th instar larvae observed in the present studies is in accordance with the studies by Mulla *et al.* (2003), who reported that percentage inhibition of emergence was 100% at concentrations of 0.25 to 1.0g/L on both 2nd and 4th instar larvae of *Ae. aegypti*. In addition, Kostyukovsky and Trostanetsky (2004) reported that novaluron reduced hatching of 3rd instar larvae of *Tribolium castaneum* by 100%. These suggest that novaluron selectively target immature insect stages by inhibiting chitin formation and so causing abnormal endocuticular deposition and abortive molting. The benzoylureas act on larval stages of most insects by inhibiting or blocking the synthesis of chitin (Soltani *et al.*, 1995), a vital and almost indestructible part of the insect exoskeleton (Chebira *et al.*, 2000). More than being the typical poisons that attack the insect nervous system, they interfere with chitin synthesis and they are more taken up by ingestion than by contact (Soltani *et al.*, 1995). Typical effects on developing larvae are the cleaving of malformed cuticle or death by starvation (Chebira *et al.*, 2000). Su *et al.* (2003) reported that Novaluron exhibited a high level of activity against *Culex* mosquitoes.

On the other hand, Moringa seed oil was also found to have promising larvicidal properties. The toxicity effect of the oil was high, as the mortality and the rate of inhibition of emergence increased with increased concentration and time. The effectiveness of plant oil in regulating growth and inhibition of many insect and fungal development has been reported previously (Jilani and Su, 1983, Mulla *et al.*, 2003, Kostyukovsky and Trostanetsky, 2004). Manas *et al.* (2005) reported that *Moringa oleifera* extracts inhibited 97.32% germination of fungal pathogen. Shema and Sexena (1994) also found that the petroleum ether extract of *Tagetes erectes* had toxic effect on larvae of *Anopheles stephensi* and on its significant growth index. Mwangi and Mukiyama (1988) observed that fraction of *Melia volkensii* fruit kernel extract had growth inhibition activity at low concentration, whereas two other fractions had acute toxic effects on the mosquito larvae. This was observed in the present study as only few surviving larvae were seen in the treated sample, which were unable to develop to the next stage. Pushpalatha and Muthukrishnam (1995) reported that leaf extracts of *Vitex negundo* at very low concentration had larvicidal activity against *Culex quinquefasciatus* and *Anopheles stepensi* and also extended the duration of larval instar pupation.

These insecticides, Novaluron and Moringa seed oil have shown to be very effective in mosquito control as seen in the present study. Novaluron action is by chitin inhibition according to Soltani *et al.* (1995), Mulla *et al.* (1974, 2003) and Kostyukovsky and Trostanetsky (2004) for novaluron and MOSO action as growth regulator and by blocking the respiratory surfaces. According to Hirashima *et al.* 1998, Enan, 2001) some plant essential oil block the octopamine neuroreceptors that regulate the movement, heart rate, behaviour, metabolism, and pupation of insects. This regulatory activity of these two insecticides was evidenced by the death recorded in both 1st and 4th instar larvae and inhibition of their emergence to pupae. In both insecticide treated water, it was obviously noticed that the larvae were weakened before death occurs.

In the present studies, the 1st instar larvae of *Ae. aegypti* were more susceptible than the 4th instar. This observation was confirmed by the previous studies that early instars of insects are more susceptible to the inhibitory and growth regulatory effects of these insecticides (Mulla *et al.*, 2003). This is however suggested to be due to the size and the tender nature of the chitin in the early instar of mosquitoes. The observation by Mulla *et al.* (2003) that 2nd instar of *A. aegypti* was more susceptible than 4th instar using novaluron and Akpa *et al.* (2003) that 3rd instar was more susceptible than 4th instar when a plant, *Phytolacca dodecandra* oil was used against *Ae. Egypti* larvae verify these findings.

References

- Abbott, W.A. (1925). A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology*, 18:265-267.
- Adendonon, A; Aveling, T.A.S, Labusehanone; N. and Tamo, M. (2006). Biocontrol agent in Combination with *Moringa Oleifera* Extract for Integrated Control of Sclerotium-Cause Cowpea Damping off and stem rot. *European Journal of Plant Pathology*, 115(4): 409-418.
- Ajayi, F. A. (2007). Effect of grain breakage and application of edible oils to protect pearl millet, *Pennisetum glaucum* (L.) R. Br., against infestation by adult *Tribolium castaneum* (Herbst.) (Coleoptera: Tenebrionidae) in Lafia, Nigeria. *Nig. J. Ent.*, 24: 107- 113.
- Akinbode, O. A. and Ikotun, T. (2008). Efficacy of certain plant extracts against seed-borne infection of *Collectotrichum destructivum* on

cowpea (*Vigna unguiculata*). *African Journal of Biotechnology*, 7 (20): 3683-3685.

- Akpa, A. U. C., Mwale, N. A., Ozumba, N. A. and Yohannes, L. W. (2003). The efficacy of *Phytolacca dodecandra* (L ‘ Herit) as a larvicide against *Aedes aegypti* (Linnaeus)- a yellow fever vector in Nigeria. *Journal of Science and Technology*, 9: 13- 22.
- Badruddoza, D. M. and Rahman, S. M. K. (2008). larvicidal activities of aqueous extracts of nineteen Indian plants against two *mosquitoes*, *Aedes albopictus* and *Culex quinquefasciatus*. *Crop protection and Toxicology*, 4:19-25.
- Bassole, I. H, Guelbeogo, W. M., Nebie, R., Constantini, R: Sagnon, N., Kabore, Z. I. and Traore, S.A. (2003). Ovicidal and larvicidal activity of essential oils extract from three spontaneous plant against *Aedes aegypti* and *An. Gambiae* complex mosquitoes of Burkina Faso. *Parasitologia*, 45(1): 23-26.
- Chebira, S. N., Soltani-Mazouni and Soltani, N. (2000). Evaluation of two insect growth regulators, Flucycloxuron and Diflubenzuron, affecting in vivo and in vitro the ecdysteroid production in *Tenebrio molitor* (Coleoptera), In Proceedings, 10th Intl. Conference of Cairo University, Pp, 75- 89.
- Clements, A. N. (1992). *The biology of mosquitoes: (Vol. 1) Development, nutrition and reproduction*. Chapman and Hall, New York. P126-135
- Donli P.V. Dauda H. Evolution of Acqueous Moringa seed extract as a seed treatment biofungicide for groundnut. *Pest Management Sci.* 2003 59: 9
- Enan, E. (2001). Insecticidal activity of essential oils: octopaminergic sites of action. *Comp. Biochem. Physiol.*, 130: 325- 337.
- Finney, D.J. (1971). *Probit analysis*. Ed. 3. Cambridge University Press, London.
- Hirashima, A. S., Hirokado, R., Tojikubo, R., Takeya, E., Taniguchi, and Eto, M (1998). Metamorphosis of the red flour beetle, *Tribolium castaneum* Freeman Hinton (Coleoptera: Tenebrionidae): alteration of octopamine content modulates activity of juvenile-hormone

- esterase, ecdysteroid level, and pupation. *Arch. Insect Biochem. Physiol.*, 37: 33- 46.
- Jilani, G. and Su, H. C. F. (1983). Laboratory studies on several plant materials as insect repellents for protection of cereal grains. *J. Econ. Entomol.*, 76: 154- 157.
- Kostyukovsky, M. and Trostanetsky .A. (2004). The effect of a new chitin synthesis inhibitor, Novaluron, on various developmental stages of *Tribolium castaneum* (Herbst). *Journal of stored Product Research*, 42 (2): 136 – 148.
- Lim G.S. Sivapragasam, A; and Ioke W.H. (1997). Crucifer Insect Pest Problems Trends, Issues and Management Strategies in: Sivapragasam, A. Ioke, W.H. Hussan, A.K. and lim G.S. (Eds). *The Management of Diamond Moth and other Crucifer Pest*. Proceeding of the Third International Workshop, Kuala Lumpur, Malaysia. Malaysia Agricultural Research and Development Institute (MARDI).
- Manas, D. M., Souman, C., Pratheesh, K., Beera, S. (2005). In vitro screening of some plant extract against fungal pathogens of mulberry (*Morus* spp). *Archives of Phytopathology and Plant Protection*, 38 (3): 157- 164.
- Mulla, M. S., Darwazeh, H. A. and Norland, R. L. (1974). Insect growth regulators: evaluation procedure and activity against mosquitoes. *Mosquito News*, 10: 329- 332.
- Mulla. M. S., Thavura, U., A. Tawatsin, J., Chomposri, M., Zaim and Su, T. (2003). Laboratory and Field Evaluation of Novaluron, a new acylclurea insect growth regulator; against *Aedes aegypti* (Diptera; Culicidae). *J. Vector Ecol.*, 28:241-254.
- Mwangi, R.W. & Mukhiama, T.K. (1988) Evaluation of *Melia volkensii* extract fractions as mosquito larvicides. *J. Amer. Mosq. Cont. Assoc.*, 4:442- 447.
- Otto, D. (1992). Insektizidresistenz in Insektenpopulation- ein Überblick. *Nachrichtenblatt des Dt. Pflanzen- Schutzdienstes*, 44 (2): 34- 39 (Abstract only seen).

- Ozumba N. A. (2005) "Health and Nutrition." An invited paper presented at a workshop on skills Enhancement for Traditional Medicine Practitioners. Organized by Nigeria National Medicine Development Agency, Federal Ministry of Science and Technology. Modotel, Enugu, 24-26 January.
- Pushpalatha, E. & Muthukrishnan, J. (1995). Larvicidal activity of few plant extracts against *Culex quinquefasciatus* and *Anopheles stephensi*. *Ind. J. Malariol.*, 32:14-23.
- Reinert, J. F. (2000). New classification for the composite genus *Aedes* (Diptera: Culicidae: Aedini) elevation of the subgenus *Ochleratatus* to generic rank, reclassification of the other subgenera, and notes on certain subgenera and subspecies. *Journal of American Mosquito Control Association*, 16:175- 188.
- Schmutterer, H. (1995). The neem Tree: source of unique natural products for the integrated pest management, medicine, industry and other purposes. Vch Verlagsgesellschaft, Weinheim.
- Sharma, M. & Saxena, R.C. (1994) Phytotoxicological activity of *Tagetes erectes* in aquatic stages of *Anopheles stephensi*. *Ind. J. Malariol.*, 31:21-26.
- Soltani, N. S., Chebira, N. Pitoizet, J. P., Delbecoq and Delachambre J., (1995). Effect of Flucyclohexuron, a novel benzoylphenylurea derivative, on the *in vivo* and *in vitro*, product of ecdysteroids in *Tenebrio molitor*; *Med. Fac. Landbowi Univ. Gent*, 607(3b) :1017-1022.
- Stephen, F. (2005). First draft on Novaluron presented to Health effects Divisions US Environmental Protection Agency, Washington DC, USA.
- Su, T. Y., Zaim, M. and Mulla, M. S. (2003). Laboratory and field evaluation of Novaluron, a new insect growth regulator (IGR) against *Culex* mosquitoes. *J. American Mosquito Control Assoc.*, 19: 408- 418.
- WHO (1981). Instructions for determining the susceptibility or resistance of adult mosquitoes to organochlorine, organophosphate and carbamate insecticides: establishment of the baseline. Geneva, World Health Organization (WHO/VBC/81.805).

WHO (2005). Review of Novaluron 10% EC, World Health Organization Communicable Disease Control, Prevention and Eradication. WHO Pesticides Evaluation Scheme (WHOPES). Report of the eighth WHOPES Working Group Meeting held in Geneva, 1- 3 December, 2004.

Table 1: The mean mortality effects of different concentrations of Novaluron on 4th instar larvae *Aedes aegypti* after 12 hours interval

Mean of Mortality (%)

Dose ($\mu\text{g/ml}$)	3(hrs)	6(hrs)	9(hrs)	12(hrs)	Mean	% Mortality
200	1	11	12	19	4.75 \pm 1.9	95
100	3	7	12	17	4.25 \pm 0.8	85
50	1	5	10	15	3.75 \pm 1.1	75
20	1	4	10	14	3.5 \pm 1.2	70
12.5	1	3	7	12	3.0 \pm 1.3	60
6.25	1	3	5	9.0	2.25 \pm 0.9	45
3.125	1	2	4	7.0	1.75 \pm 0.05	35
Total	9.0	35.0	60.0	93		
Mean	1.29 \pm 0.2	5.00 \pm 1.5	8.57 \pm 1.8	13.29 \pm 2.0		
Control	0	0	0	0		

LSD= 2.214

Means of five replicates (\pm s.e)


Table 2: The mean mortality effects of different concentrations of MOSO on 4th instar larvae of *Aedes aegypti* mosquito.

Mean of Mortality (%)						
Dose ($\mu\text{g/ml}$)	3(hrs)	6(hrs)	9(hrs)	12(hrs)	Mean	% Mortality
200	1	4	4	20	4.75 \pm 1.5	100
100	1	2	11	17	4.25 \pm 1.8	85
50	2	4	11	16	4.00 \pm 1.4	80
20	1	3	7	13	3.25 \pm 1.3	65
12.5	1	3	6	12	3.0 \pm 1.2	60
6.25	1	2	4	9	2.25 \pm 0.1	45
3.125	1	2	3	5	2.25 \pm 0.1	25
Total	8.0	20.0	64.0	91		
Mean	1.14 \pm 0.1	2.85 \pm 1.1	8.00 \pm 1.5	13.00 \pm 2.1		
Control	0	0	0	0		

LSD= 2.44

Means of five replicates (\pm s.e)

Figure 1: Probit vs log₁₀dose of Novaluron and MOSO on 4th instar larvae of *Ae. aegypti*



The regression equations are given as: Probit (Y) = 0.912X + 3.916 where X = log dose value

From the graph, for Novaluron, log LD₅₀ = 0.9046. So, LD₅₀ = 8.0280µg/ml for *Moringa Oleifera* log LD₅₀ = 0.999. So, LD₅₀ = 9.977µg/ml

Table 3: The mean mortality effects of different concentrations of Novaluron on 1st *Aedes aegypti* mosquito after 12 hours interval
Mean of Mortality (%)

Dose (µg/ml)	3(hrs)	6(hrs)	9(hrs)	12(hrs)	Mean	% Mortality
200	10	13	17	19	4.75±1.6	95
100	8	11	16	18	4.5±1.4	90
50	6	8	11	17	4.25±1.3	85
20	9	12	14	15	3.75±1.2	75
12.5	8	9	11	14	3.5±1.2	70
6.25	6	8	10	11	2.75±1.1	60
3.125	4	5	7	9	2.25±0.09	45
Total	51	66	86	103		
Mean	7.29± 2.1	16.5± 4.2	12.29±2.5	14.71±3.3		
Control	0	0	0	0		

LSD= 1.59

Means of five replicates (± s.e)


Table 4: The mean mortality effect of different concentrations of MOSO on 1st instar larvae of *Aedes aegypti* mosquito
 Mean of Mortality (%)

Dose (µg/ml)	3(hrs)	6(hrs)	9(hrs)	12(hrs)	Mean	% Mortality
200	5	14	16	18	4.5±1.5	90
100	5	10	12	15	3.75±1.1	75
50	6	8	9	13	3.25±1.3	65
20	5	7	8	10	2.5±1.1	50
12.5	5	6	8	9	2.25±1.1	45
6.25	2	5	6	8	2.0±0.07	40
3.125	2	4	5	6	1.5±0.07	30
Total	30	54	58	79		
Mean	4.28±0.8	7.71±2.1	9.14±2.2	11.28±3.3		
Control	0	0	0	0		

LSD= 1.73

Means of five replicates (± s.e)

Figure 2: graph of probit against logdose on effect of Novaluron and MOSO on 1st instar larvae of *A. aegypti*.



The regression equation is given as

$$\text{Probit (Y)} = 0.912X + 3.916 \quad \text{where } X = \text{log dose value}$$

From the graph, for Novaluron, $\log LD_{50} = 0.6578$. So, $LD_{50} = 4.546\mu\text{g/ml}$
 for *Moringa olifera* $\log LD_{50} = 1.1886$. So, $LD_{50} = 15.44\mu\text{g/ml}$