

# EXTENSION OF FORMULAS FOR PARTITION FUNCTIONS

M.S. Ladan<sup>1</sup>, D. Singh<sup>2</sup> and Y. Tella<sup>3</sup>

<sup>1</sup>Department of Statistics, Yaba College of Technology, Yaba, Lagos.

<sup>2</sup>Department of Mathematics, Ahmadu Bello University, Zaria. Nigeria.

<sup>3</sup>Department of Mathematics, Kaduna State University, Kaduna. Nigeria.

Corresponding Author's E-mail: [sm\\_ladan@yahoo.com](mailto:sm_ladan@yahoo.com)

Tel: +2348023313732

## ABSTRACT

This paper studies the elementary method for finding formulas for some partition functions. The idea dates back to Cayley and Macmahon, using partial fractions decomposition to obtain traceable power series expansions. The ideas are hereby extended with the aid of software called *maple package*.

**Keywords:** Formula, Integer Partitions, Partition functions

## INTRODUCTION

Partition of a non-negative integer  $n$  is a non-increasing sequence of positive integers  $\lambda_1, \dots, \lambda_n$ , that sum to  $n$ . For example the integer 4 has the following five partitions,  $\{4, 31, 22, 211, 1111\}$ , see Andrews (1984).

We denote by  $p_k(n)$  the number of partitions of  $n$  into at most  $k$  parts, Gupta (1970).

Andrew and Erikson (2004) also stated that  $p_k(n)$  equals the number of partitions of  $n$  into at most  $k$  parts. There is an extensive literature concerning the formula for  $p_k(n)$ , including contributions by Cayley (1856), Sylvester (1882), Glaisher (1909) and Gupta (1958). For additional references and historical notes, see Gupta (1970). Furthermore, the theory of  $q$ -partial fractions and its formula was developed in Munagi (2007).

## Some Important Definitions

### Partition

Singh *et al.* (2012) defined a partition of a positive integer  $n$  as a sequence of positive integers whose sum is  $n$ .

The order of the summands is unimportant when writing the partitions of  $n$ . For example the partitions of  $n = 4$  are given as

$$\begin{aligned} 4 &= 4 \\ &= 3 + 1 \\ &= 2 + 2 \\ &= 2 + 1 + 1 \\ &= 1 + 1 + 1 + 1 \end{aligned}$$

### Partition function $p(n)$

Andrew and Erikson (2004) stated that the partition function  $p(n)$  counts the number of unique partitions of the positive integer  $n$ .

For example, there are five unique partitions of 4. Hence,  $p(4) = 5$ .

### Intermediate function

The intermediate function denoted by  $p_k(n)$  is defined such that it counts the partitions of  $n$  with the largest added being no smaller than  $k$ , Gupta (1970).

## METHODOLOGY

In this section, we consider the formula  $p_k(n)$ , which was computed for  $k = 1, 2, 3, 4$ . by Andrews (2003). So our desire in this paper is to compute this formula for  $k = 5, 6, \dots, 11$ .

Furthermore, we implement this formula using computer program called *maple software*.

That is, Andrews (2003) deduced

$$\begin{aligned} p_1(n) &= 1, \\ p_2(n) &= \left\lfloor \frac{n+1}{2} \right\rfloor, \\ p_3(n) &= \{(n+3)^2/12\} \end{aligned}$$

Thereafter, Andrew and Erikson (2004) gave  $p_k(n) = p(n, k)$  the following generating function

$$\sum_{n=0}^{\infty} p(n, k) q^n = \frac{1}{(1-q)(1-q^2)\dots(1-q^k)} \quad (1)$$

Now, consider the case  $k = 4$ , we have maple calculate that

$$\begin{aligned} \sum_{n=0}^{\infty} p(n, 4) q^n &= \frac{1}{(1-q)(1-q^2)(1-q^3)(1-q^4)} \\ &= \frac{\frac{25}{144}}{(1-q)^2} + \frac{\frac{1}{8}}{288(1-q)^3} \\ &\quad + \frac{\frac{1}{24}}{(1-q)^4} + \frac{\frac{1}{16}}{(1-q)^2} \\ &\quad + \frac{\frac{1}{8}}{(1-q^2)(1-q)^2} \\ &\quad + \frac{1}{q(q+2)} \left( \frac{1}{1-q^3} \right) + \frac{\frac{1}{4}}{(1-q^4)} \\ &= \frac{\frac{1}{24}}{(1-q)^4} + \frac{\frac{1}{8}}{(1-q)^3} + \frac{\left(\frac{5}{12}\right)^2}{(1-q)^2} \\ &\quad + \frac{\frac{1}{8}}{(1-q^2)^2} + \frac{\frac{1}{16}}{1-q^2} + \frac{\frac{(2+q)}{9}}{1-q^3} \\ &\quad + \frac{\frac{1}{4}}{1-q^4} \\ &= \sum_{n=0}^{\infty} \left( \frac{1}{24} \binom{n+3}{3} + \frac{1}{8} \binom{n+2}{2} \right. \\ &\quad \left. + \left(\frac{5}{12}\right)^2 (n+1) \right) q^n \\ &\quad + \left( \frac{1}{8} (n+1) + \frac{1}{16} \right) q^{2n} \\ &\quad + \sum_{n=0}^{\infty} \left( -\frac{1}{16} q^{2n} + \frac{2}{9} q^{3n} + \frac{1}{9} q^{3n+1} \right. \\ &\quad \left. + \frac{1}{4} q^{4n} \right) \end{aligned}$$

Here, the coefficient of  $q^n$  is the formula for  

$$p(n, 4) = \frac{1}{24} \binom{n+3}{3} + \frac{1}{8} \binom{n+2}{2} + \frac{25}{144} (n+1) \quad (2)$$

**Case k = 5**

$$\sum_{n=0}^{\infty} p(n, 5) q^n = 1(1-q) \left( 1 - q^2 \right) \dots \left( 1 - q^5 \right) \frac{1}{(1-q)^5}$$

$$+ \frac{1}{24} \frac{1}{(1-q)^4} + \frac{31}{288} \frac{1}{(1-q)^3} + \frac{11}{64} \frac{1}{(1-q)^2}$$

$$+ \frac{11}{64} \frac{1}{1-q^2} + \frac{1}{9} \frac{1}{1-q^3} + \frac{1}{5} \frac{1}{1-q^5}$$

$$= \left( \frac{1}{120} \binom{n+4}{4} + \frac{1}{24} \binom{n+3}{3} \right. \\ \left. + \frac{31}{288} \binom{n+2}{2} + \frac{11}{64} \binom{n+1}{1} \right) q^n$$

$$+ \frac{11}{64} \sum_{n=0}^{\infty} q^{2n} + \frac{1}{9} q^{3n} + \frac{1}{5} \sum_{n=0}^{\infty} q^{5n}$$

$$p(n, 5) = \frac{1}{120} \binom{n+4}{4} + \frac{1}{24} \binom{n+3}{3} + \frac{31}{288} \binom{n+2}{2} + \frac{11}{64} \binom{n+1}{1} \quad (3)$$

**Case k = 6**

$$\sum_{n=0}^{\infty} p(n, 6) q^n = \frac{1}{(1-q)(1-q^2) \dots (1-q^6)}$$

$$= \frac{1}{720(1-q)^6} + \frac{1}{96(1-q)^5}$$

$$+ \frac{17}{432(1-q)^4} + \frac{331}{3456} \left( \frac{1}{(1-q)^3} \right)$$

$$+ \frac{777611}{518400} \left( \frac{1}{(1-q)^2} \right) + \frac{25}{256(1-q^2)}$$

$$= \frac{1}{720} \sum_{n=0}^{\infty} \binom{n+5}{5} q^n$$

$$+ \frac{1}{96} \sum_{n=0}^{\infty} \binom{n+4}{4} + \frac{17}{432} \sum_{n=0}^{\infty} \binom{n+3}{3}$$

$$+ \frac{331}{3456} \sum_{n=0}^{\infty} \binom{n+2}{2}$$

$$+ \frac{777611}{518400} \sum_{n=0}^{\infty} (n+1) + \frac{25}{256} q^{2n}$$

$$p(n, 6) = \frac{1}{720} \binom{n+5}{5} + \frac{1}{96} \binom{n+4}{4} + \frac{17}{432} \binom{n+3}{3} + \frac{331}{3456} \binom{n+2}{2} + \frac{777611}{518400} \binom{n+1}{1} \quad (4)$$

**Case k = 7**

$$\sum_{n=0}^{\infty} p(n, 7) q^n = \frac{1}{(1-q)(1-q^2) \dots (1-q^7)}$$

$$= \frac{1}{5040} \left( \frac{1}{(1-q)^6} \right) + \frac{1}{480(1-q)^5}$$

$$+ \frac{47}{4320} \left( \frac{1}{(1-q)^4} \right)$$

$$+ \frac{161}{4320} \left( \frac{1}{(1-q)^3} \right)$$

$$+ \frac{7913}{86400} \left( \frac{1}{(1-q)^2} \right)$$

$$= \frac{1}{5040} \sum_{n=0}^{\infty} \binom{n+5}{5} q^n$$

$$+ \frac{1}{400} \sum_{n=0}^{\infty} \binom{n+4}{4} q^n$$

$$+ \frac{47}{4320} \sum_{n=0}^{\infty} \binom{n+3}{3} q^n$$

$$+ \frac{161}{4320} \sum_{n=0}^{\infty} \binom{n+2}{2} q^n$$

$$+ \frac{7713}{86400} \sum_{n=0}^{\infty} \binom{n+1}{1} q^n$$

$$p(n, 7) = \frac{1}{5040} \binom{n+5}{5} + \frac{1}{480} \binom{n+4}{4} + \frac{47}{4320} \binom{n+3}{3} + \frac{161}{4320} \binom{n+2}{2} + \frac{7913}{86400} \binom{n+1}{1} \quad (5)$$

**Case k = 8**

$$\sum_{n=0}^{\infty} p(n, 8) q^n = \frac{1}{(1-q)(1-q^2) \dots (1-q^8)}$$

$$= \sum_{n=0}^{\infty} \left\{ \left( \frac{1}{40320} \right) \left( \frac{1}{(1-q)^8} \right) \right.$$

$$+ \left( \frac{1}{2880} \right) \left( \frac{1}{(1-q)^7} \right)$$

$$+ \left( \frac{83}{34560} \right) \left( \frac{1}{(1-q)^6} \right)$$

$$+ \left( \frac{25}{2304} \right) \left( \frac{1}{(1-q)^5} \right)$$

$$+ \left( \frac{24523}{691200} \right) \left( \frac{1}{(1-q)^4} \right)$$

$$+ \left( \frac{139}{1600} \right) \left( \frac{1}{(1-q)^3} \right)$$

$$+ \left. \frac{487033}{3175200} \binom{n+1}{1} \right\}$$

$$p(n, 8) = \frac{1}{40320} \binom{n+7}{7} + \frac{1}{2880} \binom{n+6}{6} + \frac{83}{34560} \binom{n+5}{5} + \frac{25}{2304} \binom{n+4}{4} + \frac{24523}{691200} \binom{n+3}{3} + \frac{139}{1600} \binom{n+2}{2} + \frac{487033}{3175200} \binom{n+1}{1} \quad (6)$$

**Case k = 9**

$$\sum_{n=0}^{\infty} p(n, 9) q^n = \frac{1}{(1-q)(1-q^2) \dots (1-q^9)} = \sum_{n=0}^{\infty} \left\{ \frac{1}{326880(1-q)^9} + \frac{1}{20160(1-q)^8} + \frac{1}{725760(1-q)^7} + \frac{1}{725760(1-q)^6} + \frac{87091200(1-q)^5}{239185} + \frac{8335199}{929377} \right\} q^n =$$

$$\frac{1}{6967296(1-q)^4} + \frac{1}{97542144(1-q)^3} + \frac{1843}{11943936(1-q)^2} \left. \right\} q^n =$$

$$\frac{1}{326880} \binom{n+8}{8} + \frac{1}{20160} \binom{n+7}{7} + \frac{319}{725760} \binom{n+6}{6} + \frac{1843}{725760} \binom{n+5}{5} + \frac{929377}{87091200} \binom{n+4}{4} + \frac{239185}{6967296} \binom{n+3}{3} + \frac{8335199}{97542144} \binom{n+2}{2} + \frac{1821389}{11943936} \binom{n+1}{1} \quad (7)$$

**Case k = 10**

$$\sum_{n=0}^{\infty} p(n, 10) q^n = \frac{1}{(1-q)(1-q^2) \dots (1-q^{10})}$$

$$= \sum \left\{ \frac{1}{362880(1-q)^{10}} + \frac{199}{161280(1-q)^9} + \frac{2867}{2903040(1-q)^8} + \frac{5806080}{2867(1-q)^7} + \frac{161111}{161111(1-q)^6} + \frac{62208000}{2600533(1-q)^5} + \frac{248832000}{1826428579(1-q)^4} + \frac{54867456000}{36618675691(1-q)^3} + \frac{438939648000}{40769204821(1-q)^2} \right\} q^n$$

$$p(n, 10) = \frac{1}{326880} \binom{n+9}{9} + \frac{1}{161280} \binom{n+8}{8} + \frac{199}{2903040} \binom{n+7}{7} + \frac{2867}{5806080} \binom{n+6}{6} + \frac{161111}{62208000} \binom{n+5}{5} + \frac{2600533}{248832000} \binom{n+4}{4} + \frac{1826428579}{54867456000} \binom{n+3}{3} + \frac{36618675691}{438939648000} \binom{n+2}{2} + \frac{40769204821}{268738560000} \binom{n+1}{1} \quad (8)$$

**Case k = 11**

$$\sum_{n=0}^{\infty} p(n, 11) q^n = \frac{1}{(1-q)(1-q^2) \dots (1-q^{11})}$$

$$= \sum_{n=0}^{\infty} \left\{ \frac{1}{39916800(1-q)^{11}} + \frac{1}{14151520(1-q)^{10}} + \frac{1}{107520(1-q)^9} + \frac{1161216}{95(1-q)^8} + \frac{114221}{114221(1-q)^7} + \frac{217728000}{2262473(1-q)^6} + \frac{870912000}{2250121541(1-q)^5} + \frac{219469824000}{14347821041(1-q)^4} + \frac{438939648000}{1099442945011(1-q)^3} + \frac{13168189440000}{17080331207(1-q)^2} \right\} q^n$$

$$p(n, 11) = \frac{1}{39916800} \binom{n+10}{10} + \frac{1}{1415120} \binom{n+9}{9} + \frac{1}{107520} \binom{n+8}{8} + \frac{95}{1161216} \binom{n+7}{7} + \frac{114221}{217728000} \binom{n+6}{6} + \frac{2262473}{870912000} \binom{n+5}{5} + \frac{2250121541}{219469824000} \binom{n+4}{4} + \frac{14347821041}{438939648000} \binom{n+3}{3} + \frac{1099442945011}{13168189440000} \binom{n+2}{2} + \frac{17080331207}{107495424000} \binom{n+1}{1} \quad (9)$$

**Example**

We illustrate the above formulas for the case n = 50, m = 10  
 Substituting n = 50

$$p(50, 10) = \frac{1}{3628800} \binom{59}{9} + \frac{1}{161280} \binom{58}{8} + \frac{199}{2903040} \binom{57}{7} + \frac{2867}{5806080} \binom{56}{6} + \frac{161111}{62208000} \binom{55}{5} + \frac{2600533}{248832000} \binom{54}{4} + \frac{1826428579}{54867456000} \binom{53}{3} + \frac{36618675691}{438939648000} \binom{52}{2} + \frac{40769204821}{268738560000} \binom{51}{1}$$

$$= 16928$$

This result agrees with the one obtained with maple via inbuilt command "numbpart" from the combination package:

>with (combinat, numbpart);

> NumbPart(50, 10); 16928

**Conclusion**

We have studied the elementary method for finding formulas for some partition functions, which used partial fractions decomposition to obtain traceable power series expansions. The ideas had been extended with the aid of Maple software package.

**REFERENCES**

Andrews, G.E., 2003, Partitions at the interface of q-series and modular forms, *Ramanujan Journal.*, Vol.7, 384-400.  
 Andrews, G.E., 1984, *The Theory of Partitions*, Cambridge University Press, New York.  
 Andrews, G.E. & Erikson, K., 2004, *Integer Partitions*, Cambridge University Press, New York.  
 Cayley, A., 1856, Researches on the partition of numbers, *Phil. Trans. Royal Soc. of London*, 146, 127-140.  
 Glaisher, J.W.L., 1909, Formulae for partitions into given elements, derived from Sylvester's theorem, *Quart. J. Pure Appl. Math.* 40, 275-348.  
 Gupta, H., 1958, *Tables of partitions*, Royal Society Mathematical Tables, vol.4, Cambridge University Press.  
 Gupta, H., 1970, Partitions - A survey, *Journal of research of the National Bureau of Standards - Mathematical Sciences*, January - March, 748(1)  
 Sylvester, J.J., 1882, On Sub-invariants, i.e. semi-invariants to binary quartics of an unlimited order, on rational fractions and partitions, *Quart. J. Math. (oxford series (2))* 2, 85-108.  
 Singh, D., Ibrahim, A.M., Singh, J.N., & Ladan, M.S. 2012, An overview of Integer partitions, *Journal of Mathematical Sciences and Mathematical Education*, 7(2), 19-31.  
 Munagi, A.O., 2007, Computation of q-partial fractions, *Integers* 7, #A25.

**APPENDIX**

**Maple Code for Generating q-Partial Fraction and Iterative Transformation Algorithm**

```
with(numtheory, cyclotomic, divisors, phi, invphi)
local a, t, i, h;
type/qfactors := proc(f, q::name)
a := primpart (f, q)*(1 + q^7)^2;
qseries:=table();
qseries[aqprod]:=proc(a,q,n)
: added else bit when n not an integer
local x,i;
if type(n,nonnegint) then
x:=1:
fori from 1 to n do
x := x * (1-a*q^(i-1));
od:
else
x:=`(a,q)[n];
fi:
RETURN(x);
end:
```

**>Partitions Function:**

```
pmn(3,n);
pmn:=proc(m, n) local a, gu, Gu, H, gu1, lu1, M, q, i, eqs, form,
unknowns, result, soln,j,k,l;
option remember;
if m = 1 then RETURN([[1]]) end if;
Gu :=mul(1/(1-q^i), i = 1 .. m);
gu := convert(Gu, parfrac);
H := [];
```

```
fori to m do
M := floor((m-i)/i);
gu1 := add(coeff(gu, numtheory:-cyclotomic(i, q),
-j)/numtheory:-cyclotomic(i, q)^j, j = 1 .. floor(m/i));
lu1 := convert(taylor(gu1, q = 0, i*(1+M)), polynomial);
unknowns := {seq(a[j], j = 0 .. -1+i*(1+M))};
eqs := {seq(seq(add((k*i+l)^j*a[j*i+l], j = 0 .. M)
= coeff(lu1, q, k*i+l), k = 0 .. M), l = 0 .. i-1)};
form := [seq(add(a[j]*l^(1-q)^j, j = 0 .. M), l = 0 .. i-1)];
soln := solve(eqs, unknowns);
result := subs(soln, form);
result:=[op(2..nops(result),result),result[1]];
H := [op(H), result]
od:
H:
end:
```

**>Polynomials of the form**  $\frac{1}{(1 - q^n)}$

```
with(numtheory, cyclotomic, divisors, phi, invphi)
type/qfactors := proc(f, q::name)
local a, t, i, h;
option
description "Tests if f is a constant multiple of factors of
polynomials of the form 1-q^n";
if not type(f, polynomial(anything, q)) then return false end if;
a := primpart (f, q)*(1 + q^7)^2;
if op(1,a) = -1 then a:= -a end if;
t :=0;
fori to nops(a) do
if type (op(i, a), { '^', '!' }) then
h :=op(1, op(i,a)^2); if abs(h) = abs(1 - q^degree (h, q))then t := t
+ 1 end if
else return false
```