

PREVALENCE OF *TRICHOMONAS VAGINALIS* AMONG PREGNANT WOMEN ATTENDING ANTENATAL CLINIC IN TWO HEALTH FACILITIES WITHIN KADUNA METROPOLIS, KADUNA, NIGERIA

*I.K. Auta, Ibrahim B. and D. Henry

Department of Biological Sciences, Kaduna State University, Tafawa Balewa Way, Kaduna, Nigeria

*Corresponding Author's Email Address: ishayakato@yahoo.com

Phone: +2348034856567

ABSTRACT

Trichomoniasis being referred to as the most prevalent sexually transmitted infection of both male and female urogenital organs. This study was aimed at determining the prevalence of *Trichomonas vaginalis* among pregnant women attending antenatal at Sabon Tasha General Hospital (STGH) and Yusuf Dantsoho Memorial Hospital (YDMH) who were not on any treatment for sexually transmitted infection for 30 days prior to the period of this research. A total of four hundred and five (405) pregnant women consisting of two hundred and one (201) and two hundred four (204) from Sabon Tasha General Hospital and Yusuf Dantsoho Memorial Hospital respectively, were involved in the study between the periods of September 2019 to January 2020. Socio-demographic information such as age and marital status were obtained from all participating women within the age range of 15-43 years. The prevalence of trichomoniasis 25(6.17%) and 9(4.48%) for YDMH and STGH respectively was observed. Although the highest rate of trichomoniasis infection was recorded in age group of sexually active women (26-30), the prevalence of trichomoniasis in this study was not significantly affected by age and marital status ($P>0.05$). However, the variation of infection rate between the two health facilities in this study is an indication that the prevalence rate of trichomoniasis varies with location, nature of hygiene, sexual habit and level of education of the individual. The study advocates for sexual and health education at every antenatal clinic and routine screening for trichomoniasis among pregnant and women of reproductive age in order to curb the outbreak of the disease.

Keywords: Prevalence, *Trichomonas vaginalis*, Pregnant Women, Antenatal, Health Facilities, Kaduna Metropolis

INTRODUCTION

Trichomonas vaginalis is a protozoan parasite and it is the most prevalent non-viral sexually transmitted infection worldwide causing the curable sexually transmitted disease called trichomoniasis (Secor *et al.*, 2014; Adegbaaju and Morenike, 2008). It has been recognized as a cosmopolitan parasite of genital tract of both male and female as reported by Jatau, *et al.* (2006) with an annual prevalence of about 180 million cases in the world (World Health Organisation, 2012). The majority of cases of trichomoniasis are localized in regions of low income particularly in African countries where the prevalence rate ranges from 15 to 37% (Swygard, *et al.*, 2004; Usanga *et al.*, 2009; Okoko, 2011; Sam -wobo *et al.*, 2012; and Chinedum, *et al.*, 2014), due to lack of adequate resources for health care (WHO,

2012). *T. vaginalis* is mainly transmitted through sexual contact but may also be transmitted through sharing of towels and under wears with infected individuals (Alcamo, 2000). Most cases of *T. vaginalis* remain undiagnosed as it is currently not a target of sexually transmitted infections control and besides, because of its asymptomatic nature in about half of infected men and women (Fouts and Kraus, 1980). The disease is reported as a major cause of pathology in obstetrics and gynecology (Hook, 1999; Hobbs *et al.*, 2006). The life cycle of *T. vaginalis* is still poorly understood like many other protozoan parasites, it is known to exist only as astrophozoite and lacks a cystic stage. It is a primitive eukaryotic organism that in its carbohydrate and energy metabolism shows remarkable similarity to anaerobic bacteria (Petrin, *et al.*, 1998). *T. vaginalis* trophozoite is an oval, parasite with five flagella and an axostyle project which may be used for attachment to surface and may also cause tissue damage noted in *T. vaginalis* infection (Swygard, *et al.*, 2012).

Historically, the presence of *T. vaginalis* has been viewed as a risk marker for other sexually transmitted agents such as *Chlamydia trachomatis*, *Neisseria gonorrhoea* or bacterial vaginosis (Wolner – Hanssen *et al.*, 1989; Petrin *et al.*, 1998). Its importance is also being reassessed in the light of recent evidence that is associated with adverse pregnancy outcome and facilitates the sexual transmission of HIV infection (Mabey *et al.*, 2006; McClelland *et al.*, 2007). Reports have also implicated *T. vaginalis* in upper reproductive tract post-surgical infection, reversible infertility, neonatal morbidity and mortality (Laga *et al.*, 1993; Draper *et al.*, 1995; Pastorek *et al.*, 1996). *T. vaginalis* is also increasingly been recognized to be associated with reproductive tract complications including sepsis that occurs after abortion and after cesarean section (Minkoff *et al.*, 1984) as well as adverse pregnancy outcome (Cotch *et al.*, 1997). It has also been reported that the disease causes discomfort and psychosocial distress in infected patients (Jatau *et al.*, 2006). Complications of *T. vaginalis* reported among pregnant women and non-pregnant patients may include premature rupture of membrane, premature labour, low birth weight, post abortion infections, pelvic inflammatory disease, urinary tract infections, bronchitis pneumonia and oral lesions (Soper, 2004) and infertility (Gookin *et al.*, 2005; Fichorova, 2009). It has also been linked to one of the predisposing factors of HIV infection, acquired immune deficiency syndrome and cervical cancer (Soper, 2004; Sobel, 2005; Mabey *et al.*, 2006; McClelland *et al.*, 2007).

The symptoms of *T. vaginalis* which is commonly observed in women than in men (Smith and Ramos, 2010) may include frothy-greenish foul-smelling vaginal discharge accompanied with


vulvovaginal irritation, postictal bleeding, frequency in micturition, dysuria and lower abdominal pains (CDC, 2006). Lower birth weight infants, preterm rupture of membranes, preterm delivery and neonatal mortality and morbidity are symptoms associated with pregnancy (Johnson *et al.*, 2011).

Multiple sex partners, poor personal hygiene, low socio-economic status and underdevelopment are factors reported to be associated with high incidence of infection (Huppert, 2009). *Trichomonas vaginalis* is detected in vaginal prostatic or urethral secretions, semen and urine of infected individuals using different laboratory methods such as wet mount, various staining methods, culture, latex agglutination, Enzyme Linked Immunosorbent Assay (ELISA) and more recently, Polymerase Chain Reaction (PCR) (Radonjic *et al.*, 2006). The most common means of diagnosis still remains microscopic visualization of the vaginal fluid (Schwebke and Burgess, 2004). Direct examination of wet mount preparation of clinical specimen is the most rapid and least expensive technique for identifying *T. vaginalis*.

MATERIALS AND METHODS

Study Area

The study was conducted in two major Government owned hospitals in Kaduna metropolis which include Sabon Tasha General Hospital with GPS coordinate of 10° 26' 56.9" N 7° 28' 43" E and Yusuf Dan Tsoho Memorial Hospital with GPS coordinate of 10°31' 15" N 7°24' 59" E in Kaduna South Local Government Area of Kaduna State. The two health facilities were established to provide secondary health services to people living in the respective wards and environs



Ethical approval

Ethical approval was obtained from Kaduna State Ministry of Health, Sabon Tasha General Hospital and Yusuf Dantsoho Memorial Hospital, Kaduna.

Specimen collection and processing

Method described by Cheesbrough (1999) was used to obtain the vaginal swab from each participating woman. The high vaginal swab was obtained by inserting a sterile cotton wool swab stick into the posterior fornix of the vagina with the aid of plastic disposable speculum and gently rotated about 2-3times in order to pick the vaginal exudate. It was immediately replaced in its casing and transported to the laboratory. Wet mount was prepared by emulsifying the vaginal exudate on a drop of normal saline placed on a grease-free microscopic slide, and viewed microscopically for the presence or absence of motile trophozoites using 10x and 40x objective lenses. Giemsa staining was also conducted as described by Manson *et al.* (1976) by making smear of the exudate on a grease-free microscopic slide, fixing it in absolute ethanol for 1minute and allowed to air-dry. Diluted Giemsa was poured on the smear and allowed to stain for 10min after which it was washed, air-dried and examined microscopically with oil immersion 100x objective lens for the presence or absence of trichomonas vaginalis.

Statistical analysis of data

Data obtained were analyzed using Chi-square and Odds ratio (OR) for the potential risk factors. P-value ≤ 0.05 was considered statistically significant.

RESULTS

Out of the 405 pregnant women examined, 16 (7.84%) and 9 (4.48%) were positive for women attending antenatal in Yusuf Dantsoho Memorial Hospital and Sabon Tasha General Hospital respectively. There was no significant difference ($P > 0.05$) between the prevalence of the infection at the two health facilities (Table 1)

Table 1: Prevalence of *Trichomonas vaginalis* among pregnant women attending antenatal clinic in YDMH and STGH within Kaduna Metropolis

Health Facility	No Examined	Positive (%)	Negative (%)
YDMH	204	16 (7.84)	188 (92.15)
STGH	201	9 (4.48)	192 (95.52)
Total	405	25 (6.17)	380 (93.82)

$P > 0.05$

Keys: YDMH = Yusuf Dan Tsoho Memorial Hospital, STGH = Sabon Tasha General Hospital

However, women between the ages of 26 and 30 years had the highest prevalence of 11 (9.32%) and 6 (5.45%) while those between ages 15 and 20 years had 1 (2.38) and 1 (1.92) for YDMH and STGH respectively. There were no significance difference between the two health facilities based on age of the women (Table 2).

Table 2: Prevalence of *Trichomonas vaginalis* among pregnant women attending antenatal clinic in STGH and YDMH in relation to age within Kaduna Metropolis

Age category (years)	YDMH		STGH		Total positive (%)
	No. examined	No. positive (%)	No. examined	No. positive (%)	
15-20	41	1 (2.38)	11	0 (0.0)	1 (1.92)
21-25	61	4 (6.55)	49	2 (4.08)	6 (5.45)
26-30	48	8 (16.6)	70	3 (4.29)	11 (9.32)
31-35	34	3 (8.82)	48	2 (4.17)	5 (6.10)
36 and above	20	0 (0)	23	2 (8.69)	2 (4.65)
Total	204	16 (7.84)	201	9 (4.48)	25 (6.17)

P> 0.05

Keys: YDMH = Yusuf Dan Tsoho Memorial Hospital, STGH = Sabon Tasha General Hospital

DISCUSSION

The findings from this study revealed the prevalence of *T. vaginalis* among pregnant women attending Yusuf Dan Tsoho Memorial Hospital and Sabon Tasha General Hospital within Kaduna Metropolis. This is similar to the findings of Aboyeji and Nwabusi (2003) who reported the prevalence of 4.7% in Ilorin. Similarly, Roger *et al.* (2019) reported the prevalence of 4.8% in Senegal. However, the prevalence of 7.84% in Yusuf Dan Tsoho memorial Hospital is similar to the finding of Oladeinde *et al.* (2016) who reported the prevalence of 7.7% among pregnant women of reproductive age in Edo state Nigeria.

The prevalence in this study is however lower than the 9.7% reported by Isiaka *et al.* (2014) in Ilorin, 13.3% reported by Obiukwu (2010) in Anambra, 17.7% reported by Okpara *et al.* (2009) in Uyo, 20.0% in Abeokuta by Ojuronbe *et al.* (2010), 18.7% reported by Jatau *et al.* (2006) in Zaria, 12.16% reported in Maryland by Coth *et al.* (1997), 16% reported by Mayaud *et al.* (1998) in Tanzania and 46.9% reported in New York by Shuter (1998). The finding is also higher than the findings of Uneke *et al.* (2005) who reported the prevalence of 2.8% in Ebonyi state South-eastern Nigeria. It is also higher than 1.4% reported by Begun *et al.* (2003) in Dakan Bangladesh, 3.3% by Adeoye and Akande (2007) in Lagos, 2.8% by Akinbo *et al.* (2017) in Benin City.

Although the prevalence rate of 4.48% and 7.84% observed in both health facilities can be considered to be within the normal range in a healthy women population. Cameron and Padian (1990) Observed infection rates of 5 to 10% in a healthy women population while it could be as high as 50% in prostitutes and female prisoners. There is a general conception that prevalence of *T. vaginalis* ranges markedly based on settings and locations (Bowden and Garnett, 2000). Hence, the variation in respect to the two health facility and findings of other authors could be due to differences in geographical location, population, personal hygiene as well as diagnostic methods used in the study.

Conclusion

Prevalence of (25) 6.17% and 9 (4.48%) were obtained for Yusuf Dan Tsoho Memorial Hospital and Sabon Tasha General Hospital respectively in the present study. The highest rate of infection was

recorded in age group of sexually active women (26-30), the prevalence of trichomoniasis in this study was not significantly affected by age and marital status (P>0.05). There was variation of infection rate between the two health facilities in the study area. There should be routine screening of trichomoniasis among pregnant women and women of reproductive age in order to curb out the disease.

Acknowledgements

We acknowledged the assistance of the laboratory technologists of Yusuf Dantsoho Memorial Hospital Tudun Wada and Sabon Tasha General Hospital for their immense contribution and Kaduna State Ministry of Health for granting the Ethical approval for the research work

REFERENCES

- Aboyeji, A.P. and Nwabuisi, C. (2003). Prevalence of sexually transmitted disease among pregnant women in Ilorin, Nigeria. *Journal of Obstetrics and Gynaecology*, 22: 637-639.
- Adegbaju, A. and Morenikeji, O.A. (2008). Cytoadherence and pathogenesis of *Trichomonas vaginalis*. *Science Research Essay*. 3 (4): 132-138.
- Adeoye, G. O. and Akande, A. H. (2007). Epidemiology of *T. vaginalis* among women in Lagos metropolis, Nigeria. *Pakistan Journal of Biological Science*. 10 (13) :2198-2201. (Pubmed)(Google Scholar).
- Akinbo, F. O., Mokobia, C. N. and Ande, A. B. (2017). Prevalence of Trichomoniasis among pregnant women in Benin City. *Sahel Medical Journal*, 20:67-71
- Alcama, I. E. (2000). *Fundamentals of Microbiology*. Boston: Jones and Bartlett Publishers. Pp. 486-487
- Amadi, A. N. and Nwangbo, A. K. (2013). *Trichomonas vaginalis* Infection among Women in Ikwuano, Abia State, Nigeria. *Journal of Applied Sciences and Environmental Management*. 17 (3) : 389-398.
- Begun, A., Nilufar, S., Akther, K., Rahman, A., Khatun, F. and Rahman, M. (2003). Prevalence of selected reproductive tract infection among pregnant women attending an urban maternal and childcare unit in Dhaka, Bangladesh. *Journal of Health Population and Nutrition*, 21: 112-116.
- Bowden, F. J. and Garnet, G. P. (2000). *Trichomonas vaginalis* epidemiology: Parameterising and analysing a model of treatment interventions. *Sexually Transmitted Infections*, 76:248-56.
- Cameron, D. W. and Padian, N. S. (1990). Sexual transmission of HIV and the epidemiology of other sexually transmitted diseases. *AIDS*, 4:S99-103.
- CDC (2006). Sexually Transmitted Diseases. Treatment Guideline Available from <http://www.cdc.gov/STD/treatment/2006/vaginaldischarge.htm#vagdis3>. (last accessed on 2007 Jul 20).
- Cheesbrough M. (1999). *District Laboratory Practice in Tropical Countries*, Part 2, Cambridge University Press.
- Chinedum, O. K., Ifeanyi, O. E., Ugwu, U. G. and Ngozi, G. E. (2014). Prevalence Of *Trichomonas vaginalis* Among Pregnant Women Attending Hospital In Irrua Specialist Teaching Hospital In Edo State. *Nigeria. Journal of Medical and Dental Sciences*. 13 (9) : 79-82.
- Cotch, M. F., Pastorek, J. G., Nugent, R. P., Hillier, S. L., Gibbs, R. S., Martin, D. H., Eschenbach, D. A., Edelman, R., Carey,

- J. C., Regan, J. A., Klebanoff, M. A., Rao, A. V. and Rhoads, G. G. (1997). *Trichomonas vaginalis* Associated with Low Birth Weight and preterm Delivery. The Vaginal Infections and Prematurity Study Group. *Sexually Transmitted Diseases*, **24**:353-360.
- Draper, D., Jones, W., Heine, R. P., Beutz, M., French, J. L. and McGregor, J. A. (1995). *Trichomonas vaginalis* Weakens Human chorion in an In Vitro Model of Premature Membrane Rupture. *Infectious Disease Obstetrics and Gynecology*, **2**:267-74.
- Fichorova, R. N. (2009). Impact of *T. vaginalis* infection on innate immune responses and reproductive outcome. *Journal of Reproductive Immunology*, **83**:185-9.
- Fouts, A. C. and Kraus, S. J. (1980). *Trichomonas vaginalis*. Reevaluation of its clinical presentation and laboratory diagnosis. *Journal of Infectious Diseases*, **141**: 137-43.
- Gookin, J. L., Birkenheuer, A. J., St John, V., Spector, M. and Levy, M. G. (2005). Molecular characterization of trichomonads from faeces of dogs with diarrhea. *Journal of Parasitology*, **91**:939-43.
- Huppert, J. S. (2009). Trichomoniasis in Teens: An Update. *Current Opinion in Obstetrics and Gynecology*, **21**: 371378
- Isiaka-Lawal, S. A, Nwabuisi, C., Fakeye, O., Saidu, R., Adesina, K. T., Ijaiya, M. A, et al., (2014). Pattern of Sexually transmitted infections in human immunodeficiency virus-positive women attending antenatal clinics in north-central Nigeria. *Sahel Medical Journal*. **17** (4): 145-150.
- Jatau, E. D., Olonitola, O. S. and Olayinka, A. T. (2006). Prevalence of *Trichomonas* infection among women attending antenatal clinics in Zaria, Nigeria. *Annals of Africa Medicine*, **5**:178-81.
- Johnson, H. L., Ghanem, K. G., Zenilman, J. M. and Erbeling, E. J. (2011). Sexually transmitted infections and adverse pregnancy outcomes among women attending inner city public sexually transmitted diseases clinics. *Sexually Transmitted Diseases*, **38**:167-71.
- Klufio, C. A., Amoa, A. B., Delamare, O., Hombhanje, M. and Kariwiga, G. I. (1995). Prevalence of vaginal infections with bacterial vaginosis *Trichomonas vaginalis* and *Candida albicans* among pregnant women at the Port moresby General Hospital antenatal clinic. *Papua and New Guinea Medical Journal*, **38**:163-171
- Laga, M., Manoka, A., Kivuvu, M., Malele, B., Tuiiza, M., Nzila, N., Gafeman, J., Behets, F., Batter, V. and Aiary, M. (1993). Non-active Sexually Transmitted Diseases as Risk Factors for HIV-1 Transmission in Women: Results From a Cohort study. *AIDS*, **7**:95-102.
- Mabey, D., Ackers, J. and Adu-Sarkodie, Y. (2006). *Trichomonas vaginalis* Infection. *Sexually Transmitted Infectio*, **82**:26-7.
- Manson, P. R., Super, H., Fripp, P. J., (1976). Comparison of four techniques for the routine diagnosis of trichomonas vaginalis infection. *Journal of Clinical Pathology*, **29** : 154-157
- Mayaud, P., Uledi, E., Cornelissen, J. et al., (1998). Risk Scores to detect cervical infections in urban antenatal clinic attenders in Mwanza, Tanzania. *Sexually Transmitted Infection*. **74** [1]: S139-146
- McClelland, R. S., Sangare, L., Hassan, W. M., Lavreys, L., Mandaliya, K., Kiari, J., Ndinya-Achola, J., Jaoko, W. and Baeten, J.M. (2007). Infection with *Trichomonas vaginalis* Increases the Risk of HIV1 Acquisition. *Journal of Infectious Diseases*, **195**:698-702.
- Minkoff, H., Grunebaum, A. N., Schwarz, R. H., Feldman, J., Cummings, M., Crombleholme, W., Clark, L., Pringle, G. and McCormack, W.M. (1984). Risk Factors for Prematurity and Premature Rupture of Membranes: A Prospective Study of the Vaginal Flora in Pregnancy. *American Journal of Obstetrics and Gynecology*, **150**:965-72.
- Obianuju, I. O. and Ogbulie, J. N. (2005). Comparative study of the prevalence of sexually transmitted disease between pregnant women and non-pregnant women in Imo State, Nigeria. *Global Journal of Pure and Applied Sciences*, **339**-342.
- Obiukwu, M. O., Onyido, A. E., Duru, J. U., Aleke, O. (2010). *Trichomonas vaginalis* infection in anambra state: Demography and behavioural predictor. *Journal of Advances in Medical and Pharmaceutical Sciences*, **4**:16-20.
- Ojuronje, O., Taiwo, B. O., Dina, B. O., Agbaje, O. R., Bolaji, O. S., Adeyeba, A. O. (2010). Prevalence of *Trichomonas vaginalis* infection among pregnant women in Abeokuta, Nigeria. *Sierra Leone Journal of Biomedical Research*, **2**: 82-86.
- Okoko, F. J. (2011). Prevalence of Trichomoniasis among Women at Effurun Metropolis, Delta State, Nigeria. *Crescent Journal of Biological Sciences*, **4** (2):45-48.
- Okpara, K., Udoidung, N., Ating, I., Bassey, E., Okon, O. and Nwabueze, A. (2009). Risk factor for vaginalis trichomoniasis among women in Uyo Nigeria. *International Journal of Health Sciences*, **9**:1528-8315.
- Oladeinde, B. H., Oladeinde, O. B., Imade, O. S. and Onifade, A. A. (2016). *Trichomonas vaginalis* infection among pregnant of reproduction age in a rural community in Nigeria. *Nigerian Journal of Experimental Clinical Biosciences*, **4**:1-5.
- Pastorek, J. G., Cotch, M. F., Martin, D. H. and Eschenbach, D. A. (1996). Clinical and Microbiological Correlates of Vaginal Trichomoniasis During Pregnancy. The Vaginal Infections and Prematurity Study Group. *Clinical Infectious Diseases*, **23**:1075-80.
- Petrin, D., Delgaty, K., Bliatt, R., Garber, G. (1998). Clinical and microbiological aspects of *Trichomonas vaginalis*. *Clinical Microbiology Review*, **11**: 300-17.
- Radonjic, I. V., Dzamic, A. M., Mitrovic, S. M., ArsicArsenijevic, V. S., Popadic, D. M. and Kranjic Zee, I. F. (2006). Diagnosis of *Trichomonas vaginalis* Infection: The Sensitivities and Specificities of Microscopy, Culture and PCR assay. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, **126**:116-20.
- Roger, C., Tine, KhadineSylla, Rougyatou Ka., Lamine Dia, Doudou Sow, et al.(2019). *Journal of Parasitology Research Volume 2019 Article ID 2069672*, 8 pages <https://doi.org/10.1155/2019/2069672>
- Sam-wobo, S. O., Ajao, O. K., Adeleke, M. O. and Ekpo, U. F. (2012). Trichomoniasis among Antenatal Attendees in a Tertiary Health Facility, Abeokuta, Nigeria. *Munis Entomology and Zoology Journal*. **7** (1) : 380-384.
- Schwebke, J. R. and Burgess, D. (2004). Trichomoniasis. *Clinical Microbiology Review*; **17**, 794-803, table of contents.
- Secor, W. E., Meites, E., Starr, M. C. and Workowski, K. A. (2014). Neglected Parasitic Infections in the United States: Trichomoniasis. *American Journal of Tropical Medicine and Hygiene*. **90** (5) :800-804.
- Shuter, J., Bell, D., Graham, D., Holbrook, K. A., Bellin, E. Y.

- (1998). Rates of and Risk Factors for trichomoniasis among pregnant inmates in New York City. *Sexually Transmitted Diseases*, **25**:303-307.
- Smith, D. A. and Ramos, N. (2010). Trichomoniasis. eMedicine Specialties; Available from: <http://www.emedicine.mediscape.com>. (Last accessed on 2016 Feb 16).
- Sobel, J. D. (2005). What's new in bacterial vaginosis and trichomoniasis? *Infectious Diseases Clinical North America*, **19**:387-406.
- Soper, D. (2004). Trichomoniasis: Under control or undercontrolled? *American Journal of Obstetrics and Gynecology*, **190**:281-90.
- Sary, A., Kocha, A and Toedorowie, Z. (2000). Detection of *Trichomonas vaginalis* as prevalent of sexually transmitted diseases. *Journal of clinical microbiology*, **40**:3277-3280.
- Swygard, H., Sena, A. C., Hobbs, M. M. and Cohen, M. S. (2004). Trichomoniasis: clinical manifestations, diagnosis, and management. *Sexually Transmitted Infection*. **80** (2) : 91-95.
- Uneke, C., Ugwuoru, C, Ali, E. and Ali, M. (2005). *Trichomonas vaginalis* infection among pregnant women in South-Eastern Nigeria; The Public Health Significance. *The International Journal of Gynecology and obstetrics*, **6**:1.
- Usanga, V. L., Abia-Basse, L., Inyang-etoh, P., Udoh, S. F., Ani, E. and Archibong, E.(2009). Trichomonas vaginalis Infection Among Pregnant Women In Calabar, Cross River State, Nigeria. *The Internet Journal of Gynecology and Obstetrics*. **14**(2): 1-7.
- Woken, G. N. (2006). *Trichomonas vaginalis* infection in some parts of Niger Delta Region, Rivers State, Nigeria. *Journal of Parasitology*, **27**:68-72.