

Case report

Amyloidosis, sarcoidosis and systemic lupus erythematosus

Amel Rezgui^{1,*}, Imene Ben Hassine¹, Monia Karmani¹, Fatma Ben Fredj¹, Chadia Laouani¹

¹Internal Medicine Department, CHU Sahloul, Tunisia

*Corresponding author: Amel Rezgui, Internal Medicine Department, CHU Sahloul, Tunisia

Key words: Amyloidosis, sarcoidosis, lupus erythematosus

Received: 14/01/2016 - Accepted: 10/04/2016 - Published: 06/05/2016

Abstract

The occurrence of renal and multiple organ Amyloidosis is currently considered exceptional in the course of systemic lupus erythematosus. We report a case of a concomitant SLE and Amyloidosis in a 57 year old female patient with hypothyroidism history, who presented with erythema nodosum, fever, arthralgia and sicca syndrome. Biological findings showed an inflammatory syndrome, renal failure, proteinuria (1g / 24h), positive auto antibodies and anti DNA. Lung radiology revealed mediastinal lymphadenopathy, pleural nodules, ground glass infiltrates and pleuritis. Bronchial biopsy showed non specific inflammation. The salivary gland biopsy showed amyloid deposits. This case report reminds us that lupus and Amyloidosis association, although exceptional remains possible. The occurrence of Lofgren syndrome in this situation make the originality of this report.

Pan African Medical Journal. 2016; 24:23 doi:10.11604/pamj.2016.24.23.8853

This article is available online at: <http://www.panafrican-med-journal.com/content/article/24/23/full/>

© Amel Rezgui et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Systemic lupus erythematosus (SLE) is characterized by a large clinical polymorphism which explains the diagnostic difficulties encountered in some cases. Amyloidosis is histologically diagnosed by the presence of insoluble protein deposits in tissues. The AA amyloid form is secondary to a chronic inflammatory condition due to systemic diseases. The occurrence of renal and multiple organ amyloidosis is exceptional in the course of systemic lupus erythematosus. We report a case of concomitant diagnosis of SLE and amyloidosis.

Patient and observation

This is a 57 year old female with hypothyroidism history for 20 years who consulted for fever, arthralgia and erythema nodosum. She presented with a 10 day history of subcutaneous nodules in lower limbs, fever, and arthritis. On physical examination, there was hepatosplenomegaly, and no lymphadenopathy. On the skin examination there was hot nodules on the front and sides of the legs. On ophthalmological examination, there was keratitis. Biological findings showed an elevated erythrocyte sedimentation rate (123 mm during the first hour), a normocytic normochromic non-regenerative anemia (9 g / dl), a positive polyclonal hyper gamma globulin rate (16 g / L), an elevated C-reactive protein rate (190 mg / L), renal failure (creatinine 200 μ mol / l, Urea 16 mmol / l, Renal clearance: 28 ml / min), anti-nuclear and anti-native DNA antibodies were positive. There was also significant proteinuria (1g/day). Calcium and phosphate balance over a 3 day periods was normal. Chest X-ray showed mediastinal enlargement suggestive of mediastinal adenomegaly with calcifications (Figure 1). On chest CT there was mediastinal lymph nodes in Baretz space and bilateral hilar lymphadenopathy which were partially calcified. There were changes in the lung parenchyma described as ground glass infiltrates with pleural nodules, pleural and pericardial effusions (Figure 2). Tuberculin intradermal test was negative, mycobacterium tuberculosis wasn't present in gastric fluid. On functional respiratory tests there was distal obstructive deficit with a normal DLCO. Bronchoscopy showed macroscopically normal bronchi. Bronchial biopsies suggested non specific inflammation lesions. Renal Ultrasound revealed poorly differentiated kidneys with a small left kidney, which didn't allow the renal biopsy. Skin biopsy was performed on healthy skin and showed no lupus band. Renal biopsy was contraindicated by the findings of the renal ultrasound. Biopsy of the salivary glands showed chronic sialadenitis the special Congo Red stain revealed amyloid deposits around the vessels and interlobular ducts (Figure 3) which are birefringent amyloid deposits under polarized light (Figure 4). Accordingly, the retained diagnosis was the association of SLE, sarcoidosis (Lofgren syndrome) and amyloidosis. Our patient has already been treated. There was a complete regression of the erythema nodosum lesions under analgesics and potassium iodide after 4 weeks. There were no indications for corticosteroid therapy because kidneys were already affected, CKD stage (Chronic kidney disease) with small dedifferentiated kidneys and there were no signs of activity of sarcoidosis. The patient was prescribed Colchicine® 1mg per day and Nivaquine® 200mg per day. The patient monitoring included a chest CT, a transthoracic echocardiography (which showed regression of the pericardial effusion), creatininemia and Proteinuria. A rectal biopsy was scheduled.

Discussion

The association SLE and Sarcoidosis has been described in the literature as a non-fortuitous association [1]. The blocking of the reticuloendothelial purification system by immune complexes excess in SLE can enhance the formation sarcoid granulomas. Hepatitis C treatment with IFN- α can induce sarcoidosis and SLE [2- 4]. There is common cytokinetic channel stimulation between the two conditions. Amyloidosis is defined by the extracellular deposition of protein agglomerates all having common tinctorial affinity, a fibrillary appearance in electron microscopy, and a so-called spatial conformation β -pleated sheet [5, 6]. There are several types of amyloidosis according to the nature of the deposits of amyloid proteins [7]. AA amyloidosis is a complication of chronic inflammation (Rheumatoid arthritis, Ankylosing spondylitis, Crohn's disease, Juvenile idiopathic arthritis) or chronic infections such as tuberculosis. Amyloidosis almost never occurs as a complication of SLE [8,9]. There are a number of reasons which make LES and amyloidosis are mutually exclusive: there is low SAA protein levels during active lupus, whereas the developed amyloidosis secondary to SLE shows primarily elevated SAA protein levels (reactive amyloidosis) [10]. There have been nearly 30 reported cases that associate SLE and amyloidosis since 1956. It is most often an AA amyloidosis type. An AL amyloidosis has been described in some lupus with monoclonal gammopathy. The elapsed time from diagnosis of SLE and amyloidosis varies from 1 year to 35 years. The diagnosis was concomitant in our patient. In half of the patients, amyloidosis appears in less than five years' time. The occurrence of amyloidosis mechanisms is not very much known. High levels of SAA proteins were found in cases where they were assayed, which showed strong correlation between elevated SAA and amyloidogenesis. The diagnosis of amyloidosis was confirmed according to histopathological findings after pulmonary, renal, gastric, cardiac and cutaneous biopsies. The association Sarcoidosis and Amyloidosis has been described in the literature. Five cases were reported in the literature with renal amyloidosis. The diagnosis of sarcoidosis in our case is to be discussed. The clinical features are highly suggestive of Lofgren's Syndrome, staged II on chest X-Ray. There is no need to histological evidence as it is not mandatory

Conclusion

Although concomitant SLE and amyloidosis is an exceptional combination it remains possible. Indeed amyloidosis can complicate any chronic inflammatory condition including SLE. It can be localized or generalized. Treatment is directly based on the management of the exacerbations of SLE

Competing interests

The authors declare no competing interest.

Authors' contributions

All the authors had contributed to this work (medical staff, discussion, therapeutic decision and bibliography). All the authors of the manuscript have read and agreed to its content. The manuscript will be freely available to any scientist wishing to use them for non-commercial purposes

Figures

Figure 1: Chest X-ray showing Mediastinal enlargement

Figure 2: Mediastinal lymph nodes in Baretz space and bilateral hilar lymphadenopathy on chest CT

Figure 3: Congo red stain: amyloid deposits around the vessels and interlobular ducts

Figure 4: Irefringent amyloid deposits under polarized light

References

1. Neville E, Mills RG, Jash DK, Mackinnon DM, Carstairs LS, James DG. Sarcoidosis of the upper respiratory tract and its association with lupus pernio. *Thorax*. 1976 ;31(6):660-4. **PubMed | Google Scholar**
2. Kikawada M, Ichinose Y, Kunisawa A, Yanagisawa N, Minemura K, Kasuga I et al. Sarcoidosis induced by interferon therapy for chronic myelogenous leukaemia. *Respirology*. 1998;3(1):41-4. **PubMed | Google Scholar**
3. Denys BG, Bogaerts Y, Coenegrachts KL, De Vriese AS. Steroid resistant sarcoidosis: Is antagonism of TNFalpha the answer?. *ClinSci*. 2007;112(5) :281-9. **PubMed | Google Scholar**
4. Ziegenhagen MW, Müller-Quernheim J. The cytokine network in sarcoidosis and its clinical relevance. *J Int Med*. 2003;253(1):18-30. **PubMed | Google Scholar**
5. Guillevin Loïc, Meyer Olivier, Hachulla Éric, Sibilia Jean. *Traité des Maladies et syndromes systémiques* . 6th edition. 2016 Lavoisier SAS. **PubMed | Google Scholar**
6. Rodney Falk H, Raymond LMD, Comenzo MD, Martha Skinner MD. *The Systemic Amyloidoses*. *N Engl J Med*. 1997; 337(13):898-909. **PubMed | Google Scholar**
7. Sipe JD, Benson MD, Buxbaum JN, Ikeda S, Merlini G, Saraiva MJM et al. Amyloid fibril protein nomenclature: 2012 recommendations from the Nomenclature Committee of the International Society of Amyloidosis. *Amyloid*. 2012;19(4):167-70. **PubMed | Google Scholar**
8. Gómez-Puerta JA, Cervera R, Moll C, Solé M, Collado A, Sanmartí R. Proliferative lupus nephritis in a patient with systemic lupus erythematosus and long standing secondary amyloid nephropathy. *Clin Rheumatol*. 2009 ;28(1):95-7. **PubMed | Google Scholar**
9. Monteiro P, Abreu P, Salvador MJ, Malcata A. Secondary amyloidosis and systemic lupus erythematosus. *Acta Reumatol Port*. 2009;34(2B):400-4. **PubMed | Google Scholar**
10. Ridley MG, Maddison PJ, Tribe CR. Amyloidosis and systemic lupus erythematosus. *Ann Rheum Dis* .1984; 43(4):649-50. **PubMed | Google Scholar**




Figure 1: Chest X-ray showing Mediastinal enlargement



Figure 2: Mediastinal lymph nodes in Baretz space and bilateral hilar lymphadenopathy on chest CT




Figure 3: Congo red stain: amyloid deposits around the vessels and interlobular ducts




Figure 4: Birefringent amyloid deposits under polarized light