

**Case Report**

*Annals of Pediatric Surgery*  
Vol. 6, No 1, January 2010, PP 39-40

## Foreign body impacted in the Jejunum

*Javad Ghoroubi, Farid Imanzadeh, Ali-Akbar Sayyari, Hazhir Javaherizadeh*

*Mofid Children's Hospital, Shahid Beheshti University, M.C., Tehran, IRAN*

### ABSTRACT

*We present a case with protracted vomiting which admitted to our hospital. After radiologic investigation, patients underwent laparotomy, were a foreign body was found impacted in the jejunum.*

*Index Word: foreign body, jejunum, impaction, intestinal obstruction.*

### INTRODUCTION

The vast majority of foreign bodies pass unimpeded through the gastrointestinal tract, often without the parent having any notion of the event. Parents may find a coin or marble in the toilet or diaper and wonder as to its origin. Foreign bodies impacted in the esophagus may initially be completely asymptomatic. Much less frequently, foreign bodies impact the duodenum, terminal ileum or rectum. Foreign body may cause damage to gastrointestinal tract with several mechanisms. Solid foreign bodies, such as coins, toy parts or sharp objects, cause damage by perforation or pressure necrosis. In addition, some coins contain a high concentration of zinc, which itself is toxic to tissue<sup>1</sup>. Some medications may also cause esophageal damage; these medications may be acid (e.g. tetracycline, ferrous sulfate), alkaline (e.g. phenytoin) or have a high osmolality when dissolved. The

mucosal damage may be deep enough to produce ulceration and subsequent scarring<sup>2,3</sup>.

### CASE REPORT

A 16 months old girl was admitted to the hospital due to protracted vomiting. Nasogastric tube was placed. Vomiting was bilious. Patient was not passing stools. Abdomen was not distended. Plain X-ray was done to verify obstruction. Laparotomy was done for intestinal obstruction. A distention was seen in the first 30 cm of jejunum. In this location a foreign body was trapped. We asked again about her history from her mother. Her mother told that her girl ingested an aquarium stone about 1 weeks ago (Figure-1). The foreign body was a gelatinous stone from aquarium which increased in size while was in the bowel.




Figure-1: An aquarium stone. After ingestion(Left). Before ingestion(Right)

## DISCUSSION

Objects that reach the stomach can usually be left to pass spontaneously down the bowel. The exception would be a long or large object that can not negotiate the pylorus, duodenal C-loop or ileocecal valve<sup>4</sup>. Neumann H et al., reported removal foreign body that had remained lodged in the jejunum for 8 days with double balloon enteroscopy<sup>5</sup>. There is some reports about role of double balloon enteroscopy of foreign body removal<sup>6,7</sup>. Traumatic intramural hematoma and bezoar may be the cause of duodenal or jejunal obstruction and must be keep in mind<sup>8</sup>. Removal of foreign body by laparoscopy was also reported by some authors<sup>9</sup>. In order to prevent such complication, in every child with history of vomiting, foreign body impaction must be considered.

## REFERENCES

- O'Hara SM, Donnelly LF, Chaung E, et al. Gastric retention of zinc-based pennies: radiographic appearance and hazards. *Radiology* 213:113-117,1999.
- Kirkendall JM, Pill esophagitis, *J Clin Gastroenterol* 28:298-305, 1999.
- Jaspersen D. Drug-induced oesophageal disorders: pathogenesis, incidence, prevention and management. *Drug Saf* 22:237-249, 2000.
- Jones A. Caustic ingestion and foreign bodies: damage to the upper gastrointestinal tract. In: Wyllie R, Hyams JS(Ed). *Pediatric Gastrointestinal and Liver Disease*. 3rd ed.; Saunders 269, 2006.
- Neumann H, Fry LC, Rickes S, et al. A 'double-balloon enteroscopy worth the money': endoscopic removal of a coin lodged in the small bowel. *Dig Dis* 26:388-9, 2008.
- Shibuya T, Osada T, Asaoka D et al. Double-balloon endoscopy for treatment of long-term abdominal discomfort due to small bowel penetration by an eel bone. *Med Sci Monit* 14: CS107-9, 2008.
- Osada T, Shibuya T, Kodani T. et al. Obstructing small bowel bezoars due to an agar diet: diagnosis using double balloon enteroscopy. *Intern Med* 47:617-20, 2008.
- Chien JH, Ho TY, Shih-Peng L, et al. Acquired duodenal obstruction in children. *Pediatr Neonatol* 49:193-6,2008.
- Chin EH, Hazzan D, Herron DM, Salky B. Laparoscopic retrieval of intraabdominal foreign bodies. *Surg Endosc* 21:1457,2007.
-