

Cardiac Surgery: One year experience of cardiac surgery at Muhimbili National Hospital, Dar es Salaam- TANZANIA.

Nyawawa ETM¹, Ussiri EV¹, Chillo P¹, Waane T¹, Lugazia E¹, Mpoki U¹, Luchemba R¹, Wandwi B¹, Nyangasa B¹, Bgoya J² Mahalu W³.

¹ Muhimbili National Hospital, Tanzania, ² visiting Cardiac Anaesthesiologist, Zimbabwe, ³Visiting Cardiac surgeon Bugando University College of Health Sciences-Tanzania

Correspondence to: Dr. Evarist T.M Nyawawa, E-mail: enyawawa@gmail.com

Background: Establishing a cardiac unit in developing countries is usually difficult as it is associated with many obstacles of both expertise and financial constraints and more alarming is the mortality rate that may be high. Even after success in the initial stage sustainability of such program is a dilemma. The aim of this study was to determine pattern of disease profile, type of cardiac surgery done and the overall outcome.

Methods: All patients who underwent cardiac operation at the centre were prospectively recruited. Patients' demography and disease characteristics as demonstrated at echocardiography and its confirmation at operation were recorded. Peri-operative factors were the measurable statistics that determined the overall patients' outcome. All data were entered and analyzed using a spss11.5 window program.

Results: A total of 105 cases of cardiac surgery were done 21% were male and 79% were females. Mean age was 19.4±12.3. The majority of cases were due to Rheumatic heart diseases (47.6%), congenital heart disease (35.2%), myxomatous valvular degeneration (16.2%) and pericardial disease 1%. Mitral valve disease was the commonest cause of cardiac disease (58.1%). Prolonged duration of aortic cross-clamp and total operation time were associated with prolonged intensive care stay and poor patients' outcome respectively (p<0.05). While, ventricular dysfunction and total cardiopulmonary bypass time were not. The overall mortality rate was 13.3%. Majority of all death (64.3%) followed mitral valve repair.

Conclusion: The majority of patients (86.7%) who underwent cardiac surgery had full recovery. The mortality of (13.3%) is probably comparable to other settings. The diversity of spectrum of cardiac disease found elsewhere is also found in our community and therefore need to increase community awareness. Mitral valve repair deserve a special entity that requires skills and expertise. The mere presence of suboptimal ventricular dysfunction is probably not a contraindication to cardiac operation. The duration of aortic cross-clamp and total operation time were determinant of postoperative outcome.

Introduction

In view of the outstanding global heart disease that is expected the world population of 6.5 billion people is at risk of heart diseases¹. While the developed countries estimate a population of one billion, the majority of people (5.5 billion) are found in developing countries. With emerging economy the backlog of patients with rheumatic heart disease, congenital heart diseases are immense. There is an epidemic rise of atherosclerotic coronary heart disease due primarily notable risk factors such as hypertension, hyperlipidemia, westernized diet, smoking, inactivity and change in life style². Fifty-seven million people die each year on the planet with over 17 million from cardiovascular disease³. Non communicable diseases are clearly a greater cause of death than communicable disease and will continue to rise⁴. To the emerging countries the situation is even worse as there are few cardiac centers when comparing number of inhabitants per centre (Africa 1/33,000,000 as compared to Asia 1/16,000,000 Europe 1/1,000,000 USA 1/120,000)⁵. To worsen the

situation in the developing countries there is lack of expertise and the frequent political conflicts.

Establishing a cardiac centre especially in a developing country is usually difficult and associated with many obstacles⁶. Pezzela⁷ classified cardiac projects in developing countries as ; nonexistent but wanting to start, previously existed but failed, small or even larger existing programs now limited by financial and political consideration, ready to start but need financial and political support, already functional but needing academic support and various combinations of these. Even after starting the reputation might be lowered with an exceedingly high mortality rate that can be anticipated that may again retards the smooth running of the unit. Sustainability of such program in emerging countries is a problem again. If not well planned even the cadres of staff might be inadequate. If there good success particularly with regards to number of operation and having the staff acquired good skills and experience, moving of staff to green pastures is all what is anticipated due to low payment⁶.⁸.With exception of south Africa and Egypt a few other African countries have far made progress in open heart Surgery if not fully established yet some are doing operations in collaboration with visiting cardiac team from else where^{9, 10}. In Tanzania earliest plan to establish cardiac unit started in early 1970s but could not mature till 2005 when strong commitment by the government was taken including sending the team of staffs to train in various institutions that had composed all various cadres of staff required to make a complete cardiac team. Further more commitment was the necessity to acquire the tools that it managed successful. Since then the team was back by 2007 and appropriate measures was taken to officially start open heart surgery that by 21st may 2008 open heart surgery started and by June 2009 a total of 105 cases had been operated. And we here present our first one year outcome of open heart surgery at Muhimbili National Hospital in Dar es Salaam.

Patients and Methods

This was a prospective study that recruited all patients who underwent cardiac operation between May 2008 and June 2009. Patients' details including age, sex and duration of symptoms were taken. The diagnosis as found from echocardiography and confirmed during operation and whether there was an associated cardiac lesion was recorded; the type of operation whether done on pump or off pump was coded and recorded. The pulmonary pressure was also determined during echocardiography and quantified to whether there was pulmonary hypertension or not. The ejection fraction combined with fraction of shortening were used to assess ventricular function simultaneously the diastolic function was assessed and whether cardiac dysfunction was present or not were recorded. During operation the time of aortic cross-clamp that was categorized into short if it was less or equal to 60 minutes and prolonged when more than 60minutes, total duration of conduct of bypass that was categorized into short if less or equal to 60 minutes, moderate when 61-100minutes and prolonged when >100minutes. Operation time that was counted from skin incision to its closure was recorded and categorized into short to average time if the procedure lasted to less or equal to 4 hours and prolonged when it was more than that. Patient was followed into the ICU while closely monitored for any complication that develops. Further noted was the total duration of intensive care stay. The duration of ICU stay was categorized as early stay if it was five days or less and prolonged when it was more. The duration of hospital stay known as post intensive care hospital stay was noted at the time of discharge of the patient from the hospital. This duration was categorized into short to normal when the patient stayed for less than or equal to two weeks and prolonged when it was more. Further noted were any complication the patient develops while in the ward. The final disposition of the patient as to

whether recovered fully, partially or died was also noted. Data were entered into a master data sheet and then into SPSS 11.5 window program and analyzed accordingly.

Results

There were 105 cardiac cases of which 22(21%) were males and 83(79%) were females making 1:4 male to female ratio. The mean age was 19.4±12.3 years, range 2-52 years. The mean duration of operation was 3.1±1.0 hours ;(range 1-5.5 hours). The mean duration of aortic cross-clamp was 64.4±25.4 minutes; (range 12-176 minutes) for 72 cases. The mean duration of cardio bypass was 94.0±33.6 minutes; (range 40-240 minutes) for 72 cases. The mean duration of intensive care stay was 4.5±2.1 days; (range 2-20 days). And mean duration of post ICU hospital stay 13.7±6.6 days ;(range 5-50 days) for 94 cases. Females (79%) had more cardiac disease as compared to males (21%). Majority of patients occurred in the ages between 7-16 years. (Table 1)

Table1. Age-sex Distribution

Age Group(years)	Male	Female	Total (%)
2-6	3(30)	7(70)	10(9.5)
7-11	4(19)	17(81)	21(20.0)
12-16	8(30.8)	18(69.2)	26(24.8)
17-21	3(23)	10(77)	13(12.4)
22-26	2(22)	7(78)	9(8.5)
27-31	2(25)	6(75)	8(7.7)
32-36	0(0)	6(100)	6(5.7)
37-41	0(0)	4(100)	4(3.8)
42-46	0(0)	3(100)	3(2.8)
47-51	0(0)	3(100)	3(2.9)
52-56	0(0)	2(100)	2(1.9)
Total	22(21)	83(79)	105(100)




Figure 1. Number of Operations per month and Year

Table 2. Pattern of group of cardiac disease

Cardiac disease	frequency	percentage	Valid percentage	Cumulative percentage
CHD	37	35.2	35.2	35.5
RHD	50	47.6	47.6	82.9
Myxomatous	17	16.2	16.2	99.0
Pericardial	1	1.0	1.0	100.0
Total	105	100.0	100.0	

Table 3. Distribution of Cardiac Diseases

Diagnosis	Frequency	Percentage	Valid percentage	Cumulative Percentage
PDA	18	17.1	17.1	17.1
ASD	6	5.7	5.7	22.9
VSD	5	4.8	4.8	27.6
Single atrium	2	1.9	1.9	29.5
Pulmonary stenosis	4	3.8	3.8	33.3
AV_Canal	2	1.9	1.9	35.2
MR	34	32.4	32.4**	67.6
TR	1	1.0	1.0	68.6
MS	27	25.7	25.7**	94.3
AR	5	4.8	4.8	99.0
Pericardial effusion	1	1.0	1.0	100.0
Total	105	100.0	100.0	




Figure 2. Cardiac Diseases Occurring in Combination

Table 4. Presence of Ventricular Dysfunction and Patients' Outcome


Perioperative factors		Total	Postoperative outcome			p-value
			Full recovered	Partial recovered	Died	
Ventricular dysfunction(105cases)	Yes	23(21.9)	19(82.6)	0(0.0)	4(17.4)	0.3 $\chi^2=2.4$ df=2,
	No	82(78.1)	65(79.3)	7(8.5)	10(12.2)	
Ischaemia time(72cases)	Short	41(56.9)	34(82.9)	4(9.8)	3(7.3)	0.0 $\chi^2=7.4$ df 2,
	Prolonged	31(43.1)	19(61.3)	2(6.5)	10(32.3)	

Table 5. Total operation time, ventricular dysfunction and postoperative complications

*= Total operation time, §= ventricular dysfunction, CCF= congestive cardiac failure, Techn.diff= technically

Perioperative factors		Total	Postoperative complications							p-Value	
			None	CCF	Techn.diff	PVS	LOS	Wound inf.	ARDS		Par'lysis
TOT*	Short	89(84.8)	65(73.0)	8(9.0)	2(2.2)	5(5.6)	3(3.4)	4(4.5)	2(2.0)	0(0.0)	0.04 $\chi^2=14.4$ df 7
	Prolonged	16(15.2)	9(56.3)	0(0.0)	0(0.0)	1(6.3)	3(18.8)	1(6.3)	1(6.0)	1(1.0)	
Ventr.dysf§	Yes	23(21.9)	18(78.3)	1(4.3)	0(0.0)	2(8.7)	2(8.7)	0(0.0)	0(0.0)	0(0.0)	0.7 $\chi^2=4.7$ df=7,
	No	82(78.1)	56(68.3)	7(8.5)	2(2.4)	4(4.9)	4(4.9)	5(6.1)	3(3.7)	1(0.0)	

difficult, PVS= paroxysmal ventricular systoles, Wound inf= wound infection, ARDS= Acute respiratory distress syndrome, Para'lysis= paralysis



Majority of patients had recovered fully and were discharged(86.7%)

Mitral valve repair accounted for more death 9(64.3%) of all deaths

Figure 3. Morbidity/Mortality with Type of Operation

Table 6. Total operation time and duration of ICU Stay

Total operation time	Duration of ICU stay			Total (%)
	Died in theatre	Short	Prolonged	
Short	0(0)	78(87.6)	11(12.4)	89(84.8)
Prolonged	2(12.5)	10(62.5)	4(25.0)	16(15.2)
Total	2(1.9)	88(83.8)	15(14.3)	105(100)

$\chi^2=13.7$; df 2, p=0.001

Table 7. Outcome of Management

Outcome	Number(Percentage)	Number of Deaths (%)
Discharged home	91 (86.7)	
Died in theatre	2 (1.9)	2(14.3)
Died in ICU	9 (8.6)	9(64.3)
Died in ward	3 (2.9)	3(21.4)
Total	105 (100)	14 13.3)

Operations started in may 2008 and we progressed to December (red bars) (Figure 1). Rheumatic heart diseases accounted for the large number of cases that were operated at the unit (47.6%) (Table 2)

Mitral valve disease accounted for the majority of cardiac diseases (58.1%) (Table 3). Majority of cardiac diseases occurred with other associated cardiac lesions (Figure 2). There was statistical significant between duration of aortic-cross clamp and postoperative outcome (p<0.05). There was no statistical significant difference between patients with ventricular dysfunction and patient outcome (p>0.05). (Table 4). There was significant statistical difference between total duration of operation and emergency of complications (p<0.05) There was no statistical significant between ventricular dysfunction and postoperative complications (p>0.05) (Table 5). There was significant statistical difference between total operation time and duration of intensive care stay (p<0.01) (Table 6)

When comparing morbidity/mortality, majority of patients had full recovery and the mortality was probably within acceptable limits in a newly established cardiac centre (Figure 3). The overall mortality rate was 13.3% with 64.3% of all deaths occurring in the ICU. Nine (64.3%) of the deaths followed mitral valve repair, 3 patients died following mitral valve replacement, 1 patient died following VSD closure and one patient died following PDA ligation.(Table 7).

Discussion

The study showed that with initial setting of a cardiac unit in our centre majority of patients (86.7%) had fully recovery and were discharged. However the mortality of 13.3% is probably comparable to other settings¹¹. The spectrum of cardiac disease found elsewhere is also found in our community⁸. Majority of our patients had Rheumatic valvular heart disease (47.6% Table 2) this high incidence noted in this study as compared to other cardiac lesions goes in parallel with the high incidence of the disease to any African country that is the result of the burden of rheumatic fever attributed to combination of lack of resources, infrastructure, political and economic instability, poverty, overcrowding and malnutrition¹². Further the

highest incidence of rheumatic heart disease is found in sub-Saharan Africa with a prevalence of 5.7 per 1000 as compared to 1.8 per 1000 in North Africa and 0.3 per 1000 in economically developed countries with established market economy^{13, 14}. Since the majority of cases were mitral valvular disease and in particular mitral regurgitation and patients who succumbed to death followed mitral valve repair it implies that mitral valve repair requires more skills and expertise. Nkomo VT, portrays that problem to valve repair or replacement for rheumatic heart disease with or without infective endocarditic in young economically disadvantaged patients are numerous¹². The fulminant course of rheumatic fever, rheumatic heart disease and infective endocarditis coupled with delayed patient presentation results in a high incidence of heart failure at presentation¹⁵, associated with high in-hospital and late mortality even with surgical intervention^{16,17,18}. Our patient had long duration of symptoms of mean 32±14 months (range 18.2-46.3) in rheumatic heart diseases and at presentation 80% were in NYHA class III and IV. Despite staying in the ward for 3 weeks to 3 months in some, for optimization of medication before surgery was undertaken, it was possible that some portions myocardial tissue could have undergone remodeling. The mere presence of suboptimal ventricular dysfunction is probably not a contraindication to cardiac operation.

The duration of aortic cross-clamp and total operation time were determinant of postoperative outcome. This finding was similar to other series where ischemia time and total operation time was found to be determinant of early extubation and postoperative complications for prolonged operations^{19, 20}. However this study could not demonstrate any association with the total pump time. Garcia-Montes et al also could not find any similarity with total pump time while determining factors associated with prolonged mechanical ventilation in paediatric patients²¹.

Conclusion

Majority of patients who underwent cardiac surgery had full recovery. The diversity of spectrum of cardiac disease found elsewhere is also found in our community and therefore need to increase community awareness. Mitral valve repair deserve a special entity that requires skills and expertise. The mere presence of suboptimal ventricular dysfunction is probably not a contraindication to cardiac operation. The duration of aortic cross-clamp and total operation time were determinant of postoperative outcome.

Acknowledgement

We thank the government through Ministry of Health and Social welfare for having realized the need of its people and training the cardiac team. We also thank the administration of Muhimbili National Hospital for having supported the unit throughout the period since its establishment. We thank members of the task force who have played a big role since its early plans and for close monitoring of the project.

References

1. World health report 2000, [http:// www. who.int/whr](http://www.who.int/whr)
2. Pezzela AT. Open heart Surgery in a developing country. *Asian Cardiovascular and Thoracic Ann.* 2006; **14**: 355-356
3. Pezzela AT, International Cardiac Surgery: A global perspective seminars in Thoracic and Cardiovascular Surgery. *Asian Journal of cardiovascular and thoracic Ann* 2002; **14**: 298-320
4. Fuster V, Voute J. Comment: MDGS: Chronic diseases are not on the Agenda. *Lancet* 2005; **366**: 1512-1514

5. Vladmir V, Michel M, Marek B, Gregory K, Xaverier M, Peter N. The development of cardiac surgery in an emerging country. *Texas Heart Inst J.* 2008;**35(3)**: 301-306
6. Ghosh p. Setting up an open heart surgical program in a developing country. *Asian Cardiovascular Thorac Ann.* 2005;**13(3)**: 299-301
7. Pezzella A T. Global Expansion of Cardiac Surgery in the New Millennium. *Asian Cardiovasc Thoracic Ann* 2001; **9**: 253-256
8. John CE and Ndubueze E. Open heart surgery in Nigeria indications and challenges. *Texas Heart Inst J.* 2007;**34**: 8-10
9. National department of health annual report 2008/2009. www.doh.gov.za
10. Akomea-Agyin C, Galukande M, Mwambu T, TtendoS, and Clarke I. pioneer human open Heart Surgery using cardiopulmonary by pass in Uganda. *African Health Sciences*, 2008; **8(4)**: 259-260
11. Luis AL, Kathy JJ, Kimberlel G. Improvement in Congenital Heart Surgery in a Developing Country: The Guatemalan Experience, *Circulation* 2007; **116**: 1882-1887.
12. Nkomo VT. Global burden of cardiovascular disease: epidemiology and prevention of valvular heart diseases and infective endocarditis in Africa. *Heart* 2007; **93**: 1510-1519
13. World Health organization. The current evidence for the burden of group A streptococcal diseases. <http://www.who.int/child-adolescent>
14. Carapetis JR, Steer AC, Mulholland EK et al. The global burden of group A streptococcal diseases. *Lancet Inf Dis* 2005; **5**: 685-694
15. Amoah AG, Kallen C. Aetiology of heart failure as seen from a National Cardiac Referral Centre in Africa. *Cardiology* 2000; **93**: 11-18
16. Touze JE, Ouattra K, Coulibaly AO et al. Infectious endocarditis surgically treated during the acute phase. 26 cases. *Presse med* 1986;**15**: 787-790
17. Louw JW, Kinsley RH, Dion RA et al . Emergency heart valve replacement: an analysis of 170 patients. *Ann Thorac Surg* 1980; **29**: 415-422
18. Fradi I, Drissa MA Cheour M, et al Retrospective study on 100 cases of infective endocarditis, Rabta University Hospital, Tunis. *Arch Mal Coeur Vaiss* 2005; **98**: 966-971
19. Steve D, Sarah W, Roger BB, Mee MB, Harrison. Factors associated with early extubation after cardiac surgery in young children. *PCCM* 2004; **5(5)**: 63-68.
20. Varro M, Gombocz K, Wrana G. Factors influencing early extubation after open heart surgery. *Orv Hetil* 2001; **142(23)**: 1217-1220.
21. Garcia-Montes JA, Calderon-Colmenero J, Casanova M, Zarco E, Fernandez de la Requera G, Buendia A. Risk factors for prolonged mechanical ventilation after surgical repair of congenital heart disease. *Arch. cardiol. Mex* 2005; **75(40)**: 402-407