



## CROWN RATIO ASSESSMENT FOR *PROSOPIS AFRICANA* (GUILLAND PERR.) TAUB SPECIES IN MAKURDI, NIGERIA

Chukwu, O.\*<sup>1</sup>, <sup>2</sup>Dau, J.H., Ogundipe, O. C.<sup>1</sup> and Wali, M. D.<sup>3</sup>

<sup>1</sup>Department of Forest Resources Management, University of Ibadan, Ibadan, Nigeria

<sup>2</sup>Department of Forest Production and Products, Federal University of Agriculture Makurdi, Benue State Nigeria

<sup>3</sup>Department of Forestry and Wildlife, University of Maiduguri, Maiduguri, Nigeria

\*Corresponding author: [onye20042000@yahoo.com](mailto:onye20042000@yahoo.com), +2348032633835

### ABSTRACT

*Information on tree crown is prerequisite for sustainable utilization of forest tree resources, as it is the site for physiological activities that lead to tree growth and development. Crown ratio (CR) is an index of tree stability; it indicates tree vigour and is a useful parameter for assessing forest condition. This study aimed at assessing crown ratio for improved silvicultural management of naturally grown Prosopis africana species within the University of Agriculture Makurdi, Nigeria. Nine temporary sample plots (TSPs) of size 100 m x 100 m were established using simple random sampling method. Diameter at breast height (Dbh), total height (Ht) and crown diameter (Cd) and height to live crown base (HCB) were measured for all living P. africana trees with Dbh > 10 cm within the 9 TSPs. Data collected were subjected to descriptive and bivariate correlation analysis. The trees were further classified based on their Crown ratio as high vigour (CR>0.50), moderate vigour (CR= 0.30-0.50) and low vigour (CR<0.30). The result revealed that CR had strong negative correlation with Ht and HCB. Tree diameter attributes showed low positive correlation with CR. However, about 69% of the trees are within moderate CR class and only 3% are of the low CR class. This implies that majority of the P. africana species in the study area are of moderate and high vigour. Hence, appropriate silvicultural Treatment such as thinning is recommended for trees with CR <0.3 to ensure moderate, better stand stability and sustainability of the species benefits.*

**Keywords:** Correlation; Crown ratio; Economic trees; Tree growth characteristics; *Prosopis africana*

### INTRODUCTION

Forests are large areas of land with varieties of tree, shrub and herb. Forest benefits to the environment and humankind, particularly the rural people who live close to the forest are enormous. These benefits include timber and non-timber products, such as fruits, pods, medicine, gum, resins, construction materials, nuts and seeds, tools, decorations, cosmetics, cultural artefacts and other needs (Oboho and Ogana, 2011). Hence, the vigour of these economic trees needs to be checked and known to ensure sustainability of its benefits.

*Prosopis africana* (Guill. and Perr.) Taub is one of the economic tree species that has both

timber and non-timber benefits. *P. africana* belongs to the family Fabaceae, subfamily Mimosoideae. The common names include; Iron wood, Ubwa (Igbo), Kiriya (Hausa), Okpehe (Idoma), Kpaaye (Tiv) and Ayan in Yoruba language (Agboola, 2004). *P. africana* is one of the only known species of its genus found in Africa; the species occurring from Senegal to Ethiopia in the zone between the Sahel and savannah forests. Abahet *al.* (2015), reported that almost all parts of the tree are medicinal and of high economic value to rural communities. The leaves and pods of *P. africana* are used for fodder and the seeds as seasoning for food (Laouali, 2016). It is used for the

planks, mortars and pestles due to its resistant to wood borers and high wood density (SoteloMontes and Weber, 2009; Dau and Chenge, 2016). The tree species is highly valued for charcoal by blacksmiths due to the high calorific value of its wood (SoteloMontes *et al.*, 2011). Furthermore, the leaves, roots and especially the bark are used in traditional medicine. According to Laouali (2016), *P. africana* is over exploited due to its numerous values, to ensure its sustainability, management practices that will aid its natural regeneration and promoting their domestication is required to reduce the pressure on natural populations.

Often times, the impact of most natural or anthropogenic stress on trees, are evident on the crown as signs of deterioration may first be observed in the tree crowns. Thus, measurement of a tree crown is often used to assist in the quantification of tree growth (Popoola and Adesoye, 2012). However, tree crown is a significant feature of stand structure; it is the centre of physiological activity, particularly gas exchange, which drives essential living processes such as photosynthesis, growth and development (Leites and Robinson, 2004). Tree crown contains the foliage, network of branches and the anchor of fruits which are source of food for animals and man. Hence, understanding crown characteristics and status, will aid sustainable utilization of its benefits.

Previous studies have identified crown ratio as an indicator of tree's vigour, wood quality and an indicator of a tree's competitiveness in a stand (Sprinz and Burkhart 1987; Temesgen *et al.*, 2005). Crown ratio is the ratio of live crown length to tree height. In other words, the percentage of crown length from the base of the live crown to the tree tip to total tree height. It is often used as an important predictor variable for tree growth and yield equation. Hence, an important input variable to estimate the mortality of individual trees and also display changes in appearance of stand over-time (Adesoye and Oluwadare, 2008).

However, despite its aforementioned importance, assessment of crown ratio is one of the most difficult and mostly imprecisely measured tasks in forestry; as it is a function of tree height and crown diameter. Furthermore, the use of crown as an index of tree vigour is highly important, as larger ratios are indicative of healthier, faster growing trees (Schomaker *et al.*, 2007). Hence, this

will aid proper decision-making. Therefore, this study was aimed at assessing crown ratio of *Prosopis africana* species found in a natural forest within the University of Agriculture, Makurdi, for sustainable and silvicultural management of the species and as baseline information for risk management.

## MATERIAL AND METHODS

### Study Area

The study was carried out in at the northern part of University of Agriculture Makurdi, Benue State, Nigeria. It is located on longitude 8° 21' E to 8° 39' E and latitude 7° 33' N to 7° 52' N in Benue State, within the southern guinea savannah ecological zone of Nigeria and covers a total land area of 7,978 km<sup>2</sup>. The topography of the study area is characterized by gentle hills. The soil is mainly sandy-loamy; the climate is characterized by distinct rainy and dry seasons. The annual rainfall ranges between 1016 mm to 1524 mm spreading over May to October. The climate of the area is tropical sub-humid with high temperatures and high humidity; the average maximum and minimum daily temperature of 35°C and 21°C in wet season, and 38°C and 16°C in dry season. The study area is bounded at the North-East by Guma Local Government Area and by River Benue in the South (Dau and Chenge, 2016).

### Sampling Procedure

Data used for this study was collected from nine (9) randomly selected Temporary Sample Plots (TSPs) of size 100 m × 100 m within the study area. Within each TSP, total enumeration living *Prosopis africana* trees with Dbh ≥ 10.0 cm in the selected plots were made. Diameter at the base (Db), middle (Dm), top (Dt) and at breast (Dbh), total (Ht) and merchantable height (Mt), height to live crown base (HCB), crown diameter (Cd), crown length (CL), basal area (BA) were measured from a total number of three hundred and twenty-three (323) *P. africana* trees found in all the nine selected plots.

### Measurement of Tree Growth Variables

#### Tree diameter

Stem diameter for all tree individuals with Dbh ≥ 10.0 cm were measured. The point of the measurement was recorded from the uphill sides of the trees and on the inside of the lean for leaning

trees. Furthermore, trees with deformations at 1.3 m (Dbh) and 0.3 m (stump diameter), measurements were made at the sound points on the stem above the abnormality. Also, for buttressed trees, a point of measurement was selected approximately 0.5 m above the convergence of the buttress (Huschet *al.*, 1982). Diameter measurements of trees were recorded using diameter tape graduated in centimeters. During the measurement, loose bark, climbers and epiphytes were lifted above the measuring tape. Spiegel relaskop was used to measure diameters at the top and middle of the trees.

**Crown diameter**

The tree crown diameter measurements were based on the assumption that the vertical projection of a tree crown is circular. Four radii were measured and, in the direction, forming equal angles (Zuhaidi, 2009).

The average crown diameter (Cd) was then calculated as such:

$$Cd = \sum \frac{r_i}{2} \dots\dots\dots (1)$$

Where; Cd = average crown diameter,  $r_i$  = projected crown radii measured on four axis and  $i=1, 2, 3$  and  $4$ .

**Height measurement**

Total height and height to base of the live crown (HCB) was measured using Spiegel relaskop. HCB was determined by identifying the point along the stem where the lowest live branch is attached to the main stem as indicated by Jiang *et al.*, (2007). Hence, height from the ground to the tip of the tree was measured as the total height.

**Data Analysis**

Individual tree derived variables were computed from the measured growth variables

**Basal area**

Basal Area for each tree was computed using:

$$BA = \frac{\pi(Dbh)^2}{4} \dots\dots\dots (2)$$

Where: BA = Basal Area (m<sup>2</sup>),  $\pi$  = Pi is constant (3.143) and Dbh = Diameter at breast height (cm)

**Volume**

Tree volume was estimated using the Newton-Simpson’s formula (Akindele, 2005) expressed as:

$$V = \frac{\pi H}{24}(Db^2 + 4Dm^2 + Dt^2) \dots\dots\dots (3) \quad 22$$

Where; V = Total volume (m<sup>3</sup>), H = total height (m),  $\pi$  = Pi is constant (3.143) Db = Diameter at the base (cm), Dm= Diameter at mid-point (cm) and Dt = Diameter at the top (cm).

**Descriptive statistics**

Descriptive and inferential statistics were used in this study. The tree growth variables were described using measures of central tendency and measures of dispersion.

**Bivariate correlation analysis**

Pearson’s product-moment correlation coefficient was used to examine the linear relationship between crown ratio and other tree variables

The correlation coefficient was computed as:

$$r = \frac{\sum XY - \frac{(\sum X)(\sum Y)}{N}}{\sqrt{[\sum X^2 - \frac{(\sum X)^2}{N}][\sum Y^2 - \frac{(\sum Y)^2}{N}]}} \dots\dots\dots (4)$$

Where, r= correlation coefficient, X = variable (1) to be compared, Y =variable (2) to be compared, N = total numbers of observations or trees measured.

**Crown ratio**

Crown ratios of individual trees were computed using the mathematical expression below:

$$CR = \frac{CL}{Ht} \dots\dots\dots (5)$$

Where CL= crown length (m) and Ht= total height (m)

**Crown ratio classification**

Crown ratio of *P. africana* species found in the study area was computed and classified into three vigour classes modified from Schütz, (2001):

- i. Low vigour (CR<0.3)
- ii. Moderate vigour (CR 0.3-0.5)
- iii. High vigour (CR> 0.5)

Frequencies of the observed numbers of tree in each class was calculated

## RESULTS

The entire data obtained from the field inventory were carefully collated and analysed to detect and represent fundamental growth patterns. The dataset comprises of tree growth variables measured from *Prosopis africana* species found

within the study area. A total of 323 tree: measured and summary statistics of the dataset used in this study are presented in Table 1. The diameter at breast height ranged from 17.27- 63.82 cm, total height ranged from 4.0 to 13 m, crown length ranged from 1.2 to 6.4m from and crown ratio ranged from 0.2 to 0.8 m.

**Table 1: Summary statistics table for growth characteristics variables.**

Variables	Descriptive Statistics		
	Minimum	Maximum	Mean $\pm$ SE
Ds (cm)	21.303	70.180	32.165 $\pm$ 0.557
Dbh (cm)	17.271	63.820	29.241 $\pm$ 0.506
Dm (cm)	12.108	51.167	24.367 $\pm$ 0.422
Dt (cm)	10.678	33.478	25.427 $\pm$ 0.440
Cd (m)	3.070	10.800	6.320 $\pm$ 0.090
Ht (m)	4.050	13.400	7.327 $\pm$ 0.092
BA (m <sup>2</sup> )	0.019	0.320	0.074 $\pm$ 0.003
V (m <sup>3</sup> )	0.095	3.388	0.446 $\pm$ 0.020
HCB (m)	1.060	9.433	4.081 $\pm$ 0.081
CL (m)	1.237	6.408	3.246 $\pm$ 0.031
CR	0.216	0.806	0.457 $\pm$ 0.005
HDR	8.558	62.526	6.771 $\pm$ 0.455

Where: Ds= stump diameter, Dbh= diameter at breast height, Dm= diameter at the middle, Dbh= diameter at the top, Ht= total height, HCB= height to live crown base, CD= crown diameter, HDR = height to diameter ratio, CL= crown length, CR= crown ratio, BA=Basal area, V = Total volume and SE = standard error. Number of trees = 323

The result of Pearson's product-moment correlation analysis between the crown ratio (CR) and other measured and derived tree growth variables was shown in Table 2. Crown ratio was significant and shows strong negatively correlation with HCB and Ht. However, the result revealed low and positive correlation with tree stem diameter (Ds and Dbh), Basal area, crown diameter, crown length and total volume (non-significant). The graphical


relationships of CR versus Ht; HCB; HDR and CR versus Dbh were presented in Figures 1-4. It can be seen that crown ratio of *Prosopis africana* trees in the study area are essentially linearly (negative) related with total height, height to live crown base and height diameter ratio. Hence, crown ratio displayed a positively linear relationship with diameter at breast height.

**Table 2: Results of correlation analysis of Tree Size Variables with CR**


Variables	Correlation								
	Ds	Dbh	Cd	Ht	BA	V	HCB	CL	HDR
CR	0.27*	0.27*	0.05*	-0.63*	0.28*	0.04 <sup>ns</sup>	-0.84*	0.29*	-0.71*

\* =Correlation is significant and ns=Correlation is not significant at 0.05 level (2-tailed).Ds= stump diameter (cm), Dbh= diameter at breast height (cm), Ht= total height (m), HCB= height to live crown base (m), Cd= crown diameter (m), HDR


= height diameter ratio, CL= crown length (m), CR= crown ratio, BA=Basal area (m<sup>2</sup>), V = Total volume (m<sup>3</sup>). 1 of trees = 323 24




**Figure 1: Relationship between crown ratio and total height**



**Figure 2: Relationship between crown ratio and height to live crown base**




**Figure 3: Relationship between crown ratio and height-diameter ratio**




**Figure 4: Relationship between crown ratio and diameter at breast height**

The result of the tree crown ratio (CR) categorization (Figures 5 and 6) for the pooled data in the study area revealed that about 224 trees/ha (69%) trees in the study area are between 0.3-0.5

(Moderate), about 91 trees/ha (28%) trees are of High CR (>0.5) and 8 trees/ha (3%) fell under low CR (<0.3)



**Figure 5: Chart showing number of trees in crown ratio classes**



**Figure 6: Chart showing percentage of trees in crown ratio classes**

## DISCUSSION

In this study, information on tree growth characteristics were collected and processed. The pooled summary of tree growth characteristics for the *P. africana* species found in a natural forest within the University of Agriculture Makurdi, Nigeria. However, the study revealed significant variations across Ds, Dbh, Cd, Ht, BA, V, CL, HDR and HCB. The low associations or relationship among tree growth attribute of Ds, Dbh, Cd, BA, V, and CL indicates low impact of these tree attribute on the CR. This implies that there is less effect of horizontal increment of growth variables on tree vigour. This is in disagreement with the report of Popoola and Adesoye (2012) that correlated crown ratio with tree for *Tectona grandis* in Osho forest, Oyo State, Nigeria; that crown ratio decreased with increasing tree size. The results of the bivariate correlation between tree growth attributes and crown ratio of *P. africana* species in the study area indicated inverse associations with total height and HCB. This implies that CR decrease with increase in tree height. This result is in congruent with the report of Temesgenet *al.* (2005), that tall and slender trees have lower CR.

The result of classification of the crown ratio of *P. africana* species in the study area reveals

that the larger percentages (69 and 28%) of the species are within the moderate and high vigour classes. Based on the CR classification considered in this study, the overall result shows the trees within the study area were vigorous, thereby; indicates that *P. africana* trees in the study are in moderate condition. This might be attributed to the location of the trees (within a University environment). Hence, there was little pressure and /or less exploitation on crown resources of the trees species investigated, because the University environment has some level of protection.

## CONCLUSION

The assessment of tree growth attributes is the fundamental and most important aspect of estimating and managing trees in the forest stand. Findings from this study showed that, there is a strong negative correlation between the tree crown ratio and stemheight. Crown ratio shows significant variation with most tree growth variables in the study area. Majority of the *P. Africana* trees in the study area have moderate vigour. Therefore, an appropriate silvicultural treatment such as thinning is recommended for trees with CR <0.3 to ensure moderate, better stand stability and sustainability of the species benefits.

## REFERENCES

- Abah, J. O., Agunu, A., Ibrahim, G., Halilu, M. E. and Abubakar, M. S. (2015). Development of quality standards of *Prosopis africana* (Guill. and Perr.) Taub. stem bark. *J. BiolAgric Healthcare*. 6: 10-17.
- Adesoye, P. O. and Oluwadare, A. O. (2008). Interim Crown Ratio Models for a mixed *Tectona grandis* and *Gmelina arborea* stand in the University of Ibadan, Nigeria. *Resource Journal of Forest*. 2 (1): 34-42.
- Agboola, D. A. (2004). *Prosopis africana* (Mimosaceae): stem, roots, and seeds in the economy of the savannah areas of Nigeria. *J Econ Bot*. 58, Suppl.: S34-S42.
- Akindele, S. O. (2005). Volume Functions for Common Timber Species of Nigeria's Tropical Rain Forests, a Technical Document: International Tropical Timber Organization (ITTO). pp. 1-85.
- Dau, J. H. and Chenge, B. I. (2016). Growth space requirements models for *Prosopis africana* (Guill& Perr) Taub tree species in Makurdi, Nigeria. *European Journal of Biological Research*, 6(3): 209-217.
- Husch, B., Miller C.I. and Beers, T. W. (1982). *Forest Measurement*. John Wiley & Sons, New York, USA.
- Jiang, L., Brooks, J. R. and Hobbs G. R. (2007). Using Crown Ratio in Yellow-poplar Compatible Taper and Volume Equations. *Northern Journal of Applied Forestry*. 24 (4): 271 275p.
- Laouali, A., Boubé, M., Tougiani, A. and Ali, M. (2016). Analysis of the structure and diversity of *Prosopis africana* (G. et Perr.) Taub. tree stands in the southeastern Niger. *J Plant Stud*. 5(1): 58-67.
- Leites, L. and Robinson, A. (2004). Improving taper equations of loblolly pine with crown

- dimensions in a mixed-effects modelling framework. *Forest Science*. 50(2): 204-212.
- Oboho, E.G. and Ogana, F.N. (2011). Effects of varying water soaking duration on the germination of *Garciniakola*(Heckel) Seeds. *Nigerian Journal of Agriculture, Food and Environment*.7 (2):57-62.
- Popoola, F. S. and Adesoye, P. O. (2012). Crown ratio models for *Tectona grandis* (Linn. f.) stands in Osho Forest Reserve, Oyo State, Nigeria. *Journal of Forest Science*. 28(2): 63-67.
- Schomaker, M. E., Zarnoch, S. J., Bechtold, W. A., Latelle, D. J., Burkman, W. G. and Cox, S. M. (2007). Crown-condition classification: a guide to data collection and analysis. Gen. Tech. Rep. SRS-102. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 78 p.
- Schütz, J.-Ph. (2001). Der Plenterwald und weitere Formenstrukturierter und gemischter Wälder. [Theselection forest and other forms of structured and mixed forests]. Parey Buchverlag. Berlin. 207p.
- Sotelo Montes, C. and Weber, J. C. (2009). Genetic variation in wood density and correlations with tree growth in *Prosopis africana* from Burkina Faso and Niger. *Ann. For. Sci.* 66: 713.
- Sotelo Montes, C., Silva, D. A., Garcia, R. A., Muñiz, G. I. B. and Weber, J. C. (2011). Calorific value of *Prosopis africana* and *Balanitesaegyptiacawood*: relationships with tree growth, wood density and rainfall gradients in the West African Sahel. *Biomass Bioenerg.* 35: 346-353.
- Sprinz, P. T. and Burkhart, H. E. (1987). Relationships between tree crown, stem, and stand characteristics in unthinned Loblolly Pine plantations. *Canadian Journal of Forest Research*. 17: 534–538.
- Temesgen, H., LeMay, V. and Mitchell, S. J. (2005). Tree crown ratio models for multi-species and multi-layered stands of South-eastern British Columbia. *The Forestry Chronicle*. 81(1):133-141.
- Zuhaidi, Y.A., (2009). Local growth model in modeling the crown diameter of plantation-grown *Dryobalanops aromatic*. *Journal of Tropical Forest Science*. 21(1): 66-71.