

International Journal of Basic, Applied and Innovative Research

ISSN: 2315 - 5388

IJBAIR, 2015, 4(4): 108 - 117

www.arpjournals.com; www.antrescentpub.com

E-ISSN: 2384 - 681X

REVIEW PAPER

A REVIEW ON THE PHARMACOLOGY AND PHYTOCHEMISTRY OF FOLKLORE MEDICINAL PLANT *HYPTIS SUAVEOLENS* (L.) POIT

Devi Priya, M.

Department of Botany, St. Thomas College, Rammi, Pathanamthitta, Kerala- 689 673, India

*Corresponding author: devi.priya.m@gmail.comReceived: 20th November, 2015Accepted: 20th December, 2015Published: 31st December, 2015

ABSTRACT

Hyptis suaveolens (L.) Poit. (Bush mint) belongs to the family Lamiaceae and commonly known as 'wilayati tulsi' in India. It is an invasive weed famous for its insecticidal property. Traditionally it is used for the treatment of respiratory tract infections, infection of uterus and skin diseases. Phytochemicals like steroids, alkaloids, phenolics, flavonoids, tannins, glycosides and essential oil isolated from the plant has medicinal and pharmaceutical importance. Extensive literature survey was made to identify the established research works on this magnificent plant and compiled it in a nutshell. The present review article is mainly focused on pharmacological and other important aspects of *H. suaveolens*.

Key-words: *Hyptis suaveolens*, essential oil, leaves, extract, phytochemicals

INTRODUCTION

Hyptis suaveolens (L.) Poit is a folklore medicinal plant used to heal wounds and treat gastrointestinal disorders, respiratory tract and uterus infections and skin diseases (Fig. 1). It is commonly found in dense clumps on roadsides, degraded moist and dry deciduous forest, waste lands and over-grazed pasture. It is a weedy alien species native to tropical America, now naturalized in the tropics and during its establishment, it upsets the recruitment pattern in the nearby occupied. The plant is commonly known as 'wilayati tulsi' because of its unique resemblance to 'tulsi leaves'. The other vernacular names are jungli tulsi, beejabandh (Hindi), ganga/gandha thulasi (Oriya), tokma, ganja tulsi, bilati tulsi, kusmai (Bengali) bhusri, dimbubha, gusumpuru (Bihari), konda thulasi, adavi tulasi (Telugu), bhustrena (Sanskrit), poothachedaiyan (Tamil) manjpatre (Kannada) naatta poochedi (Malayalam) and american mint, bush mint or the chan plant in English.

The plant is a sweet smelling woody herb with tetragonal hispid stem, leaves long petioled, pubescent, broadly elliptic-ovate, base oblique, truncate or acute, margin coarsely serrulate, apex acute, chartaceous. (Kumar *et al.*, 2005) Flowers blue in short stalked axillary cymes, calyx ovoid, campanulate and subequally 5 lobed, often accrescent. Corolla is bilabiate with lower deflexed and seccate lip and upper rounded. Stamens didynamous, anther cells confluent, glandular ovary 4 partite, gynobasic style with a shortly bifid stigma. Nut lets ovoid, smooth or rugulose (Gamble, 1921).





Fig. 1: Habit of *Hyptis suaveolens* (L.) Poit.

TRADITIONAL USE

Hyptis is used for ethnobotanical applications in rural communities in Africa (Edeoga *et al.*, 2006). Traditionally the decoction of the roots is used as appetizer and the root is chewed with betel nuts as a stomachic; leaf sap with lemon juice added is taken in Sierra Leone for stomach ache and the leaf is applied around the head for head ache or topically to maturate boils (Dalzier, 1937). A leaf poultice is applied to cancers and tumours in America (Hartwell, 1969). The plant is used by various tribal communities of Maharashtra to cure various diseases like, parasitical cutaneous, diseases, infection of uterus, and as a sudorific in catarrhal condition. The plant is stimulant carminative, antispasmodic, antirheumatic and antisporific baths. It is also used for headache, stomach-ache and snuff to stop bleeding of the nose (Nadkarni, 2005). It is a medicinal antidote to poison (Jain, 1991). In Guinea Bissau, periodic burning of leaves and flower (Palsson and Jaenson, 1999) and in Kenya, the overnight burning of leaves show repellence against mosquitoes (Seyoum *et al.*, 2002). The tribal people use the plant to repel mosquitoes in Tamil Nadu (Singh *et al.*, 2005). In Nigeria the plant used for treating respiratory tract infections, colds, pain, fever, cramps and skin diseases. Seeds extract is a remedy for menorrhagia, leucorrhoea and temporary male sterility (Prashantkumar and Vidyasagar, 2006). Seeds soaked in water are used as soothing agent by Bheel tribe (Jain *et al.*, 2010) (Ekka, 2011). In the Songkhla province, the whole plant decoction is used to cure fever and fatigue and the seeds are used to cure constipation and as anti-diarrhoeal (Neamsuvan *et al.*, 2012). Mal Paharia tribes use the root as a blood purifier and leaves as an effective wound healer (Kumar and Abbas, 2012).

CHEMICAL CONSTITUENTS

Photochemical offers by plants have many high value chemicals useful as drugs, cosmetics, bio-polymers etc which are highly beneficial to human kind. Preliminary phytochemical screening of *Hyptis* shows the presence of many photochemical like steroids, alkaloids, phenolics, flavonoids, tannins, glycosides (Ijeh *et al.*, 2007; Pachkore and Dhale, 2011^a), juglones (Jelani and Prabhakar, 1991) and saponins. The antioxidant and anti-inflammatory activity of flavonoids, anti-diarrhoeal and stomachic activity of tannins, analgesic, anti-pyretic and anti-coagulant properties of



glycosides were well established. The essential oil of aerial parts of *H. suaveolens* is predominated by β -caryophyllene (Fig. 2), 4-terpinenol, α -terpineol, α -terpinolene, α -phellandrene, α -copaene, α -thujene, α -humulene, α -bergamotene, β -pinene, β -pynene, β -elemene, γ -terpinene, γ -cadinene, δ -cadinene, linalool, fenchol, eugenol, camphene, sabinene, myrcene, 1, 8-cineole, aromadendrene, benzyl benzoate and a diterpene hydrocarbon II etc. Eucaliptol is the most abundant component in the oil in leaves, followed for γ -elemene, β -pynene, (+)-3-carene, trans- β -caryophyllene and germacrene. These essential oil can be applied in the aspergillus treatment (Peerzada, 2007) dehydroabietinol (Azevedo *et al.*, 2001), limonene, bicyclogermacrene, β -phellandrene (Ziegler *et al.*, 2002), monoterpenes, α -pinene and p-cymene (McNeil *et al.*, 2011).

Natural triterpenoid, characterized as 3β -hydroxylup-20(29)-en-27-oic acid (Triguna *et al.*, 1983), 3β -hydroxylup-12-en-28-oic acid and 3β -hydroxy lup-12-en-28-oic acid, α - and β -amyryn (Misra *et al.*, 1983). The plant derived triterpenoids possess various pharmacological properties. A large number of triterpenoids exhibits cytotoxicity against a variety of tumor cells as well as anticancer efficacy in preclinical animal models. A triterpenoid similar to betulin and betulinic acid was also identified (Sharma *et al.*, 2010). Betulin and its derivatives however possess biological effects like anti-inflammatory, antiviral, anti-HIV, hepatoprotective. The oil of stems contains dierpene hydrocarbon II and palustrol (Tonzibo *et al.*, 2009), hyptadienic acid, suaveolic acid, suaveolol, methyl suaveolate, β -sitosterol, oleanolic acid, ursolic acid, rosmarinic acid (Manchand *et al.*, 1974). Suaveolol and methyl suaveolate are responsible for anti-inflammatory activity of the plant and hence used to treat dermatological diseases. Ursolic acid occur in the form of an aglycone of triterpene saponins inhibits the inflammatory enzymes cyclooxygenase and lipoxygenase. Rosmarinic acid, a polyphenol, possesses anti-oxidant property. (2E)-1-(2-hydroxy phenyl) pent-2-en-1-one (I) and 1-[(3-hydroxy-5, 5-dimethyl cyclohex-3-en-1yl) oxy] hexan-3-one (II) (Jayakumar *et al.*, 2005), 3β -hydroxylup-12-en-28-oic acid, 3β -hydroxylup-20(29)-en-27-oic acid (Satish *et al.*, 2010) etc were isolated from various parts.




Fig. 2: Major photochemical found in *Hyptis suaveolens*

PHARMACOLOGY

The plant has antiinflammatory, antinociceptive (Santos *et al.*, 2007), antiplasmodial (Chukwujekwu *et al.*, 2005), antirheumatic, anticonvulsant (Akah and Nwambie, 1993), antiulcerogenic, carminative and lactagogue properties. It is used in catarrhal and uterine affections, parasitical cutaneous diseases, epistaxis etc (Khare, 2007). The plant is also used in nausea and infection of the gall bladder (Ahmed *et al.*, 1994). It can reduce calcium oxalate crystallization (Kumkum and Ranjana, 2012), which is comparable to that of proprietary drugs for dissolving kidney stones. The plant cures skin diseases and bronchial disorders (Koche *et al.*, 2010). Natural HIV-integrase inhibitor (Chatterjee and Pakrashi, 1997) and trypsin inhibitor (Aguirre *et al.*, 2004) were also isolated from this plant.

Antibacterial Activity: Methanolic leaf extract is active against *X. campestris* and is more effective than Kanamycin and Neomycin (De Britto and Gracelin, 2011). It shows inhibitory action on *S. aureus*, *E. coli* and *P. aeruginosa*. The volatile oil is toxic to human pathogenic bacteria (Iwu *et al.*, 1990), *B. cereus* (Asekun *et al.*, 1999), *Serratia marcescens* and *K. pneumoneae* (Pachkore *et al.*, 2011^b).

Antidiabetic Activity: The methanol extract of leaves posses anti-diabetic activity in alloxan-induced diabetic rats. The presence of alkaloids, carbohydrates, flavonoids, tannins, steroids and/or terpenes in the leaves either as single constituents or in combination may be responsible for the anti-diabetic activity (Danmalam *et al.*, 2009).

Antifungal Activity: The essential oil extracted from the leaves has antifungal and antibacterial properties (Singh *et al.*, 1992) (Pandey *et al.*, 1982) (Nantitanon *et al.*, 2007). These are active against *S. cerevisiae*, *Mucor* sp. and *F. moniliforme* (Malele *et al.*, 2003) and *F. oxysporum* (Tripathi *et al.*, 2009). Anti-aspergillus property against *A. fumigates*, *A. parasiticus* and *A. flavus* (Moreira *et al.*, 2010) is well established. The plant shows activity against *Rhizoctonia solani*, *Sclerotium rolfsii* and *Sclerotinia sclerotiorum* and the oil in combination with *Trichoderma harzianum* controlled wilt and rot diseases by *S. sclerotiorum* (Singh and Handique, 1999). Growth inhibition also noticed in *A. niger*, *C. albicans*, *Cryptococcus* and *Fursarium* species in some instances the inhibition exceeded Griseofulvin (Mbatbou *et al.*, 2010). The aqueous and ethanol extracts show inhibition of *C. albicans*, *Collectrotrichum capsici*, *F. oxysporum* *F. sp. Lycopersici* (Pachkore *et al.*, 2011).

Anti-Inflammatory Activity: *H. suaveolens* contains tannins which are used as antiinflammatory agents and also used topically for treatment of burns. Suaveolol and methyl suaveolate inhibits of croton oil-induced dermatitis of the mouse ear due to topical anti-inflammatory activity which is only two to three times lower than that of the reference drug Indomethacin (Grassi *et al.*, 2006).

Anti-Oxidant Activity: The healing properties exhibited by the ethanolic extract (Shirwaikar *et al.*, 2003) and the protective properties of free soluble polyphenols (Oboh, 2008) on Fe (II) induced lipid peroxidation in isolated Rat's brain are with a supportive role of antioxidant enzymes. It is identified that the antioxidant activity is due to the presence of flavonoids (Gavani and Paarakh, 2008).

Anti-Tumor Activity: The plant is also known for its anti-tumor and anti-cancer activities. The leaf is a potent anticancer agent in traditional medicine (Kingston *et al.*, 1979). Recently 1, 3-Propanediamine N (3- Aminopropyl)-N-Methyl isolated from *Hyptis suaveolens* has proved its anticancer potentials in Ehrlich Ascites Carcionoma Cell Line (Gurunagarajan and Pemaiah, 2011).

Gastro Protective Activity: Suaveolol obtained from ethanolic and aqueous extracts have gastro protective activity (Prabhat *et al.*, 2009) (Vera-Arzave *et al.*, 2012).

Hepatoprotective Activity: The leaf aqueous extract protects liver tissues against oxidative damages and acetaminophen induced toxicity (Babalola *et al.*, 2011).

Toxicity: The extract of leaves exhibits low acute toxicity (Santos *et al.*, 2007). The effect of water extract in Wistar rats reveals the trivial effect of the extract at given doses in rats (Attawish *et al.*, 2005).



Wound Healing Activity: Ethanolic, petroleum ether, alcoholic and chloroform water extract of leaves showed wound healing activity (Shenoy *et al.*, 2009). The activity may be due to free radical scavenging action of the plant and enhancing level of antioxidant enzymes in granuloma tissue (Shirwaikar *et al.*, 2003).

Insecticidal and Larvicidal Activity: The phytochemicals from *H. suaveolens* can act as larvicide, insect growth regulators, and repellent and ovipositor attractant. Bush mint produce a very strong aromatic mint/thyme-like smell when the leaves are crushed lead to use of the plant as a powerful insect repellent (Oparaeke *et al.*, 2002) and the fumes of dried leaves are used to repel mosquitoes and control insect pests of stored grains (Alok *et al.*, 2010; Okigbo *et al.*, 2010). It is active against cowpea weevil, *Callosobruchus maculatus* (Jayakumar and Ganesh, 2012) and cow pea borer, *Maruca testulalis*. The aqueous extract with a lower dose of insecticides can control cotton bollworms and is active against stem borer *Sesamia calamistis* (Adda *et al.*, 2011). Hyptis extracts shows mortality against *Sitophilus zeamais* and *Callosobruchus maculatus* (Iloba and Ekrakene, 2006). It can also control insects and nematodes (Oyedunmade, 1998). Dehydroabietinol revealed an *in vitro* antiplasmodial effect against *Plasmodium falciparum* (Ziegler *et al.*, 2002). Essential oil is as effective for personal protection against mosquito bites (Abagli and Alavo, 2011). The hexane leaf extracts showed larvicidal, and ovidal activity against *A. aegypti*, *An. stephensi* and *C. quinquefasciatus* (Arivoli and Tennyson, 2011; Raja *et al.*, 2012). The ethyl acetate extracts acts against *Helicoverpa armigera* and *Spodoptera litura* (Ivoke *et al.*, 2009; Musa *et al.*, 2009). The aqueous and ethanolic extract show reduced the viability in *Anopheles gambiae* (Kala *et al.*, 2006). Its application is required to control *Trogoderma granarium* and the methanol extract of seed can be a substitute to synthetic insecticides (Bernard *et al.*, 2001). The ticks of *Hyalomma* sp., *Rhipicephalus* sp. and *Haemophysalis* sp. were found to be highly susceptible to the steam distillate of *Hyptis* (Kapoor, 2011).

Other Important Activities: The plant is reported to be a potent antihelminthic and antimalaric (Chukwujekwu *et al.*, 2005). An anti-phospholipase A2 activity was observed for the aqueous extract (Berbard *et al.*, 2001). The essential oil from leaves has antifertility properties (Peerzada, 2007; Shirwaikar *et al.*, 2003). The leaf residues can be used as potent bio-herbicide to control the spread of *Parthenium* (Kapoor, 2011).

CONCLUSION:

Though *Hyptis suaveolens* is an invasive obnoxious weed, the literature survey reveals the therapeutic efficiency of the plant. The phytochemicals isolated from this medicinal has been effectively using in many health problems since a long time. The present review work provides a wide area of interest for planning and conducting research on this wonderful plant to develop novel drug molecules for the future.

REFERENCES

Abagli, A.Z. and Alavo, T.B.C. (2011). Essential oil from bush mint, *Hyptis suaveolens*, is as effective as DEET for personal protection against Mosquito bites. *The Open Entomol J*; 5:45-8.

Adda, C., Atachi, P., Hell, K. and Tamò, M. (2011). Potential use of the bushmint, *Hyptis suaveolens*, for the control of infestation by the pink stalk borer, *Sesamia calamistis* on maize in southern Benin, West Africa. *J Insect Sci*; 11:1-13.

Aguirre, C., Valdés-Rodríguez, S., Mendoza-Hernández, G., Rojo-Domínguez, A. and Blanco-Labra A. (2004). A novel 8.7 kDa protease inhibitor from chan seeds (*Hyptis suaveolens* L.) inhibits proteases from the larger grain borer *Prostephanus truncatus* (Coleoptera: Bostrichidae). *Comp Biochem Physiol Biochem Mol Biol*; 4 (138):81-89.

Ahmed, M., Scora, R.W. and Ting I.P (1994). Composition of leaf oil of *Hyptis suaveolens* (L.) Poit. *J Essen Oil Res*; 6: 571-575.

Akah, P.A. and Nwambie, A.I. (1993). Nigerian plants with anticonvulsant properties. *Fitoterapia*; 64:42-44.



Alok, S., Sanjay, J., Monika, S., Alok, M., Padmini, S. and Prabodh S. (2010). *In-vitro* evaluation of antioxidant activity of leaves of *Hyptis suaveolens* (L.) Poit. *Int J Pharm Res*; 1: 86-93.

Arivoli, S. and Tennyson, S. (2011). Mosquitocidal activity of *Hyptis suaveolens* (L.) Poit. (Lamiaceae) extracts against *Aedes aegypti*, *Anopheles stephensi* and *Culex quinquefasciatus* (Diptera: Culicidae). *Int J Recent Scientific Res*; 2:143-149.

Asekun, O.T., Ekundayo, O. and Adeniyi B.A. (1999). Antimicrobial activity of the essential oil of *Hyptis suaveolens* leaves. *Fitoterapia*; 70:440-442.

Attawish, A., Chivapat, S., Chavalittumrong, P., Phadungpat, S., Bansiddhi, J. and Chaorai B. (2005). Chronic toxicity study of *Hyptis suaveolens* (L.) Poit. in rats. *Songklanakarin J Sci Technol*; 27:27-36.

Azevedo, N.R., Campos, I.F., Ferreira, H.D., Portes, T.A., Santos, S.C., Seraphin, J.C., *et al.* (2001). Chemical variability in the essential oil of *H. suaveolens*. *Phytochem*; 57: 733-736.

Babalola, O.O., Ojo, O.E., Oloyede, F.A. (2011). Hepatoprotective activity of aqueous extract of the leaves of *Hyptis suaveolens* (L.) Poit. on acetaminophen induced hepatotoxicity in Rabbits. *Res J Chem Sci*; 1:85-88.

Bernard, P., Scior, T., Didier, B., Hibert, M. and Berthon, J.Y. (2001). Ethnopharmacology and bioinformatic combination for leads discovery: application to phospholipase A (2) inhibitors. *Phytochem*; 58: 865-874.

Chatterjee, A. and Pakrashi, S.C. (1997). The Treatise on Indian Medicinal Plants. Vol. 5, PID, New Delhi; 1997.

Chukwujekwu, J.C., Smith, P., Coombes, P.H., Mulholland, D.A. and Van Staden, J. (2005). Antiplasmodial diterpenoid from the leaves of *Hyptis suaveolens*. *J Ethnopharmacol*; 102: 295-297.

Dalzier, J.M. (1937). The useful plants of Tropical West Africa. Crown Overseas Agents Colonies, London.

Danmalam, U.H., Abdullahi, L.M., Agunu, A. and Musa, K.Y. (2009). Acute toxicity studies and hypoglycemic activity of the methanol extract of the leaves of *Hyptis suaveolens* Poit. (Lamiaceae). *Nig J Pharm Sci*; 8: 87-92.

De Britto, A.J. and Gracelin, D.H.S. (2011). Phytochemical screening and antibacterial activity of a few medicinal plants against *Xanthomonas campestris*. *Pharmacologyonline*; 2:271-277.

Edeoga, H.O., Omosun G. and Uche L.C. (2006). Chemical composition of *Hyptis suaveolens* and *Ocimum gratissimum* hybrids from Nigeria. *Afr J Biotech*; 5:892-5.

Ekka, A. (2011). Folklore claims of some medicinal plants used by tribal community of Chhattisgarh, India. *Res J Biol*; 1:16-20.

Gamble, J.S. (1921). Flora of Presidency of Madras. Vol 2, Adlard & Son Ltd., London.

Gavani, U. and Paarakh, P.M. (2008). Antioxidant activity of *Hyptis suaveolens* Poit. *Int J Pharmacol*; 4:227-229.

Grassi, P., Reyes, T.S.U., Sosa, S., Tubaro, A., Hofer, O. and Zitterl-Eglseer, K. (2006). Anti-inflammatory activity of two diterpenes of *Hyptis suaveolens* from El Salvador. *Z. Naturforsch*; 61:165-70.

Gurunagarajan, S. and Pemaiah, B. (2011). Comparative studies on cytotoxic effect of *Hyptis suaveolens* Poit. and *Leonotis nepeatefolia* R.Br. against EAC cell lines. *J Pharm Res*; 4: 222-4.

Hartwell, J.L. (1969). Plants used against Cancer: A survey. *Lloydia*; 32: 153-205.



Ijeh, I.I., Edeoga, H.O., Jimoh, M.A. and Ejeke C. (2007). Preliminary phytochemical, nutritional and toxicological studies of leaves and stems of *Hyptis suaveolens*. *Res J Pharmacol*; 1:34-6.

Iloba, B.N. and Ekrakene, T. (2006). Comparative assessment of Insecticidal effect of *Azadirachta indica*, *Hyptis suaveolens* and *Ocimum gratissimum* on *Sitophilus zeamais* and *Calosobruchus maculatus*. *J Biol Sci*; 6: 626-30.

Ivoke, N., Okafor, F.C., Owoicho L.O. (2009). Evaluation of ovicidal and larvicidal effects of leaf extracts of *Hyptis suaveolens* (L.) Poit (Lamiaceae) against *Anopheles gambiae* (Diptera: Anophelidae) complex. *Animal Res Int*; 6:1072-1076.

Iwu, M.M., Ezeugwu, C.O., Okunji C.O., Sanson D.R. and Tempesta M.S. (1990). Antimicrobial activity and terpenoids of the essential oil of *Hyptis suaveolens*. *Int J Crude Drug Res*; 28:73-6.

Jain, K.A., Vairale M.G. and Singh R. (2010). Folklore claims on some medicinal plants used by Bheel tribe of Guna district Madhya Pradesh. *Ind J Traditional Knowledge*; 9:105-7.

Jain, S.K. (1991). Dictionary of Indian folk medicine and Ethnobotany. Deep Publications, New Delhi.

Jayakumar, S.V. and Ganesh, S.M.K. (2012). New enolic type bioactive constituents from *Hyptis suaveolens* (L.) Poit. *Asian J Plant Sci & Res*; 2:403-8.

Jayakumar, M., Elumalai K., Jeyasankar A., Raja N. and Ignacimuthu, S. (2005). Biological activity of *Hyptis suaveolens* Poit (Lamiaceae) and *Melochia chorcorifolia* L. (Sterculiaceae) on cowpea weevil, *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae). *J Entomol Res*; 29:265-269.

Jelani, S. and Prabhakar, M. (1991). Pharmacognostic study of leaf of *Hyptis suaveolens*L. *Ancient Sci Life*; 11:31-37.

Kala, K.B., Praneeth, R.N., Kishore, K., Gopala, R.A. and Reddy, K.S. (2006). Evaluation of *Hyptis suaveolens* hydrodistillation an acaricide. *Vet Parasitol*, 2006; 20: 99-100.

Kapoor, R.T. (2011). Bioherbicidal Potential of Leaf-residue of *Hyptis suaveolens* on the Growth and Physiological Parameters of *Parthenium hysterophorus* L. *Curr Res J Biol Sci*; 3: 341-50.

Khare, C.P. (2007). Indian Medicinal Plants: An Illustrated Dictionary. Springer.

Kingston, D.G., Rao, M.M. and Zucker, W.V. (1979). Plant anticancer agents IX constituents of *Hyptis tomentosa*. *J Nat Prod*; 42: 496-499.

Koche, D., Shirsat, R., Imran S. and Bhadange D.G. Phytochemical screening of eight traditionally used ethnomedicinal plants from Akola district (MS) India. *Int J Pharm Bio Sci*; 1:253-256.

Kumar, K. and Abbas S.G. (2012). Ethnomedicinal composition depends on floristic composition: A case studied in Sal forests of Jharkhand. *Int J Pharm & Life Sci*; 3:1710-1719.

Kumar, N.A., Sivadasan, M. and Ravi, N. (2005). Flora of Pathanamthitta. Daya Publishing House, Delhi.

Kumkum, A. and Ranjana V. (2012). Inhibition of calcium oxalate crystallization *in-vitro* by various extracts of *Hyptis suaveolens* (L.) Poit. *Int Res J Pharm*; 3:261-264.

Malele, R.S., Mutayabarwa, C.K., Mwangi, J.W., Thoithi G.N., Lopez, A.G., Lucini, E.I. and Zygadlo, J.A. (2003). Essential oil of *Hyptis suaveolens* (L.) Poit from Tanzania: composition and antifungal activity. *J Essent Oil Res*; 15:438-440.



Manchand, P.S., White, J.D., Fayos, J. and Clardy J. (1974). Chemical constituents of *Hyptis suaveolens*. *J Org Chem*; 39:2306-8.

Mbatabou, V.C., Abdullatif, S. and Glover R. (2010). Phytochemical screening of solvent extracts from *Hyptis suaveolens* for fungal growth inhibition. *Pak J Nut*; 9:358-361.

Mc Neil, M., Facey, P. and Porter R. (2011). Essential oils from the *Hyptis* genus-a review (1909-2009). *Nat Prod Commun*; 6:1775-1796.

Misra, T.N., Singh, R.S. and Upadhyay, J. (1983). Triterpenoids from *Hyptis suaveolens* roots. *Phytochem*; 22:603-605.

Moreira, A.C.P., Lima, E.O., Wanderley, P.A., Carmo, E.S. and de Souza, E.L. Chemical composition and antifungal activity of *Hyptis suaveolens* (L.) Poit. leaves essential oil against *Aspergillus* species. *Braz J Microbiol*; 41:28-33.

Musa, A.K., Dike M.C. and Onu, I. (2009). Evaluation of Nitta (*Hyptis suaveolens* Poit.) seed and leaf extracts and seed powder for the control of *Trogoderma granarium* Everts (Coleoptera: Dermestidae) in stored Groundnut. *American- Eurasian J Agronomy*; 2:176-179.

Nadkarni, K.M. (2005). Indian plants and drugs with their medical properties and uses. Srishti Book Distributers, Delhi.

Nantitanon, W., Chowwanapoonpohn, S. and Okonogi, S. (2007). Antioxidant and antimicrobial activities of *Hyptis suaveolens* essential oil. *Sci Pharm*; 75:35-46.

Neamsuvan, O., Singdam, P., Yingcharoen, K. and Sengnon, N. (2012). A survey of medicinal plants in mangrove and beach forests from sating Phra Peninsula, Songkhla Province, Thailand. *J Med Plants Res*; 6:2421-2437.

Oboh, G. (2008). Polyphenol extracts from *Hyptis suaveolens* leaves inhibit Fe²⁺-induced lipid peroxidation in brain. *Int J Biomed Pharm Sci*; 2:41-46.

Okigbo, R.N., Okeke, J.J. and Madu, N.C. (2010). Larvicidal effects of *Azadirachta indica*, *Ocimum gratissimum* and *Hyptis suaveolens* against mosquito larvae. *J Agri Tech*; 6:703-719.

Oparaeke, A.M., Dike, M.C. and Onu L. (2002). Control of *Callosobruchus maculatus* (F.) on stored cowpea with African curry (*Ocimum gratissimum* L.) and African bush tea (*Hyptis suaveolens* Poit) leaf powders. *Nig J Entomol*; 19:99-108.

Oyedunmade, E.E.A. (1998). Control of nematode pests of cowpea with Mocap(Ethoprop), leaf residues of neem (*Azadirachta indica*), rattle weed (*Crotalaria retusa*) and nitta (*Hyptis suaveolens*). *Centrepoin Sci Edn*; 8:57-63.

Pachkore, G.L. and Dhale, D.A. (2011a). Pharmacognostic evaluation of *Hyptis Suaveolens* Poit. Lamiaceae. *J Ecobiotech*; 3:5-10.

Pachkore, G.L., Dhale, D.A. and Dharasurkar, A.N. (2011b). Antimicrobial and phytochemical screening of *Hyptis suaveolens* (L.) Poit. Lamiaceae. *Internat Multidiscipl Res J*; 1:01-03.

Palsson, K. and Jaenson T.G. (1999). Plant products used as Mosquito repellents in Guinea Bissau, W. Africa. *Acta Trop*; 72:39-52.

Pandey, D.K., Tripathi, N.N., Tripathi, R.D. and Dixit, S.N. (1982). Fungitoxic and phytotoxic properties of the essential oil of *Hyptis suaveolens*. *J Plant Diseases and Protection*; 89:344-349.



- Peerzada, N. (2007). Chemical composition of the essential oil of *Hyptis Suaveolens*. *Molecules*; 2: 165-168.
- Prabhat, K.D., Sabuj, S., Ranjan, S., Praveen, S.N., Shweta, N. and Anirudha, J. (2009). Phytochemical and pharmacological investigation of the protective effect of plant *Hyptis suaveolens* against duodenal ulceration. *J Glob Pharm Technol*; 1:82-87.
- Prashantkumar, P. and Vidyasagar, G.M. (2006). Documentation of traditional knowledge on medicinal plants of Bidar district, Karnataka. *Ind J Traditional Knowledge*; 5:295-299.
- Raja, N., Jeyasankar, A., Venkatesan, S.J. and Ignacimuthu S. (2005). Efficiency of *Hyptis suaveolens* against Lepidopteran pests. *Curr Sci*; 88: 221-222.
- Santos, T.C., Marques, M.S., Menezes, I.A., Dias, K.S. and Silva, A.B. (2007). Antinociceptive effect and acute toxicity of the *Hyptis suaveolens* leaves aqueous extract on mice. *Fitoterapia*, 2007; 78:333-6.
- Satish, V., Ravichandran, V.D., Gavani, U. and Paarakh, P.M. (2010). Antimicrobial studies on the extracts of *Cocculus hirsutus* Linn. and *Hyptis suaveolens* Poit. *Ind J Nat Prod and Resour*; 1:49-52.
- Seyoum, A., Kabiru, E.W., Lwande, W., Killeen, G.F., Hassanali, A., Knols, B.G. (2002). Repellency of live potted plants against *Anopheles gambiae* from human baits in semi-field experimental huts. *Am J Trop Med Hyg.*; 67:191-195.
- Sharma, P.P., Roy, R.K., Anurag, B., Gupta, D. (2010). Pentacyclic Triterpinoids from *Betula utilis* and *Hyptis suaveolens*. *Ind J PharmTech Res*; 2:1558-1562.
- Shenoy, C., Patil, M.B. and Kumar, R. (2009). Wound Healing Activity of *Hyptis suaveolens* (L.) Poit (Lamiaceae). *Ind J Pharm Tech Res*; 1:737-744.
- Shirwaikar, A., Shenoy, R., Udupa, A.L., Udupa, S.L. and Shetty, S. (2003). Wound healing property of ethanolic extract of leaves of *Hyptis suaveolens* with supportive role of antioxidant enzymes. *Ind J Expt Biol*; 41:238-241.
- Singh, G., Upadhyay, R.K. and Rao, GP. (1992). Fungitoxic activity of volatile oils of *Hyptis suaveolens*. *Fitoterapia*; 63:462-465.
- Singh, H.B. and Handique, A.K. (1999). Antifungal activity of the essential oil of *Hyptis suaveolens* and its efficacy in biocontrol measures in combination with *Trichoderma harzianum*. *J Essen Oil Res*; 9: 683-687.
- Singh, R.S., Uvarani, M., Mohanmarugaraja, M.K., Sureshkumar, K. and Shivakumar K.K. (2005). Some less known medicinal plants traditionally used in Dharmapuri District- Tamilnadu. *Ancient Sci Life*; 4:1-4.
- Tonzibo, Z.F., Florence, A.B., Bédi, G. and Chalchat, J.C. (2009). Chemical Composition of Essential Oil of *Hyptis Suaveolensis*(L) Poit. from Côte d'Ivoire. *Eur J Sci Res*, 2009; 38:565-571.
- Triguna, N.M., Singh, R.S. and Upadhyay, J. (1983). A natural triterpene acid from *Hyptis suaveolens*. *Phytochem*; 22:557-558.
- Tripathi, A., Sharma, N. and Sharma, V. (2009). *In vitro* efficacy of *Hyptis suaveolens* L. (Poit) essential oil on growth and morphogenesis of *Fusarium oxysporum* f.sp. gladioli (Massey) Snyder & Hansen. *WJMB*; 25:503-512.
- Vera-Arzave, C., Antonio, L.C., Arrieta, J., Cruz-Hernández, G., Velázquez-Méndez, A.M., Reyes-Ramírez, A, et al. (2012). Gastroprotection of Suaveolol, Isolated from *Hyptis suaveolens*, against Ethanol-Induced Gastric Lesions in Wistar Rats: Role of Prostaglandins, Nitric Oxide and Sulfhydryls. *Molecules*; 17:8917-8927.



Ziegler, H.L., Jensen, T.H., Christensen, J., Staerk, D., Hagerstrand, H., Sittie, A.A., *et al.* (2002). Possible artefacts in the *in vitro* determination of antimalarial activity of natural products that incorporate into lipid bilayer: apparent antiplasmodial activity of dehydroabietinol, a constituent of *H. suaveolens*. *Planta Med*; 68:547-549.

