

Tef (*Eragrostis tef*) Variety 'Tesfa'

Worku Kebede¹, Kidist Tolossa¹, Nigussu Hussein¹, Tsion Fikre¹, Yazachew Genet¹,
Abate Bekele¹, Kidu Gebremeskel², Atinkut Fentahun³, Habte Jifar⁴, Sonia Plaza-
Wüthrich⁵, Regula Blösch⁵, Solomon Chanyalew¹, Kebebew Assefa¹, and Zerihun Tadele⁵

¹Ethiopian Agricultural Research Institute, DebreZeit Agricultural Research Center, P.O. Box 32, DebreZeit, Ethiopia; ²Tigray Agricultural Research Institute, Axsum Agricultural Research Center, P.O. Box 230 Axum, Ethiopia; ³Amhara Regional Agricultural Research Institute, Adet Agricultural Research Center, P.O. Box 8, Bahir Dar, Ethiopia; ⁴Ethiopian Institute of Agricultural Research, Holetta Agricultural Research Center, P.O. Box 31 Holetta, Ethiopia; ⁵University of Bern, Institute of Plant Sciences, Altenbergrain 21, CH-3013 Bern, Switzerland

አህፅሮት

ጤፍ በኢትዮጵያ ከሚመረቱ የብርዕና አገዳ ሰብሎች ዋናው ሲሆን በየዓመቱ ቁጥር ከ6.5 ሚሊዮን የማያንስ አርሶ አደር ያመርተዋል። ይህም አጠቃላይ በብርዕና አገዳ ሰብሎች ከሚሸፈነው ማሳ 30% ድርሻ አለው። ይሁን እንጂ ከሌሎች ሰብሎች ጋር ሲነፃፀር ምርታማነቱ አነስተኛ ነው። ለዚህም ምክንያቱ በከፊል አርሶ አደሩ ያልተሻሻሉ የአካባቢ ዝርያዎችን በመጠቀሙና፣ የሰብሉ ተፈጥሯዊ የመጋሽብ ባህሪ ናቸው። የዚህ ጥናት ዓላማ በተለየ ምርምር አሰራር የተገኘን የጤፍ ዝርያን በመፈተሽ የተሻለ ምርት፣ የአገዳ ጥንካሬን አንዲሁም የዘር ቀለም ያለውና በአርሶ አደሩ ተመራጭ ዝርያ ማፍለቅ ነበር። በጥናቱ በቅርቡ የተለቀቀ አንድ ዝርያና አንድ የአካባቢ ዝርያን ጨምሮ 10 የተለያዩ የጤፍ አይነቶችን በማካተት በስድስት ወካይ ጤፍ አብቃይ ቦታዎች ላይ ተፈትሸው ተስፋ (ደዙ-ክርስ-457) ተብሎ የተሰየመውና የተለያዩ የጤፍ አይነቶች ተዳቅለው የተገኘው ዝርያ ከሌሎች ማወዳደሪያ ተፈተሽ ዝርያዎች የተሻለ ውጤት በማስመዘገቡ በብሄራዊ የዝርያ አፅዳቂ ኮሚቴ ተገምግሞ ለምርት እንዲለቀቅ ተወስኗል። ይህ ዝርያ ከሌሎች ዝርያዎች በንፅፅር መጋሽብን በመቋቋሙ፣ የተሻለ ምርት በመስጠቱ በአርሶአደሩ ተፈላጊ ከመሆኑም በሻገር ከዝርያው ባህሪ የተነሳ ለመስኖ እርሻና በሰብል መድረሻ ጊዜ የማጨጃ የእርሻ መሳሪያ መጠቀም ያስችላል።

Abstract

Tef [*Eragrostis tef* (Zucc.) Trotter] is extensively cultivated cereal crop in Ethiopia where it is annually grown by about 6.5 million smallholder farmers on about 30% of the total area allocated to cereal crops. However, the productivity of tef is very low compared to other cereals mainly due to lack of high yielding and lodging tolerant cultivars. The objective of this study was to evaluate the performance of the recently released tef variety called Tesfa and to provide unique morphological and agronomic descriptions of this new variety. Ten genotypes which were derived from three independent crosses and bred for at least eight generations plus a local and standard check varieties were tested at six tef growing sites in Ethiopia (namely, Adet, Chafe Donsa, DebreZeit black soil, DebreZeit light soil, Ginchiand Holetta) using Randomized Complete Block Design with three replications. The study found that Tesfa (DZ-Cr-457 RIL-181) which was obtained from a cross between kinde (a semi-dwarf mutant line developed at the University of Bern, Switzerland) and Kay Murri (a landrace) outperformed other genotypes and approved for release by the Ethiopian National Variety Release Committee in March 2017. The main advantages of Tesfa over other tested lines were its higher grain yield and higher lodging tolerance. In addition, Tesfa possesses unique properties for which high acceptance by growers is expected. These are its compact panicle, lack of shattering, and thick and strong culm which makes it potentially cultivated under irrigation and also mechanically harvested.

Introduction

Tef, *Eragrostis tef* (Zucc.) Trotter, is a member of the Grass or Poaceae Family and Chlorodoideae sub-family. Tef is an allotetraploid ($2n=4x=40$) with estimated genome size ranging from 648 to 926Mbp (Ayele *et al.*, 1996; Hundera *et al.*, 2000), which is approximately 60% larger than the rice genome. It is a self-pollinated with very low degree of out-crossing, that ranges from 0.2% to 1.0% (Ketema, 1997). Tef represents a unique biodiversity component in the agriculture and food security systems of millions of poor farmers in Ethiopia. In Ethiopia, tef is the most valuable source of human food (the grain) and livestock feed (the straw), cash and foreign currency earnings. It ranks first in terms of total area under cultivation where it accounts for 30% of the total acreage under cereal crops (CSA, 2015). Tef is also known to be tolerant to extreme climatic and soil conditions; hence, it is a favorite crop in the semi-arid areas with moisture limitations (Tadele and Assefa, 2012). In recent years, tef is receiving global attention for its nutritional and health-related benefits (Provost and Jobson, 2014) especially due to the absence of gluten, a cause for celiac disease, in its grain (Spaenij-Dekking *et al.*, 2005).

Despite its versatility in adapting to adverse environmental conditions and being the staple food over 60 million people in Ethiopia, the seed yield of tef is low. The national average yield is 1.57 t ha^{-1} , in contrast to 2.5 t ha^{-1} for wheat and 3.4 t ha^{-1} for maize (CSA, 2015). A major cause of low productivity of tef is lodging, the permanent displacement of the stem from the upright position. Tef has a tall and slender stem, which is susceptible to lodging caused by wind and rain. In addition, when fertilizer is applied to increase yield, stems of tef grow taller and become even more susceptible to lodging, resulting in significantly reduced quantity and quality of grain and straw. Lodging is most likely to occur when tillers have a high center of gravity and a heavy, high yielding panicle (Cheverton *et al.*, 1992). On the average, the tef yield loss due to lodging was estimated to be 11-25% (Ketema, 1993).

Since scientific research started on tef five decades ago, 42 varieties were released mainly by DebreZeit Agricultural Research Center (MoANR, 2017). Among these, recently released varieties including *Quncho* (Assefa *et al.*, 2011), *Kora* (Assefa *et al.*, 2017) and *Dagim* (Chanyalew *et al.*, 2017) showed significant yield benefits. This paper presents the performance of the recently released tef variety called *Tesfa* comparing with other candidate genotypes and standard variety. In addition, the paper describes morphological and agronomic properties of the new variety.

Methodology

Experimental sites

The experiment was carried out at six locations in Ethiopia where tef is the major crop. These locations are Adet, Chefe Donsa, Debre Zeit black soil, Debre Zeit light soil, Ginchi and Holetta. Details of the six locations in terms of altitude, climate other related parameters are shown in Table 1.

Table 1. The altitude, coordinates and climate of six locations in Ethiopia where candidate tef lines were tested

| Location | Altitude (m) | Geographic coordinate | Climate | Average temperature (Max./min.) | Annual rainfall (mm) |
|-----------------------|--------------|-----------------------|------------|---------------------------------|----------------------|
| Adet | 2240 | 11°16'0"N, 37°29'0"E | Moist-cool | 24°C/9°C | 658.6 |
| Chefe Donsa | 2450 | 8°57'15"N, 39°06'04"E | Cool-wet | 23.2°C/10.5°C | 950 |
| Debre Zeit black soil | 1800 | 8°45'0"N, 38°59'0"E | Temperate | 26.1°C/8°C | 892 |
| Debre Zeit light soil | 1800 | 8°45'0"N, 38°59'0"E | Temperate | 26.1°C/8°C | 892 |
| Ginchi | 2236 | 9°1'60"N, 38°9'0"E | Moist-cool | 24.6°C/8.4°C | 1095 |
| Holetta | 2390 | 9°06'33"N, 38°49'02"E | Cool-wet | 22°C/6°C | 1144 |

Plant materials: The source of experimental materials were two dwarf mutant tef lines, namely *kegne* (Jost *et al.*, 2015) and *kinde* (Tadele *et al.* unpublished), which were identified at the University of Bern in Switzerland from screening 5000 mutagenized tef populations. The other parental lines were a popular tef variety called *Quncho* (DZ-Cr-387 RIL355) (Assefa *et al.*, 2011) and a landrace called *Kay Murri*. *Kay Murri* was selected as a paternal parent for its very white seed color, thick culm and vigorous growth habit. The eight recombinant inbred lines used in the current study were obtained from three independent crosses. The three independent crosses were, *kinde* x *Kay Murri* (lines 2 to 6), *Kay Murri* x *kegne* (line 7) and *Quncho* x *kinde* (lines 8 and 9). From each cross, 500 F₂ populations were generated and substantially reduced to few lines with best performance after eight generations of stringent selection focusing on standing ability and grain yield. Crossing and early generation testing were done for all breeding population at Debre Zeit Agricultural Research Center from where the National Tef Breeding Program is coordinated. The crossing combinations and names of recombinant inbred lines as well as control materials used in the current study are shown in Table 2.

Results: The performance of 10 tef genotypes which includes 8 inbred lines from the three independent crosses to semi-dwarf tef lines as well as two controls (farmers' check and improved *Quncho* variety) were tested at six locations (Adet, Chefe Donsa, Debre Zeit black soil, Debre Zeit light soil, Ginchi and Holetta) using Randomized Complete Block Design with three replications. Agronomic and yield data were collected and subjected to statistical analysis in order to identify the best genotypes of the evaluated genotypes. Summary of the results of the study is shown in Table 2 and below under variety description.

Based on a two-year multi-location trial, RIL181 was selected for its high grain and biomass yield as well as additional traits indicated below. Hence, RIL 181 was given the name *Tesfa* and applied for registration as a new improved tef variety. Based on the application, the National Variety Release Committee in Ethiopia investigated the two-year performance of RIL 181 and visited several locations where the new variety was grown for evaluation. Consequently, the committee approved the release of RIL 181 as a new variety with the name '*Tesfa*'.

Table 2. Combined mean agronomic performance of candidate tef lines at the National variety Trial at Adet, Chefe Donsa, Debre Zeit black soil, Debre Zeit light soil, Ginchi and Holetta

| Candidate line | Genotypes | DTH | DTM | GFP | PH | PL | LI | SMB/HA | GY/HA |
|----------------|---------------------------------|-------------|--------------|-------------|-------------|-------------|-------------|----------------|---------------|
| <i>Quncho</i> | DZ-Cr-387 (control) | 54.3 | 105.2 | 50.9 | 104.8 | 40.5 | 80.0 | 11604.2 | 2420.9 |
| RIL-227 | <i>kindex</i> Kay Murri | 48.2 | 103.2 | 55.0 | 90.4 | 32.5 | 78.1 | 9562.5 | 2198.3 |
| RIL-181 | <i>Kinde</i> x Kay Murri | 47.1 | 103.0 | 55.9 | 91.4 | 35.2 | 78.6 | 10500.0 | 2443.7 |
| RIL-110 | <i>Kindex</i> Kay Murri | 50.8 | 103.2 | 52.4 | 88.5 | 33.6 | 79.7 | 9645.8 | 2233.8 |
| RIL-232 | <i>Kindex</i> Kay Murri | 47.7 | 103.9 | 56.2 | 87.2 | 29.3 | 76.5 | 8994.8 | 2137.5 |
| RIL-237 | <i>Kindex</i> Kay Murri | 49.3 | 104.2 | 55.0 | 93.0 | 32.8 | 74.0 | 8729.2 | 1958.2 |
| RIL-195 | <i>KayMurrrix kegne</i> | 50.8 | 104.4 | 53.5 | 94.2 | 35.8 | 77.3 | 10802.1 | 2289.6 |
| RIL-171 | <i>Qunchox kinde</i> | 48.5 | 103.5 | 55.0 | 93.0 | 33.8 | 79.2 | 9822.9 | 2342.3 |
| RIL-189 | <i>Qunchox kinde</i> | 50.5 | 103.6 | 53.1 | 96.3 | 36.1 | 78.8 | 9484.4 | 2097.8 |
| Local check | | 47.0 | 101.3 | 54.4 | 89.0 | 33.9 | 84.4 | 10380.2 | 2364.4 |
| | Mean | 49.4 | 103.5 | 54.1 | 92.8 | 34.3 | 78.7 | 9952.6 | 2248.7 |
| | R² | 0.98 | 0.98 | 0.93 | 0.84 | 0.77 | 0.77 | 0.84 | 0.68 |
| | CV | 3.1 | 2.1 | 4.8 | 5.0 | 6.7 | 8.0 | 11.7 | 14.9 |
| | LSD(0.05) | 0.8 | 1.2 | 1.5 | 2.6 | 1.3 | 3.6 | 666.78 | 191.86 |

Abbreviations. DTH: days to heading; DTM: days to physiological maturity; GFP: grain filling period; PH: plant height; PL: panicle length; LI: lodging index; SMB: shoot biomass; GY: grain yield.

In general, *Tesfa* tef variety had above average yield and the variety with non-shattering type, thick culm, and compact panicle form which makes it suitable for both rainfed and irrigated agriculture. *Tesfa* is the first variety with compact panicle to be released by tef researchers in Ethiopia while all other 41 tef varieties released in the country are with loose or semi-loose panicle type.

Description

Characters

The new tef variety named *Tesfa*(DZ-Cr-457 RIL-181) is the product of a cross between a semi-dwarf mutant line called *kinde* and a landrace called *Kay Murri*. *Tesfa* is a medium size plant with compact panicle type (Fig. 1). Other distinct characters of *Tesfa* are pink lemma color, red anther color, and a very white seed color. On the average, *Tesfa* reaches the heading stage in 47 days and physiological maturity in 103 days after sowing. From the average plant height of 91 cm, the panicle of *Tesfa* contributes to 40%.



Figure 1. The newly released Tesfa tef variety grown at DebreZeit Research Center, Ethiopia. Photo: Zerihun Tadele.

Yield Performance

Highly significant variations were observed in shoot biomass and total grain yield among tef genotypes tested in all study years and locations. The average grain yield of *Tesfa* was 2.5 t ha⁻¹ at research centers and 2.0-2.4 t ha⁻¹ in farmers' fields. In the first year of its cultivation by farmers, *Tesfa* received positive feedback from growers because of its several unique and desirable properties including its compact panicle, lack of shattering, and thick and strong culm which makes the new variety potentially cultivated under irrigation and also mechanically harvested.

Adaptation

Tesfa is released for the optimum moisture and high rainfall areas (high potential tef producing areas) of the country. It performs very well in areas having an altitude 1700-2400 m and annual rainfall of 500-1000 mm.

Variety maintenance

Breeder and foundation seeds of the variety are maintained by Debre Zeit Agricultural Research Center.

References

- Assefa K, S Aliye, G Belay, G Metaferia, H Tefera, ME Sorrells. 2011. Quncho: the first popular tef variety in Ethiopia. *International Journal of Agricultural Sustainability* 9: 25-34.
- Assefa K, S Chanyalew, Y Genet, M Asfaw, T Fikre, H Jifar, N Hussien, M Hailesellassie, G Chemedda, G Assefa. 2017 Tef (*Eragrostis tef*) variety Kora. *Ethiopian Journal of Agricultural Science* 27: 137-140.
- Ayele M, J Dolezel, M VanDuren, H Brunner, FJ Zapata Arias. 1996. Flow cytometric analysis of nuclear genome of the Ethiopian cereal Tef [*Eragrostis tef*(Zucc) Trotter]. *Genetica* 98: 211-215.
- Chanyalew S, K Assefa, M Asfaw, Y Genet, K Tolossa, W Kebede, T Fikre, N Hussien, H Jifar, A Fentahun, K Gebremeskel, G Chemedda, T Belete. 2017. Tef (*Eragrostis tef*) variety Dagim. *Ethiopian Journal of Agricultural Science* 27: 131-135.
- Cheverton M, M Pullen, F Didehvar, G Jones. 1992. Database of accessions in the *Eragrostis tef* germplasm collection at Wye, Interim Report. *In*. Wye College, University London.
- CSA. 2015. Agricultural sample survey 2014/2015. Volume I. Report on area and production of major crops. Statistical Bulletin 578. *In*. Central Statistical Agency (CSA), Addis Ababa, Ethiopia.
- Hundera F, K Arumuganathan, P Baenziger. 2000. Determination of relative nuclear DNA content of tef [*Eragrostis tef* (Zucc.) Trotter] using flow cytometry. *Journal of Genetics & Breeding* 54: 165-168.
- Jost M, K Esfeld, A Burian, G Cannarozzi, S Chanyalew, C Kuhlemeier, K Assefa, Z Tadele. 2015. Semi-dwarfism and lodging tolerance in tef (*Eragrostis tef*) is linked to a mutation in the alpha-Tubulin 1 gene. *Journal of Experimental Botany* 66: 933-944.
- Ketema S. 1993. Tef (*Eragrostis tef*): breeding, agronomy, genetic resources, utilization, and role in Ethiopian agriculture. Institute of Agricultural Research ;, Addis Abeba, Ethiopia
- Ketema S. 1997. Tef, *Eragrostis tef* (Zucc.) Trotter. *In*. Institute of Plant Genetics and Crop Plant Research, Gatersleben/International Plant Genetic Resources Institute, Rome, Italy, p 52.
- MoANR. 2017. Plant Variety Release, Protection and Seed Quality Control Directorate, Crop Variety Register, Vol Issue No. 20. Ministry of Agriculture and Natural Resources (MoANR), Addis Ababa, Ethiopia.
- Provost C, E Jobson. 2014. Move over quinoa, Ethiopia's teff poised to be next big super grain. *In* The Guardian January 23, 2014.
- Spaenij-Dekking L, Y Kooy-Winkelaar, F Koning. 2005 The Ethiopian cereal tef in celiac disease. *The New England Journal of Medicine* 353: 1748-1749.
- Tadele Z, K Assefa. 2012. Increasing food production in africa by boosting the productivity of understudied crops. *Agronomy* 2: 240-283.